

A peer-reviewed version of this preprint was published in PeerJ on 15 April 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.350) (peerj.com/articles/350), which is the preferred citable publication unless you specifically need to cite this preprint.

Kismul H, Van den Broeck J, Lunde TM. 2014. Diet and kwashiorkor: a prospective study from rural DR Congo. PeerJ 2:e350
<https://doi.org/10.7717/peerj.350>

Diet and kwashiorkor: a prospective study from rural DR Congo

Hallgeir Kismul, Jan Van den Broeck, Torleif Markussen Lunde

The etiology of kwashiorkor remains enigmatic and longitudinal studies examining potential causes of kwashiorkor are scarce. Using historical, longitudinal study data from the rural area of Bwamanda, Democratic Republic of Congo, we investigated the potential causal association between diet and the development of kwashiorkor in 5 657 preschool children followed 3-monthly during 15 months. We compared dietary risk factors for kwashiorkor with those of marasmus.

Kwashiorkor was diagnosed as pitting oedema of the ankles; marasmus as abnormal visibility of skeletal structures and palpable wasting of the gluteus muscle. A food frequency questionnaire was administered 3-monthly to record the consumption of the 41 locally most frequent food items. We specified Hanley-Miettinen smooth-in-time risk models containing potential causal factors, including food items, special meals prepared for the child, breastfeeding, disease status, nutritional status, birth rank, age, season and number of meals. Bayesian Information Criteria identified the most plausible causal model of why some children developed kwashiorkor.

In a descriptive analysis of the diet at the last dietary assessment prior to development of kwashiorkor, the diet of children who developed kwashiorkor was characterized by a low intake of sweet potatoes, papaya and “other vegetables” [0.0% , 2.3% (95%CI: 0.4, 12.1) and 2.3% (95%CI: 0.4, 12.1)] in comparison with children who did not develop kwashiorkor [6.8% (95%CI: 6.4, 7.2), 15.5% (95%CI: 15, 16.1) and 15.1% (95%CI: 14.6, 15.7)] or children who developed marasmus [4.5% (95%CI 2.6, 7.5) 11.8% (95%CI: 8.5, 16.0) and 17.6% (95%CI: 13.7, 22.5)].

Sweet potatoes and papaya have high β -Carotene content and so may some of “the other vegetables”. We found that a risk model containing an age function, length/height-for age Z-score, consumption of sweet potatoes, papaya or other vegetables , duration of this consumption and its interaction term, was the most plausible model. Among children aged 10-42 months, the risk of developing kwashiorkor increased with longer non-consumption of these foods. The analysis was repeated with only children who developed marasmus as the reference series, yielding similar results.

Our study supports that β -Carotene may play an important role in the protection against kwashiorkor development.

Diet and kwashiorkor: a prospective study from rural DR Congo

Hallgeir Kismul^{1,2}

Jan Van den Broeck^{1,2}

Torleif Markussen Lunde^{1,2}

To whom correspondence should be addressed, Phone: +47 55 58 85 56

Fax: +47 55 58 85 61 e-mail: hallgeir.kismul@cih.uib.no

¹ Centre for International Health, University of Bergen, Norway

² Current address: Centre for International Health, University of Bergen, 5009 Bergen, Norway

Introduction

Malnutrition contributes significantly to the high under-five year mortality in the world and as an underlying factor it has been estimated that it contributes to over one third of all child deaths [1]. Mortality is very high among children with marasmus, and even higher among those with kwashiorkor [2, 3] Kwashiorkor and marasmus are characterised by different metabolic response to severe undernutrition [4, 5].

Kwashiorkor has been linked to diet since its first description. Williams, who introduced the name kwashiorkor, suggested protein undernutrition as the etiology of kwashiorkor [6] . The association of kwashiorkor with low protein intake has later been questioned. So far no research has demonstrated that children with kwashiorkor consume less protein than children with marasmus. Golden and Ramdath proposed excess free radicals as the explanation of clinical findings in kwashiorkor [7]. Relations between antioxidant depletion and the occurrence of kwashiorkor have been investigated, but the role of oxidative stress as primary cause of kwashiorkor is still debated [8, 9].

To our knowledge there are only two prospective studies that have examined the relations between diet diversity and kwashiorkor. Investigating protein-calorie malnutrition, a study following Indian children from birth to 3 years, examined differences in diet between children developing kwashiorkor and children developing marasmus [10]. A more recent study examined dietary factors determining kwashiorkor by assessing dietary intake of one to three years old Malawian children [11]. The former study did not find significant differences in diet between children who developed kwashiorkor and those who developed marasmus, while the latter did not find differences between those who developed kwashiorkor and those who did not.

The overall aim of our study is to investigate, in a large longitudinal population-based study, the possible association between diet and the development of kwashiorkor. We also wanted to compare the causal influence of dietary risk factors for kwashiorkor with those of marasmus. We performed this analysis by applying the causal investigation method proposed by Miettinen and the statistical approach by Hanley and Miettinen, using a random sample of person moments from the entire dynamic population as the reference series [12, 13].

Method

The Bwamanda study

We did a secondary analysis of the historical data from the Bwamanda study, conducted from 1989 -1991 in a rural area of the northwest part of the Democratic Republic of Congo, (DRC), located at 19.2 degrees east and 3.2 degrees north. The people of Bwamanda are, up till today, predominantly subsistence farmers and the base diet consists of mainly of maize, cassava supplemented with fish, vegetables and fruits. Health care in the area is provided by a central hospital and 10 minor health centres with a few of these providing some limited nutritional rehabilitation services. With virtually unchanged living conditions in the study area, the secondary analysis was viewed to be contemporary relevant.

Study design

The Bwamanda study was a dynamic population study with follow-up including three-monthly contacts, survey rounds, making up 15 months of follow-up and 6 contacts. At the first round 4 235 preschool children were enrolled and at the last round a total of 5 657 had been enrolled. A full description of the study population can be found in Van den Broeck et al. [14]. Trained interviewers conducted interviews according to an interviewer's manual. They determined the children's age on the basis of birth date noted on children's road to health chart or on parents' identity paper or on the basis of an interview using a local events calendar.

Children were examined for kwashiorkor by using the presence of pitting oedema of the feet or ankles as a criterion. All children were examined for marasmus through inspection of abnormal visibility of skeletal structures and by absence or near-absence of palpable gluteus muscle. A locally constructed measuring board was used for measuring the length of children below 12 months, while a microtoise was used for measuring older children. In both cases length was measured to the nearest 0.1 cm. A spring scale (CMS weighting equipment) was used to weigh the children to the nearest 100 gram. We applied the WHO-MGRS Child Growth Standards for anthropometric scoring. Z-scores were calculated for weight for length/height (WHZ) and for length/height for age (HAZ) [15].

At each contact interviewers undertook face-to-face interviews with the most proximal caregiver of the child, usually the biological mother. The questionnaire included food

frequency questions with the 41 locally most consumed food items listed and interviewees providing “yes or no” answers to the questions if children had consumed the listed food items during the previous day. The food items had been identified through a pilot study. The interviewees were also asked about number of meals prepared for the families, special meals prepared for the child and breastfeeding.

Statistical method

In an initial descriptive analysis, we tabulated the percentage (95% confidence interval) of individuals eating the different items, grouped by those who developed kwashiorkor, those who did no and those who developed marasmus. We used a two-sample test for equality of proportions to test if the fractions were different.

We specified Hanley-Miettinen smooth-in-time risk models containing all potential causal factors, including food items, special meals prepared for the child, breastfeeding, disease status, nutritional status, birth rank, age, season and number of meas [13]. To select a representative sample from the study population we used the method proposed by Miettinen , and used the whole study population as reference series [12]. Results are reported as log-odds ratios (LOR) and risk reductions. In line with this method we took the dataset to consist of 35 person moments (c) where kwashiorkor was observed as the case series, and a representative sample (b) of the infinite number of person moments that constitute the 46 397 person-months in the study base. We use a c/b-ratio of 150 assuring variances and covariance have minimal errors (less than 1 percentage) compared to using the entire series. The data are summarised in table 1.

Age function and age as a risk factor

Given that risk is not changing linearly with age, as seen in figure 1, we developed an age variable that accounted for nonlinear change in risk. Accordingly age was included as an independent variable in the model. Based on visual inspection of how kwashiorkor was distributed according to age, a transformation of the age variable was done:

$$f(\text{age}) = a \exp(-b \text{ Age}) * \exp(-a \exp(-b \text{ Age})) \quad \text{Eq 1}$$

To find the parameters a and b, we optimized Eq 2 with binomial errors using logistic regression.

$$y = a + b \cdot f(\text{age}) \quad \text{Eq 2}$$

The AIC (Akaike Information criteria Information criteria) was used to compare models. We used an algorithm combining the golden section search and successive parabolic interpolation to find the parameters which minimized AIC, resulting in $a = 11.55$, and $b = 0.90$.

Other risk factors

The risk factors associated with kwashiorkor were defined in two steps; first we specified a log-linear hazard model with binomial errors where the independent variables were all food types, presence or absence of diarrhoea, and stunting and wasting at last visit. We defined time as the natural logarithm of number of months a person had, or had not, consumed a specific food item. We assumed that the food items reported at a given point in time were consumed up to the next contact, with the interview during the current visit providing data on any alterations in the consumption patterns since the previous contact.

Each variable was multiplied with the natural logarithm of time the item had been consumed or not. Next, we used the BIC (Bayesian Information Criteria) implemented in R's MASS package (stepAIC) to find the most plausible model based on our data; the posteriori most probable candidate model. The fitted candidate model corresponding to the lowest value of BIC is the candidate model corresponding to the highest Bayesian posterior probability.

Based on the selected model we address the risk of developing kwashiorkor given a prior personal profile. We report and risk reduction estimates on the basis of profiles.

To test if the model could also explain the difference between subjects who developed marasmus from those who developed kwashiorkor we applied the final selected model, with the same 35 person moments (c2) where kwashiorkor was observed as the case series, but this time with the refrecreses constituted by a sample of the infinite number of persons moments including 1 173 person-months observed in 372 new cases of marasmus.

Ethical approval for the Bwamanda study had been granted by the University of Leuven's Tropical Childcare Health Working Group. Community and individual verbal consent was obtained from children's caretakers.

Results

Table 2 reports the consumption of different food items by the children in the survey round prior to the incidence of kwashiorkor. A high proportion of the children had consumed cassava roots, maize and cassava leaves. The proportion who had consumed cassava roots and maize was non-significantly higher for those who developed kwashiorkor, but for cassava leaves the consumption was lowest for the children with kwashiorkor. The proportion of children who had consumed yam, pineapple, citrus, snails, and eggs was non-significantly higher for the children who developed kwashiorkor than for the others. On the other hand the proportion of children with kwashiorkor who had consumed okra, ground nuts, banana, squash, meat, chili, fish and other vegetables was non-significantly lower than for the rest. There were no significant differences in proportion of children who had consumed palm oil between children who developed kwashiorkor and the other children. The diet of children who developed kwashiorkor was characterized by a low intake of sweet potatoes, papaya and other vegetables [0.0% , 2.3% (95%CI: 0.4, 12.1) and 2.3% (95%CI: 0.4, 12.1)] in comparison with children who did not develop kwashiorkor [6.8% (95%CI: 6.4, 7.2), 15.5% (95%CI: 15, 16.1) and 15.1% (95%CI: 14.6, 15.7)] or children who developed marasmus [4.5% (95%CI 2.6, 7.5) 11.8% (95%CI: 8.5, 16.0) and 17.6% (95%CI: 13.7, 22.5)].

Given that the β -Carotene could be the main acting agent in sweet potatoes, papaya and “other vegetables” we constructed a variable, PaSV (papaya, sweet potatoe and “other vegetables”), which combined all these items, weighted by the β -Carotene content of 100 g of each item. The weighting of sweet potatoes equalled 1 and papaya 1/3. The variable “other vegetables” includes mostly taro including taro leaves. In addition the variable encompasses wild vegetables. Taro leaves are rich in β -Carotene and a study from DRC shows that wild vegetables are also rich in β -Carotene [16]. Since we have not been able to determine the more precise content of the other vegetables in our study, we have weighted the “other vegetables” low, with the weighing equal to 1/10. For the construction of smooth-in-time risk models we defined time for this combined variable as for the single food items.

The most probable model based on BIC, was one that included age, time, PaSV, and HAZ. Replacing HAZ with WHZ gave similar results. The two variables were correlated ($R^2 = 0.50$). As seen in Figure 1 the risk of developing kwashiorkor was highest in the age interval between 16 and 38 months.

Table 3 shows the coefficients for the non-proportional hazard model with person moments sampled from the entire population. The log-odds for the continuous variable HAZ; LOR -0.8 (CI 95% -1.1, -0.5), length/ height for age Z-score, describes an increased risk of developing kwashiorkor with lower height for age. Chronic malnourished children on average have a negative HAZ score, hence the negative log-odds. We found the log-odds for the time variable to be LOR 4.7 (CI 95% 3.4 6.1), for the combined variable for food items containing β -Carotene it was LOR -9.2 (CI 95% -21.0 -3.1), for PaSV, and their interaction it was LOR -8.1 (CI 95% -11.1 -2.1). These findings must be understood together. A child not consuming the PaSV food items will have PaSV = 0, and thus the interaction term is also zero. The risk of developing kwashiorkor therefore increases the longer the child does not consume the PaSV food items. On the other hand, as illustrated in figure 2 a child consuming PaSV food items, PaSV > 0, will reduce the risk over time. The overall model fit was good with an AIC of 251.3 and a Nagelkerke R² index of 0.44.

Table 4 shows the findings from sampling control-moments only from children who developed marasmus. The table shows that the HAZ score for those who developed marasmus is the same as the HAZ score for the children who developed kwashiorkor (LOR = 0.0, CI 95% -0.1, 0.2). Then again it shows that there is a difference with regards to consumption of products containing β -Carotene with the LOR for PaSV being -6.8 (CI 95% -17.8 -1.7) and for PaSV combined with the time variable T the LOR was -6.3 (CI 95% -9.0 -0.8). The age of children who developed kwashiorkor was also different from children who developed marasmus with LOR being 7.9 (CI 95% 4.3 12.1). The age of children with marasmus was distributed within the age of 11 to 64 months, while the age of children with kwashiorkor mainfell between 16 and 38 months, reaching a top around 26 months.

Ethical approval for the Bwamanda study had been granted by the University of Leuven's Tropical Childcare Health Working Group. Community and individual verbal consent was obtained from children's caretakers.

Discussion

Our study shows that the children who developed kwashiorkor were mainly stunted children aged 16-38 months. Their diet was characterized by a low or no consumption of sweet potatoes, papaya and “other vegetables”. The children who developed marasmus resembled the children who developed kwashiorkor by being chronically malnourished. However, the consumption of papaya, sweet potatoes and “other vegetables” were lower among children with kwashiorkor than among marasmic children. Sweet potatoes, papaya and “other vegetables” are characterised by their high β -Carotene content. We found that the risk of developing kwashiorkor increased the longer the child did not consume these food items.

We emphasize the importance of β -Carotene because it is a substance with significant antioxidant activities. Then again sweet potatoes, papaya and what we have termed other vegetables contain several other carotenoids that have antioxidant activities. As a result the various carotenoids, the mixture of carotenoids or carotenoids in association with other antioxidants in these food items can have played important roles in the protective pathway in kwashiorkor.

Children who developed kwashiorkor also consumed palm oil, with the oil being rich in β -Carotene. In Bwamanda palm oil is extracted locally and mostly stored in bottles, often being exposed to strong sunlight. Palm oil is used for preparing cassava leaves stew. The oil is mixed with cassava leaves and boiled in a pot for an hour. Studies show that carotenoids are vulnerable to degradation. It is especially their unsaturated structures that make them sensitive to heat, oxygen and light [17]. A study from Nigeria on heating of palm oil demonstrated that the amount of β -carotene declined with temperature increase and that the destruction of β -carotene was larger when the oil was heated continuously for 30 min at any given temperature [18]. Another study on heating red palm oil showed that all trans- β -Carotene were almost lost after 20 minutes [19]. Thus given that local practices expose palm oil to sunlight and long duration cooking it is unlikely that palm oil consumption would have compensated for the lack of provitamin A in the local diet.

Besides our study there are only two other observational longitudinal studies we know of that examine the relation between diet and the development kwashiorkor including Gopalan's study of Indian children up to 10 years and Lin et al.'s study on Malawian children [10, 11].

While our study analyses diet with reference to a variety of food items, Gopalan's study concentrated on differences in terms of protein-calorie consumption. Gopalan did not find evidence that there were any differences in the protein-calorie intake between cases of marasmus and kwashiorkor. The study conducted by Lin et al. actually found that the daily intake of vitamin A equivalents was low among children who developed kwashiorkor, but not significantly lower than in the control group [11]. We suspect that if the study had examined the cumulative intake of vitamin A equivalents over a longer period the difference in vitamin A equivalents, intake between children with kwashiorkor and marasmus might have been significant.

Golden suggested that kwashiorkor results from oxidative stress and a general defiance in protective mechanism that could reduce the oxidative damage with most of the protective pathways necessitating micronutrients [7]. Before target specific antibodies have been produced, immune cells generate and release reactive oxygen species [20]. These oxidative bursts have been shown to be important in the early phase of for example malaria infections. Oxidative burst can lead to haemolysis and cellular dysfunction [21]. Since carotenoids can act as an important anti-oxidants our findings support theories that relate kwashiorkor to oxidative stress and the importance of micronutrients in the protective pathways [20, 22].

A case control study examined the antioxidant hypotheses by comparing the diet in siblings of children presenting with marasmus and children with kwashiorkor [23]. It concluded that siblings of children with kwashiorkor consumed egg and tomatoes less frequent than children with marasmus. We note that the differences in consumption between the two were minor. Studies have questioned the hypothesis that antioxidant depletion causes kwashiorkor. In a controlled trial from Malawi, Ciliberto et al. assessed the efficiency of antioxidant supplementation in preventing kwashiorkor in children aged 1 to 4 years [23]. The intervention arm received antioxidant powder containing riboflavin, vitamin E, selenium and N-acetylcysteine, while the control arm received as a placebo of an identical looking powder. According to Ciliberto et al. the study showed that the supplementation of antioxidant powder did not prevent children from developing kwashiorkor. We give notice to the fact that the study did not involve supplementation of any provitamin A and it thereby does not inform about efficiency of provitamin A in preventing kwashiorkor. The study did not provide baseline data about study participants' diet. The study design does therefore not allow for

examining how dietary variables might have influenced the treatment with antioxidant powder and the influence of such variables on the development of kwashiorkor.

As far as we know the Bwamanda study is with exception Gopalan and Lin et al. the only observational longitudinal study that describes the diet of children who develop kwashiorkor and marasmus. The results from our study support the hypothesis that kwashiorkor is caused by oxidative stress, with the study supporting an important role of food containing carotenoids in the protective pathway. In Bwamanda the dietary sources of preformed vitamin A are limited and pro-vitamin A carotenoid constitutes a major source of vitamin A. The pro-vitamin A carotenoids including β -Carotene can through cleavage be converted into retinaldehyde (a form of vitamin A). Studies have shown that the bioavailability of pro vitamin A carotenoids is less than of preformed vitamin A [24, 25]. Our study proposes that the consumption of fruits plays a role in reducing the risk of developing kwashiorkor. In comparison with green leafy vegetables and carrots, fruits are also more effective in improving vitamin A status among children [25].

Studies have shown that there is a relationship between diet and infection during the development of kwashiorkor were recurrent infections contributes to hypo-albuminaemia and the development of edema [26, 27]. We also know that vitamin A modulates many types of specific and non-specific immune system and those vitamin deficiencies have a negative impact on different types of immunity functions [28, 29]. In this manner pro-vitamin A can play a role of reducing the severity of a number of types of infections and we speculate that provitamin A played a role in reducing the risk of developing edema among chronically malnourished children in Bwamanda

The strength of our study is that in a large population based study we have managed to examine the dietary diversity over a long period of children who developed kwashiorkor and marasmus. The study design has also some disadvantages. Given that information on diet was based on 24 hour recall we were not able to determine food consumed in between the follow up period. Furthermore, we have restricted information on the quantity of food consumed and we are therefore not in a position to determine the amount of pro-vitamin vitamin A consumed by the children. We also realise that our findings require confirmation preferable in a large randomised trial that examine the development of kwashiorkor in a trial of carotenoids supplementation versus placebo.

In conclusion, this is the first observational longitudinal study that demonstrates a relation between diet diversity and the development of kwashiorkor. We have suggested that the consumption of a diet that includes food items containing carotenoids reduces the risk in children aged 10-71 months of developing kwashiorkor. Our findings should be considered as a support to on-going efforts that aim at promoting a diverse agricultural and horticulture production and in this manner stimulate consumption of a more varied diet. In rural communities where there is a shortage of vitamin A rich food it is in particular important to promote increased production of vegetables and fruits rich carotenoids including sweet potatoes and papaya.

Acknowledgement

The Bwamanda study was supported by the Nutricia Research Foundation. H. K and T.M.L. conducted the analysis and H.K. wrote the first draft of the manuscript. All authors edited the manuscript and contributed in the interpretation of the results. All authors read and approved the final manuscript. The Bwamanda study was initiated and conducted by J.V.d.B. and Roger Eeckels

Accessibility of the Bwamanda dataset

As the principle investigator Jan Van den Broeck is the custodian of the Bwamanda dataset. Jan Van den Broeck supervised our study and provided Hallgeir Kismul as the first author access to the Bwamanda data. The dataset can be made available by contacting Jan Van den Broeck; Jan.Broeck@cih.uib.no

Literature cited

1. Lim, S.S., *A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010*. Lancet, 2012. **380**(9859): p. 2224-60.
2. Briend, A., B. Wojtyniak, and M.G.M. Rowland, *Arm Circumference and Other Factors in Children at High-Risk of Death in Rural Bangladesh*. Lancet, 1987. **2**(8561): p. 725-728.
3. Prudhon, C., et al., *A model to standardise mortality of severely malnourished children using nutritional status on admission to therapeutic feeding centres*. European Journal of Clinical Nutrition, 1997. **51**(11): p. 771-777.
4. Jahoor, F., et al., *Protein metabolism in severe childhood malnutrition*. Annals of Tropical Paediatrics, 2008. **28**(2): p. 87-101.
5. Badaloo, A.V., et al., *Lipid kinetic differences between children with kwashiorkor and those with marasmus*. Am J Clin Nutr, 2006. **83**(6): p. 1283-8.
6. Williams, C.D., *Kwashiorkor: a nutritional disease of children associated with a maize diet*. The Lancet, 1935. **Volume 226**(5855): p. 1151-1152.
7. Golden, M.H. and D. Ramdath, *Free radicals in the pathogenesis of kwashiorkor*. Proc Nutr Soc, 1987. **46**(1): p. 53-68.
8. Manary, M.J., C. Leeuwenburgh, and J.W. Heinecke, *Increased oxidative stress in kwashiorkor*. J Pediatr, 2000. **137**(3): p. 421-4.
9. Lenhartz, H., et al., *The clinical manifestation of the kwashiorkor syndrome is related to increased lipid peroxidation*. J Pediatr, 1998. **132**(5): p. 879-81.
10. Gopalan, C., *Kwashiorkor and marasmus: evolution and distinguishing features*. Natl Med J India, 1992. **5**(3): p. 145-51.
11. Lin, C.A., et al., *A prospective assessment of food and nutrient intake in a population of Malawian children at risk for kwashiorkor*. J Pediatr Gastroenterol Nutr, 2007. **44**(4): p. 487-93.
12. Miettinen, O.S., *Etiologic study vis-a-vis intervention study*. Eur J Epidemiol, 2010. **25**(10): p. 671-5.
13. Hanley, J.A. and O.S. Miettinen, *Fitting smooth-in-time prognostic risk functions via logistic Regression*. The International Journal of Biostatistics, 2009. **5**: p. 1- 23.
14. Van den Broeck, J., R. Eeckels, and J. Vuylsteke, *Influence of nutritional-status on child-mortality in rural Zaire*. Lancet, 1993. **341**(8859): p. 1491-1495.
15. WHO, *Global Database on Child Growth and Malnutrition*. 2012.
16. Termote, C., et al., *A Biodiverse Rich Environment Does Not Contribute to a Better Diet: A Case Study from DR Congo*. Plos One, 2012. **7**(1).
17. Leskova, E., et al., *Vitamin losses: Retention during heat treatment and continual changes expressed by mathematical models*. Journal of Food Composition and Analysis, 2006. **19**(4): p. 252-276.
18. Mudambi, R.S. and M.V. Rajagopa, *Effect of heat on the beta-carotene content of Nigerian palm oil*. Journal of Food Science and Technology-Mysore, 2006. **43**(5): p. 1414-1416.
19. Fillion, L. and C.J.K. Henry, *Nutrient losses and gains during frying: a review*. International Journal of Food Sciences and Nutrition, 1998. **49**(2): p. 157-168.
20. Strobel, M., J. Tinz, and H.K. Biesalski, *The importance of beta-carotene as a source of vitamin A with special regard to pregnant and breastfeeding women*. Eur J Nutr, 2007. **46 Suppl 1**: p. 1-20.
21. Isaksson, C., et al., *Explaining variance of avian malaria infection in the wild: the importance of host density, habitat, individual life-history and oxidative stress*. BMC Ecol, 2013. **13**: p. 15.
22. Sergio, A.R., *b-carotene and other carotenoids as antioxidants*. Journal of the American College of Nutrition, 1999. **18**(5): p. 426-433.

23. Ciliberto, H., et al., *Antioxidant supplementation for the prevention of kwashiorkor in Malawian children: randomised, double blind, placebo controlled trial*. *BMJ*, 2005. **330**(7500): p. 1-5.
24. de Pee, S., et al., *Lack of improvement in vitamin A status with increased consumption of dark-green leafy vegetables*. *Lancet*, 1995. **346**(8967): p. 75-81.
25. de Pee, S., et al., *Orange fruit is more effective than are dark-green, leafy vegetables in increasing serum concentrations of retinol and beta-carotene in schoolchildren in Indonesia*. *Am J Clin Nutr*, 1998. **68**(5): p. 1058-67.
26. Frood, J.D., *Relationship between pattern of infection and development of hypoalbuminaemia and hypo-beta-lipoproteinaemia in rural Ugandan children*. *Lancet*, 1971. **2**(7733): p. 1047-9.
27. Whitehead, R.G., *Infection and the development of kwashiorkor and marasmus in Africa*. *Am J Clin Nutr*, 1977. **30**(8): p. 1281-4.
28. Stephensen, C.B., *Vitamin A, infection, and immune function*. *Annu Rev Nutr*, 2001. **21**: p. 167-92.
29. Villamor, E. and W.W. Fawzi, *Effects of vitamin A supplementation on immune responses and correlation with clinical outcomes*. *Clinical Microbiology Reviews*, 2005. **18**(3): p. 446-64.

Table 1 Age distribution, length/height-for-Age Z-score (HAZ), and Weight for Length Z-score (WLZ) and for children with different clinical nutritional status. Z-scores based on the WHO-2006 Child Growth Standards[15].

	Age in month:			HAZ:			WLZ:		
	Q10	Q50	Q90	Q10	Q50	Q90	Q10	Q50	Q90
Kwashiorkor	15.9	26.5	38.4	-4.3	-2.3	-0.8	-2.5	-1.0	0.1
Reference population	7.4	35.9	66.5	-2.9	-1.4	-0.1	-1.4	-0.1	1.2
Marasmus	10.8	28.8	64.0	-4.5	-2.7	-1.2	-2.8	-1.3	0.1

Table 2 Consumption of different food items in the survey round preceding the development of kwashiorkor (n = 37) and non-development of kwashiorkor (n = 8108) and development of marasmus (n=374) in children between 6 and 50 months of age. * denotes p-value < 0.05, while** denotes p-value<0.01 with the value estimated using 2-sample test for equality of proportions with continuity correction as implemented in the prop test in R

Food items	Children with Kwashiorkor % (95%CI)	Children without kwashiorkor % (95%CI)	Children with Marasmus % (95%CI)	Food item	Children with Kwashiorkor % (95%CI)	Children without kwashiorkor % (95%CI)	Children with Marasmus % (95%CI)
African pear	0.0 (0.0, 8.2)	0.0 (0.0, 0.1)	0.0 (0.0, 1.3)	Okra	2.3 (0.4, 12.1)	4.5 (4.2, 4.8)	5.2 (3.2, 8.4)
Amaranth	7.0 (2.4, 18.6)	1.8 (1.7, 2.1)	3.1 (1.6, 5.8)	Palm oil	86.0 (72.0, 93.4)	88.8 (88.3, 89.3)	80.3 (75.4, 84.5)
Aubergine	0.0 (0.0, 8.2)	0.8 (0.7, 0.9)	0.0 (0.0, 1.3)	Papaya	2.3 (0.4, 12.1)	15.5 (15.0, 16.1)*	11.8 (8.5, 16.0)*
Avocado	0.0 (0.0, 8.2)	0.2 (0.1, 0.2)	0.0 (0.0, 1.3)	Pineapple	4.7 (1.3, 15.5)	1.4 (1.2, 1.6)	1.7 (0.7, 4.0)
Banana	9.3 (3.7, 21.6)	19.1 (18.5, 19.7)	16.3 (12.5, 21.0)	Powder milk	0.0 (0.0, 8.2)	0.0 (0.0, 0.1)	0.0 (0.0, 1.3)
Beans	31.1 (30.4, 31.8)	0.4 (0.3, 0.5)	0.0 (0.0, 1.3)	Rice	2.3 (0.4, 12.1)	0.6 (0.5, 0.8)	0.0 (0.0, 1.3)
Breadfruit	0.0 (0.0, 8.2)	1.1 (0.9, 1.2)	1.7 (0.7, 4.0)	Sesame	0.0 (0.0, 8.2)	0.1 (0.1, 0.2)	0.3 (0.1, 1.9)
Cassava leaves	76.7 (62.3, 86.8)	79.2 (78.6, 79.9)	70.7 (65.2, 75.6)	Shrimp	0.0 (0.0, 8.2)	0.1 (0.1, 0.2)	0.0 (0.0, 1.3)
Caterpillars	2.0 (1.8, 2.2)	2.0 (1.8, 2.2)	1.0 (0.4, 3.0)	Snails	2.3 (0.4, 12.1)	1.4 (1.3, 1.6)	1.4 (0.5, 3.5)
Cassava Roots	76.7 (62.3, 86.8)	72.6 (71.9, 73.3)	72.3 (66.9, 77.2)	Soya	4.7 (1.3, 15.5)	5.2 (4.9, 5.5)	5.9 (3.7, 9.2)
Chili pepper	4.7 (1.3, 15.5)	8.9 (8.4, 9.3)	4.5 (2.6, 7.5)	Spinach	2.3 (0.4, 12.1)	2.8 (2.6, 3.1)	2.1 (1.0, 4.5)
Egg	4.7 (1.3, 15.5)	0.7 (0.5, 0.8)**	1.0 (0.4, 3.0)	Squash	0.0 (0.0, 8.2)	4.9 (4.6, 5.2)	5.2 (3.2, 8.4)
Fish	18.6 (9.7, 32.6)	31.1 (30.4, 31.8)	25.3 (20.6, 30.6)	Sugar cane	0.0 (0.0, 8.2)	0.7 (0.6, 0.9)	0.3 (0.1, 1.9)

Fruit (others)	0.0 (0.0, 8.2)	1.8 (1.6, 2.0)	1.0 (0.4, 3.0)	Sweet potato	0.0 (0.0, 8.2)	6.8 (6.4,7.2)	4.5 (2.6, 7.5)
Ground nuts	18.6 (9.7, 32.6)	28.6 (27.9, 29.3)	23.9 (19.3, 29.1)	Termites	0.0 (0.0, 8.2)	0.3 (0.3,0.4)	0.0 (0.0, 1.3)
Maize	97.7 (87.9, 99.6)	93.5 (93.1, 93.8)	91.7 (88.0, 94.4)	Tomatoes	0.0 (0.0, 8.2)	1.3 (1.1,1.5)	0.0 (0.0, 1.3)
Mango	0.0 (0.0, 8.2)	0.9 (0.7, 1.0)	0.7 (0.2, 2.5)	Wheat	0.0 (0.0, 8.2)	0.6 (0.5,0.7)	0.3 (0.1, 1.9)
Meat	0.0 (0.0, 8.2)	4.7 (4.4, 5.0)	5.5 (3.4, 8.8)	Other vegetables	2.3 (0.4, 12.1)	15.1 (14.6,15.7)*	17.6 (13.7, 22.5)**
Milk	0.0 (0.0, 8.2)	0.1 (0.1, 0.1)	0.0 (0.0, 1.3)	Yam	2.3 (0.4, 12.1)	1.3 (1.1,1.4)	0.7 (0.2, 2.5)
Mushroom	0.0 (0.0,8.2)	2.3 (2.1, 2.5)	1.7 (0.7, 4.0)				

Table 3 The coefficients for the non-proportional hazard model with person moments sampled from the entire population. The Age variable is a transformation based on the distribution of kwashiorkor across age. T is a variable that describes the duration of consuming a food item containing β -Carotene. PaSV is a variable that combines papaya, other vegetables and sweet potatoes and weighted by the β -Carotene content of 100 g of the item. The Height-for-Age Z-score (HAZ) are based on the WHO-2006 Child Growth Standards[15].

Term	Log odds - Estimate	Confidence interval, 95%
Intercept	-15.5	-18.1, -13.4
Age	10.1	6.1, 14.8
T	4.7	3.4 , 6.1
Var X	-9.2	-21.0, -3.1
HAZ	-0.8	-1.1, -0.5
T * PaSV	-8.1	-11.1, -2.1

PeerJ Preprints

Table 4 The coefficients for the non-proportional hazard model with reference persons moments drawn from persons developing marasmus. The Age variable is based on the distribution of kwashiorkor across age. T is a variable that describes the duration of consuming a food item containing β -Carotene. PaSV is a variable that includes papaya, other vegetables and sweet potatoes weighted by the β -Carotene content of 100 g of the item. Height-for-Age Z-score (HAZ) are based on the WHO-2006 Child Growth Standards[15].

Term	Log odds - Estimate	Confidence interval, 95%
Intercept	-9.7	-12.1, -7.7
Age	7.9	4.3, 12.1
T	4.2	3.2, 5.4
Var X	-6.8	-17.8, -1.7
HAZ	0.0	-0.3, 0.2
T * PaSV	-6.3	-9.0, -0.8




Figure 1 Three months incidence of kwashiorkor according to age in months in intervals of six months.

Incidence of kwashiorkor (y-axis) against age in months (x-axis). Black dots indicate age of those children which developed kwashiorkor.




Figure 2: Risk reduction for developing kwashiorkor reduction of consuming β -Carotene rich products according to age in months. Dotted line is risk reduction after two months, dashed line after four months, and solid line after six months. Panel a) shows risk reduction for a child with a Height-for-Age Z-score (HAZ) of minus five, b) for a child with HAZ of minus three, and c) a child with HAZ of zero. HAZ-scores are based on the WHO-2006 Child Growth Standards[15].