

1

2 **Actively restored ecosystems as a refuge for biological diversity: A case study**  
3 **from eelgrass (*Zostera marina* L.)**

4 Jonathan S. Lefcheck<sup>1†</sup>, Scott R. Marion<sup>2</sup>, Robert J. Orth<sup>1</sup>

5  
6 <sup>1</sup>Virginia Institute of Marine Science, The College of William & Mary, Gloucester Point, VA

7 <sup>2</sup>Oregon Department of Fish & Wildlife, Marine Resources Program, Newport, OR

8  
9 <sup>†</sup>Corresponding author. Email: [jslefche@vims.edu](mailto:jslefche@vims.edu). Phone: +1 804 684 7150. Mailing address:  
10 PO Box 1346, Rt 1208 Greate Rd, Gloucester Point, VA 23062, USA

11  
12 Running head: Eelgrass restoration and biodiversity

**Abstract:**

As nearshore ecosystems are increasingly degraded by human activities, active restoration is a critical element in ensuring the continued provision of goods and services by coastal habitats. Since 1997, over 1800 ha of the foundational species eelgrass (*Zostera marina* L.) has been reestablished in the coastal bays of the mid-western Atlantic. Here, we follow the functional recovery of the epifaunal invertebrate community associated with the restored eelgrass habitat from 2001-2013. After less than a decade, the invertebrate community in the restored bed was richer, more even, and exhibited greater variation in functional traits than a nearby reference bed. Analysis of environmental and physical data using random forests revealed the primary drivers of these differences was the increasing area and density of eelgrass directly attributable to ongoing restoration efforts. We also infer based on existing paradigms that the diverse epifaunal communities observed may have positive consequences for critical ecosystem services, including removal of fouling epiphytes from eelgrass blades and transfer of energy to higher trophic levels. Given that restored eelgrass now make up 32% of total seagrass cover in the mid-Atlantic coastal bays, this restoration may foster regional biodiversity by providing new habitat, particularly given the general decline of natural eelgrass in this region and globally.

**Keywords:** biodiversity, seagrass, restoration, functional traits, grazers

## Introduction

Coastal habitats are becoming increasingly impacted through human activities (Halpern et al. 2008), leading to loss of important and valuable services such as food production, pollution control, shoreline buffering, and carbon storage (Barbier et al. 2011). At the same time, anthropogenic impacts are driving the rapid and irreversible loss of marine biological diversity (McCauley et al. 2015). As high biodiversity has been generally shown to promote ecosystem functioning (Cardinale et al. 2012) – including in marine systems (Gamfeldt et al. 2015) – species extinctions may further erode the ability of nearshore habitats to provide goods and services. Thus, there is strong incentive within the conservation movement to protect and manage for high biodiversity in marine realms (Palumbi et al. 2008), in addition to protecting habitat itself.

More recently, researchers have broadened their definition of biodiversity to include the individual roles of species within an ecosystem, represented by their *functional traits*: measurable characteristics of individual organisms relating to their morphology, physiology, ecology, behavior, and life history (Díaz et al. 2013). Functional diversity, then, captures the breadth of variation in functional traits within an assemblage. Traits can quantify the degree of redundancy with an assemblage (Rosenfeld 2002), and thus be used to refine conservation priorities by identifying areas of particular vulnerability (Micheli & Halpern 2005), or that harbor unique or functionally varied assemblages (Devictor et al. 2010, Stuart-Smith et al. 2013). By specifying how organisms perform and interact with one another, traits can be also be used to mechanistically link patterns and processes (Díaz & Cabido 2001), generating more accurate and generalizable predictions for the ecosystem consequences of species losses (Mouillot, Bellwood, et al. 2013).

PeerJ PrePrints

In addition to conserving current coastal habitats and their biological diversity, there have been increasing efforts to actively restore lost or degraded systems, including seagrasses (Orth et al. 2006), marshes (Zedler 2000), mangroves (Ellison 2000), and oyster (Beck et al. 2011) and coral reefs (Mumby & Steneck 2008). A recent meta-analysis revealed that aquatic restorations have, on average, recovered 86% of their biodiversity and 80% of their ecosystem services relative to unrestored reference systems (Benayas et al. 2009). These values, however, reveal that restored systems have generally failed to reach levels of structure and functioning exhibited in their pre-disturbance states (Lotze et al. 2011, Duarte et al. 2015). Moreover, many assessments of recovery have focused on quantifying the cover or abundance of the restored habitat itself (Lotze et al. 2011, Duarte et al. 2015), and not the abundance or diversity the organisms that use it, even though it is these communities that underpin many of the services provided by coastal ecosystems (but see Fonseca, Meyer & Hall 1996; Coen *et al.* 2007). Finally, only a handful of studies have integrated functional trait information into their assessment of faunal communities associated with restored ecosystems, which may better reflect the functional recovery of the restored systems by focusing on what constituent organisms are doing, as opposed to replicating some historical community composition (Tullos et al. 2009, Barnes et al. 2014, Nordström et al. 2015).

One of the more successful examples of coastal restorations has been eelgrass (*Zostera marina* L.) in the Delmarva coastal bays of the mid-western Atlantic (Orth & McGlathery 2012). Eelgrass is a foundational marine angiosperm distributed across the northern hemisphere and provides a number of valuable ecosystem services, including nursery habitat, shoreline protection, and carbon storage (Barbier et al. 2011). Eelgrass habitat supports a variety of epifaunal invertebrates, including amphipods, isopods, decapods, and gastropods, which are an

important link between primary producers and secondary consumers (Sobocinski & Latour 2015). The diversity of these epifauna has been implicated in the functioning of temperate seagrass ecosystems, with greater species richness generally leading to increased grazing of fouling epiphytes in experimental mesocosms and field exclusions, and more recently, across 15 field sites in the northern hemisphere (Duffy et al. 2015). Emerging evidence has also shown that variation in the functional traits of these organisms better predicted epifaunal biomass, algal consumption, and transfer to higher trophic levels than species richness alone (Best et al. 2013, Lefcheck & Duffy 2015), suggesting that the functioning of eelgrass systems can, in part, be inferred through functional trait diversity of the grazer community.

Here, we present long-term data from 2001-2003 and again from 2010-2013 on the recovery of associated epifaunal invertebrate communities alongside the successful restoration of eelgrass habitat in the Delmarva coastal bays. Many of these bays had been unvegetated since the early 1930s, after the pandemic decline of eelgrass due to hurricanes and disease (Orth et al. 2010). In the late 1990s, seed-based restoration efforts began to reestablish eelgrass to the region, which at present now supports over 1800 hectares, with more than half occurring in a single area, South Bay (Orth et al. 2013). Despite the rapid reestablishment of this foundational habitat, it remains unclear whether the epifaunal community, which plays a critical role in the maintenance and functioning of these systems, has experienced similar recovery. To address this question, we compared temporal trends in the epifaunal community in South Bay to a nearby persistent bed in Chincoteague Bay, which served as a baseline or reference bed for assessing restoration success. Drift macroalgae was also collected at each site to understand potential sources of recruitment to the restored bed. We coupled these survey data with information on nine functional traits representing differences in morphology, diet, habitat use, and life history (Table A1). We

hypothesized that the epifauna in South Bay would initially reflect macroalgal assemblages, representing the primary recruitment vector, and over time to become more like the reference bed as the communities matured.

## Methods

### *Survey Sites*

Two coastal lagoons were chosen for comparison: a restored bed in South Bay (37.26 N, -75.84 W), and a reference bed in Chincoteague Bay (38.04 N, -75.31 W). Eelgrass had been absent in South Bay since 1933, until seed-based restoration efforts began in 1997. As of 2013, South Bay contained nearly 1500 ha of dense, continuous eelgrass. In contrast, eelgrass recovered naturally after the pandemic decline in Chincoteague Bay, and by the late 1990s constituted a significant area of our study region. By the early 2000s, Chincoteague represented the largest natural eelgrass bed remaining in this region by at least a factor of 10 (Orth et al. 2013), making it the default candidate for comparison. The two are generally expected to share fauna based on their geographic proximity (<100 km) and known ranges of target invertebrates (Pollock 1998).

### *Survey Methods*

Faunal sampling occurred in two distinct periods: (1) 2001-2003 when eelgrass was initially colonizing South Bay and beds were sparse and patchy, covering only 15 ha of bottom; and (2) 2010 -2013 after the restored bed had increased to nearly 1500 ha. Sampling was conducted quarterly from May to November each year. In each bay, 2-22 random samples (contingent on the area sampled) were collected using suction sampling (Orth & van Montfrans 1987). At low tide, a weighted 0.33-m<sup>2</sup> metal cylinder was placed over a continuous stand of

eelgrass. The suction head was then inserted into the top of the sampling frame and the contents in the cylinder were suctioned into a 0.8-mm mesh bag. Sampling continued for two minutes after which collection bags were sealed, returned to the laboratory, and frozen. In the laboratory, mobile epifauna were enumerated, identified to lowest possible taxon, and preserved in 90% ethyl alcohol. Additionally, we sampled epifauna of drift macroalgae to investigate potential vectors of recruitment. Macroalgal samples were taken by gently enclosing drift algae in the general vicinity of the sampling location in fine mesh bags, and storing and processing the samples identically to the suction samples. The number of algal samples taken during each sampling date ranged from 5-38, with the exact number dependent on the availability of drift macroalgae, which declined in beds that were denser and more continuous.

### *Functional Traits*

We identified nine functional traits based on their relevance to colonization potential, competitive interactions, and effects on ecosystem functioning. Traits included indicators of defense and morphology (exoskeleton material and body plan), trophic ecology (trophic group and specific diet), body size (maximum length), habitat use (mobility and position in the water column), and life history (egg retention and development mode). These traits have been shown to discriminate among epifaunal invertebrates, and to describe and predict benthic community structure and function (Bremner et al. 2003, 2006, Best et al. 2013, Lefcheck & Duffy 2015). The trait data were collected directly from peer-reviewed literature and verified using expert knowledge of the organisms. A detailed description of these traits, including their units and functional interpretations, is given in Table A1. The trait data themselves and a bibliography of references from which they were extracted are found in Supplement 1.

### *Environmental Covariates*

Water quality data from 2001-2003 for South Bay was sampled monthly by the Virginia Coastal Long-Term Ecological Reserve, and from 2010-2013 by the Virginia Estuarine and Coastal Monitoring System program at the Virginia Institute of Marine Science. Physical variables were measured using a YSI 6600 EDS sensor array (YSI Inc., Yellow Springs, Ohio). Additional water samples were taken at eight separate sampling stations to quantify total suspended solids, chlorophyll-*a*, and phosphate concentrations. Further details can be found in (Orth et al. 2012, McGlathery & Christian 2014). Water quality data from 2001-2013 for Chincoteague Bay was collected by the National Park Service at Assateague Island National Seashore (NPS) and the Maryland Department of Natural Resources (DNR) Water Quality Monitoring Programs. Both programs recorded on-site water quality parameters, and collected samples to send to laboratories for nutrient and chlorophyll-*a* analyses. Further details can be found in (Wazniak et al. 2007).

Eelgrass bed area and density in both bays were derived from aerial photography acquired annually from 2001 through 2013, except for 2005, when weather conditions prohibited photography. Black and white photography was acquired at a scale of 1:24,000 along two flight lines covering all shorelines and adjacent shoal areas of the two bays. Aerial photography was scanned from negatives, and *Z. marina* bed boundaries were then directly photo-interpreted on-screen while maintaining a fixed scale using ESRI ArcMap GIS software (ESRI, Redlands CA). *Z. marina* beds were categorized as very sparse (1-10% cover), sparse (11-40% cover), moderate (41-70% cover), or dense (70-100% cover) based on a visual estimate of the percent cover. Ground surveys were conducted in the bays each year to corroborate the occurrence of *Z. marina* identified in the photography outside the boundaries of the seeded plots. Further details can be found in (Orth et al. 2010).



167 *Statistical Analysis*

168 To adjust for varying sampling effort through time, we averaged all observations for a  
169 particular quarterly sampling period so that each period constituted a single replicate (e.g.,  
170 Dornelas et al. 2014). Then, for each replicate, we calculated a number of community metrics.  
171 First, we summed the total number of mean individuals observed in each averaged replicate to  
172 yield a measure of mean total abundance. Next, we calculated species richness ( $S$ ), as the mean  
173 number of species observed, and Simpson diversity ( $D$ ), which further incorporates information  
174 on species' (mean) relative abundances. For Simpson diversity, we converted the resulting  
175 probability values to 'effective number' of species using the transformation from (Jost 2006):

$$D_{eff} = \frac{1}{(1 - D)} \quad (1)$$

176 which obey the 'doubling property' and scales the values comparably with species richness. We  
177 used these two indices to construct an index of relative evenness ( $J$ ), which is calculated as:

$$J = \frac{\ln(D)}{\ln(S)} \quad (2)$$

178 Unlike other indices of evenness, this index of evenness has been shown to independent of both  
179 richness and Simpson diversity (Jost 2010).

180 For functional diversity (FD) we used the index of Rao's quadratic entropy (Rao 1982).  
181 This index has been shown to be independent of species richness (Botta-Dukát 2005), and is  
182 calculated using the following equation:

$$Q = \sum_{i=1}^{S-1} \sum_{j=i+1}^S d_{ij} p_i p_j \quad (3)$$

183 where  $d_{ij}$  is the difference between species  $i$  and  $j$  based on their functional traits,  $p_i$  is the  
184 relative abundance of species  $i$ , and  $p_j$  is the relative abundance of species  $j$ . The differences  $d_{ij}$

are derived from any combination of functional traits using Gower's dissimilarity measure (Gower 1971), which allows both continuous and categorical trait data to be collapsed into a single continuous distance measure. Because Rao's  $Q$  can be maximized when fewer than all functional types are present when Gower's distances are not ultrametric (Pavoine et al. 2005), we generated ultrametric distances using the procedure described in (Mouchet et al. 2008, M  rigot et al. 2010). In the event where species are either the same or different,  $d_{ij}$  collapses to 0 or 1 and  $Q = D$ , or Simpson diversity. Thus, Rao's  $Q$  can also be transformed using Equation (1) to yield units that are on the same scale as Simpson diversity. We calculated two separate indices of FD: one weighted by relative abundances, which we interpret as the functional analogue to Simpson diversity, and one based on presence-absence, or the functional analogue to species richness.

Finally, to further account for variable effort, we generated rarefied estimates of species richness and Simpson Diversity using fixed-coverage subsampling (Chao & Jost 2012). This method generates a value of sample coverage (i.e., how complete the sample is) based on the number of observed singletons (species for which only 1 individual was observed), and estimates of diversity are then rarefied down to a fixed level of coverage, as opposed to a number of individuals or samples. This method results in less data loss than traditional rarefaction, and more efficiently ranks communities based on their true diversities (Chao & Jost 2012). Rarefied estimates were calculated using the *iNEXT* package in R (Hsieh et al. 2014).

### *Statistical Analysis*

Differences among the community metrics for each bay and sampling period were assessed using general linear mixed effects models with an autoregressive 1 (AR1) correlation structure corresponding to each sampling period. Assumptions of normality of errors and homogeneity of variance were assessed visually. Models were constructed using the *nlme*

package in R (Pinheiro et al. 2013). The role of environmental covariates in driving the observed patterns was then estimated using random forests (RF, (Breiman 2001)). We built an ensemble of 50 trees, after which we observed no appreciable decrease in mean squared error, with ‘out-of-bag’ (OOB) permutation to estimate importance partitioned based on Pearson correlation threshold  $\geq 0.5$ . Each community metric (abundance, richness, etc.) was regressed against the following variables that had near complete coverage across all sampling locations and periods: year, bed location, area, and density, temperature, salinity, dissolved oxygen, pH, and water column chl-*a* and phosphates (as a proxy for suspended organic material). We then evaluated the importance of each predictor by quantifying the mean percent decrease in accuracy, derived from the increase in mean error across all trees in the ensemble when a given variable is randomly permuted and then challenged with the OOB data. Model parameters were optimized using the *caret* package (Kuhn 2015), and RFs were constructed using the *randomForest* package in R (Liaw & Wiener 2002). For the top predictors, we plotted bivariate correlations and tested whether the observed correlation was significantly different from zero by comparing values to a *t*-distribution with  $n - 2$  degrees of freedom.

We used non-metric multidimensional scaling (NMDS) to visualize differences in multivariate community structure across bays, algal vs. eelgrass samples, and through time. Because algal samples were collected differently than eelgrass samples, we used a sample-by-species incidence matrix to downplay the influence of the amount of habitat or area sampled. We also ran a separate NMDS for each habitat type using relative abundances, applying a Wisconsin square-root transformation to reduce the impact of highly abundant species. To statistically test for differences among species composition through time, we conducted PERMANOVA, which partitioned variance in the community dissimilarity matrix as a function of sampling period, and

used random permutations of the data to assess significance (McArdle & Anderson 2001). We conducted the NMDS and PERMANOVA using the *vegan* package (Oksanen et al. 2013).

All statistical analyses were conducted in R version 3.1.2 (R Development Core Team 2015), and we held a study-wide  $\alpha = 0.05$  for assessing statistical significance. All data and R scripts used to conduct the analyses are provided in Supplement 1.

## Results

In contrast to our expectations that the restored bed would appear more like the reference bed through time, the two were not significantly different in terms of most community metrics even after only one year (Fig. 1,  $P > 0.05$  based on output from general linear mixed effects models). The two exceptions were average abundance, which was significantly lower in South Bay during the period from 2001-2003 (Fig. 1a,  $P = 0.027$ ), and functional diversity based on presence-absence, which was, on average, significantly higher (Fig. 1f,  $P < 0.001$ ). By the early 2010s, all but one community metric were actually higher in the restored bed (Fig. 1b-f,  $P < 0.001$ ). Functional diversity weighted by relative abundance in particular was 15-32% higher in South Bay than in Chincoteague Bay from 2010-2013 (Fig. 1e). Average abundance was still significantly lower in South Bay during the later period ( $P < 0.001$ ), although we note the expected difference between the two was approximately 25% less than in the earlier period (based on parameter estimates from mixed models).

To understand the factors driving the rapid recovery and ultimate transcendence of the restored bed, we fit random forests for each community metric to a suite of environmental and physical covariates. Random forests consistently identified bed location, total area and mean density to be the top three most important predictors across all community metrics (Fig. A1).

PeerJ PrePrints

Bivariate plots revealed strong and highly significant positive correlations between area and density for almost all metrics of diversity in South Bay (Fig. 2), particularly estimated richness and Simpson diversity (Fig. 2b, c) and functional diversity weighted by relative abundance (Fig. 2f). Chincoteague Bay showed generally weaker or even opposite trends in these bivariate correlations, which may at first appear to be at odds with the output from the random forest analysis. However, it is important to recall that random forests account for collinearity and interactions among variables, and thus the ranking reflects the importance independent of other confounding factors, while the bivariate plots do not.

Exploration of individual trait values revealed that assemblages in the South Bay bed generally exhibited greater variation in traits related to habitat use, life history, and morphology, particularly during the later stages of restoration (Fig. A2). Specifically, South Bay had proportionally more benthic-dwelling crawlers, with calcium carbonate exoskeletons, and external release of eggs with subsequent planktonic dispersal (Figs. A3-7). Thus, it appears that gastropods are responsible for a majority of the difference in functional diversity between the restored and reference beds. Even within this taxonomic group, however, richness, Simpson, and functional diversity weighted by relative abundance were all higher in South Bay during the late period (Fig. A8).

Analysis of multivariate community data using non-metric multidimensional scaling and PERMANOVA revealed that the composition of the restored bed changed significantly from the early period of restoration to the late period (Fig. 3), particularly when factoring in relative abundances ( $P < 0.001$  based on 9,999 random permutations of the data, Fig. A9). Shifts in community composition were principally driven by declines in the relative abundances of the grazing amphipod *Batea catharinensis*, the grazing snail *Astyris lunata*, and the shrimp

276 *Hippolyte pleuracanthus*, and by the appearance of the snail *Costoanachis avara*. In contrast,  
277 there were no significant changes in community structure across years in Chincoteague Bay  
278 based on relative abundances ( $P > 0.05$ , Fig. 3, Fig. A9). Further, the changes in community  
279 structure through time did not appear to be a consequence of recruitment via drift macroalgae,  
280 with the macroalgal communities in each bay being significantly distinct from the corresponding  
281 grass beds ( $P < 0.001$ , Fig. 3).

282 Of the 42 species captured over the entire survey, five were found exclusively in South  
283 Bay, including three gastropod species in the genus *Costoanachis*, as well as the grazing snail  
284 *Marshallora nigrocincta*, and the detritivorous amphipod *Photis macrocoxa* (although this  
285 species was found only on drift macroalgae). All five of these species occupied unique parts of  
286 functional trait space, based on principal coordinates analysis of their trait values (Fig. A10).  
287 Only two species – the grazing isopod *Paracerceis caudata* and the detritivorous amphipod  
288 *Lysianopsis alba* – were found exclusively in Chincoteague Bay, and they were nearly  
289 functionally equivalent with other species (Fig. A10).

## 290 Discussion

291 In this study, we found that the diversity of epifaunal invertebrates and their functional  
292 traits in a restored eelgrass bed matched and then exceeded values exhibited by a nearby  
293 reference bed in less than a decade (Fig. 1). These invertebrates, and particularly their diversity,  
294 are known to play a critical role in the maintenance, structure, and functioning of eelgrass  
295 ecosystems (Duffy et al. 2014). Further, the rapid recovery of the epifaunal community was  
296 shown to be a direct consequence of restoration actions that encouraged the growth and  
297 expansion of the restored bed, represented by bed area and density (Fig. 2). These findings may  
298 be of particular consequence as eelgrass cover has generally declined in the region (Orth et al.

299 2013) and worldwide (Waycott et al. 2009), and is likely to continue this downward trajectory  
300 without further intervention. The area of the reference bed, for example, declined 49% over the  
301 12 years of the survey (Fig. 4a), presumably as a consequence of decreasing water quality and  
302 thermal stress known to drive eelgrass decline in the nearby Chesapeake Bay (Wazniak et al.  
303 2007, Moore et al. 2012). In contrast, the area of the restored bed increased by 910% over the  
304 same time period (Fig. 4b). Thus, if these trends continue, then restored beds in an environment  
305 not challenged by poor water quality may play an increasingly vital role in providing habitat for  
306 a variety of taxa and functional forms present in the region. Indeed, restored beds now make up  
307 roughly one-third of total seagrass cover in the Delmarva coastal bays (Orth et al. 2013).

308 If many aquatic restorations fail to achieve pre-disturbance levels of diversity (Benayas et  
309 al. 2009), including in other seagrasses (Paling et al. 2009), then why has this particular effort  
310 been so successful? Low human impacts in this region are undoubtedly a major factor. Long-  
311 term monitoring by the Virginia Coastal LTER has shown that nutrient loading and water  
312 column chlorophyll-*a* have been low and relatively stable across all of Virginia's coastal bays  
313 over the past two decades (Orth & McGlathery 2012). The bays also experience regular flushing  
314 with cooler ocean water, alleviating temperature stress that is a major driver of eelgrass die-  
315 backs in nearby regions, such as the Chesapeake Bay (Moore et al. 2012). Moreover, as the  
316 restored bed has increased in size and density, it has more effectively captured particulates,  
317 leading to increased water clarity and greater light availability for photosynthesis (Moore et al.  
318 2012, Orth et al. 2012). This self-facilitation has been identified as a key predictor of the  
319 persistence of restored seagrass ecosystems (van Katwijk et al. 2009).

320 Altogether, the factors that promote the growth and expansion of the restored bed should,  
321 in theory, facilitate the recruitment and survival of epifaunal invertebrates, principally by

increasing the availability of habitat and substrate for algal resources. While the positive relationship between area sampled and species richness has been established generally, there are few examples from seagrass systems (Boström et al. 2006). Our study is also unique in the sense that increasing area does not simply represent a shift in geographic focus, but a product of ongoing efforts to actively increase the total habitat in the same location, and the natural spread of eelgrass from the initial effort to restore this habitat. Moreover, the presence and diversity of these grazers has been shown to promote their eelgrass habitat both in experiments and observational studies, generating a positive feedback where habitat begets diversity, and so on. Thus, the abundant and functionally diverse grazer community in the restored bed may play an important role in the self-facilitation mentioned above.

The reproductive biology and dispersal abilities of the focal organisms may have also played a large role in the rapid recovery of the epifaunal community in the restored bed. Recent reviews have suggested that ‘successful restorations’ – those that have achieved similar levels of biodiversity as before a disturbance – have concerned communities of fast-reproducing organisms with high dispersal ability (Jones & Schmitz 2009, Duarte et al. 2015). While there did not appear to be a change in the proportion of individuals belonging to high vs. low dispersal species throughout the course of restoration – those that brooded or released their eggs, and those with planktonic vs. non-planktonic larval forms (Figs. A6-7) – most of the organisms in this system have generation times on the scale of weeks to months. The amphipod *Gammarus mucronatus*, for instance, can reach reproductive maturity in as few as three weeks during hotter months (Fredette & Diaz 1986). The quick generation and turnover of populations certainly contributed to the ability of these small invertebrates to reach and exploit this new habitat.



344 A complementary hypothesis is that, as the bed expanded, it also recruited mesopredators  
345 of epifauna, including various fishes and crabs, which also utilize the eelgrass as habitat.  
346 Predator trawls conducted in South Bay the early 2010s revealed a diverse and abundant predator  
347 community (Schmitt et al., *submitted*). A recent meta-analysis produced strong evidence  
348 supporting the long-standing hypothesis that predation enhances prey diversity by relieving  
349 competition among prey species (Katano et al. 2015), a trend which has been seen in mesocosm  
350 experiments using several of the species observed in our surveys (Duffy et al. 2005, Douglass et  
351 al. 2008). Consistent with an active and increasingly abundant predator community, epifaunal  
352 abundance actually had an inverse relationship with bed area and density in South Bay (Fig. 2a).  
353 An alternative explanation is that epifauna are resource-limited, as suggested by their remarkably  
354 invariant mean abundances observed in the later stages of restored bed (Fig. 1a). Thus,  
355 increasing the density of habitat but not resources could have diluted the average abundance per  
356 unit sampling area.

357 One of the more striking differences between the restored and reference beds was the  
358 higher functional diversity in South Bay compared to Chincoteague Bay (Fig. 1e-f). Habitat  
359 complexity and resource availability, proxied by mean density and total phosphates in our  
360 random forest analysis, may have played a large role here as well (Fig. 2e-f, Fig. A1). Increasing  
361 both should theoretically open niche space and facilitate coexistence, leading to a wider variety  
362 of observed functional morphologies, behaviors, and life histories. Classical successional theory  
363 would predict that this would be especially true during the later stages of restoration, as biotic as  
364 opposed to abiotic filters play an increasingly larger role and functionally similar species are  
365 weeded out as consequence of competitive interactions (Connell & Slatyer 1977). Given these  
366 predictions, it is surprising that we observed significantly higher functional diversity in the

367 restored bed compared to the reference bed in the early stages of restoration. Once again, the  
368 relatively fast recruitment and generation times of the organisms may play a role, allowing such  
369 interactions to play out on much shorter time scales than might be expected in other systems,  
370 such as terrestrial plants (Purschke et al. 2013).

371 A recent publication also suggested that it is not competitive interactions, but continued  
372 colonization, that drives the functional trait structure of successional communities (Li et al.  
373 2015). We observed significant differences in community composition through time, particularly  
374 in the restored bed (Fig. 3), which was a consequence of the arrival of new species or shifts in  
375 dominance. Along similar lines, the demography of these epifaunal invertebrates is notoriously  
376 unpredictable. In nearby systems in the Chesapeake Bay, which have been well studied for  
377 decades, species appear and then disappear regularly, generally in response to loss of habitat  
378 (Douglass et al. 2010). Similar stochastic processes may also be at play here, with high trait  
379 diversity in the early years simply an outcome of chance processes that favored recruitment of  
380 functionally varied individuals to the sudden appearance of uncolonized habitat.

381 Finally, we must acknowledge the possibility that we are still observing some transitional  
382 state, and that the restored bed is still progressing towards the stable state exhibited by the  
383 reference bed (Fig. 3). Several lines of evidence suggest this is not the case. First, inter- and  
384 intra-annual variability was similar between both bays (Fig. 1), suggesting that the reference bed  
385 had not achieved a more stable, and thus mature, state than the restored bed. In fact, the error  
386 around most of the means in the restored bed is actually lower than in the reference bed (Fig. 1).  
387 Second, we observed significant compositional differences between the beds, suggesting  
388 potentially different avenues for colonization (Fig. 3). Finally, given the contrasting trajectories  
389 of the two beds (Fig. 4), it may be unreasonable to suspect these two communities to ever

converge despite their proximity and the known ranges of the epifaunal species considered in this analysis.

Aggregating at broad temporal scales emphasizes overall trajectories of these beds, but ignores some of the finer scale variation. One particularly notable instance occurred in June 2012, with the recruitment of juvenile pinfish (*Lagodon rhomboides*) to these coastal bays. During early life history stages, pinfish are known to be voracious predators of epifaunal invertebrates (Luczkovich & Stellwag 1993), and were indisputably responsible for the steep declines in epifaunal abundance and diversity during this month (Fig. A11). Generally, pinfish do not range this far north, but with increasing temperatures, we may be witness to a northward range expansion. Such a shift may have drastic consequence for the eelgrass: first, by removing the critical top-down control provided by epifaunal grazers, and second, by directly consuming eelgrass, particularly later during their ontogeny (Luczkovich & Stellwag 1993). Thus, while local conditions over the last decade have favored the growth and expansion of eelgrass in Virginia's coastal bays, climate change may introduce a new variable in the form of pinfish with uncertain consequences for the maintenance of eelgrass diversity and function in South Bay.

Within the larger context of restoration ecology, our study stands out in that it documents the transition from a near complete lack of vegetation to dense, contiguous beds (Fig. 4, Orth & McGlathery 2012). This state-shift stands in contrast to many other restoration projects, which have focused on recovery of degraded, but not totally destroyed, systems. The drastic transition from a near unvegetated to a vegetated system may explain some of our results by widening the scope for detectable effects. We note, however, that the greatest differences occurred during the later stages of restoration, when the restored and reference beds were comparable in terms of their total coverage (Fig. 4). Further, even in degraded eelgrass beds, remediation of

environmental stressors has been shown to enhance the functional trait diversity of epifaunal invertebrates (Dolbeth et al. 2013). As with our results (Fig. 1e), functional recovery was partly a consequence of the redistribution of biomass among different functional types (Dolbeth et al. 2013). Similarly rapid recovery of functional diversity has been observed in other invertebrate communities, such as beetles in restored wetlands (Watts & Mason 2015). These studies, along with ours, suggest that functional traits can be a powerful tool for assessing the response and subsequent recovery of communities to disturbance (Mouillot, Graham, et al. 2013).

It is worth noting a few limitations of this study. First, we are comparing two bays, even though these bays combined account for 88% of the submersed aquatic vegetation in the Delmarva coastal bays. While our results may not be representative of eelgrass systems in general, or even restored systems, we do demonstrate that restored ecosystems at least have the potential to greatly exceed the diversity of reference systems, which is a relatively rare outcome in restoration ecology (Benayas et al. 2009). Second, we had variable sampling effort through time, largely as a consequence of available habitat to sample. To further explore the potential for bias, we additionally weighted our regression analyses by the inverse of sample size ( $1 / N$ ) to explore the relative influence of less intensively sampled periods (see Supplement 1). This exercise yielded nearly identical results, with the one exception that functional diversity weighted by relative abundance in the early period was now significantly higher in the restored bed than the reference bed ( $P = 0.017$ ), where it was previously non-significant in the unweighted regression (Fig. 1e). Finally, we did not have robust and consistent estimates of the local density of the habitat to use in our analysis. Future monitoring efforts should focus on a method that is not only standard to the area of bottom sampled, but also yields some value of habitat availability (biomass, etc.).

PeerJ PrePrints

In this study, we show that epifaunal community diversity in a restored eelgrass bed was significantly higher than an unrestored bed in the region, particularly when considering the breadth of organismal functional traits. Given the wealth of evidence linking both epifaunal species and trait diversity to enhanced ecosystem functioning (Best et al. 2013, Duffy et al. 2015, Lefcheck & Duffy 2015), it is reasonable to infer that the restored bed may also experience higher levels and greater stability in ecosystem functioning than unrestored beds. Indeed, sediment organic content and ammonium exchange have been shown to be higher during the later stages of restoration (McGlathery et al. 2012). While these functions were attributed to the recovery of the eelgrass itself, previous mesocosm experiments have shown that epifaunal richness also impacts these properties as well, irrespective of eelgrass biomass (Duffy et al. 2003, Spivak et al. 2007). Future work, however, will have to directly quantify processes related to these grazers, such as secondary production, to demonstrate their independent contributions to the eelgrass community function. In the interim, the restored systems in Virginia's coastal bays may serve an important role in conserving regional biological diversity in the face of large-scale declines in existing beds.

## Acknowledgements

We foremost thank the past and current members of the VIMS SAV lab for the collection and sampling of the hundreds of benthic samples taken over the course of this survey, but especially C Holbert, S Sumoski, A Johnson, and P Trivett. We are also indebted to B Lusk and B Truitt of the Nature Conservancy for their assistance for the duration of the project. Funding for all faunal work and later water quality data in South Bay was provided by the grants from numerous agencies notably: the Coastal Programs of the Virginia Department of Environmental Quality funded by Coastal Zone Management Act of 1972, as amended, administered by NOAA's Office

of Ocean and Coastal Resource Management, the Virginia Recreational Fishing License Fund, the American Recovery and Reinvestment Act with funding to NOAA, Grant NA09NMF4630308, The Nature Conservancy, U.S. Army Corps of Engineers, as well as private grants from the Allied-Signal Foundation, Norfolk-Southern, and the Keith Campbell Foundation for the Environment. Early South Bay water quality data were provided J Porter and K McGlathery at Virginia Coast Reserve Long-Term Ecological Research project, supported by the National Science Foundation under Grants No. BSR-8702333-06, DEB-9211772, DEB-9411974, DEB-0080381, DEB-0621014 and DEB-1237733. Chincoteague Bay water quality data were provided by C Wazniak at the Maryland Department of Natural Resources Tidewater Ecosystem Assessment Division, funded by the State of Maryland. We thank C Patrick for comments on a previous version of this manuscript. This is contribution no. ##### of the Virginia Institute of Marine Science.

## References

- Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR (2011) The value of estuarine and coastal ecosystem services. *Ecol Monogr* 81:169–193
- Barnes AD, Emberson RM, Krell F (2014) The Role of Species Traits in Mediating Functional Recovery during Matrix Restoration. *PLoS One*:1–19
- Beck MW, Brumbaugh RD, Airoidi L, Carranza A, Coen LD, Crawford C, Defeo O, Edgar GJ, Hancock B, Kay MC, Lenihan HS, Luckenbach MW, Toropova CL, Zhang G, Guo X (2011) Oyster Reefs at Risk and Recommendations for Conservation, Restoration, and Management. *Bioscience* 61:107–116
- Benayas JMR, Newton AC, Diaz A, Bullock JM (2009) Enhancement of Biodiversity and Ecosystem Services by Ecological Restoration: A Meta-Analysis. *Science* (80- ) 325:1121–1124
- Best RJ, Caulk NC, Stachowicz JJ (2013) Trait vs. phylogenetic diversity as predictors of competition and community composition in herbivorous marine amphipods. *Ecol Lett* 16:72–80

- 486 Boström C, Jackson EL, Simenstad CA (2006) Seagrass landscapes and their effects on  
487 associated fauna: A review. *Estuar Coast Shelf Sci* 68:383–403
- 488 Botta-Dukát Z (2005) Rao's quadratic entropy as a measure of functional diversity based on  
489 multiple traits. *J Veg Sci* 16:533–540
- 490 Breiman L (2001) Random forests. *Mach Learn* 45:5–32
- 491 Bremner J, Paramor OAL, Frid CLJ (2006) Developing a methodology for incorporating  
492 ecological structure and functioning into designation of Special Areas of Conservation  
493 (SAD) in the 0-12 nautical mile zone. :1–158
- 494 Bremner J, Rogers SI, Frid CLJ (2003) Assessing functional diversity in marine benthic  
495 ecosystems: a comparison of approaches. *Mar Ecol Prog Ser* 254:11–25
- 496 Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM,  
497 Tilman D, Wardle DA, Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A,  
498 Srivastava DS, Naeem S (2012) Biodiversity loss and its impact on humanity. *Nature*  
499 486:59–67
- 500 Chao A, Jost L (2012) Coverage-based rarefaction and extrapolation: standardizing samples by  
501 completeness rather than size. *Ecology* 93:2533–2547
- 502 Coen LD, Brumbaugh RD, Bushek D, Grizzle R, Luckenbach MW, Posey MH, Powers SP,  
503 Tolley SG (2007) Ecosystem services related to oyster restoration. *Mar Ecol Prog Ser*  
504 341:303–307
- 505 Connell JH, Slatyer RO (1977) Mechanisms of succession in natural communities and their role  
506 in community stability and organization. *Am Nat* 111:1119–1144
- 507 Devictor V, Mouillot D, Meynard C, Jiguet F, Thuiller W, Mouquet N (2010) Spatial mismatch  
508 and congruence between taxonomic, phylogenetic and functional diversity: the need for  
509 integrative conservation strategies in a changing world. *Ecol Lett* 13:1030–1040
- 510 Díaz S, Cabido M (2001) Vive la difference: plant functional diversity matters to ecosystem  
511 processes. *Trends Ecol Evol* 16:646–655
- 512 Díaz S, Purvis A, Cornelissen JHC, Mace GM, Donoghue MJ, Ewers RM, Jordano P, Pearse  
513 WD (2013) Functional traits, the phylogeny of function, and ecosystem service  
514 vulnerability. *Ecol Evol* 3:2958–2975
- 515 Dolbeth M, Cardoso P, Grilo T, Raffaelli D, Pardal MÂ (2013) Drivers of estuarine benthic  
516 species distribution patterns following a restoration of a seagrass bed: A functional trait  
517 analyses. *Mar Pollut Bull* 72:47–54



- 518 Dornelas M, Gotelli NJ, McGill B, Shimadzu H, Moyes F, Sievers C, Magurran AE (2014)  
519 Assemblage time series reveal biodiversity change but not systematic loss. *Science* (80- )  
520 344:296–299
- 521 Douglass JG, Duffy JE, Bruno JF (2008) Herbivore and predator diversity interactively affect  
522 ecosystem properties in an experimental marine community. *Ecol Lett* 11:598–608
- 523 Douglass JG, France KE, Paul Richardson J, Duffy JE (2010) Seasonal and interannual changes  
524 in a Chesapeake Bay eelgrass community: Insights into biotic and abiotic control of  
525 community structure. *Limnol Oceanogr* 55:1499–1520
- 526 Duarte CM, Borja A, Carstensen J, Elliott M, Krause-Jensen D, Marbà N (2015) Paradigms in  
527 the Recovery of Estuarine and Coastal Ecosystems. *Estuaries and Coasts* 38:1202–1212
- 528 Duffy JE, Hughes AR, Moksnes P-O (2014) Ecology of Seagrass Communities. In: Bertness  
529 MD, Silliman BR, Bruno JF, Stachowicz JJ (eds) *Marine Community Ecology and*  
530 *Conservation*. Sinauer, Sunderland, MA, p 271–297
- 531 Duffy JE, Paul Richardson J, France KE (2005) Ecosystem consequences of diversity depend on  
532 food chain length in estuarine vegetation. *Ecol Lett* 8:301–309
- 533 Duffy JE, Reynolds PL, Boström C, Coyer JA, Cusson M, Donadi S, Douglass JG, Eklöf JS,  
534 Engelen AH, Eriksson BK, Fredriksen S, Gamfeldt L, Gustafsson C, Hoarau G, Hori M,  
535 Hovel K, Iken K, Lefcheck JS, Moksnes P-O, Nakaoka M, O'Connor MI, Olsen JL,  
536 Richardson JP, Ruesink JL, Sotka EE, Thormar J, Whalen MA, Stachowicz JJ (2015)  
537 Biodiversity mediates top-down control in eelgrass ecosystems: A global comparative-  
538 experimental approach. *Ecol Lett*:n/a–n/a
- 539 Duffy JE, Richardson JP, Canuel EA (2003) Grazer diversity effects on ecosystem functioning in  
540 seagrass beds. *Ecol Lett* 6:637–645
- 541 Ellison AM (2000) Mangrove restoration: Do we know enough? *Restor Ecol* 8:219–229
- 542 Fonseca MS, Meyer DL, Hall MO (1996) Development of planted seagrass beds in Tampa Bay,  
543 Florida, USA. II. Faunal components. *Mar Ecol Prog Ser* 132:141–156
- 544 Fredette TJ, Diaz RJ (1986) Life history of *Gammarus mucronatus* Say (Amphipoda:  
545 Gammaridae) in warm temperate estuarine habitats, York River, Virginia. *J Crustac Biol*  
546 6:57–78
- 547 Gamfeldt L, Lefcheck JS, Byrnes JEK, Cardinale BJ, Emmett Duffy J, Griffin JN (2015) Marine  
548 biodiversity and ecosystem functioning: what's known and what's next. *Oikos* 124:252–265
- 549 Gower JC (1971) A general coefficient of similarity and some of its properties. *Biometrics*  
550 27:857–874



- 551 Halpern BS, Walbridge S, Selkoe KA, Kappel C V., Micheli F, D'Agrosa C, Bruno JF, Casey  
552 KS, Ebert C, Fox HE, Fujita R, Heinemann D, Lenihan HS, Madin EMP, Perry MT, Selig  
553 ER, Spalding M, Steneck R, Watson R (2008) A Global Map of Human Impact on Marine  
554 Ecosystems. *Science* (80- ) 319:948–952
- 555 Hsieh TC, Ma KH, Chao A (2014) iNEXT: iNterpolation and EXTrapolation for species  
556 diversity.
- 557 Jones HP, Schmitz OJ (2009) Rapid recovery of damaged ecosystems. *PLoS One* 4
- 558 Jost L (2006) Entropy and diversity. *Oikos* 113:363–375
- 559 Jost L (2010) The relation between evenness and diversity. *Diversity* 2:207–232
- 560 Katano I, Doi H, Eriksson BK, Hillebrand H (2015) A cross-system meta-analysis reveals  
561 coupled predation effects on prey biomass and diversity. *Oikos*:n/a–n/a
- 562 Katwijk MM van, Bos a. R, Jonge VN de, Hanssen LS a M, Hermus DCR, Jong DJ de (2009)  
563 Guidelines for seagrass restoration: Importance of habitat selection and donor population,  
564 spreading of risks, and ecosystem engineering effects. *Mar Pollut Bull* 58:179–188
- 565 Kuhn M (2015) caret: Classification and Regression Training.
- 566 Lefcheck JS, Duffy JE (2015) Multitrophic functional diversity predicts ecosystem functioning  
567 in experimental assemblages of estuarine consumers. *Ecology*:n/a–n/a
- 568 Li S, Cadotte MW, Meiners SJ, Hua Z, Jiang L, Shu W (2015) Species colonisation, not  
569 competitive exclusion, drives community overdispersion over long-term succession. *Ecol*  
570 *Lett*:n/a–n/a
- 571 Liaw A, Wiener M (2002) Classification and Regression by randomForest. *R News* 2:18–22
- 572 Lotze HK, Coll M, Magera AM, Ward-Paige C, Airoidi L (2011) Recovery of marine animal  
573 populations and ecosystems. *Trends Ecol Evol* 26:595–605
- 574 Luczkovich JJ, Stellwag EJ (1993) Isolation of cellulolytic microbes from the intestinal tract of  
575 the pinfish, *Lagodon rhomboides*: size-related changes in diet and microbial abundance.  
576 *Mar Biol* 116:381–388
- 577 McArdle BH, Anderson MJ (2001) Fitting multivariate models to community data: a comment  
578 on distance-based redundancy analysis. *Ecology* 82:290–297
- 579 McCauley DJ, Pinsky ML, Palumbi SR, Estes JA, Joyce FH, Warner RR (2015) Marine  
580 defaunation: Animal loss in the global ocean. *Science* (80- ) 347:1255641

- 581 McGlathery KJ, Christian RR (2014) Water Quality Sampling - integrated measurements for the  
582 Virginia Coast, 2004-2014. Virginia Coast Reserv Dataset VCR15243
- 583 McGlathery KJ, Reynolds LK, Cole LW, Orth RJ, Marion SR, Schwarzschild A (2012)  
584 Recovery trajectories during state change from bare sediment to eelgrass dominance. Mar  
585 Ecol Prog Ser 448:209–221
- 586 Mérigot B, Durbec J-P, Gaertner J-C (2010) On goodness-of-fit measure for dendrogram-based  
587 analyses. Ecology 91:1850–1859
- 588 Micheli F, Halpern BS (2005) Low functional redundancy in coastal marine assemblages. Ecol  
589 Lett 8:391–400
- 590 Moore KA, Shields EC, Parrish DB, Orth RJ (2012) Eelgrass survival in two contrasting  
591 systems: Role of turbidity and summer water temperatures. Mar Ecol Prog Ser 448:247–258
- 592 Mouchet M, Guilhaumon F, Villéger S, Mason NWH, Tomasini J-A, Mouillot D (2008)  
593 Towards a consensus for calculating dendrogram-based functional diversity indices. Oikos  
594 117:794–800
- 595 Mouillot D, Bellwood DR, Baraloto C, Chave J, Galzin R, Harmelin-Vivien M, Kulbicki M,  
596 Lavergne S, Lavorel S, Mouquet N, Paine CET, Renaud J, Thuiller W (2013) Rare species  
597 support vulnerable functions in high-diversity ecosystems. PLoS Biol 11:e1001569
- 598 Mouillot D, Graham NAJ, Villéger S, Mason NWH, Bellwood DR (2013) A functional approach  
599 reveals community responses to disturbances. Trends Ecol Evol 28:167–177
- 600 Mumby PJ, Steneck RS (2008) Coral reef management and conservation in light of rapidly  
601 evolving ecological paradigms. Trends Ecol Evol 23:555–563
- 602 Nordström MC, Demopoulos AWJ, Whitcraft CR, Rismondo A, McMillan P, Gonzalez JP,  
603 Levin LA (2015) Food web heterogeneity and succession in created saltmarshes.
- 604 Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos  
605 P, Stevens MHH, Wagner H (2013) vegan: Community Ecology Package.
- 606 Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, James W, Heck Jr. KL, Hughes AR,  
607 Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M, Williams SL (2006) A  
608 global crisis for seagrass ecosystems. Bioscience 56:987–996
- 609 Orth RJ, Marion SR, Moore KA, Wilcox DJ (2010) Eelgrass (*Zostera marina* L.) in the  
610 Chesapeake Bay region of mid-Atlantic coast of the USA: Challenges in conservation and  
611 restoration. Estuaries and Coasts 33:139–150
- 612 Orth RJ, McGlathery KJ (2012) Eelgrass recovery in the coastal bays of the Virginia Coast  
613 Reserve, USA. Mar Ecol Prog Ser 448:173–176

- 614 Orth R, Montfrans J van (1987) Utilization of a seagrass meadow and tidal marsh creek by blue  
615 crabs *Callinectes sapidus*. I. Seasonal and annual variations in abundance with emphasis on  
616 post-settlement juveniles . Mar Ecol Prog Ser 41:283–294
- 617 Orth RJ, Moore KA, Marion SR, Wilcox DJ, Parrish DB (2012) Seed addition facilitates eelgrass  
618 recovery in a coastal bay system. Mar Ecol Prog Ser 448:177–195
- 619 Orth RJ, Wilcox DJ, Whiting JR, Nagey L, Kenne AK, Smith ER (2013) 2013 Distribution of  
620 Submerged Aquatic Vegetation in the Chesapeake Bay and Coastal Bays.
- 621 Paling EI, Fonseca M, Katwijk MM van, Keulen M van (2009) Seagrass Restoration. In: Perillo  
622 GME, Wolanski E, Cahoon DR, Brinson M (eds) Coastal Wetlands - An Integrated  
623 Ecosystem Approach. Elsevier, Amsterdam, p 687–713
- 624 Palumbi SR, Sandifer PA, Allan JD, Beck MW, Fautin DG, Fogarty MJ, Halpern BS, Incze LS,  
625 Leong J-A, Norse E, Stachowicz JJ, Wall DH (2008) Managing for ocean biodiversity to  
626 sustain marine ecosystem services. Front Ecol Environ 7:204–211
- 627 Pavoine S, Ollier S, Pontier D (2005) Measuring diversity from dissimilarities with Rao's  
628 quadratic entropy: Are any dissimilarities suitable? Theor Popul Biol 67:231–239
- 629 Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2013) nlme: Linear and Nonlinear  
630 Mixed Effects Models.
- 631 Pollock LW (1998) A Practical Guide to the Marine Animals of Northeastern North America.  
632 Rutgers University Press, New Brunswick, NJ
- 633 Purschke O, Schmid BC, Sykes MT, Poschlod P, Michalski SG, Durka W, Kühn I, Winter M,  
634 Prentice HC (2013) Contrasting changes in taxonomic, phylogenetic and functional  
635 diversity during a long-term succession: insights into assembly processes (J Fridley, Ed.). J  
636 Ecol 101:857–866
- 637 R Development Core Team (2015) R: A Language and Environment for Statistical Computing.
- 638 Rao CR (1982) Diversity and dissimilarity coefficients: A unified approach. Theor Popul Biol  
639 21:24–43
- 640 Rosenfeld JS (2002) Functional redundancy in ecology and conservation. Oikos 98:156–162
- 641 Sobocinski KL, Latour RJ (2015) Trophic transfer in seagrass systems : estimating seasonal  
642 production of an abundant seagrass fish , *Bairdiella chrysoura* , in lower Chesapeake Bay.  
643 Mar Ecol Prog Ser 523:157–174
- 644 Spivak AC, Canuel EA, Duffy JE, Richardson JP (2007) Top-down and bottom-up controls on  
645 sediment organic matter composition in an experimental seagrass ecosystem. Limnol  
646 Oceanogr 52:2595–2607

- 647 Stuart-Smith RD, Bates AE, Lefcheck JS, Duffy JE, Baker SC, Thomson RJ, Stuart-Smith JF,  
648 Hill N a., Kininmonth SJ, Airoidi L, Becerro MA, Campbell SJ, Dawson TP, Navarrete SA,  
649 Soler GA, Strain EMA, Willis TJ, Edgar GJ (2013) Integrating abundance and functional  
650 traits reveals new global hotspots of fish diversity. *Nature* 501:539–542
- 651 Tullos DD, Penrose DL, Jennings GD, Cope WG (2009) Analysis of functional traits in  
652 reconfigured channels: implications for the bioassessment and disturbance of river  
653 restoration. *J North Am Benthol Soc* 28:80–92
- 654 Watts CH, Mason NWH (2015) If we build - they mostly come: partial functional recovery but  
655 persistent compositional differences in wetland beetle community restoration. *Restor*  
656 *Ecol:n/a–n/a*
- 657 Waycott M, Duarte CM, Carruthers TJB, Orth RJ, Dennison WC, Olyarnik S, Calladine A,  
658 Fourqurean JW, Heck KL, Hughes a R, Kendrick G a, Kenworthy WJ, Short FT, Williams  
659 SL (2009) Accelerating loss of seagrasses across the globe threatens coastal ecosystems.  
660 *PNAS* 106:12377–81
- 661 Wazniak CE, Hall MR, Carruthers TJB, Sturgis B, Dennison WC, Orth RJ (2007) Linking water  
662 quality to living resources in a mid-Atlantic lagoon system, USA. *Ecol Appl* 17:S64–S68
- 663 Zedler JB (2000) Progress in wetland restoration ecology. *Trends Ecol Evol* 15:402–407

## Figure Captions

**Figure 1:** Times series plotting the annual mean  $\pm$  1 SE for various community properties. Light grey circles represent the mature bed in Chincoteague Bay. Black triangles represent the restored bed in South Bay. FD = functional diversity, calculated as Rao's quadratic entropy from all nine functional traits.

**Figure 2:** Plots of total bed area (in hectares, left column) and mean density (as the average density bin based on comparison of aerial footage against a crown density scale, ranging from 0 for no cover to 1 for full cover) for South Bay. Points represent each year and month of the survey. The black line represents a simple linear regression between the two variables. The Pearson correlation ( $r$ ) from this regression is given in the bottom of each panel along with the following indicator of statistical significance: \*  $P = 0.05$ , \*\*  $P = 0.01$ , \*\*\*  $P = 0.001$ .

**Figure 3:** Non-metric multidimensional scaling conducted on relative abundance data for both eelgrass (green) and drift macroalgal (brown) communities. Small open points represent each year and month of the survey. Large points represent the centroid for each year, graded from early years (dark grey) to later years (light grey). Arrows represent the trajectories from one year to the next. Stress – a measure of concordance between the multivariate data and the two-dimensional representation – is given in the lower right.

**Figure 4:** Total area (in hectares) of eelgrass beds in (a) Chincoteague Bay and (b) South Bay through time. Bars are divided into estimated density bins (% cover) based on comparison of aerial footage to a crown density scale. Note: aerial photography could not be conducted in 2005 due to poor weather conditions.




Figure 1




Figure 2




Figure 3






Figure 4