

A peer-reviewed version of this preprint was published in PeerJ on 4 February 2016.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1666) (peerj.com/articles/1666), which is the preferred citable publication unless you specifically need to cite this preprint.

Plaete J, De Bourdeaudhuij I, Verloigne M, Crombez G. 2016. The use and evaluation of self-regulation techniques can predict health goal attainment in adults: an explorative study. PeerJ 4:e1666
<https://doi.org/10.7717/peerj.1666>

The use and evaluation of self-regulation techniques can predict health goal attainment in adults: an explorative study

Jolien Plaete, Ilse De Bourdeaudhuij, Maite Verloigne, Geert Crombez

Background. Self-regulation tools are not always used optimally, and implementation intention plans often lack quality. Therefore, this study explored participants' use and feasibility evaluation of self-regulation techniques and their impact on goal attainment.

Methods. Data were obtained from 452 adults in a proof of concept (POC) intervention of 'MyPlan', an eHealth intervention using self-regulation techniques to promote three healthy behaviours (physical activity(PA), fruit intake, or vegetable intake). Participants applied self-regulation techniques to a self-selected health behaviour, and evaluated the self-regulation techniques. The quality of implementation intentions was rated by the authors as a function of instrumentality (instrumental and non-instrumental) and specificity (non-specific and medium to high specific). Logistic regression analyses were conducted to predict goal attainment.

Results. Goal attainment was significantly predicted by the motivational value of the personal advice (OR:1.86), by the specificity of the implementation intentions (OR:3.5), by the motivational value of the action plan (OR:1.86), and by making a new action plan at follow-up (OR:4.10). Interaction-effects with behaviour showed that the specificity score of the implementation intention plans (OR:4.59), the motivational value of the personal advice (OR:2.38), selecting hindering factors and solutions(OR:2.00) and making a new action plan at follow-up (OR:7.54) were predictive of goal attainment only for fruit or vegetable intake. Also, when participants in the fruit and vegetable group made more than three plans, they were more likely to attain their goal (OR:1.73), whereas the reverse was the case in the PA group (OR:0.34).

Discussion. Feedback on goal feasibility, coping implementation intentions, further research to investigate the optimal number of plans for different behaviours, the optimal frequency and timing of follow-up modules and new ways to incorporate social support in eHealth interventions, are recommended.

1 **The use and evaluation of self-regulation techniques can predict health goal**
2 **attainment in adults: an explorative study**

3 Jolien Plaete¹, Ilse De Bourdeaudhuij¹, Maïté Verloigne¹, Geert Crombez²

4

5 ¹ Department of Movement and Sports Sciences, Ghent University, Belgium

6 ² Department of Experimental-Clinical and Health Psychology, Ghent University, Belgium

7

8 Corresponding author:

9 Jolien Plaete¹

10 Watersportlaan 2, 9000 Gent, Belgium. Tel.: +32 9 264 6363

11 Email address: jolien.plaete@ugent.be

12

13 **Abstract**

14 **Background.** Self-regulation tools are not always used optimally, and implementation intention
15 plans often lack quality. Therefore, this study explored participants' use and feasibility
16 evaluation of self-regulation techniques and their impact on goal attainment.

17 **Methods.** Data were obtained from 452 adults in a proof of concept (POC) intervention of
18 'MyPlan', an eHealth intervention using self-regulation techniques to promote three healthy
19 behaviours (physical activity(PA), fruit intake, or vegetable intake). Participants applied self-
20 regulation techniques to a self-selected health behaviour, and evaluated the self-regulation
21 techniques. The quality of implementation intentions was rated by the authors as a function of
22 instrumentality (instrumental and non-instrumental) and specificity (non-specific and medium to
23 high specific). Logistic regression analyses were conducted to predict goal attainment.

24 **Results.** Goal attainment was significantly predicted by the motivational value of the personal
25 advice (OR:1.86), by the specificity of the implementation intentions (OR:3.5), by the
26 motivational value of the action plan (OR:1.86), and by making a new action plan at follow-up
27 (OR:4.10). Interaction-effects with behaviour showed that the specificity score of the
28 implementation intention plans (OR:4.59), the motivational value of the personal advice
29 (OR:2.38), selecting hindering factors and solutions(OR:2.00) and making a new action plan at
30 follow-up (OR:7.54) were predictive of goal attainment only for fruit or vegetable intake. Also,
31 when participants in the fruit and vegetable group made more than three plans, they were more
32 likely to attain their goal (OR:1.73), whereas the reverse was the case in the PA group (OR:0.34).

33 **Discussion.** Feedback on goal feasibility, coping implementation intentions, further research to
34 investigate the optimal number of plans for different behaviours, the optimal frequency and
35 timing of follow-up modules and new ways to incorporate social support in eHealth
36 interventions, are recommended.

37

38 Background

39 Physical activity (PA) and a varied diet with fruits and vegetables are associated with decreased
40 risk of cardiovascular diseases and cancer(1-3). Therefore, adults are recommended to perform
41 at least 30 minutes of PA at moderate- to vigorous intensity on most, preferably all days of the
42 week, and to consume at least 400 g of fruit and vegetable per day(4). However, many people do
43 not meet these recommendations(5). Despite the efforts to promote these health behaviours in
44 adults, fruit and vegetable intake have been decreasing, and PA levels have remained the same
45 since 2008 in Belgium(6). A recent meta-analysis focusing on these health behaviours indeed
46 stated that changing unhealthy lifestyle is difficult, and there is room for improvement(7).

47

48 Interventions most often targeted motivational determinants that are important during the early
49 stages of behaviour change, such as attitude and knowledge. However, interventions based upon
50 theories of intentions are often more effective in changing intentions than in changing
51 behaviour(8, 9), resulting in a so-called intention-behaviour “gap”. This gap can be targeted by
52 adopting self-regulation techniques. One useful framework is the Health Action Process
53 Approach model that includes both pre-intentional processes that lead to a behavioural intention
54 and post-intentional processes that lead to the actual health behaviour(10). The model states that
55 individuals first have to become conscious of their own health behaviour and have to become
56 motivated to change their behaviour, where after they have to initiate the new health behaviour to
57 bridge the gap between intentions and behaviour. This can be achieved by defining specific
58 action plans on ‘when’, ‘where’, and ‘how’ to perform the health behaviour, and by stating
59 implementation intentions in which strategies to initiate the action are stated (i.e. “If situation Y
60 is encountered, then I will initiate goal-directed behaviour X”)(11). Finally, in order to be able to
61 adopt health behaviours people may make coping plans in which they cope with anticipated

62 barriers and problems that may hinder goal attainment(10, 12, 13). Research has shown that
63 interventions that applied self-regulation techniques (i.e. specific goal setting, implementation
64 intentions, providing feedback on performance, prompting review of behaviour goals, social
65 support and self-monitoring) were more effective in changing health behaviour than other
66 interventions(14-17). The meta-analysis of Michie et al. (2009) offers support to prompt
67 intention formation, prompt goal setting, provide feedback on performance, and prompt review
68 of goal progress and self-monitoring(14). Another meta-analysis of Gollwitzer and
69 Sheeran(2006) also showed that implementation intention plans were effective in goal
70 achievement(18). Based on the findings we developed an eHealth intervention ‘MyPlan’, which
71 included self-regulation techniques (e.g. goal setting, prompting monitoring of behaviour,
72 implementation intentions), guided by the Health Action Process Approach (HAPA) theory(10).
73 ‘MyPlan’ provides the opportunity to select one out of three health behaviours (fruit, vegetables
74 and PA), helps adults to set personal goals for that health behaviour and guides them to
75 anticipate barriers and hindering situations during goal pursuit. To help participants to attain their
76 goals, the following self-regulation techniques were implemented: tailored feedback, action
77 planning, implementation intentions, problem solving, sharing action plans with friends/family
78 for social support, stimulating self-monitoring and goal evaluation and adjustment.

79 A limitation of self-regulation interventions is that many tools (e.g. action planning,
80 implementation intentions) may be used suboptimally by participants (19-21). For example,
81 Michie et al. (2004) found that more than one-third of participants did not formulate
82 implementation intentions despite being prompted to do so(22). Furthermore, when self-
83 regulation tools are used, participants may not optimally apply them. Van Osch et al. (2010)
84 reported that plans that are relatively broad and non-specific resulted in less successful

85 behavioural change(20). Ziegelmann, Lippke and Schwarzer (2006) also found that plans that
86 were incomplete (lacking action planning or coping planning) were associated with less physical
87 activity during rehabilitation at 6 months post-test(23).

88 The first aim of this study was to evaluate whether the use of several self-regulation techniques
89 (e.g. selecting hindering factors and solutions, monitor behaviour) and feasibility evaluation of
90 the self-regulation techniques (e.g. the difficulty experienced when making an action plan, the
91 feasibility of the action plan) predicted goal attainment. Second, we rated the quality of
92 implementation intention plans and evaluated if the coded total number of instrumental plans
93 and the specificity score of the implementation intention plans predicted goal attainment. Finally,
94 the moderating effect of the selected behaviour (fruit intake, vegetable intake, PA) on the
95 predictions of use and feasibility rating of the self-regulation techniques on goal attainment was
96 examined, as previous research showed that the effect of behaviour change techniques varies for
97 different behaviours(12).

98 **Methods**

99 *Participants and procedure*

100 Data were obtained from participants in a proof of concept (POC) intervention of ‘MyPlan’.
101 ‘MyPlan’ provides personal feedback and helps adults to set and monitor personal and attainable
102 health goals in order to increase either PA level, fruit or vegetable intake. Participants were
103 recruited by distributing flyers to parents of adolescents in secondary schools, by using Facebook
104 and Twitter advertisements and by recruiting university students. Eligible participants were over
105 18 years, were able to understand Dutch, and had access to Internet. Potential participants were
106 invited to visit the website. A computer log in system was used to allocate adults to the control or
107 intervention condition. The present study only used data from participants in the intervention

108 group who applied at least one of the self-regulation tools of ‘MyPlan’. In the ‘MyPlan’
109 intervention programme, adults themselves chose a health behaviour that they wanted to change
110 (fruit, vegetables or PA), where after they filled in online questions about demographic variables
111 (age, gender, socio economic status) and questions about the selected health behaviour. Next,
112 adults had access to the computer-tailored intervention module (T1). After one week (T2) and
113 one month (T3), adults received an email with an invitation for the follow-up modules. These
114 follow-up models evaluated whether they had reached their health goals and whether they
115 attained the recommended health norms. Figure 1 shows the flow of the participants through the
116 intervention modules as a function of the selected health behaviour. The study was approved by
117 the Ghent University Ethics Committee (approval number of the Ghent University Ethics
118 Committee: 670201319313), and an informed consent statement was obtained from each
119 participant.

120 [Insert figure 1 (flow chart) here]

121 ***‘MyPlan’ intervention***

122 ‘MyPlan’ is informed by self-regulation and Health Action Process Approach theory. After
123 logging in at the website (www.mijnactieplan.be), participants selected a behaviour of interest
124 (fruit intake, vegetable intake, or PA) and completed the first module for that behaviour, which
125 consisted of several components.

126 *Tailored feedback* is based upon the answers provided on a questionnaire about the
127 selected behaviour. For PA, the International Physical Activity Questionnaire (IPAQ) was
128 used(24). Tailored feedback consisted of reporting the actual level of PA in different domains
129 (i.e. leisure time PA, active transportation, PA at work, house hold PA), providing feedback
130 about these levels taking into account the health norms, and suggestions to increase PA. For Fruit

131 intake, the average portion of fruit per day was calculated using the Flemish ‘Fruit Test’(25).
132 Participants were asked to indicate how many pieces of each type of fruit that they ate during the
133 previous week. The average portion of fruit per day was calculated. Participants received a report
134 of this average portion and a comparison of this portion with the health norms in the personal
135 feedback. For vegetables, the average grams of vegetables was calculated by means of the
136 ‘Vegetable Test’(25). Participants were asked to indicate the amount of portions of each type of
137 vegetable they ate during the previous week. Average grams of vegetables per day was reported
138 and compared with health norms in the personal feedback.

139 *Action plans* were formulated by answering a series of questions. For example,
140 participants were asked *what* they want to do (e.g. being more physically active by walking),
141 *when* they want to do this (e.g. every Monday evening), *where* they want to do this (e.g. local
142 park), *how long* they want to do this (e.g. 60 minutes) and *with whom* they want to do this (e.g.
143 friends). For PA, adults chose in which domain they wanted to increase their PA level (i.e.
144 leisure time, active life style or both), and defined their goal by selecting activities (e.g. walking,
145 swimming, biking) and by indicating the frequency (days per week) and time (minutes per
146 activity) they wanted to spend on the chosen activity. For fruit and for vegetables, participants
147 indicated the number of days and portions of vegetables they wanted to eat.

148 Next, *implementation intentions* were stated. Participants were guided to formulate their
149 action plan into an implementation intention plan format (e.g. If it is Monday evening, then I will
150 go to the aerobic lessons, in the local gym).

151 *Problem solving* was prompted by indicating hindering factors from a predefined list, or
152 –when not listed- to write down the hindering factors in an open-ended question format.
153 Participants had to reflect upon solutions to overcome these difficulties. This was also done by

154 providing a predefined list of solutions for each hindering factor that could be selected, or –when
155 not listed- adults could write down their own solutions in an open-ended question format.

156 *Sharing action plans* was made possible by providing the participants the opportunity to
157 send their action plan to family or friends for social support.

158 *Stimulating self-monitor of behaviour* was done by motivating participants to monitor
159 their behaviour between the follow-up modules and by providing a list of possibilities (e.g.
160 tracking behaviour change in an agenda) to do so.

161 *Evaluating feedback* was provided in the follow-up modules. After one week (T2) and
162 one month (T3), participants had access to follow-up modules, in which it was evaluated whether
163 their behaviour changed and goals were reached. After the evaluation feedback, participants
164 could adapt or maintain their action plan. Action plans could be adapted by stating new goals
165 (easier goals or more difficult goals) and by selecting new difficult situations, hindering factors
166 and solutions, in the same way as in the first module. Both follow-up modules were similarly
167 constructed. An overview of the intervention programme is given in figure 2.

168 [Insert figure 2 here]

169 **Measures**

170 *Demographics*

171 Participants provided information on age, gender and educational level. Participants with a
172 university or college degree were classified as having a ‘high educational level’ whereas
173 participants with a secondary school degree or lower were classified as having a ‘low educational
174 level’. Age was dummy coded into younger adults (≤ 40 years) and older adults (> 40 years).

175 *Outcome variables*

176 Goal attainment at T2 and at T3 was operationalised in terms of whether participants attained at
177 least their goal set at T1.

178 *Use and feasibility evaluation of behaviour change techniques*

179 Participants indicated whether they used particular techniques (selecting hindering factors and
180 solutions, selecting different domains and activities (for PA only), sharing the action plan,
181 monitor behaviour and making a new action plan at T2) (See table 1). These variables were
182 dummy code into used (1) or not used (0) the technique (See table 1).

183 To evaluate the feasibility of the self-regulation techniques, additional questions were assessed at
184 the end of the questionnaire in T1. All variables regarding the feasibility of the self-regulation
185 techniques were dummy coded for the analyses. Table 1 provides an overview of the predictors
186 about the use and feasibility evaluation of the self-regulation techniques.

187 [Insert Table one here.]

188 *Quality of implementation plans*

189 Plan quality of implementation intentions (if-then plans) was evaluated by rating plan
190 instrumentality and specificity. We used the rating method of van Osch et al. (2010), plans were
191 rated as (1) instrumental or (0) non-instrumental. Plans were rated as instrumental when they
192 could facilitate the chosen behaviour (fruit intake or vegetable intake, PA) and when they were
193 applicable in the situation that was mentioned. The total number of instrumental plans was used
194 for the analysis by dummy coding it into (0) one or two instrumental plans and (1) more than two
195 instrumental plan. Frequent reasons for scoring a plan as not instrumental were nonsense plans,
196 or plans that did not target the chosen behaviour. Non-instrumental plans were not rated for
197 specificity. Specificity was only scored for plans considered instrumental, and was coded as (0)
198 non-specific, (1) medium specific, and (2) highly specific. 'Non-specific plans' were vague

199 plans, which were often applicable to various behaviours, e.g. “When it is Friday, I am going to
200 sport”. Plans that were described with a certain amount of detail and direction, but that were still
201 general and applicable to several actions and/or lacked one of the following elements (when,
202 how long and where) were rated as ‘medium specific’, e.g. “When I come home after work, I
203 will go playing basket”. Plans were coded as ‘highly specific’ if a sufficient amount of precision
204 and direction of time (Monday evening 8 am) and place (the local swimming pool) was used and
205 if all elements (when, how long and where) were included, e.g. “When it is Monday evening 8
206 am, I go swimming for 45 minutes in the local Swimming pool”. Participants had the possibility
207 to make several implementation plans. ‘The mean specificity score’ of all plans was calculated
208 and used in the analysis by dummy coding it into (0) low specific plans and (1) medium/high
209 specific plans. Two researchers independently evaluated all plans on instrumentality and
210 specificity. The interrater reliability was high for instrumentality (Cohen’s κ 0.89) and
211 substantial for specificity (Cohen’s κ =0.76)(26).

212 *Statistical analyses*

213 Baseline characteristics of participants were described using descriptive statistics. Logistic
214 regression analyses were performed to predict whether participants reached their goal (= goal
215 attainment) at T2 and T3. Various predictors were taking into account. These included several
216 self-ratings of the feasibility of the self-regulation techniques: the awareness of own behaviour,
217 the motivational value of the personal advice, the instructive value of the personal advice, the
218 motivational value of the action plan, the feasibility of the action plan and the difficulty
219 experienced when making an action plan. Also, selecting hindering factors and solutions,
220 selecting different domains and activities for PA, sharing the action plan, monitor behaviour and
221 making a new action plan at T2, were added as predictors to take into account the use of these

222 self-regulation techniques. Furthermore, the coded total number of instrumental plans and the
223 mean specificity score of the implementation intention plans were taking into account. All
224 predictors were dummy coded (See Table 1).

225 First we evaluated whether the evaluation of the self-regulation techniques, use of particular self-
226 regulation techniques, and plan quality of implementation intentions could predict whether health
227 goals were attained across the three groups. Next, interaction terms (predictor X behaviour) were
228 included to investigate whether the predictors of goal attainment differed as a function of the
229 chosen behaviour ('PA' or 'fruit and vegetables') of participants. Fruit and vegetables were taken
230 together in one category. In case of a significant interaction effect, the estimated predictive main
231 effect of the predictor only applies to the group that was indicated as the reference category (0).
232 For ease of interpretation, we reported odds ratio's and confidence intervals for PA indicated as
233 reference category and also for fruit and vegetables indicated as reference category (See Table 3
234 and 4). Statistical significance was set at a level of 0.05, p-values between 0.05 and 0.1 were
235 considered borderline significant.

236 **Results**

237 *Baseline characteristics*

238 In the intervention condition, 225 participants started the intervention module for fruit, 84 for
239 vegetables and 267 for PA. Table 2 presents the baseline characteristics for the sample that
240 completed the intervention programme at baseline (T1). Descriptive percentages regarding the
241 use and evaluation of the behaviour change methods are given in Table 1.

242 Logistic regression analysis revealed that younger participants (OR = 0.38; 95% CI = 0.24-0.62;
243 $p < 0.001$), men (OR = 0.63; 95% CI = 0.43-0.92; $p = 0.018$) and participants with high SES (OR

244 = 1.75; 95% CI = 1.15-2.66; $p = 0.009$) had a significant higher probability of responding at
245 follow-up.

246 ***Goal attainment***

247 For all predictors, odds ratios and confidence intervals of the logistic regression analyses are
248 shown in table 3 and 4. In what follows, only significant and borderline significant predictors are
249 reported.

250 *Tailored feedback*

251 *The motivational value of the tailored feedback* was a borderline significant predictor of health
252 goal attainment at T2. There was also a borderline significant interaction-effect with behaviour
253 ($p=0.090$), possibly indicating that this only applied for participants in the fruit or vegetable
254 group. Participants in the fruit or vegetable group who perceived the personal advice about fruit
255 or vegetables as motivating were two times more likely to attain their goal at T2 compared to
256 participants in the fruit or vegetable group who did not perceive the personal advice as
257 motivating (OR=2.38, 95% CI = 1.15-4.94; $p < 0.10$).

258 *Action planning*

259 Borderline significance was found for *the motivational value of the action plan* for health goal
260 attainment at T2. Participants who perceived making their action plan as motivating were more
261 likely to attain their goal at T2 than participants who did not perceive this as motivating
262 (OR=2.25, 95% CI = 1.08-4.69; $p < 0.05$).

263 *Problem solving*

264 *Selecting hindering factors and solutions* was a significant predictor, after including the
265 interaction term with behaviour ($p=0.019$). Participants in the fruit or vegetable group who
266 selected hindering factors and solutions, were two times more likely to reach their goal at T2

267 compared to participants in the fruit or vegetable group who did not select hindering factors and
268 solutions.

269 *Implementation intentions*

270 No significant main effects were found for *the coded total number of instrumental plans*.

271 However, a significant interaction effect was found with behaviour ($p < 0.001$). Indicating that

272 participants in the fruit or vegetable group who made more than two instrumental

273 implementation plans for fruit or vegetable intake were three times more likely to attain their

274 goals compared to participants in the fruit and vegetable group that made one or two

275 implementation plans (OR=1.73, 95% CI = 0.92-3.27; $p < 0.10$). In contrast, participants in the

276 PA group who made one or two instrumental implementation plans for PA were three times

277 more likely to attain their goals compared to participants that made more than two

278 implementation plans for PA (OR=0.34, 95% CI = 0.15-0.73; $p < 0.05$). Furthermore, separate

279 analysis solely in the PA group indicated that *stating goals in different PA domains* (e.g. free

280 time and active living style) was also a significant predictor for PA goal attainment. Participants

281 who stated goals in different PA domains (i.e. goals for their free time and goals for an active

282 living style (e.g. at work) and goals for active transport) were less likely to attain their PA goals

283 compared to participants that stated goals for only one PA domain (i.e. goals for their leisure

284 time only or for active transport only) (OR=8.07, 95% CI = 2.20-29.55; $p < 0.05$). *The amount of*

285 *activities selected for goals in the free time* could also predict PA goal attainment. Participants

286 who chose two different physical activities (e.g. walking and swimming) were less likely to

287 attain their PA goals at T2 compares to participants who chose only one activity (OR=0.21, 95%

288 CI = 0.08-0.59; $p < 0.05$). At T3, *the coded total number of instrumental plans* was a significant

289 predictor of health goal attainment, when adjusting for behaviour ($p = 0.003$). Participants in the

290 PA group who made more than two instrumental plans for PA were less likely to succeed in their
291 health goal than participants in the PA group who made one or two instrumental plans (OR=0.40,
292 95% CI = 0.16-1.03; $p<0.10$).

293 Borderline significance was found for *the mean specificity score of the implementation intention*
294 *plans* for goal attainment at T2. However, the significant interaction-effect with behaviour
295 ($p=0.016$) indicates that the estimated effect only counts for participants in the fruit or vegetable
296 group. Participants in the fruit or vegetable group who made specific plans were five times more
297 likely to attain their health goal at T2 (OR=4.59, 95% CI = 1.26-16.79; $p<0.05$).

298 *Adapting plans*

299 *Stating new goals at T2* was found to be a significant predictor. However, the significant
300 interaction-effect with behaviour indicates that the estimated effect only counts for participants
301 in the fruit or vegetable group. Participants in the fruit and vegetable group who did not state
302 new health goals at T2 were more likely to attain their health goal at T3 than participants who
303 stated new goals at T2 (OR=7.54, 95% CI = 1.96-28.99; $p<0.05$).

304 **Discussion**

305 The results of this study provide further information on how the design, feasibility and
306 applicability of health promotion interventions can be improved to promote optimal behaviour
307 outcomes for different health behaviours. Based on the results, feasible behaviour change
308 techniques can be identified and the content of self-regulation interventions can be improved by
309 further including and optimizing the different behaviour change techniques that can predict goal
310 attainment.

311 Our study revealed several significant predictors of health goal attainment. Goal attainment of
312 health behaviours was positively predicted by the motivational value of the personal advice, the

313 motivational value of the action plan, selecting hindering factors and solutions, the specificity
314 score of the implementation intention plans, the coded total number of instrumental plans,
315 selecting different PA activities and making a new action plan after one week.

316 Our results also showed that the efficacy of particular behaviour change techniques varies as a
317 function of type of health behaviour. Some predictors were only significant for fruit and
318 vegetable intake, and other predictors only for PA. The estimated effect of the specificity score
319 of the implementation intention plans, the motivational value of the personal advice, selecting
320 hindering factors and solutions and making a new action plan after one week to attain the health
321 goal was only applicable for participants who chose for fruit or vegetable intake and not for those
322 who chose PA. In line with our results the meta-analysis of Bélanger-Gravel et al. (2013)
323 revealed ‘small-to-medium’ effect size of implementation intentions on PA compared to
324 ‘medium-to-large’ effect sizes reported by Gollwitzer and Sheeran (2006) on a variety of health-
325 related behaviours(12, 18).

326 Moreover, there was one predictor (i.e. the coded total number of instrumental plans) that was
327 positively related to goal attainment for one behaviour, and inversely related for the other health
328 behaviour. The results for fruit intake, are in line with those of Wiedemann, Lippke and
329 Schwarzer (2012), who found that forming a large number of plans may be more effective in
330 changing fruit intake than forming few plans(27). However, our results also implicate that ‘the
331 more plans, the better’ cannot be generalised for all health behaviours, as our study showed
332 opposite results for PA goals. The study of Wiedemann et al. (2011) also showed that better
333 intervention effects were associated with two rather than three PA action plans(28). Due to our
334 small sample group, we could not investigate the optimal number of plans. Therefore, future

335 research may further focus on the optimal number of plans for different behaviours, especially
336 for PA.

337 In our study, we also conducted separate analysis for the PA group, because participants who
338 choose the PA module had also the opportunity to make plans for the different domains.
339 However, we found that participants who stated goals in different PA domains (i.e. goals for
340 their free time and goals for active transport) were less likely to attain their PA goals compared
341 to participants that stated goals for only one PA domain (i.e. goals for their free time only or
342 goals for active transport only). The amount of activities selected also negatively predicted PA
343 goal attainment. Participants who chose two different physical activities (e.g. walking and
344 swimming) were less likely to attain their PA goals at T2 compared to participants who chose
345 only one activity. This could perhaps be attributed to the fact that PA is a rather complex
346 behaviour to change (12, 29). This also shows that the feasibility of PA goals and plans may be
347 important. However, the feasibility of the action plan was not a significant predictor of PA goal
348 attainment. It may be that adults have difficulties to formulate feasible PA plans, due to the
349 complexity of incorporating PA goals(12, 29). Therefore, it would be beneficial to incorporate
350 computerized feedback that gives advice on the feasibility by comparing the current health
351 behaviour with goals (especially for PA). For example, adults who never ran before and who
352 state a plan to run every day for one hour, may better receive prompt feedback about the
353 unfeasibility of their plan, if this results in lower physical activity levels. It seems that a small
354 group was already aware that their plans were not feasible, as they indicated this in their self-
355 report. It may make no sense to pursue a such goal, and in such situations adults are probably
356 better prompted to adapt their goals. It may be a good idea to implement an evaluation of the
357 feasibility of plans by participants in eHealth interventions. Another idea is to give participants

358 advice to start with only one or two plans in one PA domain, and to make repeated and/or
359 additional plans after the first goal is achieved.

360 Making implementation intentions of medium to high quality predicted goal attainment. The
361 mean specificity score of the implementation intention plans could only predict goal attainment
362 at short-term (at T2, after one week) in the fruit and vegetable group. In our study,
363 implementation intentions were used to let adults make action plans. Bélanger-Gravel et al.
364 (2013) stated that using implementation intentions for PA only for action planning and not for
365 coping planning (i.e. management of barriers) can decrease the efficacy of implementation
366 intentions(12). This may explain why the use of implementation intentions could not predict goal
367 achievement for PA goals and after a longer period. We did let participants select
368 difficulties/barriers/hindering factors and solutions (i.e. problem solving) but this was not applied
369 in an implementation intention format (i.e. if-then plans) and could also only predict goal
370 attainment in the fruit and vegetable group at short-term. By using the implementation intention
371 format, critical cues in coping plans will be linked to the goal-directed behaviour, which creates
372 a strong and automatic cue-response. Previous studies observed the 'if-then' format to yield
373 better behaviour outcomes (30, 31). Thus, our results strengthen the suggestion of Bélanger-
374 Gravel et al. (2013) to incorporate coping plans into an implementation intention format(12).

375 The timing and frequency of follow-up modules might be important as well. Adults in the fruit
376 and vegetable group who already adapted their plan after one week, had less chance to achieve
377 their goals after one month compared to those who did not adapt their plans yet. In the PA
378 group, adapting plans after one week could not predict goal attainment. This indicates that giving
379 people the possibility to adapt their goals after one week, is maybe too early. To our knowledge,
380 no studies have already investigated the optimal frequency and timing of follow-up modules in

381 self-regulation interventions. So, it is important to further investigate this to develop an optimal
382 follow-up system for eHealth interventions that target health behaviour change.

383 Only a small group sent their action plan for social support as part of 'MyPlan', and this could
384 not predict goal attainment. This result shows that further investigation on how to include social
385 support in eHealth interventions is warranted. Morrison et al. (2012) reported in their review
386 study that social context and support mediates eHealth intervention outcomes(17). To increase
387 intervention effectiveness, they suggest to provide social support by using automated dialogue,
388 peer-to-peer-mediated communication, or information about other real users. Ziegelmann,
389 Lippke, & Schwarzer (2006) reported more complete action plans and a longer duration of
390 physical activities when participants were assisted by an interviewer trained in motivational
391 interviewing(23). This suggests that additional personal support by health counsellors trained in
392 motivational interviewing could also lead to additional effects of future planning interventions
393 (23, 29).

394 There are some limitations to this study. Our results should be interpreted with caution due to the
395 small sample. The choice options (e.g. choosing to only form plans for PA in leisure time) have
396 led to some small sample groups, making it not possible to perform moderator analyses. Also,
397 the high drop-out led to a low response at T3, which may have influenced the results for the
398 impact on goal attainment at T3. Studies with larger samples are needed to confirm our results.
399 Next, a longer trial to measure the long-term impact on goal attainment (e.g. after 6 months and 1
400 year) is needed. Furthermore, it is also important to note that other factors (e.g., quality of
401 theoretical content, combination of behaviour change techniques, participants characteristics)
402 and the combination of factors might also have important effects on intervention outcome and
403 needs further investigation(14, 17). The impact of the behaviour change techniques on

404 intervention effectiveness should also be further evaluated by conducting experimental studies
405 with different conditions that do and do not include specific behaviour change techniques(1414).

406 **Conclusion**

407 In conclusion, we recommend to integrate feedback on the feasibility of adults' goals (especially
408 for PA goals) and to incorporate coping planning into a implementation intention format. We
409 also suggest further research to investigate the optimal number of plans for different behaviours,
410 the optimal frequency and timing of follow-up modules and ways to incorporate social support in
411 eHealth interventions.

412

413

414 **REFERENCES**

- 415 1. WHO. Global recommendations on physical activity for health. Geneva: World Health
416 Organisation. 2010.
- 417 2. WHO. Global Health Risks Mortality: and burden of disease attributable to selected major risks.
418 2009.
- 419 3. Lock K, Pomerleau J, Causer L, Altmann D, McKee M. The global burden of disease attributable
420 to low consumption of fruit and vegetables: implications for the global strategy on diet. Bulletin
421 of the World Health Organisation. 2005;83(2):100-8.
- 422 4. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public
423 health: updated recommendation for adults from the American College of Sports Medicine and
424 the American Heart Association. Med Sci Sports Exerc. 2007;39(8):1423-34.
- 425 5. WHO. Diet, Nutrition and the prevention of Chronic diseases, Report of Joint WHO/FAO Expert
426 Consultation. 2003.
- 427 6. Gisle L, Demarest S. Gezondheidsenquête 2013. Rapport 2: Gezondheidsgedrag en leefstijl. WIV-
428 ISP, Brussel: 2014.
- 429 7. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels:
430 surveillance progress, pitfalls, and prospects. The Lancet. 2012;380(9838):247-57.
- 431 8. Sheeran P, Milne S, Webb TL, Gollwitzer PM. Implementation intentions and health behaviour.
432 In: Conner M & Norman P, editors. Predicting health behaviour. New York: Open University
433 Press; 2005. p. 276-323.
- 434 9. Hagger MS, Lonsdale A, Koka A, Hein V, Pasi H, Lintunen T, et al. An intervention to reduce
435 alcohol consumption in undergraduate students using implementation intentions and mental
436 simulations: a cross-national study. Int J Behav Med. 2012;19(1):82-96.
- 437 10. Schwarzer R. Modeling Health Behavior Change: How to Predict and Modify the Adoption and
438 Maintenance of Health Behaviors. Applied Psychology. 2008;57(1):1-29.

- 439 11. Gollwitzer P, Sheeran P. Implementation Intentions and Goal Achievement: A Meta-analysis of
440 effects and processes. *Advances in Experimental Social Psychology*. 2006;38:69-119.
- 441 12. Bélanger-Gravel A, Godin G, Amireault S. A meta-analytic review of the effect of implementation
442 intentions on physical activity. *Health Psychology Review*. 2013;7(1):23-54.
- 443 13. Sniehotta FF, Scholz U, Schwarzer R. Action plans and coping plans for physical exercise: A
444 longitudinal intervention study in cardiac rehabilitation. *British journal of health psychology*.
445 2006;11(Pt 1):23-37.
- 446 14. Michie S, Abraham C, Whittington C, McAteer J. Effective Techniques in Healthy Eating
447 and Physical Activity Interventions: A Meta-Regression. *Health Psychology*. 2009; 28 (6): 690–
448 701.
- 449 15. Lara J, Evans HE, O'Brien N, Moynihan PJ, Meyer DT, Adamson AJ, Errington L, Sniehotta FF,
450 White M, Mathers JC. Association of behaviour change techniques with effectiveness of dietary
451 interventions among adults of retirement age: a systematic review and meta-analysis of
452 randomised controlled trials. *BMC Medicine*. 2014;12:177.
- 453 16. Broekhuizen K, Kroeze W, van Poppel MN, Oenema A, Brug J. A systematic review of randomized
454 controlled trials on the effectiveness of computer-tailored physical activity and dietary behavior
455 promotion programs: an update. *Ann Behav Med*. 2012;44(2):259-86.
- 456 17. Morrison LG, Yardley L, Powell J, Michie S. What design features are used in effective e-health
457 interventions? A review using techniques from Critical Interpretive Synthesis. *Telemedicine
458 journal and e-health : the official journal of the American Telemedicine Association*.
459 2012;18(2):137-44.
- 460 18. Gollwitzer PM & Sheeran P. Implementation intentions and goal achievement: A
461 meta-analysis of effects and processes. *Advances in Experimental Social Psychology*. 2006;38:
462 70–110.
- 463 19. Springvloed L, Lechner L, Oenema A. Planned development and evaluation protocol of two
464 versions of a web-based computer-tailored nutrition education intervention aimed at adults,
465 including cognitive and environmental feedback. *BMC Public Health*. 2014;14:47.
- 466 20. van Osch L, Lechner L, Reubsat A, De Vries H. From theory to practice: An explorative study into
467 the instrumentality and specificity of implementation intentions. *Psychology & health*.
468 2010;25(3):351-64.
- 469 21. van Genugten L: Prevention of weight gain among overweight adults: development and
470 evaluation of a computer-tailored self-regulation intervention. PhD thesis. Erasmus University,
471 Department of Public Health; 2011.
- 472 22. Michie S, Dormandy E, Marteau TM. Increasing screening uptake amongst those intending to be
473 screened: the use of action plans. *Patient education and counseling*. 2004;55(2):218-22.
- 474 23. Ziegelmann JP, Lippke S, Schwarzer R. Adoption and maintenance of physical activity: Planning
475 interventions in young, middle-aged, and older adults. *Psychology & health*. 2006;21(2):145-63.
- 476 24. Vandelanotte C, De Bourdeaudhuij I, Philippaerts R, Sjöström M, Sallis J. Reliability and Validity of
477 a Computerized and Dutch Version of the International Physical Activity Questionnaire (IPAQ).
478 *Journal of Physical Activity and Health*. 2005; 2: 6-75.
- 479 25. VIG: Gezondheidsbevordering op het werk. Resultaten en verslag van de productevaluatie
480 Voeding: fruittest, groentetest en vochttest. [Health promotion at work. Results and report of
481 product evaluation Nutrition: fruit and vegetable test] 2004.
- 482 26. Landis JR, Koch GG: The measurement of observer agreement for categorical data. *Biometrics*.
483 1977;33:159–174.
- 484 27. Wiedemann AU, Lippke S, Schwarzer R. Multiple plans and memory performance: results of a
485 randomized controlled trial targeting fruit and vegetable intake. *Journal of behavioral medicine*.
486 2012;35(4):387-92.

- 487 28. Wiedemann AU, Lippke S, Reuter T, Ziegelmann JP, Schüz B. The More the Better? The Number
488 of Plans Predicts Health Behaviour Change. *Applied Psychology: Health and Well-Being*.
489 2011;3(1):87-106.
- 490 29. De Vet E, Oenema A, Sheeran P, Brug J. Should implementation intentions interventions be
491 implemented in obesity prevention: the impact of if-then plans on daily physical activity in Dutch
492 adults. *The international journal of behavioral nutrition and physical activity*. 2009;6:11.
- 493 30. Chapman J, Armitage CJ, Norman P. Comparing implementation intention interventions in
494 relation to young adults' intake of fruit and vegetables. *Psychology & health*. 2009;24(3):317-32.
- 495 31. Armitage CJ, Arden MA. A volitional help sheet to increase physical activity in people with low
496 socioeconomic status: A randomised exploratory trial. *Psychology & health*. 2010;25(10):1129-
497 45.

498

499

500

501

502

503

504 TABLES

505 Table 1

Behaviour change technique	Predictor	Question	Values (dummy coded)	n (%)
Tailored feedback (feasibility evaluation)	The motivational value of the personal advice	“I Think the personal advice is motivating”	Personal advice perceived as motivating (1) Personal advice not perceived as motivating (0)	141 (63.2) 82 (36.8)
	The awareness of own behaviour	“Did you expect the result of the personal advice in advance?”	Aware of their behaviour (1) Not aware of their behaviour (0)	129 (57.3) 96 (42.7)
	The instructive value of the personal advice	“I Think the personal advice is instructive”	Personal advice perceived as instructive (1) Personal advice not perceived as instructive (0)	142 (63.7) 81 (36.3)
Problem solving (use)	Selected barriers and hindering situations	“Select those barriers or hindering situations you want to apply or formulate it yourself”	No barriers or hindering situations (0) Selected/formulated barriers or hindering situations (1)	126 (31.3) 277 (68.7)
Action planning (feasibility evaluation) (use)	Perceived difficulty of making an action plan	“I think it is difficult to make an action plan”	Perceived making an action plan as difficult (1) Perceived making an action plan not as difficult (0)	82 (37.3) 138 (62.7)
	The motivational value of the action plan	“The action plan motivates me to pursue my goals”	Action plan perceived as motivating (1) Action plan not perceived as motivating (0)	139(62.9) 82 (37.1)
	The feasibility of the action plan	“My action plan is feasible”	Action plan perceived as feasible (1) Action plan not perceived as feasible (0)	217 (98.2) 4 (1.8)
	Selecting different domains for PA	“How do you want to improve you physical activity level?”	By being more active in my free-time (1) By choosing an active life style (0)	99 (54.1) 84 (45.9)
	Selecting different activities for PA	“ Do you want to select a second activity for your free time plan?”	Yes, I want to perform a second activity (1) No, I do not want to perform a second activity (0)	84 (54.5) 70 (45.5)
Stimulating self-monitoring (use)	Monitoring behaviour	“Did you monitor your behaviour the past week?”	Did monitor behaviour (1) Did not monitor behaviour (0)	89 (39.6) 136 (60.4)
Sharing action plan for social support (use)	Sharing the action plan	“Select to share your action plan with friends and family and fill out their email address”	Sent action plan to family/friends (0) Did not sent action plan to family/friends (1)	57 (25.3) 168 (74.7)
Goal evaluation and adjustment (use)	Making a new plan at T2	Do you want to make a new plan?	Yes, I want to make a new plan (0) No, I want to keep the same plan (1)	28 (20.4) 109 (79.6)

506

507 **Table 2: Baseline characteristics for the total sample and the four conditions separately.**

	Total intervention group (n=452)	Intervention Physical Activity (n=158)	Intervention Fruit intake (n=166)	Intervention Vegetable intake (n=50)
Age (years)	30.5 ± 12.5	30.5 ± 12.6	28.1 ± 10.9	33.8 ± 13.4
Gender (% male)	39.2	44.5	47.8	33.3
Education level (% high university or college)	72.1	73.6	75.8	66.6
Instrumentality n (%)				
No instrumental plan (N=6)	6 (1.7)	3 (1.9)	2 (1.4)	1 (2.3)
One instrumental plan (N=159)	159 (45.7)	57 (36.3)	60 (40.5)	42 (97.7)
Two instrumental plans (N=102)	102 (29.3)	54 (34.3)	48 (32.4)	0 (0)
Three instrumental plans (N=68)	68 (19.5)	30 (19.1)	38 (25.7)	0 (0)
Four instrumental plans (N=8)	8 (2.3)	8 (5.1)	0 (0)	0 (0)
Five instrumental plans (N=3)	3 (0.9)	3 (1.9)	0 (0)	0 (0)
Six instrumental plans (N=2)	2 (0.6)	2 (1.3)	0 (0)	0 (0)
Specificity n (%)				
Low specificity (N=28)	28 (8.0)	21 (13.0)	3 (2.0)	4 (9.5)
Medium specificity (N=219)	219 (62.2)	87 (53.7)	98 (66.2)	34 (81.0)
High specificity (N=105)	105 (29.8)	54 (33.3)	47 (31.8)	4 (9.5)

508

509

510

511

512

513

514

515

516 **Table 3 predicting goal attainment at T2**

Predictor	Goal attainment T2 (n=274)			
	Main effect ^a Predictor	Interaction effect predictor X behaviour	Main effect ^b Predictor	Main effect ^c Predictor
	OR(95%CI)	(p-value)	OR(95%CI)	OR(95%CI)
The motivational value of the personal advice	1.86(1.06-3.27)*	0.090	2.38(1.15,4.94)**	1,16(0.48,2.78)
The awareness of own behaviour	1.22(0.64-2.31)	0.077	1.65(0.80-3.40)	0.77(0.33-1.76)
The instructive value of the personal advice	0.89(0.47-1.70)	0.045	1.20(0.59-2.42)	0.49(0.20-1.19)
Selecting hindering factors and solutions	1.45(0.80-2.65)	0.019	2.00(1.04-3.85)**	0.89(0.43-1.86)
The coded total number of instrumental plans	0.89(0.52-1.55)	<0.001	1.73(0.92-3.27)*	0.34(0.15-0.73)**
The mean specificity score of the implementation intention plans	3.50(0.97-12.57)*	0.016	4.59(1.26-16.79)**	2.20(0.58-8.37)
The difficulty experienced when making an action plan	1.22(0.63-2.34)	0.058	1.68(0.81-3.49)	0.48(0.15-1.60)
The motivational value of the action plan	1.86(1.06-3.27)*	0.210	2.25(1.08-4.69)**	1.34(0.57-3.13)
The feasibility of the action plan	1.30(0.62-2.74)	0.516	1.06(0.40-2.81)	1.63(0.59-4.51)
Sharing the action plan	1.66(0.89-3.08)	0.394	1.97(0.94-4.11)*	1.20(0.46-3.16)

517 ^aNo interaction term included for behaviour, with fruit and vegetables as reference category (0);

518 ^bWith included interaction term (predictorXbehaviour), with fruit and vegetables as reference category (0);

519 ^cWith included interaction term (predictorXbehaviour), with physical activity as reference category (0)

520 ** $p < 0.05$: significant predictor; * $p < 0.1$: borderline significant predictor

521 CI, confidence interval; OR, odds ratio

522

523

524

525

526

527 **Table 4 predicting goal attainment at T3**

Predictor	Goal attainment T3 (n=137)			
	Main effect ^a Predictor OR(95%CI)	Interaction effect predictor X behaviour (p-value)	Main effect ^b Predictor OR(95%CI)	Main effect ^c Predictor OR(95%CI)
The motivational value of the personal advice	1.24(0.55,2.78)	0.230	1.52(0.63-3.68)	1.88(0.67,5.30)
The awareness of own behaviour	1.09(0.49,2.40)	0.188	1.41(0.57,3.45)	0.70(0.26,1.93)
The instructive value of the personal advice	0.68(0.29,1.59)	0.101	0.38(0.14,1.05)	0.35(0.12,1.04)
Selecting hindering factors and solutions	0.97(0.44,2.17)	0.019	1.44(0.60,3.47)	0.486(0.18,1.29)
The coded total number of instrumental plans	0.99(0.46,2.10)	0.003	1.70(0.70,4.11)	0.40(0.16, 1.031)*
The mean specificity score of the implementation intention plans	1.91(0.41,8.95)	0.035	2.57(0.53,12.41)	1.10(0.22,5.57)
The difficulty experienced when making an action plan	0.76(0.34,1.69)	0.327	0.458(0.096,2.179)	0.41(0.09,1.78)
The motivational value of the action plan	1.05(0.46,2.36)	0.228	1.29(0.53,3.17)	0.70(0.25,1.96)
The feasibility of the action plan	0.66(0.26,1.62)	0.994	0.65(0.20,2.18)	0.66(0.20,2.18)
Sharing the action plan	1.73(0.74,4.03)	0.243	0.40(0.09,1.86)	0.94(0.26,3.36)
Monitoring between T1 and T2	1.18(0.57,2.45)	0.618	0.74(0.23,2.39)	0.96(0.32,2.84)
Making a new action plan at T2	4.10(1.33,12.64)**	0.022	7.54(1.96,28.99)**	1.35(0.34,5.36)

528

^aNo interaction term included for behaviour, with fruit and vegetables as reference category (0);

529

^bWith included interaction term (predictorXbehaviour), with fruit and vegetables as reference category (0);

530

^cWith included interaction term (predictorXbehaviour), with physical activity as reference category (0)

531

** $p < 0.05$: significant predictor; * $p < 0.1$: borderline significant predictor

532

CI, confidence interval; OR, odds ratio

533

534

535

Figure 1 (on next page)

Figure 1: Flow chart response rate

Figure 1: Flow chart response rate




Figure 2 (on next page)

Figure 2: An overview of the intervention programme

Figure 2: An overview of the intervention programme