

# Biodiversity collapse in a phase transition between neutral and niche communities

Leonardo A. Saravia<sup>1 2</sup>, Fernando R. Momo<sup>1</sup>

1. Instituto de Ciencias

Universidad Nacional de General Sarmiento

J.M. Gutierrez 1159 (1613), Los Polvorines

Buenos Aires, Argentina.

2. E-mail: [lsaravia@ungs.edu.ar](mailto:lsaravia@ungs.edu.ar)

## Abstract

The dynamics of ecological communities can be described by two contrasting models: the first assumes that the individuals of all species are identical and do not have competitive interactions. The second assumes that species are different, adapted to particular habitat conditions, and have strong interactions. These represent extremes of a continuum: the neutral and the niche models of community organization. Real communities are actually a mixture of both types of dynamics. Here we study the simplest model of neutral-niche communities where niche dynamics is represented as a competitive hierarchy. The competition intensity is defined as a parameter that modulates the transition between these extremes. We use a stochastic cellular automata to show that there is a phase transition between the neutral and niche model with a spanning patch formed by the most abundant species. We measure the diversity as the Shannon index and the richness as the number of species. The transition implies a sharp fall of species diversity but the richness shows a gentle decline with increasing competitive intensity. As this kind of multi-species critical transition have not been described previously, we suggest new early warning signals: the rate of exponential decay in the patch distribution of the non-dominant species. This rate decreases when the community approaches the critical point and increases when the community crosses it. As an example we apply the early warnings to the Barro Colorado Tropical forest, which as expected, result to be far from a critical transition. The model shows that the critical point occur at a very low value of competitive intensity. Low values of competitive intensity were also reported for different high diversity real communities suggesting the possibility that this kind of communities are located near the critical point. This transition could happen before habitat destruction or degradation

affect the community in response to changes in environmental conditions like the ones produced by climate change or exotic species invasions.

*Short title:* A spatial phase transition between neutral and niche communities

*keywords:* Continuous critical transitions; Percolation; Neutral-niche communities; spatial stochastic models; Biodiversity

*Type of article:* Letters

*Abstract no. words:* 317

*Main text no. words:* 5747

*No. references:* 66

*No. figures & tables:* 6

*Corresponding author:*

Dr. Leonardo A. Saravia

Área Biología y Bioinformática

Instituto de Ciencias

Universidad Nacional de General Sarmiento

J.M. Gutierrez 1159 (1613) Los Polvorines

Buenos Aires Argentina

Tel: 54 11 4469-7533

# Introduction

Much effort has been devoted to understand the mechanisms of community assembly and dynamics. In principle, the emphasis was on deterministic processes based on niche differences between species; the niche theory assumes that different species are regulated by different environmental factors and infer that diversity originates from spatial and temporal environmental heterogeneity (Tilman 1982; Chesson 2000). More recently, the emphasis shifted to stochastic mechanisms in the form of the Neutral Theory of Biodiversity and Biogeography (Hubbell 2001). The neutral theory assumes that individuals of all species are equivalent and it proposes that diversity originates from a balance between immigration, speciation, and extinction. The neutral theory has been proposed as a parsimonious formulation that can provide new insight into the patterns of community assembly (Hubbell 2005), besides this simplification it can predict some community metrics very well (Volkov *et al.* 2007; Rosindell *et al.* 2012), mainly the species abundance distribution (SAD).

Finally it arises a unified view that accepts that both kinds of mechanisms are present at the same time and the focus shift to quantify the relative importance of these in natural communities (Leibold & McPeck 2006; Vergnon *et al.* 2009; Kalyuzhny *et al.* 2014; Martorell & Freckleton 2014). The main point is to understand which species level traits are important for community dynamics and which ones are unimportant (Matthews & Whittaker 2014), and this is related to the scale of observation. The problems of pattern and scale are critical in ecology (Levin 1992; Chave 2013), because patterns that seem stochastic at one scale may reveal structure at another scale. The concept of pattern is related to some sort of repetition that our brain can detect, when this pattern repeats at different scales we talk about scale invariance or self-similarity, characterized by power laws. These patterns could be produced by critical phase transitions described by percolation theory (Stauffer & Aharony 1994). This kind of spatial phase transitions were first introduced in ecology in the framework of landscape ecology (Loehle *et al.* 1996) and habitat fragmentation (Bascompte *et al.* 1996).

Percolation is characterized by the presence of two phases defined by some macroscopic features, e.g. the presence of vegetation or a desert in arid ecosystems (Kéfi *et al.* 2007). These phases are linked by a critical point where a sudden transition happens and a large spatial pattern emerges. In a two dimensional landscape where each site is connected to the nearest neighbors with some probability  $p$ , if you start with a small  $p$  there will be some connected sites that form a cluster or patch. When  $p$  increases, there is a point at which appears a cluster of sites spanning the entire landscape. This spanning cluster has a self-similar structure and is produced by local interactions (Solé & Bascompte 2006).

Several different ecological spatial models exhibit critical behavior related to the degree of disturbance (Pascual & Guichard 2005). Some of these models showed robust criticality: a particular kind of criticality discovered for ecological systems, where self-similarity is present for a wide range of parameters and does not necessarily involve drastic changes in the biological variables of interest (Roy & Pascual 2003). This kind of criticality has been documented for arid ecosystems (Solé 2007); here the sudden shift towards a desert condition might occur when rainfall decreases (Scanlon *et al.* 2007) or also with more intense grazing (Kéfi *et al.* 2007). The mechanism producing self-similarity is the positive effect produced by local facilitation, the chance of a new seedling to become established is higher near the parent plant.

Another example of an ecosystem exhibiting criticality are savannas, where the transition occurs between tree and grass cover (Abades *et al.* 2014). In critical phenomena, the transition is produced by the capacity of the system to transmit some signal or information, for example in savannas the proportion of 60% grass 40% trees is linked to the threshold needed for fire to spread. If there is not enough grass to act as a combustible, fire can not spread across the landscape (Staver & Levin 2012). Thus the increase in the proportion of trees, due to a change in environmental conditions, the can create positive feedback mechanisms resulting in the encroachment of savanna ecosystems (Abades *et al.* 2014).

Neutral models can produce species patches with power law distributions without been near a critical state (Houchmandzadeh & Vallade 2003), so the detection of this kind of patterns does not imply a phase transition. Moreover phase transitions are also observed in non-spatial models, some studies detected the presence of a sharp transition between neutral and niche dynamics in spatially implicit models (Chisholm & Pacala 2011) and Fisher *et al.* (2014) demonstrated the presence of a phase transition for this kind of neutral-niche models. Here we study a different kind of ecological spatial phase transition that is not related to disturbance or fire: it is the transition between a neutral and a niche community. We used the simplest model of neutral-niche communities where niche dynamics is represented as a competitive hierarchy (Saravia 2015). This spatially explicit model unifies Tilman's model of hierarchical competition with the classical neutral model using one parameter: competition intensity. This parameter is represented as the probability that one species replaces another and it modulates the transition between the neutral phase and niche phase.

Our first aim is to demonstrate the existence of a spatial phase transition in neutral-niche models, our second objective is to explore the dependence of the critical point with the parameters that are more important for the spatial pattern and to suggest some possible early warnings of the transition. Finally we will apply this new kind of early warnings to the repeated censuses of the Barro Colorado Island 50-ha forest plot in Panama (BCI).

# Methods

First, we define the spatial explicit neutral-hierarchical model, then we explain how we characterized its critical behavior in terms of percolation theory and how simulations are performed. We analyze early warnings for this critical transition and apply the same techniques for BCI plot data. We refer to interested readers to more extensive introductions to percolation theory in an ecological context (Solé & Bascompte 2006; Oborny *et al.* 2007).

## The spatial stochastic model

This model represents a continuum between hierarchical and neutral model in the same spirit as Gravel *et al.* (2006), and others (Zhou & Zhang 2008; Chisholm & Pacala 2010). The model is a stochastic cellular automata (CA) or also called interactive particle system (Durrett & Levin 1994). In these kind of models space is discretized into a grid and only one individual can occupy a particular position. Each position represents an area fixed by the investigator to mimic the real system. Time is continuous so the update of the model is asynchronous. Sites are chosen at random to be updated and to perform one complete time interval  $cJ$  sites have to be updated, where  $c$  is a constant that describes the overall rate at which transitions are occurring and  $J$  is the size of the grid (Durrett & Levin 1994).

We use periodic boundary conditions, which makes the landscape a torus. It means that sites on the top edge of the grid are neighbors of those on the bottom edge, and sites on the right edge are neighbors of those on the left. With this choice we avoid edge effects, this is equivalent to thinking that the grid is embedded in a large community. The size of the community is given by  $J = \dim X \times \dim Y$ , where  $\dim X$  and  $\dim Y$  are the dimension of the grid. Thus  $J$  is the maximum number of individuals in the simulated area. As in a classical neutral model there is a metacommunity i.e. a regional species pool assumed very large and invariant in ecological time scales (Hubbell 2001). All individuals have the same parameters, besides they should belong to different species (Hubbell 2001), and each species is assigned with a number. There are only two possible differences between species:

- They may have a different frequency  $X_i$  in the metacommunity and also different abundances in the local community.
- Hierarchical competition: species with lower numbers has a probability to replace species with higher numbers as in (Tilman 1994). Thus a species with number 1 have a probability to replace species with number 2 and greater. The species with number 2 can replace species starting from 3. The probability

of replacement ( $\rho$ ) is a parameter, when it is 0 there is no replacement and the model behaves like a neutral model without competitive hierarchy.

The colonization-competition and other possible trade-offs are not explicitly included in the model. But a colonization-competition trade-off can be established if species numbering is arranged in inverse order as it's frequency  $X_i$  in the metacommunity. Hence the most competitive species (with number 1) will have the lowest migration rate and the less competitive will have the highest migration rate.

There are three processes included in the model: death, local dispersal, and migration, starting with an empty site the following events can happen:

(1) With probability  $m$  an individual of a species  $i$  can migrate from the metacommunity at a rate proportional to its frequency  $X_i$  in the metacommunity.

(2) When the grid is not full, individuals give birth with rate 1 to a new individual that disperse to the neighborhood with a dispersal kernel, here we use an inverse power kernel (Marco *et al.* 2011):

$$d(x) = \frac{\alpha-1}{x_{min}} \left( \frac{x}{x_{min}} \right)^{-\alpha} \text{ with } mean = \frac{\alpha-1}{\alpha-2} x_{min} \text{ where } \alpha > 1 \text{ and } x \geq x_{min}.$$

where  $d(x)$  is the probability that an individual disperse a distance  $x$  from the parent. In all cases I used  $x_{min} = 1$ .

(3) Individuals die at a rate  $\mu$

(4) When an individual dies, it is replaced by a migrant from metacommunity with probability  $m$  and with probability  $1 - m$  by an individual from the neighborhood. The neighborhood is established using the dispersal kernel with average distance  $d$ . Once the grid is full it keeps full, because when an individual dies it is immediately replaced by another. This is called the zero-sum assumption in neutral models.

(5) If the individual does not die it can be replaced by an individual from the metacommunity or neighborhood as in (4), but an individual of species with number  $k$  can replace an individual of a species  $k + 1$  with probability  $\rho$ . Thus a hierarchical ordering of species is established. When this probability is zero, the model behavior becomes neutral.

The model was developed using the C++ programming language and its source code is available at <https://github.com/lasaravia/neutral> and figshare <http://dx.doi.org/10.6084/m9.figshare.969692>.

# 1 Percolation and simulations

2 To characterize our model in terms of percolation theory, we need to define an order parameter that depends  
3 on an external control or tuning parameter that can be continuously varied. Thus, we defined as tuning  
4 parameter the replacement probability  $\rho$ , and the order parameter as the probability that a patch of one  
5 species connects the landscape, this is called the spanning cluster probability  $SC_p$ .

6 In our model, percolation is produced when there is at least one patch of one species that spans from one  
7 edge of the system to the opposite edge. To detect species patches we used a modified Hoshen–Kopelman  
8 cluster labeling algorithm (Hoshen & Kopelman 1976) with a neighborhood defined by the four nearest sites  
9 (Von Neumann neighborhood) available at github <https://github.com/laravia/Clusters>. The percolation  
10 point is defined as the value of the tuning parameter  $\rho$  at which  $SC_p$  is 0.5. We used one snapshot of the  
11 spatial pattern to make our results more compatible with field studies, hence we measure the patch size  
12 distributions after 5000 time iterations.

13 The size of the lattice affects the value of the critical point  $\rho_c$  at which the transition occurs; in small lattices  
14  $SC_p$  is non-zero for values of  $\rho$  below the  $\rho_c$ , because patches that connect the entire lattice appear only  
15 by chance. Therefore, in order to obtain an asymptotic estimate for  $\rho_c$  we performed a finite size scaling  
16 analysis. For this, we run simulations for different lattice sizes (Side = 100, 150, 256, 512) and obtained  
17 asymptotic values by regressing  $\rho_c$  against  $1/N$ , the intercept becomes an estimate for a lattice of infinite  
18 size ( $\rho_c^\infty$ ) (Stauffer & Aharony 1994; Sornette 2000).

19 We determined critical points for two different metacommunities: a) One with a logseries species abundance  
20 distribution, the most common distribution that fits experimental data (White *et al.* 2012). With this  
21 metacommunity we included a competition-colonization trade-off by arranging species numbers in inverse  
22 order as it's frequency  $X_i$  in the metacommunity. b) A uniform species distribution, all species have the same  
23 probability to colonize the local community, this is analogous to simulate the apparition of a new species by  
24 evolution thus the migration parameter  $m$  would represent a speciation rate. The values of the  $m$  parameter  
25 (Table 1) were at least two orders of magnitude higher than realist speciation rates (Rosindell & Cornell  
26 2009) as it is not our aim to interpret the results in an evolutionary framework, only to show the influence of  
27 different metacommunities.

28 All simulations started with an empty lattice that is colonized by migrants mimicking the assembly of a  
29 new community. For each parameter combination we made 30 simulations. Thus, we calculate the  $SC_p$  as  
30 the number of times we observed a spanning cluster divided by the number of simulations. The range of  
31 parameters used were compatible with published results in tropical forest (Condit *et al.* 2002; Etienne 2007;

Anand & Langille 2010), It was suggested that fat-tail dispersal kernels give more realistic results (Rosindell & Cornell 2009; Seri *et al.* 2012) so we used an inverse power law distribution with an exponents always greater than two—so the mean exist (Table 1). The parameter  $\rho$  is varied in all the range between 0 and 1 to determine the critical point, in the region where we suspect the  $\rho_c$  will be located (near 0) the steps were very small (0.0001) and greater (0.1) in the region where we don't expect the  $\rho_c$ . The analysis of the model output were done in the R statistical language (R Core Team 2015) and the scripts are available at github <https://github.com/lisaravia/CriticalTransition> and figshare <http://dx.doi.org/10.6084/m9.figshare.2007537>. We calculated the sensitivity of critical point for infinite lattices ( $\rho_c^\infty$ ) varying three parameters of the model, one at a time: the migration  $m$ , the dispersal distance and the number of species on the metacommunity. To change the dispersal distance we vary the power exponent of the dispersal kernel  $\alpha$ . The range of parameters we used is described in table 1. We used as a base the first row of the table, so if we vary  $m$ , the other parameters are fixed at  $\alpha = 2.04$  and the metacommunity number of species to 64.

## Communities and early warnings signals

To compare communities change near and far from the  $\rho_c$  we measured the species abundance distribution (SAD) and calculated the rank abundance diagrams (RAD). These diagrams are equivalent to cumulative distributions and thus are a robust way to visualize the SAD without losing information (Newman 2005). We also calculated richness ( $S$ ) and the Shannon diversity index ( $H$ ) for each simulation.

The patch size distribution has been suggested as an early warnings for ecological transitions (Kéfi *et al.* 2014), besides a more in-depth study is necessary to reliably identify early warnings we present these as a first step in this new kind of models. The difference with previous models of phase transition in ecology (Kéfi *et al.* 2011), is that they have one or two species patches and we have from 16 to 256 different species. We already know that the patch size distributions of species in neutral models follow a power law (Houchmandzadeh & Vallade 2003), besides we added to the neutral model a competitive interaction we expect that the patch size distribution follows a power law ( $f(x) = x^{-\alpha}$ ) or due to the finite size of the simulations a power law with exponential cutoff ( $f(x) = x^{-\alpha}e^{-\lambda x}$ ) (Pueyo 2011; Weerman *et al.* 2012). As an alternative model we fitted an exponential distribution ( $f(x) = e^{-\lambda x}$ ). Near the critical point, and after it, we have a species with a continuous patch that spans all over the area: the spanning species. Thus we fitted the patch distribution of the spanning species and the distribution of all other species aggregated. Before the critical point we don not have a spanning species thus we fitted the patch distribution of the species that has the biggest patch and also the distribution of all other species excluding the one that has the biggest patch.



For each of these species or set of species, we fitted the mentioned 3 discrete distributions using maximum likelihood methods (Clauset *et al.* 2009). After that we calculated the Akaike information criteria corrected for small samples ( $AIC_c$ ), to select the best model (Burnham & Anderson 2002; Burnham *et al.* 2011). To characterize the variation of fitted parameters against  $\rho$ , we used quantile regression (Cade & Noon 2003). All the analysis were made using the R statistical language (R Core Team 2015), the fitting of patch distributions was made using mainly the package `powerlaw` (Gillespie 2015) and quantile regressions with R package `quantreg` (Koenker 2013).

We measured the patch size distribution from simulations in a range of  $\rho$  from neutral to niche communities, using a smaller set than the ones we used to estimate the critical point:  $\rho = \{0.0000, 0.0005, 0.0010, 0.0020, 0.0030, 0.004, 0.005, 0.01\}$ , the other parameters were from the first row of table 1 and a *Side* of 256 sites. The simulations run up to time 5000 and then we collected the patches every 40 time intervals for 400 time intervals more each. In some simulations the spanning cluster occupies a great proportion of the landscape in consequence very few patches remain to estimate a distribution. We only fitted a model when there are at least 20 patches and 5 different patch sizes.

## Application of early warnings to BCI forest data

The Barro Colorado Island forest plot is a 50 hectare rectangle (1000 m x 500 m) of tropical forest located at Panama and managed by the Smithsonian Tropical Research Institute (Condit 1998). In this plot all individuals  $\geq 1$  cm diameter at breast height (dbh) of free standing woody tree species have been measured and identified. Since the first census at 1982-1983, there was a second census in 1985, and then each five years, there are seven publicly available censuses. We will estimate the species patch distributions for each year to check if there is a tendency.

To analyze the patch distribution of BCI plot we first discretized the positions of the trees to fit them in a lattice (Appendix 1). After that we fit the same models that we described above to the patch size distributions for each year. To check for a tendency in the fitted parameters we quantile regressions but only with the 0.5 quantile (median) because we have a smaller  $n$ .

## Results

We observed a typical pattern of a second order continuous phase transition (Figure 1) which means that at the critical point  $\rho_c$  one species percolates through the lattice: a mono-specific patch spreads from side

to side, this patch is called the spanning patch or spanning cluster. As expected from percolation theory (Stauffer & Aharony 1994; Sornette 2000) the probability of a spanning cluster ( $SC_p$ ) is greater than 0 for  $\rho < \rho_c$  and small lattice sizes, for bigger lattice sizes  $SC_p$  is 0 for  $\rho < \rho_c$  and jumps quickly to 1 for  $\rho > \rho_c$  and this clearly define the two phases. Consequently the critical point for smaller lattices is lower than for bigger lattices (Table 2). The size of the spanning cluster is between 0.28-0.37 in proportion of the lattice for logseries metacommunity, and 0.34-0.36 for uniform metacommunity. Communities with a species patch around these areas could be near a critical point.

When the competitive intensity surpasses the critical point, the space left by the spanning species quickly diminish so the Shannon diversity ( $H$ ) collapses (Figure 2), but some individuals can escape the competitive displacement thus richness shows a more gentle fall (Figure 3). We can observe the effect of the competition-colonization trade-off in the logseries metacommunities: a very small degree of competition produces more diverse communities than purely neutral ones. This is because in the long term the SAD from a neutral community will match the metacommunity SAD (Houchmandzadeh & Vallade 2003), thus a small degree of competition lowers the density of colonizers and raises  $H$  before the critical point. The effect of the trade-off also can be observed in richness, logseries communities have in general less species and the decrease in richness is slower than with uniform communities (Figure 3).

The differences between logseries and uniform communities are also observed looking at the RADs (Figure S1). Before the critical point logseries communities have the effect of the trade-off: neutral ones ( $\rho = 0$ ) have a greater negative slope and less species, then the slope is more flat ( $\rho$  around 0.0005), and approaching the critical point it began to fall again but the shape of the RADs are different, the curvature is inverted. For uniform communities, the shape of the RAD is constant while the dominance of competitive species raises, the number of species keep constant until we reach  $\rho=0.1$ . This explains why we observed a sharp drop in  $H$  and a gentle decrease in  $S$  when we raise  $\rho$ . The change in the RAD with  $\rho$  also suggest a method to find communities where the trade-offs are important or not.

The critical point is the value of the intensity of competition where the phase transition occurs  $\rho_c$ . We observed that in general the transition happened at very low  $\rho$  values (Figure S2). That means that low levels of competition are needed to change the phase and sharply decrease diversity, this is observed for both logseries and uniform metacommunities. The effect of the colonization-competition trade-off in communities with logseries metacommunities is to slightly increase  $\rho_c^\infty$ . Thus logseries communities can have slightly higher levels of competition than uniform communities before the transition to the niche phase.

We analyzed how the critical point changes varying model parameters in ranges suggested by previous studies, thus the validity of our findings is restricted to these ranges (Table 1). The migration parameter  $m$  had the

widest range and the biggest relative influence (0.9) on the critical point  $\rho_c^\infty$  (Appendix Table 2). When  $m$  is greater we have more influence of the metacommunity on the local community, this lowers the effect of competition intensity and makes the  $\rho_c^\infty$  higher (Figure S2). The effect becomes important when  $m = 0.016$  which results in the biggest  $\rho_c^\infty$ , approximately 0.01 which is an order of magnitude greater than all the other cases (Appendix Table 2). The number of species in the metacommunity have a strong influence on  $\rho_c^\infty$  and we observed different relative responses for logseries (0.29) and uniform (0.39) metacommunities. For higher number of species in the metacommunity we obtained higher  $\rho_c^\infty$ , this means that when we have less species the effect of competition is stronger, the community changes from the neutral to niche phase earlier. The competition-colonization trade-off, present in logseries communities, diminish the influences of competition so the variation in  $\rho_c^\infty$  is smaller (Appendix Table 2). Finally dispersal distance has a weak effect in the range we used (relative variation of 0.1).

The patch size distributions have been used to detect the closeness of the critical point when the spatial patterns are non periodic or irregular as in our case (Kéfi *et al.* 2014). In our simulations the most frequent best model—with lower  $AIC_c$ —was the power law with exponential cutoff (70%), the pure power law was found best in 8% of the cases and the exponential model never was found the best model (Appendix Table 3). The rest of the cases (22%) correspond to simulations with  $\rho$  greater than  $\rho_c$  where one big spanning patch exist and there are few other patches of the same species, so a distribution model can not be fitted following the criteria we stated in methods. Thus we adopted as a possible indicators of the phase transition the parameters of the power law with exponential cutoff model:  $\alpha$  the power exponent and  $\lambda$  the exponential decay rate. We fitted quantil regressions using three quantiles: 0.25, 0.50 and 0.75.

For the biggest patch and the spanning species  $\alpha$  has an increasing tendency but slopes are non-significative at 5% level in near 60% of the cases (Appendix Table 4 and appendix Figure S3), The  $\lambda$ 's slopes are also non-significative in near 60% and do not have consistent patterns for all the cases: a decreasing pattern for logseries communities; for uniform communities most abundant species have negative slopes and spanning species have positive slopes (Appendix Figure S4). The most clear and consistent patterns are observed for the patches of other species clumped, which have for  $\lambda$  only 1 case non-significative (8%) but for  $\alpha$  also have 58% of cases non-significative (Appendix table 4). Before the critical point the  $\lambda$  have a negative slope and after the critical point have a positive slope, we also observed that the median of  $\lambda$  reaches a minimum right after the critical point ( $\rho_c$  between 0.0011-0.0013). Thus the slope of  $\lambda$  could be used as an indicator of the critical point. If the community is approaching the critical point the slope will be negative and if it is past the critical point the slope will be positive.

For BCI data we found that there is no spanning patch, the relatively small scale we chose for discretization

produced small patches so the biggest patch have in average a size of 0.006 % of the lattice. The best models were the power law with exponential cutoff for the biggest patch species, and the power law for the other species, but in this last group the power law with exponential cutoff is also a plausible model because the differences in  $AIC_c$  are around 2, and the fitted power exponents are the same (up to two decimals). Thus we regress the parameter  $\lambda$  against year to detect any tendency that would indicate an approaching or a moving away from a critical point. The slopes we obtained were all non-significative (Appendix table 5) thus there is no indication that the BCI is near a critical point.

## Discussion

We described for the first time a spatial phase transition between neutral and niche ecological communities. The power laws of patch sizes distributions observed in this model are not only produced at the critical point but are present over the whole range of the control parameter  $\rho$ . There are some ecological models that display this kind of behavior and it has been termed robust criticality, because of the permanence of scaling laws (Pascual & Guichard 2005). These previous models all included some kind of disturbances or stress, and with the increase in disturbance levels an increase in the exponential decay ( $\lambda$ ) of patch sizes is observed. This increase can be the result of a switch from a power law to a power law with exponential cutoff model (Kéfi *et al.* 2011), or by an increase of  $\lambda$  in the power law with exponential cutoff model (Weerman *et al.* 2012). In our model model, the competition intensity  $\rho$  can play the same role of stress; a very small degree of competition produce a critical transition from a neutral phase to a niche phase, and the most competitive species invades a great portion of the landscape.

The sequence of an increasing  $\lambda$  when the critical point is approached is not observed in our model but a decrease of it when the system goes towards the critical point, and then an increase. This coincides with theoretical predictions from percolation theory (Stauffer & Aharony 1994), at the critical point the exponential decay in the patch distribution vanishes and the patch distribution becomes a pure power-law. We did not observe the pure power-law because the size of the simulations is finite, but as  $\lambda$  becomes very small the power-law and power-law with exponential cutoff become almost identical.

When the increase of  $\lambda$  starts we already have a biodiversity collapse, thus the pattern that can be used to detect the closeness to the collapse is the decrease in  $\lambda$  observed for the patches of all species excluding the one with biggest patch. The existence of a patch of one dominant species that covers near 30% of the area can also point us that the system is near the critical point. This patch that extends from side to side of the system —also called the spanning cluster— has a variable size depending on the size of the system

(from 28% to 38%). Thus it can not be used alone as an early warning, the size of the biggest mono-specific patch should be used in conjunction with the decrease of  $\lambda$ , if the size of the biggest patch is around 10% the system might not be close to the critical point, but if its size is around 30% or more the probability that the system is near a critical transition is greater.

We calculated  $\lambda$  for the patch distribution of the Barro Colorado Island in Panama (BCI), the method could be applied because the power-law with exponential cutoff is a plausible model for the patch distributions. We did not find any tendency with time in  $\lambda$  and the biggest mono-specific patches are very small, besides we may be underestimating these sizes. There are contradicting results about the BCI regarding if it is in a steady-state, some point out BCI as been in non-equilibrium state (Fort & Inchausti 2013) and another points that it is in a steady state (Azaele *et al.* 2006). We found that  $\lambda$  is not changing with time, this is compatible with a steady-state of the forest, moreover the forest seems not to be close —by now—to a critical point.

A commonly used pattern to characterize ecological communities is the species abundance distribution, that we used in the form of rank abundance diagrams (RAD). We found that there is a change in the RAD when the critical point is approached but these changes are small and will probably go undetected (Saravia 2015). The shape of the RAD is highly dependent on the metacommunity, which is difficult to estimate as a baseline, thus there is no RAD characteristic of a community near the critical point. The species richness and Shannon diversity of the community are both resumed versions of the species abundance distribution. Richness keeps fluctuating around a more or less constant value when the community goes through the critical transition. Shannon diversity depend on the metacommunity and on the existence of trade-off: in some cases increases before the critical point and in other cases starts to decrease. Thus these two indices can be a misleading indicators of the proximity of a critical transition.

In any case the use of several indicators is recommended (Kéfi *et al.* 2014), the indicators developed in this study have the advantage of combining the spatial and temporal information thus they should be more robust than indicators based in only one kind of information (Martín *et al.* 2015). Other kinds of early warnings signals developed for spatial systems like spatial variance or autocorrelation should be adapted for multispecific communities and a detailed study comparing all of them should be done.

Percolation transitions like the one observed in this model are second order or continuous critical transitions. This means that unless the system becomes degraded and change its internal dynamics, this transitions are reversible. Much of the ecological literature was dedicated to study other kinds transitions: first order or discontinuous transitions that produce hysteresis —also called regime shifts—and rely on the understanding of deterministic equations (Solé & Bascompte 2006). These kinds of transitions are practically irreversible but in real ecosystems the presence of noise and spatial heterogeneities can convert irreversible transitions into

second order transitions (Martín *et al.* 2015), this enhances the importance of second order phase transitions like the ones detected here.

A recently published model related the savanna-forest transition to a phase transition between persistent species in a neutral model without including any niche effect (Abades *et al.* 2014). The spanning cluster that at the critical point is theoretical infinite (Stauffer & Aharony 1994; Sornette 2000). But percolation is studied in finite systems as a consequence, in small lattices the spanning cluster has a positive probability to appear. In our model, we observed the apparition of a spanning cluster in the neutral phase only at the smallest lattice sizes (100-150 sites), but a spanning cluster never appears at greater sizes. Thus we did not observe a critical transition for neutral communities, this is in contrast with Abades *et al.* (2014) that reported a critical phase transition for a neutral model using a lattice with sides from 10 to 100. In both cases the same order parameter was used —the probability of a spanning cluster— but we used a different control parameter. They used the proportion of sites with a stable population of a given species as a control parameter. It seems that their results are related to static percolation, not to the neutral model. In static percolation occupied sites are located at random in a lattice and there is no time dependence or change, when the proportion of occupied sites reach a threshold percolation is observed. This is analogous to Abades *et al.* procedure, so they obtained percolation because they varied the proportion of occupied sites (and the populations within these sites were stable), not as a result the use of neutral model.

Using an spatially implicit model Fisher & Metha (2014) described a phase transition between a neutral and niche communities. They used a stochastic Lotka-Volterra model for niche communities where neutral dynamics was added as Gaussian noise. Their mechanism is similar to ours but generalized to different kinds of interactions. Their model predicts that under stress a community will suffer a biodiversity collapse produced by a shift towards neutrality. This means that disturbed less diverse communities should have neutral dynamics. This prediction is contrary to most of the models and experimental data that suggest that niche dynamics dominate low-diversity communities while neutral dynamics will be more common in high diversity communities (Chisholm & Pacala 2011). Our results are in accordance with this last prediction: the critical point  $\rho_c$  is lower for low diversity communities. This implies that they are more sensitive to changes in environmental conditions, a small change can make them to shift to a niche phase.

The results our model and similar works in spatially implicit models implies that most ecosystems will exhibit patterns of diversity that are either strongly niche-structured or indistinguishable from neutral (Chisholm & Pacala 2011; Fisher & Mehta 2014). Anyway there is another option, several field studies have demonstrated that weak interactions is a general phenomenon for species rich communities (Martorell & Freckleton 2014, Volkov *et al.* (2009)) which is also observed for natural food webs (Wootton & Emmerson

2005), so it is possible that these communities live near the critical point between neutrality and niche. This suggests the existence of a mechanism similar to critical self organization as hypothesized by Solé *et al.* (2002) called self-organized instability. In this general mechanism the immigration of new species increase diversity and connectivity, we argue that it also might increase interaction strength; this will happen until the critical point is reached and then diversity diminishes. Thus, only communities with weak interactions can maintain high diversity in the long term. The mechanisms included in this model are generic, the only differences between species are their competitive ability and their frequency in the metacommunity. The habitat is homogeneous so niche partition mechanisms are not included but the final effect could be the same at the end: the lowering of the intensity of interactions.

Habitat fragmentation produce more isolated communities (Haddad *et al.* 2015), where less effective space is available to species thus disregarding the effect of the spatial pattern of habitat destruction this is equivalent to a community composed of less sites. Then it is probable than fragmentation shifts the critical point to lower values making the communities more sensitive to environmental changes. At the same time these communities would have a stronger niche effect and a high probability of biodiversity collapse. As we have previously mentioned, the distribution of habitable and non-habitable sites produced by fragmentation could result per se in percolation and critical phenomena. The interplay between these two critical transitions can be more complex than was previously thought (Oborny *et al.* 2007). Thus the combination of these two critical phenomena should be thoroughly studied, also the extension to different kinds of interactions including food webs, mutualistic communities and intransitive competition (Soliveres *et al.* 2015) is a fundamental next step, but great care should be given to the method of simulate the interaction coefficients to recreate realistic communities (Rohr *et al.* 2014).

The novelty of this critical transition is that it occurs previously than the produced by habitat loss, deforestation, land use changes, increased grazing, fragmentation, etc., and it only needs a change in the environmental conditions that change the strength of the interaction between species. This also could be produced if habitat connectivity and species that previously did not interact have the possibility to compete, in the same line an invading species could produce the same effect and trigger a phase transition in a formerly neutral community. A great portion of biosphere's ecosystems are under the pressures produced by human-induced activities, but human activities also produce global-scale forcings —as climate change—that can reach relatively pristine ecosystems besides they are not under a direct influence of human activities (Barnosky *et al.* 2012). This kind of changes can be represented by the phase transition described here before the ecosystem is under a direct degradation, and the detection of this kind of transition could be used as a signal to detect a global state shift in the biosphere.



# Acknowledgments

We are grateful to the National University of General Sarmiento for financial support, and to Graeme Ruxton for their English revision. This work was partially supported by a grant from CONICET (PIO 144-20140100035-CO). We thank Jordi Bascompte and Juan Bonachela for their comments and discussions on earlier versions of the manuscript. The BCI forest dynamics research project was made possible by National Science Foundation grants to Stephen P. Hubbell: DEB-0640386, DEB-0425651, DEB-0346488, DEB-0129874, DEB-00753102, DEB-9909347, DEB-9615226, DEB- 9615226, DEB-9405933, DEB-9221033, DEB-9100058, DEB- 8906869, DEB-8605042, DEB-8206992, DEB-7922197, support from the Center for Tropical Forest Science, the Smithsonian Tropical Research Institute, the John D. and Catherine T. MacArthur Foundation, the Mellon Foundation, the Celera Foundation, and numerous private individuals, and through the hard work of over 100 people from 10 countries over the past two decades. The plot project is part of the Center for Tropical Forest Science, a global network of large-scale demographic tree plots.

# References

1. Abades, S.R., Gaxiola, A. & Marquet, P.A. (2014). Fire, percolation thresholds and the savanna forest transition: a neutral model approach. *Journal of Ecology*, 102, 1386–1393.
2. Anand, M. & Langille, A. (2010). A model-based method for estimating effective dispersal distance in tropical plant populations. *Theoretical Population Biology*, 77, 219–226.
3. Azalee, S., Pigolotti, S., Banavar, J.R. & Maritan, A. (2006). Dynamical evolution of ecosystems. *Nature*, 444, 927–928.
4. Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H. & Fortelius, M. *et al.* (2012). Approaching a state shift in Earth’s biosphere. *Nature*, 486, 52–58.
5. Bascompte, J., Solé, R.V. & Sole, R.V. (1996). Habitat fragmentation and extinction thresholds in spatially explicit models. *Journal of Animal Ecology*, 65, 465–473.
6. Burnham, K. & Anderson, D.R. (2002). *Model selection and multi-model inference: A practical information-theoretic approach*. 2nd. Springer-Verlag, New York.
7. Burnham, K., Anderson, D. & Huyvaert, K. (2011). AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology*, 65, 23–35.



- 1 8.Cade, B.S. & Noon, B.R. (2003). A gentle introduction to quantile regression for ecologists. *Frontiers in*  
2 *Ecology and the Environment*, 1, 412–420.
- 3 9.Chave, J. (2013). The problem of pattern and scale in ecology: what have we learned in 20 years? *Ecology*  
4 *Letters*, 16, 4–16.
- 5 10.Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and*  
6 *Systematics*, 31, 343–366.
- 7 11.Chisholm, R.A. & Pacala, S.W. (2010). Niche and neutral models predict asymptotically equivalent species  
8 abundance distributions in high-diversity ecological communities. *Proceedings of the National Academy of*  
9 *Sciences*, 107, 15821–15825.
- 10 12.Chisholm, R.A. & Pacala, S.W. (2011). Theory predicts a rapid transition from niche-structured to neutral  
11 biodiversity patterns across a speciation-rate gradient. *Theoretical Ecology*, 4, 195–200.
- 12 13.Clauset, A., Shalizi, C. & Newman, M. (2009). Power-Law Distributions in Empirical Data. *SIAM Review*,  
13 51, 661–703.
- 14 14.Condit, R. (1998). *Tropical Forest Census Plots: Methods and Results from Barro Colorado*. Springer.
- 15 15.Condit, R., Pitman, N., Jr., E.G.L., Chave, J., Terborgh, J. & Foster, R.B.*et al.* (2002). Beta-Diversity in  
16 Tropical Forest Trees. *Science*, 295, 666–669.
- 17 16.Durrett, R. & Levin, S.A. (1994). Stochastic spatial models: a user’s guide to ecological applications.  
18 *Philosophical transactions of the Royal Society of London. Series B*, 343, 329–350.
- 19 17.Etienne, R.S. (2007). A neutral sampling formula for multiple samples and an “exact” test of neutrality.  
20 *Ecology Letters*, 10, 608–618.
- 21 18.Fisher, C.K. & Mehta, P. (2014). The transition between the niche and neutral regimes in ecology.  
22 *Proceedings of the National Academy of Sciences*, 111, 13111–13116.
- 23 19.Fort, H. & Inchausti, P. (2013). Tropical Forests Are Non-Equilibrium Ecosystems Governed by Interspecific  
24 Competition Based on Universal 1/6 Niche Width. *PLoS ONE*, 8, e82768.
- 25 20.Gillespie, C.S. (2015). Fitting Heavy Tailed Distributions: The powerLaw Package. *Journal of Statistical*  
26 *Software*, 64, 1–16.
- 27 21.Gravel, D., Canham, C.D., Beaudet, M. & Messier, C. (2006). Reconciling niche and neutrality: the  
28 continuum hypothesis. *Ecology letters*, 9, 399–409.

- 22.Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A. & Holt, R.D.*et al.* (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, 1, e1500052—e1500052.
- 23.Hoshen, J. & Kopelman, R. (1976). Percolation and cluster distribution. I. Cluster multiple labeling technique and critical concentration algorithm. *Physical Review B*, 14, 3438–3445.
- 24.Houchmandzadeh, B. & Vallade, M. (2003). Clustering in neutral ecology. *Physical Review E*, 68, 1–7.
- 25.Hubbell, S.P. (2001). *The unified neutral theory of biodiversity and biogeography*. Princeton University Press.
- 26.Hubbell, S.P. (2005). Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional Ecology*, 19, 166–172.
- 27.Kalyuzhny, M., Seri, E., Chocron, R., Flather, C.H., Kadmon, R. & Shnerb, N.M. (2014). Niche versus neutrality: a dynamical analysis. *The American Naturalist*, 184, 439–446.
- 28.Kéfi, S., Guttal, V., Brock, W.A., Carpenter, S.R., Ellison, A.M. & Livina, V.N.*et al.* (2014). Early Warning Signals of Ecological Transitions: Methods for Spatial Patterns. *PLoS ONE*, 9, e92097.
- 29.Kéfi, S., Rietkerk, M., Alados, C.L., Pueyo, Y., Papanastasis, V.P. & ElAich, A.*et al.* (2007). Spatial vegetation patterns and imminent desertification in Mediterranean arid ecosystems. *Nature*, 449, 213–217.
- 30.Kéfi, S., Rietkerk, M., Roy, M., Franc, A., Ruiter, P.C. de & Pascual, M. (2011). Robust scaling in ecosystems and the meltdown of patch size distributions before extinction. *Ecology Letters*, 14, 29–35.
- 31.Koenker, R. (2013). *quantreg: Quantile Regression*.
- 32.Leibold, M.A. & McPeck, M.A. (2006). Coexistence of the Niche and Neutral Perspectives in Community Ecology. *Ecology*, 87, 1399–1410.
- 33.Levin, S.A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73, 1943–1967.
- 34.Loehle, C., Li, B.-L. & Sundell, R.C. (1996). Forest spread and phase transitions at forest-prairie ecotones in Kansas, U.S.A. *Landscape Ecology*, 11, 225–235.
- 35.Marco, D.E., Montemurro, M.A. & Cannas, S.A. (2011). Comparing short and long-distance dispersal: modelling and field case studies. *Ecography*, 34, 671–682.
- 36.Martín, P.V., Bonachela, J.A., Levin, S.A. & Muñoz, M.A. (2015). Eluding catastrophic shifts. *Proceedings of the National Academy of Sciences*, 112, E1828–E1836.
- 37.Martorell, C. & Freckleton, R.P. (2014). Testing the roles of competition, facilitation and stochasticity on community structure in a species-rich assemblage. *Journal of Ecology*, 102, 74–85.

- 1 38. Matthews, T.J. & Whittaker, R.J. (2014). Neutral theory and the species abundance distribution: recent  
2 developments and prospects for unifying niche and neutral perspectives. *Ecology and Evolution*, 4, 2263–2277.
- 3 39. Newman, M.E.J. (2005). Power laws, Pareto distributions and Zipf's law. *Contemporary Physics*, 46,  
4 323–351.
- 5 40. Oborny, B., Szabó, G. & Meszéna, G. (2007). Survival of species in patchy landscapes: percolation in  
6 space and time. In: *Scaling Biodiversity*. Cambridge University Press, pp. 409–440.
- 7 41. Pascual, M. & Guichard, F. (2005). Criticality and disturbance in spatial ecological systems. *Trends in*  
8 *Ecology & Evolution*, 20, 88–95.
- 9 42. Pueyo, S. (2011). Desertification and power laws. *Landscape Ecology*, 305–309.
- 10 43. R Core Team. (2015). *R: A Language and Environment for Statistical Computing*. R Foundation for  
11 Statistical Computing, Vienna, Austria.
- 12 44. Rohr, R.P., Saavedra, S. & Bascompte, J. (2014). On the structural stability of mutualistic systems.  
13 *Science*, 345, 1253497.
- 14 45. Rosindell, J. & Cornell, S.J. (2009). Species-area curves, neutral models, and long-distance dispersal.  
15 *Ecology*, 90, 1743–1750.
- 16 46. Rosindell, J., Hubbell, S.P., He, F., Harmon, L.J. & Etienne, R.S. (2012). The case for ecological neutral  
17 theory. *Trends in ecology & evolution*, 27, 203–208.
- 18 47. Roy, M. & Pascual, M. (2003). Broad scaling region in a spatial ecological system. *Complexity*, 8, 19–27.
- 19 48. Saravia, L.A. (2015). A new method to analyse species abundances in space using generalized dimensions.  
20 *Methods in Ecology and Evolution*, 6, 1298–1310.
- 21 49. Scanlon, T.M., Caylor, K.K., Levin, S.A. & Rodriguez-iturbe, I. (2007). Positive feedbacks promote  
22 power-law clustering of Kalahari vegetation. *Nature*, 449, 209–212.
- 23 50. Seri, E., Maruvka & Shnerb, N.M. (2012). Neutral Dynamics and Cluster Statistics in a Tropical Forest.  
24 *The American Naturalist*, 180, E161—E173.
- 25 51. Solé, R. (2007). Scaling laws in the drier. *Nature*, 449, 151–153.
- 26 52. Solé, R.V. & Bascompte, J. (2006). *Self-organization in complex ecosystems*. Princeton University Press,  
27 New Jersey, USA.
- 28 53. Solé, R.V., Alonso, D. & McKane, A. (2002). Self-organized instability in complex ecosystems. *Philosophical*  
29 *transactions of the Royal Society of London. Series B*, 357, 667–681.

- 1 54.Soliveres, S., Maestre, F.T., Ulrich, W., Manning, P., Boch, S. & Bowker, M.A.*et al.* (2015). Intransitive  
2 competition is widespread in plant communities and maintains their species richness. *Ecology Letters*, 18,  
3 790–798.
- 4 55.Sornette, D. (2000). *Critical Phenomena in Natural Sciences - Chaos, Fractals, Selforganization and*  
5 *Disorder: Concepts and Tools*. Springer, Berlin / Heidelberg.
- 6 56.Stauffer, D. & Aharony, A. (1994). *Introduction To Percolation Theory*. Taylor & Francis, London.
- 7 57.Staver, A.C. & Levin, S.A. (2012). Integrating Theoretical Climate and Fire Effects on Savanna and Forest  
8 Systems. *The American Naturalist*, 180, 211–224.
- 9 58.Tilman, D. (1982). *Resource Competition and Community Structure*. Monographs in Population Biology.  
10 Princeton University Press, Princeton, New Jersey.
- 11 59.Tilman, D. (1994). Competition and biodiversity in spatially structured habitats. *Ecology*, 75, 2–16.
- 12 60.Vergnon, R., Dulvy, N.K. & Freckleton, R.P. (2009). Niches versus neutrality: uncovering the drivers of  
13 diversity in a species-rich community. *Ecology Letters*, 12, 1079–1090.
- 14 61.Volkov, I., Banavar, J.R., Hubbell, S.P. & Maritan, A. (2007). Patterns of relative species abundance in  
15 rainforests and coral reefs. *Nature*, 450, 45–49.
- 16 62.Volkov, I., Banavar, J.R., Hubbell, S.P. & Maritan, A. (2009). Inferring species interactions in tropical  
17 forests. *Proceedings of the National Academy of Sciences*, 106, 13854–13859.
- 18 63.Weerman, E.J., Van Belzen, J., Rietkerk, M., Temmerman, S., Kéfi, S. & Herman, P.M.J.*et al.* (2012).  
19 Changes in diatom patch-size distribution and degradation in a spatially self-organized intertidal mudflat  
20 ecosystem. *Ecology*, 93, 608–618.
- 21 64.White, E.P., Thibault, K.M. & Xiao, X. (2012). Characterizing species-abundance distributions across  
22 taxa and ecosystems using a simple maximum entropy model. *Ecology*.
- 23 65.Wootton, J.T. & Emmerson, M. (2005). Measurement of interaction strength in nature. *Annual Review of*  
24 *Ecology, Evolution, and Systematics*, 36, 419–444.
- 25 66.Zhou, S.-R. & Zhang, D.-Y. (2008). A nearly neutral model of biodiversity. *Ecology*, 89, 248–258.

## Tables

Table 1: Parameters values used in the simulations of the neutral-hierarchical model. The parameter  $\mu$  is the mortality rate;  $\alpha$  is exponent of the inverse power law dispersal kernel, between brackets is the mean dispersal distance; and  $m$  is the migration from the metacommunity.

Side	Metacomm. No. Species	$\mu$	$\alpha$ (mean dist.)	$m$
100	64	0.2	2.04 (26.66)	0.00016
150	128		2.08 (13.34)	0.0016
256	16		2.18 (6.67)	0.016
512				

Table 2: Critical points  $\rho_c$  and critical cluster size for the phase transition of a neutral/niche model. We used a range of simulation lattice sizes and two metacommunity types. The metacommunities where L: logseries, U: uniform; *Side* was the side of the simulation lattice and the total size= $Side^2$ . The other parameters were migration  $m=0.00016$ , mean dispersal distance= $26.66$ .

Metacomm. Type	Side	$\rho_c$	Critical Cluster
L	100	0.00018	0.3093
L	150	0.00102	0.3777
L	256	0.00132	0.3521
L	512	0.0015	0.2862
U	100	0.00065	0.3564
U	150	0.00094	0.3609
U	256	0.0011	0.3583
U	512	0.00136	0.3448

# Figures




Figure 1: Probability of Spanning cluster for a spatial neutral/niche model as a function of the intensity of competition  $\rho$ . The columns represent two different metacommunity types: Logseries, a metacommunity with logseries species abundance distribution (SAD); Uniform, a metacommunity with a uniform SAD. The columns represent the side of the simulation lattice, the total size is  $side^2$ . The vertical green line is the critical point, the value for parameter  $\rho$  where a phase transition between neutral and niche phases occurs. The critical point was determined as the point where the spanning probability is 0.5, the other parameters used were  $m=0.00016$ , dispersal distance = 26.66




Figure 2: Shannon diversity index and critical point for a spatial neutral/niche model as a function of the intensity of competition  $\rho$ . Columns represent with metacommunity types: Logseries is a metacommunity with logseries species abundance distribution (SAD), and the Uniform metacommunity have a uniform SAD, both with 64 species. Rows represent different lattice sizes. Points are independent simulations of the model. The parameter  $\rho$  representing the intensity of competition is the control parameter and the vertical green line is the critical point where the phase transition occurs. Other parameters used were  $m=0.00016$ , dispersal distance = 26.66.






Figure 3: Richness and critical point for simulated neutral/niche model communities as a function of the intensity of competition  $\rho$ . The columns represent different metacommunity types and the rows different grid \*side\*, where the total size of the grid is \*side x side\*. Points are repeated simulations of a the spatial model (n=30) and the vertical line is the critical point were the phase transition occurs




Figure 4: Exponential decay rate  $\lambda$  for patch size distributions as a function of the intensity of competition  $\rho$ . We fitted a power law with exponential cutoff to patch size distribution of species that does not have the biggest patch (Other MaxPatch) or species that are not the spanning species (Other Spanning), the points are the value of  $\lambda$  and the continuous lines are fitted median regressions. The dashed line connects the  $\lambda$  median for each  $\rho$ . We made 10 simulations for each  $\rho$ , metacommunities have 64 species and two different species abundance distributions (SAD): \*L\*, logseries SAD; and \*U\*, uniform SAD; the critical point for logseries is 0.0015, for uniform metacommunities is 0.0014. The size of the grid was 256\*256 sites and the other parameters used are migration=0.00016, dispersal distance=26.66.