

A peer-reviewed version of this preprint was published in PeerJ on 10 April 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.345) (peerj.com/articles/345), which is the preferred citable publication unless you specifically need to cite this preprint.

Carrillo-Larco RM, Bernabé-Ortiz A, Miranda JJ, Rey de Castro J. 2014. Peruvians' sleep duration: analysis of a population-based survey on adolescents and adults. PeerJ 2:e345 <https://doi.org/10.7717/peerj.345>

Peruvians' Sleep Duration: Analysis of a Population-Based Survey on Adolescents and Adults

Sleep duration, either short or long, has been associated with diseases such as obesity, type-2 diabetes and cardiovascular diseases. Characterizing the prevalence and patterns of sleep duration at the population-level, especially in resource-constrained settings, will provide informative evidence on a potentially modifiable risk factor. The aim of this study was to explore the patterns of sleep duration in the Peruvian adult and adolescent population, together with its socio-demographic profile. This is a cross-sectional study, secondary analysis of the Use of Time National Survey conducted in 2010. Weighted means and proportions were used to describe sleep duration according to socio-demographic variables. Poisson regressions, taking into account the multistage sampling design of the survey, were used to calculate crude and adjusted prevalence ratios (PR) and 95% confidence intervals (95% CI). Main outcomes were short- (<6 hours) and long-sleep duration (≥ 9 hours). A total of 12,424 observations, mean age 35.8 years (SD ± 17.7), 50.6% males, were included in the analysis. On average, Peruvians slept 7.7 hours (95% CI 7.4-8.0) on weekdays and 8.0 hours (95% CI 7.8-8.1) during weekends. The proportions of short- and long-sleep, during weekdays, were 4.3% (95% CI 2.9%-6.3%) and 22.4% (95% CI 14.9%-32.1%), respectively. Regarding urban and rural areas, a much higher proportion of short-sleep was observed in the former. On the multivariable analysis, compared to regular-sleepers (≥ 6 to <9 hours), short-sleepers were twice more likely to be older and to have higher educational status, and 50% more likely to be currently employed. Similarly, relative to regular-sleep, long-sleepers were more likely to have a lower socioeconomic status as per educational attainment. In this nationally representative sample, the sociodemographic profile of short-sleep contrasts the long-sleep. These scenarios in Peru, as depicted by sleeping duration, differ from patterns reported in other high-income settings and could serve as the basis to inform and to improve sleep habits in the population.

1 **Running Title:** Sleep Duration in Peru

2 Rodrigo M. Carrillo-Larco¹

3 Antonio Bernabé-Ortiz¹

4 J. Jaime Miranda^{1, 2}

5 Jorge Rey de Castro^{3, 4}

6 1. CRONICAS Center of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia,
7 Lima, Peru.

8 2. Department of Medicine, School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru.

9 3. Universidad Peruana Cayetano Heredia, Lima, Peru.

10 4. Grupo de Investigación en Sueño (GIS), Lima, Peru.

11 **Corresponding Author:**

12 Rodrigo M. Carrillo-Larco

13 CRONICAS Center of Excellence in Chronic Diseases

14 Universidad Peruana Cayetano Heredia

15 Address: Av. Armendáriz 497, 2do Piso, Miraflores, Lima 18, Peru

16 Telephone: +511 241 6978

17 E-mail: rodrigo.carrillo@upch.pe

Introduction

Over the last decades, sleep duration has changed leading to people sleeping less or more than they used to. A recent study analyzed data from the 1970s to the 2000s of ten industrialized countries and concluded that long-sleep duration, over nine hours, was more prevalent (Bin et al. 2013). However, a systematic review of cross-sectional studies conducted between the 1960s and 2000s in 15 countries reported a mixed trend: whilst seven countries had an increased sleep duration, six had a reduced one (Bin et al. 2012).

Inappropriate sleep duration, either in excess or defect, has been associated with cardiovascular diseases and other non-communicable diseases. A meta-analysis of prospective studies reported that both short- and long-sleep duration was a risk factor for dying of coronary heart disease or stroke (Cappuccio et al. 2011); whereas Guo X *et al.* reported that short-sleep duration was associated with hypertension (Guo et al. 2013). Of the different known risk factors for cardiovascular diseases, type-2 diabetes has been associated with short- and long-sleep (Cappuccio et al. 2010a); whilst obesity shows a positive association with short-sleep, long-sleep, or both (Marshall et al. 2008). In general, individuals with short- or long-sleep patterns are at higher risk of all cause mortality (Cappuccio et al. 2010b; Gallicchio & Kalesan 2009); yet, the evidence is not conclusive on this matter (Kurina et al. 2013).

Determining how much time a person sleeps is important for elucidating newer avenues for prevention as it could provide a practical target of a risk factor amenable to modification. Unfortunately there is limited data published on this matter in Latin American countries (LAC). Recent systematic reviews or meta-analysis on sleep patterns, sleep duration and its association with socio-demographic variables have not included any LAC (Bin et al. 2013; Galland et al. 2012), with the exception of one effort that included Brazil with a study that targeted adolescents (Olds et al. 2010).

There are important reasons to determine sleep duration profiles in the Peruvian population as well as in other emerging countries given their context-specific environments. First, Peru is undergoing an epidemiological transition with an increasing prevalence of NCDs (Huicho et al. 2009), and this phenomena, paired with economic development, will certainly impact the profile of sleeping pattern as well as its relationship with other diseases. Second, another context-specific characteristic from resource-constrained settings, road traffic injuries have been linked to tiredness or sleepiness (Rey de Castro & Rosales-Mayor 2010), which is very common among Lima's public transportation drivers (Risco et al. 2013), thus, having a broader picture at the population-level of sleeping-related factors would inform the potential burden of this public health problem. Third, and linked to the previous argument, home injuries could be preventable events, especially among the elderly population, as some reports suggest that people aged 75 years and over and who were long-sleepers had higher odds to suffer falls (Mesas et al. 2011). Finally, from a different and yet related angle, maternal and child health remain as an area of public health priority in the developing world and sleep duration may play a role as it may be associated with pre-term delivery and post-partum depression (Chang et al. 2010). These are some examples of how approaching sleeping patterns in low- and middle-income settings, where information at a general population level is lacking, could well inform and close existing knowledge gaps.

The aim of this study was to characterize the patterns of sleep duration in the Peruvian population, and to describe the socio-demographic profile of those considered short- and long-sleepers using a nationally representative cross-sectional survey.

58 **Methods**

59 **Study Design and Participants**

60 This is a secondary analysis of a population-based survey. The data came from the Use of Time National
61 Survey (*Encuesta Nacional de Uso del Tiempo-ENUT* in Spanish) conducted by the Peruvian National
62 Institute of Statistics and Informatics (INEI) in the year 2010. Information about this survey is publicly
63 available online (Instituto Nacional de Estadística e Informática. ENCUESTA NACIONAL DE USO DEL
64 TIEMPO 2010 FICHA TÉCNICA [Internet]. 2010 [Cited 2013, Aug. 10]. Available from:
65 <http://www.inei.gob.pe/srienaho/Descarga/FichaTecnica/286-Ficha.pdf>).

66 The original survey had a random sample of participants, drawn using standard probabilistic two-step
67 procedures: clusters (primary sample units) and households (secondary sampling units). The final sample
68 included 4580 households grouped up in 510 clusters: 3080 houses were from urban and 1500 from rural
69 areas.

70 The study population consisted of all permanent residents and those living in the selected household at the
71 moment of the survey. Information on personal needs, including sleep duration, was recorded in
72 participant's aged 12 or above. People living in institutionalized collective residences (e.g. hospitals or
73 jails) were excluded.

74 **Questionnaire & Proceedings**

75 Details of the ENUT questionnaire, sections and contents are available elsewhere . The following
76 information was included in this secondary analysis: (a) household characteristics (type of house, main
77 wall material, main floor material, total number of rooms, total number of bedrooms, water source;
78 sewage management, and sharing bathroom), and assets (iron, blender, radio, television, washing machine,
79 dryer machine, computer, telephone, Internet, car); (b) household members' characteristics (sex, age,
80 marital status, and educational attainment); (c) activities for the household that include personal needs
81 (sleep duration); and (d) job status (employment status the week prior to the survey).

82 A trained fieldworker, who visited each of the selected households, administered the survey. The
83 interviewer contacted the participant, then explained the aim of the study and ensured the confidentiality
84 of the survey. In order to avoid comprehension bias, fieldworkers read the questions as they were written.
85 All the participants were asked about the activities they performed in a 24-hour period taking as a
86 reference the week before the interview; that is, the last Monday-Friday and Saturday-Sunday period. The
87 survey was conducted between November 15th and December 30th, 2010 .

88 **Variables**

89 The outcome variable for this study was the total number of hours the participant self-reported or slept
90 during the week before the survey, assessed through the question: "*how many hours did you sleep from*
91 *Monday to Friday?*" For analysis purposes, and assuming that participants had similar sleeping hours
92 every day, to calculate the average number of hours the participant slept daily we divided the total number
93 of hours the participant slept during the previous week by five. For weekends, we proceeded in the same
94 way, dividing total hours by two. Afterwards, these variables were categorized as follows: short-sleep (<6
95 hours per day), regular-sleep (from 6 to less than 9 hours per day), and long-sleep (9 and more hours). The
96 rationale for choosing these cut-off points was based on a recent critical review on sleep duration and all-
97 cause mortality that included a study that reported an elevation in mortality risk, for men and women and
98 in a U-shaped curve, using the chosen sleep categories (Kurina et al. 2013). Furthermore, according to the
99 National Sleep Foundation an adult needs 7-9 hours of sleep, while teens (10-17 years old) need 8.5-9.25
100 hours .

The ENUT survey inquired about several types of daily activities, with the premise that all of them should add up to 24 hours. We conducted verification analysis of this sum procedures using the whole dataset and the main results presented in the ENUT's final report . After adding all the activities presented in the final report, the result was on average 31.95 hours (per day); though this number is most likely to be overestimated, as in this calculation activities that are not performed in a daily basis nor every week (e.g. buying new clothes/shoes, or buying spare parts for home appliances) were included. Furthermore, after considering only activities that are more likely to be done during a regular day or week (e.g. sleep, eat, or work) the result was 23.90 hours per day. In so doing, we believe the estimates on sleep duration are accurate enough for the purposes of this study.

Additional variables considered for the socio-demographic characteristics of the participants were area (rural or urban); region (Lima, rest of the Coast, Highlands, and the Amazon Region); gender (male, female); age (12-19, 20-35, 36-64, ≥ 65 years); education (none/primary school, high school, higher); asset index (in tertiles); job status (yes or no depending upon the participant had worked the week previous to the survey), and marital status (single, cohabiting partner/married, separated/widow/divorced).

The indicator asset index was constructed from the module of the survey comprising household characteristics and assets, according to Gordon's proposed methodology (Gordon & Pantazis 1997). The variables included in the index (Cronbach's $\alpha > 0.80$) were: type of house; main wall material; main floor material; total number of rooms; total number of bedrooms; household water source; sewage service at bathroom; if the bathroom was shared or not; and assets (iron, blender, radio, television, washing machine, dryer machine, computer, telephone, Internet, car).

Statistical Methods

The analysis was conducted with STATA 11.0 (StataCorp, College Station, TX, USA). For all calculations and estimations (results and all three tables) reported, we used the *SVY* command provided the multistage design, based upon area and region variables of the ENUT. Appropriate techniques were used for estimating results in subpopulations of interest, to guarantee accurate calculation of standard errors and, hence, inference of our findings we used the *SUBPOP* command in the Poisson regression models . We calculated means and standard deviations and percentages for continuous and categorical variables, respectively. T-test and Chi-squared test were conducted to assess differences between continues and categorical variables. To assess associations with the outcomes of interest, we used Poisson regression and report prevalence ratios (PR) and 95% confidence intervals (95% CI). For the multivariable model we utilized a stepwise backward technique (all variables were included in a model, those with a p-value > 0.05 for the Wald Test were dropped out the model) and report variables independently associated with the outcomes of interest. Throughout the analysis a $p < 0.05$ was considered to be statistical significant.

Ethics

This is a secondary-data analysis of a publicly-available dataset stored at a public national repository ; so, approval from an Institutional Review Board was not considered mandatory. The dataset used does not provide any kind of information that might have allowed us, or any other researcher, to identify participants of the study, ensuring confidentiality.

Results

Sample Characteristics

There were 18,412 observations in the original dataset and 5,988 (32.5%) were excluded due to missing values in the outcome of interest. A detailed comparison of those participants with missing data and those included in the analysis is shown in Table 1 (complete data was found for marital status); a significant difference was found with the variables region and sex. The mean age was 35.8 years (SD: ± 17.7) and there were almost a similar proportion of men and women. Details on the sample characteristics are also shown in Table 1.

Sleep Duration

Peruvians reported to sleep 7.7 hours (95% CI 7.4-8.0) on average during weekdays and 8.0 hours (95% CI 7.8-8.1) during weekends ($p < 0.001$). After categorizing this variable, during weekdays 4.3% (95% CI 2.9%-6.3%), 73.4% (95% CI 65.8%-79.8%), and 22.4% (95% CI 14.9%-32.1%) would qualify as short-, regular-, and long-sleep respectively. For weekend periods these proportions were 4.1% (95% CI 3.2%-5.4%), 65.4% (95% CI 59.4%-70.9%), and 30.5% (95% CI 24.3%-37.5%) for short-, regular, and long-sleep, respectively. Given similar sleep duration in weekdays and weekends further analyses were conducted only with the weekdays' data. Table 2 shows socio-demographic variables according to sleep duration categories.

Sub-National Analysis of Sleep Duration

The proportion of short-sleepers among men was 4.5% (95% CI 3.4%-6.0%) and with regard to women it was 4.0% (95% CI 2.3%-6.8%). However, a different trend was seen for long-sleep: 22.7% (95% CI 15.5%-32.0%) and 22.0 (95% CI 14.3%-32.3%) for women and men, respectively.

Regarding urban and rural areas, a much higher proportion of short-sleep was observed in the former: 5.2% (95% CI 3.8%-7.0%) for urban and 1.4 (95% CI 1.0%-2.1%) for rural areas. However, the proportion of long-sleep was almost the double in rural versus urban areas: 36.2% (95% CI 30.1%-42.9%) for the former and 17.9% (95% CI 11.3%-27.6%) for the latter.

Sleep Duration Profile

The socio-demographic profile of short- and long-sleep is presented in Table 3. All point estimates were attenuated, became closer 1, in all calculations following adjustment. Variables independently associated with short-sleep in the multivariable model were: age, education, and work. The higher the age the higher probability of being short-sleeper and the same was found with currently being employed. However, those with high school or no education had lower probability compared to those with higher education. On the other hand, variables independently associated with long sleep were: sex, age, education, asset index, work, and marital status.

The indicators of socioeconomic disadvantage used in the analysis did not yield a unified direction in the relationships of interest. Different markers of socioeconomic status showed different directions of association with both outcomes, i.e. worse assets index and unemployment had opposite relationships with sleeping duration outcomes compared to the estimates obtained with lower education.

Discussion

Main findings

Few studies have assessed sleep duration at the population level in developing countries, and our study aimed to characterize the patterns of sleep duration in the Peruvian population taking advantage of a nationally representative cross-sectional survey. Our results indicate that the Peruvian population has appropriate sleep patterns: sleep duration was, on average, close to 8 hours, similar during weekdays and weekends. Relatively, there was five times more short-sleep and almost half long-sleep in urban areas compared to rural ones. The socio-demographic profile characterizing short-sleepers differed from long-sleepers providing almost a mirror patterns between these two profiles, albeit with different magnitudes of association, in particular for factors such as age, education, assets and job status.

Comparison with other studies

Average duration of sleep calculations were similar to those reported previously in international (Santos-Silva et al. 2010; Steptoe et al. 2006) and national (Calderón et al. 2010; Rey de Castro et al. 2004; Rosales et al. 2009) studies. On average, sleep duration in other LAC range from 7.2 hours among Colombians (Steptoe et al. 2006), 7.3 in Venezuelans (Steptoe et al. 2006), and 7.5 in Brazilians (Santos-Silva et al. 2010). Previous Peruvian studies have reported sleep duration ranging from 6.8 to 7.5 hours among bus drivers (Rey de Castro et al. 2004; Rosales et al. 2009). Another study applied the Pittsburgh Sleep Quality Index in a small sample of people from the Andes, and reported mean sleep duration of 7 hours (Calderón et al. 2010). Our observations expand the estimations available to-date to a population-based level.

When sleep duration was approached in short- and long-sleep categories, our results markedly differ from findings in other developed and developing countries. The prevalence of short-sleep during weekdays in the USA, despite using a lower cut-off (≤ 5 hours), was 7.8% (Krueger & Friedman 2009; Nunes et al. 2008). Studies from Finland (Kronholm et al. 2006) and Korea (Ryu et al. 2011) report greater proportions of short-sleep, varying from 14.5% to 37.2%, respectively. Our study found a frequency of short-sleep duration of 4.3% during weekdays, much lower than the reported literature.

On the other hand, the prevalence of long-sleep in this study (22.4%) was much greater than equivalent estimates reported in the USA (8.5%) (Krueger & Friedman 2009), Finland (13.5%) (Kronholm et al. 2006), and Korea (4.0%) (Ryu et al. 2011). A seasonal effect has been posited to explain some of the differences observed between countries, e.g. longer sleep duration in autumn compared with summer as suggested by *Bin et al.* (Bin et al. 2011) and by a study with children (Hjorth et al. 2013). However, Peru is situated near the Equator and daylight variations during the year are not substantial. As such seasonality would not affect our calculations of sleeping categories, and therefore it does not explain the differences observed between our estimates and other studies.

The socio-demographic profile of short- and long-sleep characterized in our study is also different from those previously reported with regards to employment status and educational attainment. *Krueger et al.* (Krueger & Friedman 2009), also in a population-based study in the USA, found that those not working had increased odds of both short- and long-sleep, whereas in our study we observed such similar pattern for long-sleep only and the opposite for short-sleep. In the present study, those less educated were less likely to be short-sleepers; in contrast, studies in the USA (Krueger & Friedman 2009) and Australia (Magee et al. 2009) reported the opposite, those with completed high-school or higher education had lower probability of being short-sleep. These observations from contexts of rapid emerging countries, such as Peru, depict the complexities of addressing socioeconomic assessments (Howe et al. 2012), in relation to health outcomes that would otherwise remain unobserved in studies from more developed and established societies.

Strengths and Limitations

The study benefits from the population-based nature and the use of data from a large sample size. However, this study has limitations that must be pointed out. First, the analysis was based on data collected through self-reports and prone to recall bias, a frequent limitation in large surveys. Nevertheless, good correlation between subjective and objective measurements of sleep duration has been described both in adults (Lauderdale et al. 2008) and adolescents (Wolfson et al. 2003). Second, the methodology followed to calculate the daily sleep duration (total sleep duration in a given week divided by five) could have biased the results; nonetheless, the fact that the results are comparable to previous local reports may account for appropriate internal validity. Third, the cross-sectional design can show only association instead of causality, a limitation shared by all surveys. Fourth, a great number of missing values might bias our results and reduce possibility of inferring them at the population level; additionally there were differences when comparing some variables (area, age, education and assets index) between participants with complete and missing data for the outcome of interest. Finally, the ENUT did not provide information about other important variables that have been reported to be associated with either short or long sleep such as smoking status, alcohol consumption, ethnicity, or physical activity (Krueger & Friedman 2009; Magee et al. 2009; Ryu et al. 2011; Stranges et al. 2008). Future research would benefit from an intensive exploration of the sleeping patterns reported and important health-related outcomes, including sleeping problems, e.g. obstructive sleep apnea. Also, given the rapid socio-demographic transitions occurring in many low- and middle-income countries, variations of sleep patterns over time at the national level and their relationship with health outcomes deserve further monitoring and scrutiny.

Relevance for Public Health Policy

Translating epidemiologic research into health policy could be tough; and this could be particularly sensitive with sleep duration provided a policy could be seen as a restriction in anyone's free use of time. Peru is going through an epidemiological transition, and so are other developing countries. In this vein, there is a change in the population demographical distribution leading to a higher proportion of adults and elderly. Both scenarios have led to a higher prevalence of non-communicable diseases. Consequently, further efforts should be taken to address modifiable risk factors, including sleep duration and other sleep problems.

Describing the sociodemographic profile of the Peruvian population with higher probability of short- or long-sleep may be useful to inform and to develop potential interventions. Possible strategies might include the education of people about the benefits of adequate sleep duration and of good quality, which could raise their awareness about their sleep health. As people work more hours, ideally we could anticipate that such shift should not occur in detriment of their sleep duration, as previously reported (Basner et al. 2007; Kronholm et al. 2006). These principles have been acknowledged in the USA through their National Prevention Strategy: American's Plan for Better Health and Wellness (National Prevention Council, National Prevention Strategy, Washington, DC: U.S. Department of Health and Human Services, Office of the Surgeon General, 2011). Sleep health has been included among the topics and objectives of Healthy People 2020, a set of 10-year objectives to improve USA citizens' health, and this study sets a baseline scenario to consider correspondent prevention avenues for Peru and related contexts.

Conclusions

Peruvian population sleeps around 8 hours during weekdays and weekends. There is a much higher frequency of long-sleep in contrast to short-sleep, though the majority was regular sleepers. The socio-demographic profile of short- and long-sleeping patterns is different, not only within our study but also when compared to other settings. The profile description provided by this study might be useful to develop strategies to protect and improve advantageous sleeping habits in people with short—e.g. older people and those in the highest asset index—or long—e.g. people with no formal education or just having completed high school—sleep duration.

268 **Acknowledgments**

269 CRONICAS Center of Excellence in Chronic Diseases is supported by the National Heart, Lung, and
270 Blood Institute Global Health Initiative under the contract Global Health Activities in Developing
271 Countries to Combat Non-Communicable Chronic Diseases (Project Number 268200900033C-1-0-1). The
272 authors are grateful to Walter Mendoza, Suzy Pollard, Edmundo Rosales-Mayor and Maria Kathia
273 Cardenas for their feedback on earlier drafts of the manuscript.

References

- Instituto Nacional de Estadística e Informática. Consulta por Censos, Encuestas y Registros [Internet]. 2010 [Cited 2013, Oct. 01]. Available from: http://www.inei.gob.pe/srienaho/Consulta_por_Encuesta.asp.
- Instituto Nacional de Estadística e Informática. Encuesta Nacional de USO del TIEMPO 2010 Principales Resultados [Internet]. 2010 [Cited 2013, Aug. 10]. Available from: <http://www.inei.gob.pe/biblioineipub/bancopub/Est/Lib0960/libro.pdf>.
- Instituto Nacional de Estadística e Informática. Microdatos-Base de Datos [Internet]. 2013 [Cited 2013, Oct. 23]. Available from: <http://iinei.inei.gob.pe/microdatos/>.
- National Sleep Foundation. How Much Sleep Do We Really Need? [Internet]. [Cited 2013, Oct. 28]. Available from: <http://www.sleepfoundation.org/article/how-sleep-works/how-much-sleep-do-we-really-need>.
- The STATA Journal. A closer examination of subpopulation analysis of complex-sample survey data [Internet]. 2013 [Cited 2013, Oct. 03]. Available from: <http://www.stata-journal.com/sjpdf.html?articlenum=st0153>.
- U.S. Department of Health and Human Services. Healthy People 2020 [Internet]. 2013 [Cited 2013, Sept. 07]. Available from: <http://www.healthypeople.gov/2020/default.aspx>.
- Instituto Nacional de Estadística e Informática. ENCUESTA NACIONAL DE USO DEL TIEMPO 2010 FICHA TÉCNICA [Internet]. 2010 [Cited 2013, Aug. 10]. Available from: <http://www.inei.gob.pe/srienaho/Descarga/FichaTecnica/286-Ficha.pdf>.
- National Prevention Council, National Prevention Strategy, Washington, DC: U.S. Department of Health and Human Services, Office of the Surgeon General, 2011.
- Basner M, Fomberstein KM, Razavi FM, Banks S, William JH, Rosa RR, and Dinges DF. 2007. American time use survey: sleep time and its relationship to waking activities. *Sleep* 30:1085-1095.
- Bin YS, Marshall NS, and Glozier N. 2012. Secular trends in adult sleep duration: a systematic review. *Sleep Med Rev* 16:223-230.
- Bin YS, Marshall NS, and Glozier N. 2013. Sleeping at the limits: the changing prevalence of short and long sleep durations in 10 countries. *Am J Epidemiol* 177:826-833.
- Bin YS, Marshall NS, and Glozier NS. 2011. Secular changes in sleep duration among Australian adults, 1992-2006. *Med J Aust* 195:670-672.
- Calderón R, Quiroz A, Rosales E, and Rey de Castro J. 2010. [Sleep quality of andean inhabitants at 3200 m. Ancash-Peru]. *Rev Med Hered* 21:65-69.
- Cappuccio FP, Cooper D, D'Elia L, Strazzullo P, and Miller MA. 2011. Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *Eur Heart J* 32:1484-1492.
- Cappuccio FP, D'Elia L, Strazzullo P, and Miller MA. 2010a. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care* 33:414-420.

- 313 Cappuccio FP, D'Elia L, Strazzullo P, and Miller MA. 2010b. Sleep duration and all-cause
314 mortality: a systematic review and meta-analysis of prospective studies. *Sleep* 33:585-
315 592.
- 316 Chang JJ, Pien GW, Duntley SP, and Macones GA. 2010. Sleep deprivation during pregnancy
317 and maternal and fetal outcomes: is there a relationship? *Sleep Med Rev* 14:107-114.
- 318 Galland BC, Taylor BJ, Elder DE, and Herbison P. 2012. Normal sleep patterns in infants and
319 children: a systematic review of observational studies. *Sleep Med Rev* 16:213-222.
- 320 Gallicchio L, and Kalesan B. 2009. Sleep duration and mortality: a systematic review and meta-
321 analysis. *J Sleep Res* 18:148-158.
- 322 Gordon D, and Pantazis C. 1997. *Breadline Britain in the 1990s*, Aldershot: Ashgate.
- 323 Guo X, Zheng L, Wang J, Zhang X, Zhang X, Li J, and Sun Y. 2013. Epidemiological evidence
324 for the link between sleep duration and high blood pressure: a systematic review and
325 meta-analysis. *Sleep Med* 14:324-332.
- 326 Hjorth MF, Chaput JP, Michaelsen K, Astrup A, Tetens I, and Sjodin A. 2013. Seasonal variation
327 in objectively measured physical activity, sedentary time, cardio-respiratory fitness and
328 sleep duration among 8--11 year-old Danish children: a repeated-measures study. *BMC*
329 *Public Health* 13:808.
- 330 Howe LD, Galobardes B, Matijasevich A, Gordon D, Johnston D, Onwujekwe O, Patel R, Webb
331 EA, Lawlor DA, and Hargreaves JR. 2012. Measuring socio-economic position for
332 epidemiological studies in low- and middle-income countries: a methods of measurement
333 in epidemiology paper. *Int J Epidemiol* 41:871-886.
- 334 Huicho L, Trelles M, Gonzales F, Mendoza W, and Miranda J. 2009. Mortality profiles in a
335 country facing epidemiological transition: an analysis of registered data. *BMC Public*
336 *Health* 9:47.
- 337 Kronholm E, Harma M, Hublin C, Aro AR, and Partonen T. 2006. Self-reported sleep duration in
338 Finnish general population. *J Sleep Res* 15:276-290.
- 339 Krueger PM, and Friedman EM. 2009. Sleep duration in the United States: a cross-sectional
340 population-based study. *Am J Epidemiol* 169:1052-1063.
- 341 Kurina LM, McClintock MK, Chen JH, Waite LJ, Thisted RA, and Lauderdale DS. 2013. Sleep
342 duration and all-cause mortality: a critical review of measurement and associations. *Ann*
343 *Epidemiol* 23:361-370.
- 344 Lauderdale DS, Knutson KL, Yan LL, Liu K, and Rathouz PJ. 2008. Self-reported and measured
345 sleep duration: how similar are they? *Epidemiology* 19:838-845.
- 346 Magee CA, Iverson DC, and Caputi P. 2009. Factors associated with short and long sleep. *Prev*
347 *Med* 49:461-467.
- 348 Marshall NS, Glozier N, and Grunstein RR. 2008. Is sleep duration related to obesity? A critical
349 review of the epidemiological evidence. *Sleep Med Rev* 12:289-298.
- 350 Mesas AE, Lopez-Garcia E, and Rodriguez-Artalejo F. 2011. Self-reported sleep duration and
351 falls in older adults. *J Sleep Res* 20:21-27.
- 352 Nunes J, Jean-Louis G, Zizi F, Casimir GJ, von Gizycki H, Brown CD, and McFarlane SI. 2008.
353 Sleep duration among black and white Americans: results of the National Health
354 Interview Survey. *J Natl Med Assoc* 100:317-322.

- 355 Olds T, Blunden S, Petkov J, and Forchino F. 2010. The relationships between sex, age,
356 geography and time in bed in adolescents: a meta-analysis of data from 23 countries.
357 *Sleep Med Rev* 14:371-378.
- 358 Rey de Castro J, Gallo J, and Loureiro H. 2004. [Tiredness and sleepiness in bus drivers and road
359 accidents in Peru: a quantitative study]. *Rev Panam Salud Publica* 16:11-18.
- 360 Rey de Castro J, and Rosales-Mayor E. 2010. [Tiredness and sleepiness in rural bus drivers
361 during their job performance: Peruvian experience and proposals]. *Rev Peru Med Exp*
362 *Salud Publica* 27:237-242.
- 363 Risco J, Ruiz P, Marinos A, Juarez A, Ramos M, Salmavides F, Vega J, Kruger H, and Vizcarra D.
364 2013. Excessive sleepiness prevalence in public transportation drivers of a developing
365 country. *Traffic Inj Prev* 14:145-149.
- 366 Rosales E, Egoavil MT, Durand IS, Ccaccro NM, Herrera RF, Garcia SR, Cueva CA, Baquerizo
367 LM, and Castro JRd. 2009. [Traffic accidents and his relationship with bus drivers'
368 sleepiness and fatigue]. *Rev Med Hered* 20:48-59.
- 369 Ryu SY, Kim KS, and Han MA. 2011. Factors associated with sleep duration in Korean adults:
370 results of a 2008 community health survey in Gwangju metropolitan city, Korea. *J*
371 *Korean Med Sci* 26:1124-1131.
- 372 Santos-Silva R, Bittencourt LR, Pires ML, de Mello MT, Taddei JA, Benedito-Silva AA,
373 Pompeia C, and Tufik S. 2010. Increasing trends of sleep complaints in the city of Sao
374 Paulo, Brazil. *Sleep Med* 11:520-524.
- 375 Steptoe A, Peacey V, and Wardle J. 2006. Sleep duration and health in young adults. *Arch Intern*
376 *Med* 166:1689-1692.
- 377 Stranges S, Dorn JM, Shipley MJ, Kandala NB, Trevisan M, Miller MA, Donahue RP, Hovey
378 KM, Ferrie JE, Marmot MG et al. . 2008. Correlates of short and long sleep duration: a
379 cross-cultural comparison between the United Kingdom and the United States: the
380 Whitehall II Study and the Western New York Health Study. *Am J Epidemiol* 168:1353-
381 1364.
- 382 Wolfson AR, Carskadon MA, Acebo C, Seifer R, Fallone G, Labyak SE, and Martin JL. 2003.
383 Evidence for the validity of a sleep habits survey for adolescents. *Sleep* 26:213-216.

Table 1 (on next page)

Participants' characteristics according to complete data for the outcome variable.

Table 1: Participants' characteristics according to complete data for the outcome variable.

Variable	Missing (%)	Complete (%)	p
Area	n=5,988	n=12,424	
Urban	80.0	75.8	0.50
Rural	20.0	24.2	
Region	n=5,988	n=12,424	
Highlands	20.9	31.7	<0.001
Coast (except Lima)	16.5	24.0	
Amazon	10.4	12.1	
Lima	52.2	32.2	
Sex	n=5,066	n=12,424	
Male	59.2	50.1	0.01
Female	40.9	49.9	
Age	n=873	n=12,424	
12 - 19	13.8	20.6	0.08
20 - 35	42.2	34.1	
36 - 64	33.0	37.2	
≥ 65	11.0	8.1	
Education	n=3,929	n=12,424	
None/Primary	29.4	30.9	0.13
High School	42.3	45.8	
Higher	28.3	23.3	
Assets Index	n=5,988	n=12,424	
Lowest	19.3	23.8	0.06
Middle	25.1	34.4	
Highest	55.6	41.9	

Table 2(on next page)

Distribution of self-reported sleep duration on weekdays by socio-demographic variables. ENUT Peru 2010.

*p-value for comparison between short-sleep vs. regular

**p value for comparison between long-sleep vs. regular.

Table 2: Distribution of self-reported sleep duration on weekdays by socio-demographic variables. ENUT Peru 2010.

Variable	Sleep Duration (%)			p*	p**
	Short-sleep	Regular sleep	Long-sleep		
	n=470	8,877	n=3,077		
Area					
Urban	92.0	79.4	60.8	<0.001	0.004
Rural	8.0	20.6	39.2		
Region					
Highlands	20.8	29.3	41.5	0.09	0.03
Coast (except Lima)	28.7	24.3	21.9		
Amazon	6.2	10.9	17.2		
Lima	44.3	35.5	19.4		
Sex					
Male	53.1	50.2	49.3	0.46	0.51
Female	46.9	49.8	50.7		
Age					
12 - 19	7.1	16.4	37.0	0.02	<0.001
20 - 35	32.2	36.0	28.4		
36 - 64	53.4	40.6	23.0		
≥ 65	7.3	7.0	11.6		
Education					
None/Primary	22.0	26.9	45.6	0.006	0.001
High School	38.4	46.9	43.9		
Higher	39.6	26.2	10.5		
Assets Index					
Lowest	12.9	20.3	37.0	0.04	<0.001
Middle	32.1	33.9	36.6		
Highest	55.0	45.8	26.4		
Marital Status					
Single	30.8	36.4	51.7	0.04	0.002
Married/Living together	30.1	30.8	23.6		
Separate/Divorced/Widowed	39.1	32.7	24.7		
Job status					
No	22.3	35.2	59.0	0.009	0.001
Yes	77.7	64.8	41.1		

*p-value for comparison between short-sleep vs. regular

**p value for comparison between long-sleep vs. regular.

Table 3(on next page)

Associations between socio-demographic variables and self-reported sleep duration.
ENUT Peru 2010*.

* Multivariable models were created using backward elimination technique; variables for which there is no PR value in the adjusted model were dropped during the backward elimination process. Statistical significant results ($p < 0.05$) are in bold.

† The initial model included all the variables, sex, assets index and marital status were dropped because their p-value (Wald Test) was > 0.05 .

£ The initial model included all the variables and none were dropped because all were statistical significant for the Wald Test.

Table 3: Associations between socio-demographic variables and self-reported sleep duration. ENUT Peru 2010*.

Variables	Crude Short- vs. regular-sleep PR (95%CI)	Multivariable [†] Short- vs. regular-sleep PR (95% IC)	Crude Long- vs. regular-sleep PR (95% IC)	Multivariable [‡] Long- vs. regular-sleep PR (95% IC)
Sex				
Male	1 (Reference)		1 (Reference)	1 (Reference)
Female	0.89 (0.62 – 1.28)		1.03 (0.93 – 1.13)	0.88 (0.82 – 0.94)
Age				
12 - 19	1 (Reference)	1 (Reference)	1 (Reference)	1 (Reference)
20 - 35	2.03 (1.26 – 3.27)	1.43 (0.98 – 2.08)	0.48 (0.37 – 0.62)	0.79 (0.71 – 0.87)
36 - 64	2.92 (1.39 – 6.16)	2.18 (1.30 – 3.67)	0.36 (0.24 – 0.55)	0.60 (0.50 – 0.72)
≥ 65	2.32 (1.33 – 4.04)	2.24 (1.22 – 4.11)	0.82 (0.56 – 1.21)	0.99 (0.77 – 1.28)
Education				
Higher	1 (Reference)	1 (Reference)	1 (Reference)	1 (Reference)
High School	0.56 (0.48 – 0.66)	0.53 (0.32 – 0.86)	2.04 (1.88 – 2.22)	1.42 (1.34 – 1.51)
None/Primary	0.56 (0.39 – 0.80)	0.63 (0.57 – 0.70)	3.14 (2.27 – 4.34)	2.17(1.72 – 2.73)
Assets Index				
Lowest	1 (Reference)		1 (Reference)	1 (Reference)
Middle	1.47 (0.79 – 2.75)		0.69 (0.54 – 0.89)	0.77 (0.63 – 0.94)
Highest	1.84 (1.03 – 3.28)		0.42 (0.36 – 0.49)	0.54 (0.48 – 0.60)
Job status				
No	1 (Reference)	1 (Reference)	1 (Reference)	1 (Reference)
Yes	1.84 (1.25 – 2.69)	1.50 (1.09 – 2.06)	0.48 (0.34 – 0.67)	0.59 (0.46 – 0.75)

Marital Status			
Single	1 (Reference)	1 (Reference)	1 (Reference)
Living Together/Married	1.15 (0.86 – 1.53)	0.63 (0.51 – 0.76)	0.80 (0.71 – 0.90)
Separate/Widow/Divorced	1.39 (1.02 – 1.89)	0.62 (0.46 – 0.84)	0.75 (0.66 – 0.85)

* Multivariable models were created using backward elimination technique; variables for which there is no PR value in the adjusted model were dropped during the backward elimination process. Statistical significant results ($p < 0.05$) are in bold.

[†] The initial model included all the variables, sex, assets index and marital status were dropped because their p-value (Wald Test) was > 0.05 .

[£] The initial model included all the variables and none were dropped because all were statistical significant for the Wald Test.