

1 **Potential for primary poisoning of a Critically Endangered endemic land bird during**
2 **rodent eradication operations at Gough Island, Tristan da Cunha**

3 Alexander L. Bond¹, Michelle M. Risi², Christopher W. Jones², Peter G. Ryan²

4 ¹ RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, Sandy,
5 Bedfordshire, United Kingdom

6 ² Percy FitzPatrick Institute of African Ornithology, DST/NRF Centre of Excellence, University
7 of Cape Town, Rondebosch, South Africa

8

9 Corresponding Author:

10 Alexander L. Bond¹

11 RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, The Lodge
12 Sandy, Bedfordshire, SG19 2DL United Kingdom

13 Email address: alex.bond@rspb.org.uk

14 Abstract

15 Eradicating introduced rodents from islands restores these communities, but operations must
16 mitigate bait uptake by non-target species to ensure adequate bait coverage, and minimize
17 mortality of non-target species. Ingestion of toxic bait is a recognised risk for scavenging birds,
18 but is also a concern for generalist feeders. Gough Island, Tristan da Cunha, in the South Atlantic
19 Ocean, has introduced house mice (*Mus musculus*) that negatively affect the island ecosystem. It
20 is also home to the endemic globally threatened Gough bunting (*Rowettia goughensis*), a
21 generalist that may be affected by primary poisoning. We presented 26 wild individuals with
22 non-toxic bait pellets and observed their reactions for up to 30 min, or until they flew away.
23 While 23% of Gough buntings did not react to bait pellets, 77% showed some level of interest.
24 Generalist feeders, such as Gough bunting, may also be at risk of primary poisoning during
25 rodent eradication operations.

26

27

28 Introduction

29 Introduced rodents can cause significant damage to island ecosystems (Atkinson, 1985;
30 Jones et al., 2008; Kurle et al., 2008; Angel et al., 2009), affecting the entire community,
31 including plants, invertebrates, and vertebrates (Croll et al., 2005; Kurle et al., 2008; Mulder et
32 al., 2009; Towns et al., 2011). Since the 1970s, hundreds of island ecosystems have been
33 restored by eradicating introduced vertebrates (Towns & Broome, 2003; Howald et al., 2007;
34 Russell & Holmes, 2015), to the benefit of island communities, especially their avifauna (Jones,
35 2010; Lavers et al., 2010). Operationally, this is often accomplished by helicopter-based delivery
36 of rodenticide in cereal bait pellets (Towns & Broome, 2003; McClelland, 2011; Keitt et al.,
37 2015).

38 Such rodent eradication operations are not without their challenges. Native species may
39 compete for bait pellets, which reduces bait availability for target species (Cuthbert et al., 2012;
40 Engeman et al., 2013), and has the potential to cause non-target mortality (Wanless & Wilson,
41 2007; Wanless et al., 2010; Parkes et al., 2011), although the exposure pathways of non-target
42 species to rodent anticoagulant can be poorly known (Rattner et al., 2014). Risks of primary and
43 secondary poisoning can be mitigated in part by bringing a population into captivity during the
44 eradication operations (Wilkinson & Priddel, 2011).

45 Introduced house mice (*Mus musculus*) were introduced to Gough Island, Tristan da
46 Cunha in the 19th century, and cause significant negative effects on the island's avifauna
47 (Cuthbert & Hilton, 2004; Wanless et al., 2007; Davies et al., 2015; Dilley et al., 2015). The
48 6500 ha island presents several operational challenges owing to its remoteness, rugged terrain,
49 and potential for non-target mortality among scavengers (e.g., Subantarctic Skuas, *Stercorarius*
50 *antarcticus*) and land birds (Cuthbert et al., 2011a; Cuthbert et al., 2011b; Rexer-Huber &
51 Parker, 2011; Cuthbert et al., 2014).

52 Gough buntings (*Rowettia goughensis*) are endemic to the island, and are classified as
53 Critically Endangered by the IUCN, with a population of approximately 1000-1400 individuals,

54 and are negatively affected by the mice (Cuthbert & Hilton, 2004; Ryan & Cuthbert, 2008;
55 Birdlife International, 2015b). Preliminary work suggested that Gough buntings were unlikely to
56 be susceptible to primary poisoning by ingesting cereal bait pellets, but was based on a relatively
57 small sample size (Wanless et al., 2010). Here, we examined how wild Gough buntings reacted
58 with non-toxic cereal pellets, particularly during the winter when other food sources are likely to
59 be scarce, and the likely time of any mouse eradication attempt.

60

61 **Materials and methods**

62 From 4 April-19 July 2015, we encountered Gough buntings opportunistically in
63 southeastern Gough Island (40° 21' S, 9° 55' W), and presented them with 2-11 non-toxic cereal
64 bait pellets (Pestoff 20R; Cuthbert et al., 2011a) pale green in colour, and weighing
65 approximately 1 g. These are identical (other than lacking the rodent anticoagulant) to the bait
66 that will be used during the proposed mouse eradication. Any bunting that allowed close
67 approach was tested; this created a bias towards juvenile birds, which tend to be more
68 approachable. Pellets were tossed to within 1-2 m from birds, and birds were observed until they
69 flew away from the vicinity. We scored the reactions of buntings to the bait on a scale of 0-3,
70 where 0 represented no reaction, 1 a minimal reaction (pecking at, but not eating any of the
71 pellet), 2 a moderate reaction (picking up the pellet, but apparently not eating any), and 3 a large
72 reaction (ingesting a portion of a pellet). Most trials used dry pellets, but two trials were
73 conducted with soft, moist pellets.

74

75 **Results**

76 We recorded the reactions of 26 Gough buntings to bait pellets: 22 juveniles and 4 adults
77 (3 males and 1 female). Six (23%) ignored the bait pellets entirely, but 20 (77%) inspected the
78 pellets, with 5 (19%) scored as minimal reaction, 13 (50%) as moderate reactions, and 2 (8%) as
79 large reactions. Only juveniles were observed to eat pellets. Both trials with moist pellets were
80 presented to juveniles; one showed no interest, while the other ingested part of a pellet, and
81 carried another away.

82

83 **Discussion**

84 Wanless et al. (2010) suggested that Gough buntings would not be at risk of primary
85 poisoning through cereal bait ingestion, but only tested the reactions of 6 wild and 4 captive
86 individuals of unknown age. We found that while both adult and juvenile birds showed interest
87 in bait pellets, only juveniles were seen to consume them. This could be because of the low
88 number of adults encountered opportunistically during our trials, which were often in lowlands
89 where the breeding population is lower (Ryan & Cuthbert, 2008).

90 Gough buntings have a mixed diet consisting of invertebrates, seeds, and berries, and are
91 often seen digging and pulling up vegetation (Ryan & Cuthbert, 2008). Birds with such a general
92 diet are probably more likely to ingest bait pellets than more specialised feeders. Brodifacoum in
93 invertebrates can also cause secondary poisoning in land birds (Masuda et al., 2014). This

94 demonstrates that during rodent eradication operations, generalist feeders, in addition to
95 predatory or scavenging species, have the potential for both primary and secondary poisoning.

96 The eradication of mice from Gough Island will be a complex and highly challenging
97 endeavour. The island is > 2800 km from land, is surrounded by high cliffs, deep valleys, and
98 has complex cave areas (Cuthbert et al., 2011a; Cuthbert et al., 2011b; Cuthbert et al., 2014).
99 Gough Moorhens (*Gallinula comeri*) are also endemic, are listed as Vulnerable by the IUCN
100 (Birdlife International, 2015a), and their well-known scavenging behaviour and preliminary trials
101 suggest they will ingest bait pellets (Wanless & Wilson, 2007; Wanless et al., 2010),
102 necessitating a captive population while bait pellets remain available (Rexer-Huber & Parker,
103 2011). Our results show that a similar approach will be required for Gough bunting.

104 Planning for addressing non-target mortality prior to eradication operations is important,
105 though often inadequate, as is monitoring mortality and tissue concentrations of rodenticides
106 following eradication operations (Keitt et al., 2015; Pitt et al., 2015). Gough buntings are present
107 on at least one mouse-free offshore islet, Penguin Island, which may act as a partial refugium
108 (Ryan & Cuthbert, 2008; Wanless et al., 2010). Though the risk may be relatively low, as
109 Wanless et al. (2010) suggested, as a precautionary measure, a population of Gough buntings
110 will likely be kept captive during the eradication of house mice from Gough Island.

111

112 **Acknowledgements**

113 We thank the Administrator and Island Council of Tristan da Cunha for permission to work on
114 Gough Island, and the South African Department of Environmental Affairs' South African
115 National Antarctic Programme (SANAP) for transport to the island on the S.A. *Agulhas II* and
116 logistical support. The RSPB, the UK partner in BirdLife International, funded this research.

117

118 **References**

- 119 Angel, A., Wanless, R. M. & Cooper, J., 2009. Review of impacts of the introduced house mouse
120 on islands in the Southern Ocean: are mice equivalent to rats? *Biological Invasions*, 11:
121 1743-1754.
- 122 Atkinson, I. A. E., 1985. The spread of commensal species of *Rattus* to oceanic islands and their
123 effects on island avifaunas. In: *Conservation of Island Birds*: 35-81 (P. J. Moors, Ed.).
124 ICBP Technical Publication No. 3, Cambridge.
- 125 Birdlife International, 2015a. Species factsheet: *Gallinula comeri*. Downloaded from
126 <http://www.birdlife.org> on 14/09/2015. Birdlife International, Cambridge, UK.
- 127 Birdlife International, 2015b. Species factsheet: *Rowettia goughensis*. Downloaded from
128 <http://www.birdlife.org> on 14/09/2015. Birdlife International, Cambridge, UK.
- 129 Croll, D. A., Maron, J. L., Estes, J. A., Danner, E. M. & Byrd, G. V., 2005. Introduced predators
130 transform subarctic islands from grassland to tundra. *Science*, 307: 1959-1961.
- 131 Cuthbert, R. J., Brooke, M. d. L. & Torr, N., 2012. Overcoming hermit-crab interference during
132 rodent-baiting operations: a case study from Henderson Island, South Pacific. *Wildlife*
133 *Research*, 39: 70-77.

- 134 Cuthbert, R. J., Broome, K. G., Bradley, J. & Ryan, P. G., 2014. Evaluating the effectiveness of
135 aerial baiting operations for rodent eradications on cliffs on Gough Island, Tristan da
136 Cunha. *Conservation Evidence*, 11: 25-28.
- 137 Cuthbert, R. J. & Hilton, G. M., 2004. Introduced house mice *Mus musculus*: a significant
138 predator of threatened and endemic birds on Gough Island, South Atlantic Ocean?
139 *Biological Conservation*, 117: 483-489.
- 140 Cuthbert, R. J., Visser, P., Louw, H., Rexer-Huber, K., Parker, G. & Ryan, P. G., 2011a.
141 Preparations for the eradication of mice from Gough Island: results of bait acceptance
142 trials above ground and around cave systems. In: *Island invasives: eradication and
143 management: 47-50* (C. R. Veitch, M. N. Clout, & D. R. Towns, Eds.). IUCN, Gland,
144 Switzerland.
- 145 Cuthbert, R. J., Visser, P., Louw, H. & Ryan, P. G., 2011b. Palatability and efficacy of rodent
146 baits for eradicating house mice (*Mus musculus*) from Gough Island, Tristan da Cunha.
147 *Wildlife Research*, 38: 196-203.
- 148 Davies, D., Dilley, B. J., Bond, A. L., Cuthbert, R. J. & Ryan, P. G., 2015. Trends and tactics of
149 mouse predation on Tristan Albatross *Diomedea dabbenena* chicks at Gough Island,
150 South Atlantic Ocean. *Avian Conservation and Ecology*, 10: 5.
- 151 Dilley, B. J., Davies, D., Bond, A. L. & Ryan, P. G., 2015. Effects of mouse predation on
152 burrowing petrel chicks at Gough Island. *Antarctic Science*: doi:
153 10.1017/S0954102015000279.
- 154 Engeman, R. M., Pitt, W. C., Berentsen, A. R. & Eisemann, J. D., 2013. Assessing spatial
155 variation and overall density of aerially broadcast toxic bait during a rat eradication on
156 Palmyra Atoll. *Environmental Science and Pollution Research*, 20: 480-487.
- 157 Howald, G. R., Donlan, C. J., Galván, J. P., Russell, J. C., Parkes, J., Samaniego, A., Wang, Y.,
158 Veitch, D., Genovesi, P., Pascal, M., Saunders, A. & Tershy, B., 2007. Invasive rodent
159 eradication on islands. *Conservation Biology*, 21: 1258-1268.
- 160 Jones, H. P., 2010. Seabird islands take mere decades to recover following rat eradication.
161 *Ecological Applications*, 20: 2075-2080.
- 162 Jones, H. P., Tershy, B., Zavaleta, E. S., Croll, D. A., Keitt, B. S., Finkelstein, M. E. & Howald,
163 G. R., 2008. Severity of the effects of invasive rats on seabirds: a global review.
164 *Conservation Biology*, 22: 16-26.
- 165 Keitt, B. S., Griffiths, R., Boudjelas, S., Broome, K. G., Cranwell, S., Millett, J., Pitt, W. &
166 Samaniego-Herrera, A., 2015. Best practice guidelines for rat eradication on tropical
167 islands. *Biological Conservation*, 182: 17-26.
- 168 Kurle, C. M., Croll, D. A. & Tershy, B. R., 2008. Introduced rats indirectly change marine rocky
169 intertidal communities from algae- to invertebrate-dominated. *Proceedings of the
170 National Academy of Sciences of the United States of America*, 105: 3800-3804.
- 171 Lavers, J. L., Wilcox, C. & Donlan, C. J., 2010. Bird demographic responses to predator removal
172 programs. *Biological Invasions*, 12: 3839-3859.

- 173 Masuda, B. M., Fisher, P. & Jamieson, I. G., 2014. Anticoagulant rodenticide brodifacoum
174 detected in dead nestlings of an insectivorous passerine. *New Zealand Journal of*
175 *Ecology*, 38: 110-115.
- 176 McClelland, P. J., 2011. Campbell Island – pushing the boundaries of rat eradications. In: *Island*
177 *invasives: eradication and management: 204-207* (C. R. Veitch, M. N. Clout, & D. R.
178 Towns, Eds.). IUCN, Gland, Switzerland.
- 179 Mulder, C. P. H., Grant-Hoffman, M. N., Towns, D. R., Bellingham, P. J., Wardle, D. A.,
180 Durrett, M. S., Fukami, T. & Bonner, K. I., 2009. Direct and indirect effects of rats: does
181 rat eradication restore ecosystem functioning of New Zealand seabird islands? *Biological*
182 *Invasions*, 11: 1671-1688.
- 183 Parkes, J., Fisher, P. & Forrester, G., 2011. Diagnosing the cause of failure to eradicate
184 introduced rodents on islands: brodifacoum versus diphacinone and method of bait
185 delivery. *Conservation Evidence*, 8: 100-106.
- 186 Pitt, W. C., Berentsen, A. R., Shiels, A. B., Volker, S. F., Eisemann, J. D., Wegmann, A. S. &
187 Howald, G. R., 2015. Non-target species mortality and the measurement of brodifacoum
188 rodenticide residues after a rat (*Rattus rattus*) eradication on Palmyra Atoll, tropical
189 Pacific. *Biological Conservation*, 185: 36-46.
- 190 Rattner, B. A., Lazarus, R. S., Elliott, J. E., Shore, R. F. & Van den Brink, N., 2014. Adverse
191 outcome pathway and risks of anticoagulant rodenticides to predatory wildlife.
192 *Environmental Science & Technology*, 48: 8433-8445.
- 193 Rexer-Huber, K. & Parker, G., 2011. Captive husbandry of the Gough Island bunting and
194 moorhen. RSPB Research Report No. 42. Sandy, UK.
- 195 Russell, J. C. & Holmes, N. D., 2015. Tropical island conservation: Rat eradication for species
196 recovery. *Biological Conservation*, 185: 1-7.
- 197 Ryan, P. G. & Cuthbert, R. J., 2008. The biology and conservation status of Gough Bunting
198 *Rowettia goughensis*. *Bulletin of the British Ornithologists' Club*, 128: 242-253.
- 199 Towns, D. R. & Broome, K. G., 2003. From small Maria to massive Campbell: forty years of rat
200 eradications from New Zealand islands. *New Zealand Journal of Zoology*, 30: 377-398.
- 201 Towns, D. R., Byrd, G. V., Jones, H. P., Rauzon, M. J., Russell, J. C. & Wilcox, C., 2011.
202 Impacts of introduced predators on seabirds. In: *Seabird islands: ecology, invasion, and*
203 *restoration: 56-90* (C. P. H. Mulder, W. B. Anderson, D. R. Towns, & P. J. Bellingham,
204 Eds.). Oxford University Press, Oxford.
- 205 Wanless, R. M., Angel, A., Cuthbert, R. J., Hilton, G. M. & Ryan, P. G., 2007. Can predation by
206 invasive mice drive seabird extinctions? *Biology Letters*, 3: 241-244.
- 207 Wanless, R. M., Cooper, J., Slabber, M. & Ryan, P. G., 2010. Risk assessment of birds foraging
208 terrestrially at Marion and Gough Islands to primary and secondary poisoning by
209 rodenticides. *Wildlife Research*, 37: 524-530.
- 210 Wanless, R. M. & Wilson, J. W., 2007. Predatory behaviour of the Gough Moorhen *Gallinula*
211 *comeri*: conservation implications. *Ardea*, 95: 311-315.

212 Wilkinson, I. S. & Priddel, D., 2011. Rodent eradication on Lord Howe Island: challenges posed
213 by people, livestock, and threatened endemics. In: Island invasives: eradication and
214 management: 508-514 (C. R. Veitch, & M. N. Clout, Eds.). IUCN, Gland, Switzerland.

215

216