

A peer-reviewed version of this preprint was published in PeerJ on 8 March 2016.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1762) (peerj.com/articles/1762), which is the preferred citable publication unless you specifically need to cite this preprint.

Xiang R, Oddy VH, Archibald AL, Vercoe PE, Dalrymple BP. 2016. Epithelial, metabolic and innate immunity transcriptomic signatures differentiating the rumen from other sheep and mammalian gastrointestinal tract tissues. PeerJ 4:e1762
<https://doi.org/10.7717/peerj.1762>

Epithelial, metabolic and innate immunity transcriptomic signatures differentiating the rumen from other sheep and mammalian gastrointestinal tract tissues

Ruidong Xiang, Victor Hutton Oddy, Alan L. Archibald, Phillip E. Vercoe, Brian P. Dalrymple

Background. Ruminants are successful herbivorous mammals, in part due to their specialized forestomachs, the rumen complex, which facilitates the conversion of feed to soluble nutrients by micro-organisms. Is the rumen complex a modified stomach expressing new epithelial (cornification) and metabolic programs, or a specialised stratified epithelium that has acquired new metabolic activities, potentially similar to those of the colon? How has the presence of the rumen affected other sections of the gastrointestinal tract (GIT) of ruminants compared to non-ruminants? **Methods.** Transcriptome data from 11 tissues covering the sheep GIT, two stratified epithelial and two control tissues, was analysed using principal components to cluster tissues based on gene expression profile similarity. Expression profiles of genes along the sheep GIT were used to generate a network to identify genes enriched for expression in different compartments of the GIT. The data from sheep was compared to similar data sets from two non-ruminants, pigs (closely related) and humans (more distantly related). **Results.** The rumen transcriptome clustered with the skin and tonsil, but not the GIT transcriptomes, driven by genes from the epidermal differentiation complex, and genes encoding stratified epithelium keratins and innate immunity proteins. By analysing all of the gene expression profiles across tissues together 16 major clusters were identified. The strongest of these, and consistent with the high turnover rate of the GIT, showed a marked enrichment of cell cycle process genes ($P=1.4E-46$), across the whole GIT, relative to liver and muscle, with highest expression in the caecum followed by colon and rumen. The expression patterns of several membrane transporters (chloride, zinc, nucleosides, amino acids, fatty acids, cholesterol and bile acids) along the GIT was very similar in sheep, pig and humans. In contrast, short chain fatty acid uptake and metabolism appeared to be different between the species and different between the rumen and colon in sheep. The importance of nitrogen and iodine recycling in sheep was highlighted by the highly preferential expression of *SLC14A1*-urea (rumen), *RHBG*-ammonia (intestines) and *SLC5A5*-iodine (abomasum). The gene encoding a poorly characterized member of the maltase-glucoamylase family (*MGAM2*), predicted to play a role in the degradation of starch or glycogen, was highly expressed in the small and

large intestines. **Discussion.** The rumen appears to be a specialised stratified cornified epithelium, probably derived from the oesophagus, which has gained some liver-like and other specialized metabolic functions, but probably not by expression of pre-existing colon metabolic programs. Changes in gene transcription downstream of the rumen also appear have occurred as a consequence of the evolution of the rumen and its effect on nutrient composition flowing down the GIT.

**Epithelial, metabolic and innate immunity transcriptomic signatures
differentiating the rumen from other sheep and mammalian gastrointestinal
tract tissues**

Ruidong Xiang¹, V. Hutton Oddy², Alan L. Archibald³, Phillip E. Vercoe⁴, Brian P. Dalrymple^{1,*}

¹CSIRO Agriculture, 306 Carmody Road, St Lucia, QLD 4067, Australia

²NSW Department of Primary Industries, Beef Industry Centre, University of New England,
Armidale, NSW 2351, Australia

³The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh,
Easter Bush, Midlothian EH25 9RG, UK.

⁴School of Animal Biology and Institute of Agriculture, The University of Western Australia, 35
Stirling Highway Crawley WA 6009, Australia.

Email addresses:

Ruidong Xiang: ruidong.xiang@csiro.au

Hutton Oddy: hutton.oddy@dpi.nsw.gov.au

Alan Archibald: alan.archibald@roslin.ed.ac.uk

Philip E. Vercoe: philip.vercoe@uwa.edu.au

19 *Brian P. Dalrymple: brian.dalrymple@csiro.au

20 *: corresponding author

21 Abstract

22 **Background.** Ruminants are successful herbivorous mammals, in part due to their specialized
 23 forestomachs, the rumen complex, which facilitates the conversion of feed to soluble nutrients by
 24 micro-organisms. Is the rumen complex a modified stomach expressing new epithelial
 25 (cornification) and metabolic programs, or a specialised stratified epithelium that has acquired
 26 new metabolic activities, potentially similar to those of the colon? How has the presence of the
 27 rumen affected other sections of the gastrointestinal tract (GIT) of ruminants compared to non-
 28 ruminants?

29 **Methods.** Transcriptome data from 11 tissues covering the sheep GIT, two stratified epithelial
 30 tissues and two control tissues, was analysed using principal components to cluster tissues based
 31 on gene expression profile similarity. Expression profiles of genes along the sheep GIT were
 32 used to generate a network to identify genes enriched for expression in different compartments of
 33 the GIT. The data from sheep was compared to similar data sets from two non-ruminants, pigs
 34 (closely related) and humans (more distantly related).

35 **Results.** The rumen transcriptome clustered with the skin and tonsil, but not the GIT
 36 transcriptomes, driven by genes from the epidermal differentiation complex, and genes encoding
 37 stratified epithelium keratins and innate immunity proteins. By analysing all of the gene
 38 expression profiles across tissues together 16 major clusters were identified. The strongest of
 39 these, and consistent with the high turnover rate of the GIT, showed a marked enrichment of cell
 40 cycle process genes ($P=1.4E-46$), across the whole GIT, relative to liver and muscle, with
 41 highest expression in the caecum followed by colon and rumen. The expression patterns of
 42 several membrane transporters (chloride, zinc, nucleosides, amino acids, fatty acids, cholesterol

and bile acids) along the GIT was very similar in sheep, pig and humans. In contrast, short chain fatty acid uptake and metabolism appeared to be different between the species and different between the rumen and colon in sheep. The importance of nitrogen and iodine recycling in sheep was highlighted by the highly preferential expression of *SLC14A1*-urea (rumen), RHBG-ammonia (intestines) and *SLC5A5*-iodine (abomasum). The gene encoding a poorly characterized member of the maltase-glucoamylase family (MGAM2), predicted to play a role in the degradation of starch or glycogen, was highly expressed in the small and large intestines.

Discussion. The rumen appears to be a specialised stratified cornified epithelium, probably derived from the oesophagus, which has gained some liver-like and other specialized metabolic functions, but probably not by expression of pre-existing colon metabolic programs. Changes in gene transcription downstream of the rumen also appear have occurred as a consequence of the evolution of the rumen and its effect on nutrient composition flowing down the GIT.

INTRODUCTION

The ruminants, of which sheep, cattle, buffalo and goats are the major domesticated species, are now the most numerous large herbivores on earth. Their success is largely due to their specialized forestomachs, the rumen complex (the rumen, reticulum and omasum), and to rumination, the process of recycling the partially digested material via the mouth to reduce particle size and increase rate of fermentation (Hofmann, 1989). The forestomachs follow the oesophagus and precede the abomasum (the equivalent of the stomach of non-ruminants) (Hofmann, 1989). The evolutionary origin of the rumen is the subject of debate with out-pouching of the oesophagus, or of the stomach, as the most likely origins (Beck et al., 2009; Langer, 1988). The primary chambers of the rumen facilitate the action of a complex mixture of micro-organisms to ferment a portion of the plant polysaccharides (including starch, xylan and cellulose) and lipids to short chain volatile fatty acids (SCFAs), principally acetate, butyrate and propionate (Bergman, 1990). The SCFAs are the primary energy source in carbon of ruminants, and the rumen is the major site of their uptake.

From the rumen, partially processed plant material, nutrients, and micro-organisms pass through the omasum and enter the conventional gastrointestinal system: the abomasum, and the small and large intestines for further digestion and fermentation (in the large intestine). The abomasum is primarily a digestive organ lowering the pH of the rumen fluid and facilitating the first step of proteolysis prior to more extensive degradation in the duodenum and absorption of amino acids and small peptides. Pancreatic RNAses degrade microbial RNA in the small intestine contributing to nitrogen availability. On pasture, roughage or grass diets only small amounts of starch escape fermentation in the rumen and the remaining starch is generally digested in the small intestine, providing limited amounts of glucose (Deckardt et al., 2013).

Depending on the dietary source larger amounts of starch may escape fermentation in the rumen (Huntington, 1997). As a consequence glucose is not a major source of carbon in ruminants, and the liver is not a major site of (fatty acids) FA synthesis (Ingle et al., 1972). Biohydrogenation processes in the rumen (Van Nevel & Demeyer, 1996) increase the saturation of fatty acids (Jenkins et al., 2008; Van Nevel & Demeyer, 1996), and lipids that escape fermentation in the rumen are taken up in the small intestine. Fermentation of the remaining carbohydrates, lipid etc. occurs in the large intestine/hindgut. The hindgut is responsible for 5-10% of the total digestion of carbohydrates (Gressley et al., 2011) and for 8 to 17% of total production of SCFAs (Hoover, 1978). This contribution of hindgut fermentation may be altered on high grain diets (Fox et al., 2007; Mbanzamihigo et al., 1996). The overlap in functions of the rumen and the hindgut raises the question of whether the equivalent processes in the two tissues are undertaken by the same proteins and pathways; that is co-option of the hindgut program by the rumen, or by different proteins and pathways resulting from convergent evolution.

Unlike the stomach and subsequent segments of the GIT the rumen surface is a stratified squamous epithelium that is cornified and keratinized to protect the rumen from physical damage from the ingested plant material (Scocco et al., 2013). Due to the large numbers of microorganisms in the rumen it is also exposed to colonization of surfaces and potential attack from these organisms. The nature of the defences and the interaction between the surface of the rumen and the microbial populations has not been investigated in detail.

Herein, we utilised the latest sheep genome and transcriptome data (Jiang et al., 2014) to further dissect gene expression features of the ruminant GIT. We analyze the transcriptomes of six GIT tissue/cell types covering the majority of the sheep GIT in the context of reference samples from two other tissues with stratified squamous epithelium (skin and tonsil), another

component of the immune system (spleen), and two non-epithelial tissues (liver and muscle). Further, we systematically compared our results with existing transcriptome data from the human and pig gastrointestinal tracts and with relevant literature using candidate gene/protein based approaches. Our major aims were to identify: i) the distinctive features of ruminant GIT, ii) the common features shared between ruminant and mammalian GIT and iii) the developmental origin of the rumen.

METHODS

Data acquisition and statistical analysis

No new primary datasets were generated in this work, the major secondary datasets are included in the supplementary material. The sample preparation procedures and sequencing of the RNA are described in (Jiang et al., 2014) and experimental animal information were specified in Additional file 1: Table S1. Briefly, tissue samples were obtained from a trio of Texel sheep, i.e., ram (r), ewe (e) and their lamb (l). RNA was prepared and sequenced using stranded Illumina RNA-Seq with a yield of 70-150 million reads per tissue sample. 26 files of RNA sequence alignment data in the BAM format for 11 tissue/cell types, including skin (n=3), tonsil (n=1r), ventral rumen (n=3), abomasum (n=3), duodenum (n=1r), caecum(n=2, r and l), colon(n=3), rectum (n=3), spleen (n=2, r and l), liver (n=2, r and e) and muscle (n=3), was downloaded from the Ensembl sheep RNA sequencing archive, Oar_v3.1 (Huttenhower et al., 2009; Jiang et al., 2014). Detailed animal and gender distribution can be found in Supplementary Figure S1. Detailed raw RNA sequencing data from the same samples was also retrieved from the European Nucleotide Archive (ENA), study accession PRJEB6169. The raw mapping counts for each gene were calculated from the downloaded BAM files and the Ensembl sheep gene models (Ensembl), with additional gene models for genes at the EDC locus not included in the Ensembl sheep gene

models (Jiang et al., 2014), using HTSeq in the Python environment (Anders et al., 2015). The raw count data was normalized and clustered with DEseq2 (Love et al., 2013) to produce PCA plots and variance-stabilizing transformed gene expression values for network analysis described below. DEseq2 produced PCA sample clustering was further tested for significance using a k-means method and bivariate t-distributions based on the eigenvalues of the principle components. Calculation was performed using the stat_ellipse package (2012) and the raw outputs were presented in ggplot2 in R. EdgeR (Robinson et al., 2010) in Bioconductor in R v3.1.3 was used to analysis gene differential expression. After filtering for transcripts with at least 1 count per million in at least one of the 11 tissues, data was analysed using the Analysis of Variance-like procedure (special feature in EdgeR) and fitted to a simple model:

$y = tissue_i + animal_j + e_{ij}$. Where y is raw transcript counts, $tissue_i$ ($i=11$) is 11 types of tissues and $animal_j$ ($j=3$) is the adjustment of types of animal (lamb, ram and ewe). Transcripts with significance levels (P) < 0.01 and false discovery rate (FDR) < 0.01 for tissue effects and differentially expressed in at least one of the 11 tissues were identified.

Co-expression network analysis

Variance-stabilizing transformed RNA sequencing expression values have properties similar to normalized microarray expression values in terms of network analysis (Giorgi et al., 2013) and raw counts of differentially expressed (FDR<0.01) transcripts were variance-stabilizing transformed (Durbin et al., 2002) using DEseq2. Transformed expression values were analyzed for co-expression using PCIT (Hudson et al., 2012; Reverter & Chan, 2008) in R v3.1.3 (Watson-Haigh et al., 2010). To reduce the complexity of the network the PCIT output was filtered for pairs of genes with a correlation coefficient > 0.9 and visualized in Cytoscape v3.1.2 (Shannon et al., 2003). The network cluster algorithm ‘community cluster’ within the GLaY

plugin (Su et al., 2010) of cytoscape was used to subdivide the large network and identify explanatory sub-networks in an iterative manner until no obvious sub-network was observed in the large network. Pig genes assigned to 10 clusters showing differential expression in the GIT (Freeman et al., 2012) were mapped to sheep genes based on their gene symbols. The probability of over or under representation of pig GIT genes in a sheep GIT gene cluster was calculated using the hypergeometric distribution (Andrews et al., 1999). Functional enrichment of shared sets of genes within sheep clusters was analyzed using GOrilla (Eden et al., 2009) to identify biological pathways.

Gene expression pattern clustering

The transcripts present in the gene networks described above, and with an ANOVA $P < 0.01$ and a FDR < 0.01 , were included in k-mean clustering in R v3.1.3 based on $\log_2$ fold change across 11 tissues with abomasum being the reference. The k-mean analysis aimed to identify expression patterns to represent transcript groups showing elevated expression levels for the following sets of tissues v. the remaining tissues: 1) all GIT tissues, i.e., rumen, abomasum, duodenum, caecum, colon and rectum, 2) rumen and abomasum, 3) rumen and intestinal tissues, 4) abomasum and intestinal tissues, 5) rumen, 6) abomasum, 7) intestinal tissues, 8) rumen and skin, 9) rumen and tonsil, 10) rumen, skin and tonsil, 11) spleen, duodenum, caecum, and colon. The transcript names are determined based on the tissue(s) where included transcripts showed the highest expression. We filtered these identified transcript clusters with the criteria that 1) the average absolute expression of the transcript at the highest expressed tissue > 3 count per million, 2) the $\log_2$ expression fold difference of expression of the transcript from the tissue within the reference tissue group with the highest expression to the tissue within the elevated expressed tissue group with the highest expression, be > 0.5 , and 3) from the tissue with the

highest expression to the tissue with the lowest expression within the elevated tissue group be < 0.5. The final expression of each transcript is presented in the format of log₂ Fragments Per Kilobase of exon per Million fragments mapped (FPKM). Selected gene members and associated pathways were presented in heat maps based on their log₂ FPKM values using GENE-E (Gould).

To understand the GIT associated SLC family genes, we performed a network analysis of expression as above. The PCIT output of network matrix was filtered for correlation coefficient > 0.7, clustered by GLayer (Su et al., 2010) and visualized in Cytoscape v3.1.2 (Shannon et al., 2003).

Comprehensive transcript annotation

To complement the sheep genome annotation, we used multiple annotation sources and software to identify the function of the products encoded by the identified transcripts. Firstly, the transcripts of interest, both with or without a gene symbol, were validated for existence on the sheep genome, using comparisons of the sheep gene along within the locus with its ortholog(s) of human and bovine from Ensembl and NCBI. Secondly, GO was used to annotate genes. Thirdly, the functions and annotations of the genes were searched in Ensembl and NCBI, if no available description or gene information were identified, the biomedical literature was searched with GenCLiP 2.0 (Wang et al., 2014). When multiple biomedical functions were listed, functions related to gastrointestinal activity were prioritized for annotation. Fourthly, for a subset of genes Unigene (McGrath et al., 2010) and Genevestigator (Hruz et al., 2008) were used to identify transcript expression patterns in cattle and humans respectively. Protein sequences analysis was performed using Radar (Heger & Holm, 2000), to identify amino acid sequence repeats, and NetOGlyc 4.0 (Steentoft et al., 2013), to identify glycosylation sites.

Data access

No new primary datasets were generated in this work, the major secondary datasets are included in the supplementary material.

RESULTS AND DISCUSSION

Clustering of sheep GIT tissue transcriptomes

We performed principal component analysis (PCA) using RNA-Seq data from six GIT (rumen, abomasum, duodenum, caecum, colon and rectum), two epithelial (skin and tonsil), an immune (spleen) and two reference (liver and muscle) tissue/cell types from a trio of Texel sheep (ram, ewe and lamb (Jiang et al., 2014)). We included a total of 26 tissue samples, a similar tissue sample coverage to a previous transcriptomic study of the pig GIT (Freeman et al., 2012) to which the results of this analysis will be compared below. Three clusters of tissues were identified at the 95% confidence interval: cluster 1, skin, tonsil and rumen, cluster 2, muscle, and cluster 3, liver, spleen and the remaining GIT tissues (Figure 1A, Additional file: Figure S1A).

Identification of common and specific GIT and epithelial transcriptomic signatures

To identify the genes driving the clustering of the tissues we identified those transcripts with an ANOVA $P < 0.01$ and a false discovery rate (FDR) < 0.01 , for differential expression in at least one tissue versus the other tissue types. This multi-tissue comparison reduced the impact of the small sample size for some tissues, in particular the duodenum (one tissue sample). Secondly, for a conservative gene network cluster analysis, the pair-wise gene correlation coefficient cut-off was set to 0.9 and we further filtered transcripts based on relative (fold change) and absolute (count per million) expression levels. We identified 16 major gene expression patterns,

217 representing common and specific transcriptomic signatures of the epithelial and GI tissues,
 218 accounting for 639 different transcripts (Figure 2A). A full list of the expression of the genes
 219 across the tissues with assignment to clusters is available (Additional file 1: Table S2, S3). Gene
 220 Ontology enrichment analysis of the clusters identified a number of significantly enriched terms
 221 (Table 1). A full list of the genes contributing to the enrichments is available (Additional file 1:
 222 Table S4). Most notable was the highly significant enrichment of the genes in the epithelia-
 223 intestine cluster for the GO-term, “cell cycle process”. The higher expression of the majority of
 224 these genes in the epithelial and GIT tissues (Figure 2, Supplementary Table S2) is consistent
 225 with the much higher turnover rate of these tissues compared to liver and muscle (Milo et al.,
 226 2010) and may contribute to the structural adaptability of the rumen epithelia to different diets
 227 and health conditions (Dionissopoulos et al., 2012; Penner et al., 2011). Epithelia structure
 228 related pathways including ‘cell junctions’ showed significant enrichment in genes highly
 229 expressed in the rumen and the large intestine (Table 1). Gene members involved in cell junction
 230 functions have been reported to be important for the rumen epithelia to maintain pH homeostasis
 231 (Dionissopoulos et al., 2012; Steele et al., 2011a). Two other very significant enrichments were
 232 observed, “flavonoid biosynthetic process” in the rumen-intestine-liver cluster and “regulation of
 233 chloride transport” in the large intestine cluster (Table 1). The mammalian Epidermal
 234 Development Complex (EDC) locus is a cluster of up to 70 adjacent genes encoding proteins
 235 with roles in the development and the structure of stratified epithelia (Kypriotou et al., 2012).
 236 Although no significant enrichment of genes in the rumen cluster was identified by GO analysis
 237 several genes in the EDC region were very significantly overrepresented in the cluster (Table 1).
 238 This is consistent with our previous identification of several ruminant specific genes at the EDC
 239 locus highly preferentially expressed in the rumen (Jiang et al., 2014). The genes in the epithelia-

rumen-tonsil cluster were also very significantly enriched for EDC genes (Table 1). Thus the clustering of the rumen with the skin and tonsil appears to have been driven by genes involved in the development and structure of the stratified epithelium.

The stratified squamous rumen epithelium expression signature

The EDC locus genes are not the only genes encoding proteins involved in the synthesis of the cornified surface of the rumen and we looked for additional genes involved in cornification preferentially expressed in the rumen compared to skin and tonsil. The cross linking of the proteins of the cornified surface is mediated by transglutaminases (TGMs) (Eckert et al., 2005). Multiple TGMs are expressed in the rumen in this study, TGM1 and TGM3 appear to be the major rumen transglutaminases, but are also highly expressed in the skin (Figure 3). Keratins are major components of the cornified layers so we asked the question, are there keratin genes highly preferentially expressed in the rumen? Although no *KRT* genes showed expression as exclusive to the rumen as some of the EDC locus genes in our data, *KRT36* was grouped in the rumen expression cluster (Figure 3, Additional file 1: Table S3, Additional file 2: Figure S2), with significantly elevated expression in rumen, compared to the other studied tissues, and limited expression in skin. *KRT36* was previously identified as a novel keratin gene only expressed in sheep hair cortex (Yu et al., 2011) and its rumen expression showed significant responses to dietary changes in cattle (Li et al., 2015). However, in humans the highest expression of *KRT36* was in the tongue (Genevestigator (Hruz et al., 2008) analysis). Overall the transglutaminases and keratins do not appear to be as preferentially expressed in the rumen as some of the EDC locus genes.

262 Kallikrein-related peptidases are involved in the turnover of the cornified layers of the stratified
263 epithelia, and deficiencies can lead to altered turnover of the surface layers of the epithelia
264 (Hovnanian, 2013). In our study, *KLK12* is the only *KLK* family member preferentially
265 expressed in the rumen (Figure 3, Additional file 1: Table S2). Members of the SPINK (serine
266 peptidase inhibitor, Kazal type) family are inhibitors of the *KLK* family peptidases (Hovnanian,
267 2013), *SPINK5* is the only member of the family that is highly expressed in the rumen (Figure 3,
268 Additional file 1: Table S2) in our data, but is also highly expressed in the tonsil and skin.
269 *KLK12* and *SPINK5* may be involved in the regulation of the turnover and thickness of the
270 cornified surface of the rumen epithelium, but may not form a rumen specific system.

271 **Rumen micro-organism interactions**

272 The rumen is the site of frequent interaction between the host and very dense populations of
273 micro-organisms. In our study, *DUOX2* and *DUOXA2* encoding subunits of dual oxidase were
274 preferentially expressed in the rumen (Figure 3), while *DUOX1* showed rumen-biased expression
275 (Figure 3) and *DUOXA1* was highly expressed in all epithelia tissues (Figure 3). This
276 observation is in line with the findings in the pig where the highest expression of *DUOXA1* and
277 *DUOX1* was in the epithelial tissues. e.g., tongue and lower oesophagus (Freeman et al., 2012).
278 In humans, the *DUOXA1* and *DUOX1* genes are also most highly expressed in epithelia tissues
279 exposed to air, whilst *DUOX2* and *DUOXA2* are most highly expressed in a different set of
280 tissues including the GIT (Genevestigator (Hruz et al., 2008) analysis). Thus, our findings
281 suggest that the *DUOX1*s are active in general epithelial tissues, while *DUOX2*s are probably
282 active specifically in rumen to play a major role in controlling microbial colonization. Previously
283 in sheep, the highest expression levels of *DUOX1* and *DUOX2* were reported in the bladder and

abomasum, respectively, but the rumen and epithelial tissues were not included in the tissues surveyed (Lees et al., 2012).

PIP, encoding prolactin-induced protein (an aspartyl protease), was preferentially expressed in the rumen (Figure 3). In humans, *PIP* is also highly expressed in epidermal (Genevestigator (Hruz et al., 2008) analysis) and exocrine tissues, and in pigs in the salivary gland. Although *PIP* has been reported to be involved in regulation of the cell cycle in human breast epithelial cells (Cassoni et al., 1995; Naderi & Vanneste, 2014), its expression pattern in sheep (not part of the cell cycle cluster) is more consistent with a role in mucosal immunity (Hassan et al., 2009). Also highly expressed in the rumen were members of the *SERPINB* family of peptidase inhibitors (Figure 3), which are involved in the protection of epithelial surfaces in humans (Wang et al., 2012) and mice (Sivaprasad et al., 2011). EDC locus genes *PGLYRP3* and *PGLYRP4* encode peptidoglycan recognition proteins in the N-acetylmuramoyl-L-alanine amidase 2 family, which bind to the murein peptidoglycans of Gram-positive bacteria as part of the innate immune system. Additional EDC locus genes, *S100A8*, *S100A9* and *S100A12* (calgranulins A, B and C), encode key players in the innate immune function (Funk et al., 2015; Tong et al., 2014).

Rumen steroid metabolism

Amongst the genes preferentially expressed in the rumen (and often the liver) we identified a number of aldo/keto-reductases (Figure 4). *AKR1C1* can catalyze the conversion of progesterone to 20- α -hydroxy-progesterone (PGF2 α) (Penning, 1997), retinals to retinols and bioactivates and detoxifies a range of molecules (El-Kabbani et al., 2011). Intravenous injection of PGF2 α in goats has been shown to increase contraction of rumen smooth muscle, which leads to a reduction in the contraction rate of the rumen (van Miert & van Duin, 1991; Veenendaal et al., 1980). *AKR1C1* has also been reported to be preferentially expressed in the rumen of cattle (Kato

et al., 2015). The exact role of AKR1C1 in the rumen is unknown,. In addition, the gene encoding the related enzyme AKR1D1 (catalyzes the reduction of progesterone, androstenedione, 17-alpha-hydroxyprogesterone and testosterone to 5-beta-reduced metabolites) is highly expressed in the rumen and the liver and the gene encoding ARK1C4 in the rumen, liver and duodenum (Figure 4). The products of these genes are also likely to be involved in the metabolism of steroids in the rumen epithelium. In addition, we observed marked pathway enrichment of flavonoid biosynthetic process due to the identification of five members of the UDP-glucuronosyltransferase (UGT) gene family [29], with the highest expression levels in the rumen and liver (Additional file 1: Table S2). Flavonoids are only produced by plants, but UGT enzymes are highly active in mammals and catalyze the glucuronidation of a diverse chemical base including steroids, bile acids and opioids [29]. The functions of the products of these genes in the rumen require further investigation. However, results discussed here suggest important interactions between the rumen wall and activity of steroids.

Comparison of the sheep and pig GIT transcriptomes

To compare the ruminant and a closely related non-ruminant mammal GIT transcriptomes (Jiang et al., 2014), we mapped those transcripts previously reported to show specific expression patterns in the pig GIT (Freeman et al., 2012) to the sheep gene network (Figure 2B). Pig is the genomically closest non-ruminant to the ruminants (Groenen et al., 2012; Jiang et al., 2014) for which sufficient GIT transcriptome data is available. The overall overlap of the 639 genes in the sheep GIT network and the 2634 mappable pig GIT genes is 179, which is highly significant (Table 2). The smaller number of genes showing differential expression in our study versus the pig study is due to the application of stringent statistical filtering thresholds to minimize the

impact of the small number of samples per tissue. However, the overlap of 627 genes between the set of 2475 sheep genes identified using relaxed filtering criteria and the 2634 pig genes was also highly significant ($P < 10E-20$), supporting the robustness of the approach. The set of 179 overlapping genes was highly significantly enriched for the GO-term “cell cycle process” (Table 2). The overlap of genes between the pig intestine clusters and the sheep epithelia-intestine cluster was highly significant and the overlap genes were again very highly significantly enriched for the GO-term “cell cycle process” (Table 2). A full list of the genes in the overlap and assignment to the pig and sheep gene clusters is available (Additional file 1: Table S5). Furthermore, pig genes preferentially expressed in the tongue and oesophagus have a highly significant overlap with sheep genes with high expression in the rumen and epithelial tissues (Figure 2B), enriched for the GO-term “epidermis development” (Table 2). Our results emphasises the contribution of cell cycle to the renewal of mammalian GIT epithelial surfaces (Crosnier et al., 2006).

Ruminant specific pathways for SCFA uptake and GIT metabolism?

SCFAs are the major source of energy in ruminants, with the primary sources of SCFAs being the rumen, and to a much lesser extent the large intestine. Carbonic anhydrases, which hydrate CO_2 to bicarbonate, are thought to play a significant role in the uptake of SCFAs by an SCFA/bicarbonate antiporter, and by providing protons at the rumen epithelium to neutralize the SCFAs and promote their diffusion into the ruminal epithelium (Bergman, 1990; Wang et al., 1996). There are many members of the carbonic anhydrase gene family (Tashian, 1989), several of which are expressed in mammalian gastrointestinal tissues (Freeman et al., 2012; Kivel et al., 2005; Parkkila et al., 1994; Tashian, 1989). In ruminants, *CAI* has previously been reported to encode a rumen specific carbonic anhydrase with low activities in the blood (unlike in other

mammals) and in the large intestines (Carter, 1971). Consistent with this, compared to all of the other tissues in our dataset, *CA1* is highly expressed in the rumen and, albeit with lower but significant expression, in the large intestine (Figure 4). *CA2* and *CA7* appear to encode the major carbonic anhydrases in the large intestines (Figure 4). In humans *CA1*, *CA2* and *CA7* are highly expressed in the colon (Genevestigator (Hruz et al., 2008) analysis). In contrast in pigs, whilst *CA2* is highly expressed in the stomach, it is not highly expressed in the large intestine and *CA1* and *CA7* were not reported to be differentially expressed across the GIT (Freeman et al., 2012).

The apical membrane SCFA/bicarbonate antiporter exchanges intracellular bicarbonate with intra-ruminal SCFA and consistent with previous publications, *SLC4A9*, preferentially expressed in the rumen in our dataset (Figure 4), encodes the most likely antiporter. The proposed basolateral membrane SCFA/bicarbonate antiporter gene *SLC16A1* (exchanges intracellular SCFA with blood bicarbonate), which has highest expression in the rumen in our dataset, followed by the colon and rectum, has a much more general expression across the tissues than *SLC4A9* (Figure 4). These expression patterns are consistent with previous findings in cattle (Connor et al., 2010). *SLC16A1* is also likely to be involved in the transport of ketone bodies into the blood supply to the basolateral surface of the rumen epithelium (van Hasselt et al., 2014).

HCO_3^- -independent apical uptake of acetate in the rumen has also been observed (Aschenbach et al., 2009). However, the transporter has not been identified, with candidates proposed in the *SLC4A*, *SLC16A*, *SLC21A*, *SLC22A* and *SLC26A* families (Aschenbach et al., 2009). Members of the *SLC21A* and *SLC22A* families showed generally low expression in the rumen in our study (Additional file 1: Table S2). In addition to *SLC16A1* and *SLC4A9* discussed above, *SLC26A2* and *SLC26A3* are highly expressed in the rumen in our dataset (Figure 4). Both genes encode

376 apical anion exchangers confirming them as candidates for encoding the apical HCO_3^- -
 377 independent acetate uptake transporter. SLC26A3 is a $\text{Cl}^-/\text{HCO}_3^-$ exchanger (see fluid and
 378 electrolyte balance section below) and therefore is unlikely to be an HCO_3^- -independent acetate
 379 transporter. However, SLC26A2 is a $\text{SO}_4^{2-}/\text{OH}^-/\text{Cl}^-$ exchanger (Ohana et al., 2012) and remains a
 380 candidate for the proposed apical HCO_3^- -independent acetate transporter. An HCO_3^- -independent
 381 basolateral maxi-anion channel for SCFA^- efflux to blood has also been proposed without an
 382 assigned transporter (Georgi et al., 2014). A survey of ABC (ATP-binding cassette) family
 383 transporters identified *ABCC3* as the most preferentially expressed in the rumen in our dataset
 384 and with the second highest expression in the large intestine (Figure 4). *ABCC3* is an organic
 385 anion transporter with a possible role in biliary transport and intestinal excretion (Rost et al.,
 386 2002). Therefore, *ABCC3* may be involved in the efflux transport of SCFA^- from the rumen
 387 epithelium to blood.

388 In most mammals, including humans, the liver is the major site of the synthesis of ketone bodies
 389 (acetoacetate and beta-hydroxybutyrate), but in ruminants the epithelium of the rumen is a major
 390 site of *de novo* ketogenesis (Lane et al., 2002). *HMGCS2* encodes an HMG-CoA synthase (3-
 391 hydroxy-3-methylglutaryl-CoA Synthase 2) in the ketogenesis pathway (Figure 5). This gene is
 392 significantly associated with bovine butyrate metabolism (Baldwin et al., 2012) and the encoded
 393 enzyme was predicted to be the rate limiting enzyme in sheep ruminal ketone body synthesis
 394 (Lane et al., 2002). As expected, in our data *HMGCS2* is highly expressed in the rumen
 395 compared to the other GIT tissues and the liver (Figure 4). *ACADS*, *HMGCL* and *BHDI*, which
 396 encode other enzymes involved in the ketone body pathway (Figure 5), are also highly expressed
 397 in the rumen relative to most of the other tissues studied (Figure 4). *HMGCS1* and *ACAT2* may
 398 also contribute to the ketone body pathway in the rumen, but their highest expression levels are

in the liver (Table S1). However, their expression in the rumen has been reported to actively respond to different diets (Steele et al., 2011b) and acidosis conditions (Steele et al., 2012) in cattle. Whilst *HMGCS2* is quite highly expressed in the colon, in contrast *ACADS*, *HMGCL* and *BHDI* are not highly expressed (Figure 4), consistent with the colon not being a major contributor to ketone body synthesis. Genes encoding enzymes for other steps in the pathways from acetate and butyrate to ketone bodies are much more generally expressed across the tissues, although expression of *ECHS1* and *ACAT1* are significantly higher in the rumen than in other GIT tissues (Figure 4). In humans, in addition to the liver, *HMGCS2* also has high expression in the intestine, including the jejunum and colon (Genevestigator (Hruz et al., 2008) analysis). In contrast, the only enzyme in the pathway (Figure 4, 5) reported to be preferentially expressed in the pig GIT was *BDHI*, in the fundus of the stomach (Additional file 1: Table S5). Thus the rumen, abomasum, duodenum, caecum, colon and rectum in sheep all appear to have subtly different SCFA transport and metabolism systems, and in the equivalent compartments of the GIT appear to be different between sheep, humans and pigs.

Long chain fatty acids (LCFAs) uptake, cholesterol homeostasis and bile acid recycling

Due to the activity of the microbial populations of the rumen and the production of SCFAs ruminants have less reliance on dietary LCFAs than non-ruminants. Does this reduced importance lead to detectable differences in the transcriptome? The small intestine is the principal site of uptake of LCFA and cholesterol homeostasis, and consistent with this the genes encoding the well characterized components of the intestinal fatty acid uptake (*CD36*, *SLC27A2/4/5* and *FABP2* (Wang et al., 2013)) and cholesterol homeostasis (*NPC1L1* and *ABCG5/8* (Wang et al., 2013)) systems are expressed in the sheep small intestine (Figure 4), as they are in humans and most are in the pig (Freeman et al., 2012). *FABP2* and *ABCG5* are

particularly preferentially expressed in the sheep small intestine relative to other GIT tissues (Figure 4). However, it is thought that the major route of LCFA uptake at the apical membrane of the GIT epithelium is by passive diffusion (Abumrad & Davidson, 2012). Bile acids secreted by the liver and stored in the gall bladder before being released into the small intestine play a major role in the uptake of LCFAs. Bile acids are recycled in the intestine. SLC10A2 in the apical membrane and SLC51A and SLC51B in the basolateral membrane are proposed to constitute the uptake systems in the human small intestine (Ballatori et al., 2013). *SLC10A2* is also preferentially expressed in the small intestines of the pig, but preferential expression of *SLC51A/B* has not been reported (Freeman et al., 2012). In sheep *SLC10A2* is preferentially expressed in the small intestine, albeit at a low level (Figure 4). Whilst *SLC51B* is highly expressed in the duodenum in sheep, the highest expression of the two subunits together in sheep (*SLC51A/B*) is in the caecum and the colon (Figure 4), where they are also expressed in humans and mice (Genevestigator (Hruz et al., 2008) analysis). Although described as subunits of a complex, *SLC51A* and *SLC51B* have also been reported to be regulated differently (Ballatori et al., 2013), thus the balance between expression of *SLC10A2* and *SLC51A* and *SLC51B* may indicate differences in the bile acid uptake pathways in the duodenum, large intestines and liver of sheep.

Overall despite the reduced importance of LCFAs sheep appear to have a very similar systems to human and pigs for LCFA uptake and bile acid recycling.

Saccharide metabolism

Again as a consequence of the activity of the rumen microbes in mature ruminants the uptake of dietary glucose may be less than 10% of glucose requirements (Young, 1977). The dietary

glucose comes primarily from the degradation of polysaccharides, in particular in the small intestine of starch that has escaped degradation by the rumen microbial population. The primary source of alpha-amylase required to digest the long polymers is the pancreas, which was not investigated in this study. Genes encoding three enzymes likely to contribute to the digestion of starch and other alpha-glycans, *MGAM* (maltase-glucoamylase), *MGAM2* (maltase-glucoamylase 2) and *SI* (sucrase-isomaltase) (Nichols et al., 2003), were preferentially expressed in the tissues studied here. *SI* was preferentially expressed in the intestine-low in rectum gene cluster, *MGAM2* was highly expressed in all intestinal tissues, while *MGAM* was also preferentially expressed in the intestine (primarily the duodenum), but at a much lower level (Figure 4). In humans (Genevestigator (Hruz et al., 2008) analysis) and pigs (Freeman et al., 2012), both *MGAM* and *SI* are preferentially expressed in the small intestine. Expression of the orthologues of *MGAM2* has not been reported in the GIT of humans (Genevestigator (Hruz et al., 2008) analysis) or pigs (Freeman et al., 2012).

The mammalian *MGAM* and *MGAM2* genes appear to have arisen by tandem duplication of a single ancestral gene at the base of the mammals (Nichols et al., 2003; Nichols et al., 1998). *MGAM2* genes are present in most mammals, but have been annotated as possible pseudogenes in a number of species, including man (NCBI LOC93432). *MGAM2* is not well characterized in any species. Comparative analysis of the protein sequences of *MGAM* and *MGAM2* showed that *MGAM2* has additional sequence at the carboxy-terminus comprised of multiple copies of a 40 amino acid repeat not present in *MGAM* (Figure 6). The repeat unit is enriched in serine and threonine, with similar sequences in the predicted sheep, cattle, pig and to a much lesser extent human proteins (Figure 6). The repeat unit of *MGAM2* is predicted to be heavily glycosylated (Steentoft et al., 2013) to form a mucin-like domain. As in the rumen the microbial population in

the large intestine ferments plant material, contributing up to 10% of the total carbohydrate fermentation and conversion to SCFAs in the ruminant GIT (Gressley et al., 2011). Whilst the role MGAM2 is unclear it appears to represent a contribution from the host to the breakdown of plant polysaccharides by the bacterial population in the large intestine. MGAM produces glucose from maltose and MGAM2 may have a similar functionality, and therefore contribute to the uptake of the scarce supply of glucose in ruminants. Alternatively the high expression of MGAM2 and low expression of MGAM may reflect the reduced availability of glucose in the rumen GIT. Further investigation of this gene and the activity and function of its encoded protein will improve our understanding of carbohydrate metabolism in the large intestine of ruminants.

In humans the major uptake of glucose in the GIT occurs in the small intestine via SLC5A1 (aka SGLT1) in the apical membrane, and SLC2A2 (aka GLUT2) in the basolateral membrane (Roder et al., 2014). The expression pattern of these two genes in sheep (Figure 4) and pigs (Freeman et al., 2012) is consistent with a similar process in all three species.

Nitrogen acquisition and recycling

A high level of nitrogen recycling in the GIT is a characteristic of ruminants. Urea is the major input from the animal (primarily via the saliva and the rumen epithelium) and anabolic-N sources (in the small intestine) and ammonia (in the rumen, small and large intestines) are the major uptake molecules from the GIT (Lapierre & Lobley, 2001). *SLC14A1* (Figure 4), encoding SLC14A1 which mediates the basolateral cell membrane transport of urea, a key process in nitrogen secretion into the GIT (Abdoun et al., 2010), is highly preferentially expressed in the rumen in our dataset (Figure 4). However, in cattle expression of *SLC14A1* was not affected by differences in dietary N (Rojen et al., 2011) and doubts remain about the role of SLC14A1 in increasing rumen epithelial urea permeability at low dietary N. Urea is also thought to be

released by the epithelium of the small and large intestines (Lapierre & Lobley, 2001), but our analysis did not identify a potential transporter.

Urea is converted to ammonia by microbial ureases and is used by rumen microorganisms to synthesize microbial proteins (75-85% of microbial N) and nucleic acids (15-25% of microbial N) (Fujihara & Shem, 2011) which are subsequently digested by the host in the intestines, thus recovering the majority of the secreted nitrogen (Abdoun et al., 2006). Consistent with this, *SLC3A1* (neutral and basic amino acid transporter) in our study is preferentially expressed in the duodenum (Figure 4), as is *SLC28A2* (concentrative nucleoside transporter) the product of which plays in an important role in intestinal nucleoside salvage and energy metabolism (Huber-Ruano et al., 2010). Both genes were also highly expressed in the small intestine of pigs (Freeman et al., 2012) and humans (Genevestigator (Hruz et al., 2008) analysis). *RHBG* (*SLC42A2*), an ammonia transporter, is preferentially expressed in the sheep small and large intestines and the liver (Figure 4) and is a candidate for an intestinal ammonia transporter. However, *RHBG* is not expressed at particularly high levels in the human GIT (Genevestigator (Hruz et al., 2008) analysis) relative to many other tissues, and was not reported to be preferentially expressed in the pig GIT (Freeman et al., 2012). In humans uptake of ammonia in the large intestine is thought to most likely occur (mainly) by passive non-ionic diffusion (Wrong & Vince, 1984). However, *RHCG* (apical membrane) and *RHBG* (basolateral membrane) have also been proposed to constitute an ammonium uptake pathway in the human GIT (Handlogten et al., 2005). The expression profile of *RHCG* in sheep (Figure 4) is not consistent with such a pathway in sheep.

In addition to the secretion of urea into the rumen (a ruminant specific process) the increased importance of nitrogen recycling in ruminants may have led to the apparent increased expression of *RHBG* in the GIT of sheep.

513 Iodine recycling

514 *SLC5A5*, member 5 of solute carrier family 5, encoding a sodium iodide symporter is highly
515 preferentially expressed in the abomasum in our study (Figure 4). *SLC5A5* also has higher
516 expression in human (Genevestigator (Hruz et al. 2008) analysis) and rat stomach (Kotani et al.,
517 1998) than in other digestive tissues. The latter authors reported that the distribution of *SLC5A5*
518 transcripts in the stomach epithelium was consistent with a role of *SLC5A5* in the import or
519 export of iodine, from or to the stomach contents. In the rat, iodine is actively transported into
520 the gastric lumen and this transport is at least partly mediated by a sodium-iodide symporter
521 (Josefsson et al., 2006). In cattle the rate of iodine export by the abomasum epithelium into the
522 abomasum is much greater than the import of iodine from the abomasum (Miller et al., 1975),
523 suggesting that the role of *SLC5A5* in sheep abomasum is to export iodine into the stomach
524 contents. In contrast, *SLC5A5* was not reported to be significantly more expressed in the pig
525 stomach versus other components of the GIT (Freeman et al., 2012). The specific physiological
526 role of iodine in the stomach/GIT is unknown, but a number of possibilities have been suggested:
527 iodine-conserving mechanisms to deal with low iodine concentrations in the diet (Miller et al.,
528 1975), antioxidative activity (Venturi & Venturi, 1999) and antimicrobial activity (Spitzweg et
529 al., 1999). The majority of the secreted iodine is thought to be recovered in the lower intestines.
530 Another member from the same transporter family *SLC5A6*, a sodium/multivitamin and iodide
531 co-transporter (de Carvalho & Quick, 2011), encoded by a gene showing expression in all
532 studied tissues, with the highest expression sheep large intestine (Figure 4) is a likely candidate
533 for the iodine importer. In humans, *SLC5A6* is also expressed in a wide range of tissues with
534 intestinal tissues being close to the top of the list (de Carvalho & Quick, 2011). In pigs, *SLC5A6*
535 is preferentially expressed in the small intestine (Freeman et al., 2012). The high expression of

SLC5A5 in the abomasum suggests that ruminants may have retained a higher dependence on iodine in the GIT than other mammals.

Zinc homeostasis

SLC39A4 encodes a transporter protein essential for zinc uptake in the mouse intestine (Dufner-Beattie et al., 2003) and stomach (Martin et al., 2013). *SLC39A4* is highly expressed in stomach and intestines in sheep (Figure 4) and humans (Genevestigator (Hruz et al., 2008) analysis), and showed the highest expression in pig small intestine (Freeman et al., 2012). Another zinc transporter encoding gene, *SLC39A5*, has a similar expression profile to *SLC39A4* in sheep (Figure 4), humans and pigs. However, *SLC39A5* is located in the basolateral membrane and is involved in the secretion of zinc. In mouse gastrointestinal tract cells the two zinc transporters are reciprocally regulated (Weaver et al., 2007), together controlling the influx and efflux of zinc at the intestinal epithelium. It appears likely that sheep have a similar mechanism for zinc homeostasis to other mammals.

Fluid and electrolyte balance

Maintaining salt and water balance is an important function of the mammalian GIT. In the large intestine significant GO term enrichment was identified for regulation of chloride transport, due to the inclusion of *CA2*, *7* and *CFTR* (Table 1, Figure 4). This is in agreement with the reported critical chloride secretory mechanism in intestinal epithelial cells, associated with mucosal hydration (Barrett & Keely, 2000). *SLC26A3*, which is a $\text{Cl}^-/\text{HCO}_3^-$ antiporter, imports Cl^- ions driven by bicarbonate, thus linking the activity of carbonic anhydrases and the leakage of Cl^- out of the cells by *CFTR*. *SLC26A3* is preferentially expressed in the large intestine of sheep (Figure

4) and the colon of pigs. Thus the expression of genes involved in fluid and electrolyte balance is similar between all three species.

Conclusions

As a significant event in the evolution of the true ruminants, the evolutionary origin of the rumen is the subject of debate, with out-pouching of the oesophagus, or of the stomach, as the two most likely origins (Beck et al., 2009; Langer, 1988). The cornification of the epithelia surface, tissue clustering analysis based on gene expression (driven by the epidermal structural proteins and innate immunity genes) and the relative lack of metabolic overlap with the abomasum strongly favours an oesophageal origin. Metabolically the rumen has many similarities with the liver, especially for SCFA metabolism and even though there are functional similarities with the large intestine, the complements of genes involved are not highly similar.

We have identified a small number of highly rumen specific metabolic processes, in particular the roles of SLC14A1 (urea secretion), SLC4A9 (SCFA uptake) and AKR1C1 (uncertain function). Overall our analysis has enabled gene expression data to be married up with decades of physiological and other research to link transport and enzymatic activities and the most likely genes encoding products with the activities. Nitrogen and iodine recycling have been identified as processes with a much greater importance in the sheep than in humans or pigs. These metabolic functions are protected by strong immune functions and stratified epidermis-like epithelium. The major rumen immune players are DUOX and SERPINB gene families and *DUOXA2*, *DUOX2s* and *SERPINB3/4-like 1* appear to be preferentially expressed in the rumen. These findings will bring novel insights into biomedical research on mammalian digestive and gastrointestinal systems.

579

580 Additional files

581 **Additional file 1: Table S1.** Sheep experimental design information. **Table S2.** Gene expression
582 values with P and FDR <0.01 across tissues. **Table S3.** Assignment of genes to tissue clusters.
583 **Table S4.** Gene members in each significant GO Term enrichment. **Table S5.** Genes overlapped
584 between sheep and pig GIT clusters.

585 **Additional file 2: Figure S1.** Raw output of PCA clustering of tissues. **Figure S2.** Expression
586 profile of GIT-related keratin genes. **Figure S3.** Clustering of transporter genes.

587

588 Abbreviations

589 **GIT: gastrointestinal tissue**

590 Acknowledgements

591 We would like to thank Richard Talbot for supervision of the generation of the RNA-Seq data.

592

593 References

- 594 2012. stat_ellipse <https://github.com/JoFrhwld/FAAV/blob/master/r/stat-ellipse.R>) (accessed
595 03072015.
596 Abdoun K, Stumpff F, and Martens H. 2006. Ammonia and urea transport across the rumen
597 epithelium: a review. *Animal Health Research Reviews* 7:43-59.
598 10.1017/s1466252307001156
599 Abdoun K, Stumpff F, Rabbani I, and Martens H. 2010. Modulation of urea transport across
600 sheep rumen epithelium in vitro by SCFA and CO₂. *Am J Physiol Gastrointest Liver*
601 *Physiol* 298:G190-G202. 10.1152/ajpgi.00216.2009

- 602 Abumrad NA, and Davidson NO. 2012. Role of the gut in lipid homeostasis. *Physiological*
- 603 *Reviews* 92:1061-1085. 10.1152/physrev.00019.2011
- 604 Anders S, Pyl PT, and Huber W. 2015. HTSeq--a Python framework to work with high-
- 605 throughput sequencing data. *Bioinformatics* 31:166-169. 10.1093/bioinformatics/btu638
- 606 Andrews GE, Askey R, and Roy R. 1999. *Special functions - Encyclopedia of Mathematics*.
- 607 Cambridge: Cambridge University Press.
- 608 Aschenbach JR, Bilk S, Tadesse G, Stumpff F, and Gabel G. 2009. Bicarbonate-dependent and
- 609 bicarbonate-independent mechanisms contribute to nondiffusive uptake of acetate in the
- 610 ruminal epithelium of sheep. *American Journal of Physiology Gastrointestinal and Liver*
- 611 *Physiology* 296:G1098-1107. 10.1152/ajpgi.90442.2008
- 612 Baldwin RL, Wu S, Li W, Li C, Bequette BJ, and Li RW. 2012. Quantification of
- 613 Transcriptome Responses of the Rumen Epithelium to Butyrate Infusion using RNA-seq
- 614 Technology. *Gene Regulation and Systems Biology* 6:67-80. 10.4137/grsb.s9687
- 615 Ballatori N, Christian WV, Wheeler SG, and Hammond CL. 2013. The heteromeric organic
- 616 solute transporter, OSTalpha-OSTbeta/SLC51: a transporter for steroid-derived molecules.
- 617 *Molecular Aspects of Medicine* 34:683-692. 10.1016/j.mam.2012.11.005
- 618 Barrett KE, and Keely SJ. 2000. Chloride secretion by the intestinal epithelium: molecular basis
- 619 and regulatory aspects. *Annual Review of Physiology* 62:535-572.
- 620 10.1146/annurev.physiol.62.1.535
- 621 Beck DC, Jiang H, and Zhang L. 2009. Elucidating the Evolutionary Relationships among Bos
- 622 taurus Digestive Organs Using Unigene Expression Data. *International Journal of*
- 623 *Evolutionary Biology*:Article ID 803142. 10.4061/2009/803142
- 624 Bergman EN. 1990. Energy contributions of volatile fatty acids from the gastrointestinal tract in
- 625 various species. *Physiological Reviews* 70:567-590.
- 626 Carter MJ. 1971. The carbonic anhydrase in the rumen epithelial tissue of the ox. *Biochimica et*
- 627 *Biophysica Acta* 235:222-236.
- 628 Cassoni P, Sapino A, Haagenen DE, Naldoni C, and Bussolati G. 1995. Mitogenic effect of the
- 629 15-kDa gross cystic disease fluid protein (GCDFF-15) on breast-cancer cell lines and on
- 630 immortal mammary cells. *International Journal of Cancer* 60:216-220.
- 631 Connor EE, Li RW, Baldwin RL, and Li C. 2010. Gene expression in the digestive tissues of
- 632 ruminants and their relationships with feeding and digestive processes. *animal* 4:993-1007.
- 633 10.1017/s1751731109991285
- 634 Crosnier C, Stamataki D, and Lewis J. 2006. Organizing cell renewal in the intestine: stem cells,
- 635 signals and combinatorial control. *Nature Reviews Genetics* 7:349-359. 10.1038/nrg1840
- 636 de Carvalho FD, and Quick M. 2011. Surprising substrate versatility in SLC5A6: Na⁺-coupled I⁻
- 637 transport by the human Na⁺/multivitamin transporter (hsmvt). *Journal of Biological*
- 638 *Chemistry* 286:131-137. 10.1074/jbc.M110.167197
- 639 Deckardt K, Khol-Parisini A, and Zebeli Q. 2013. Peculiarities of enhancing resistant starch in
- 640 ruminants using chemical methods: opportunities and challenges. *Nutrients* 5:1970-1988.
- 641 10.3390/nu5061970
- 642 Dionissopoulos L, Steele M, AlZahal O, and McBride B. 2012. Adaptation to high grain diets
- 643 proceeds through minimal immune system stimulation and differences in extracellular
- 644 matrix protein expression in a model of subacute ruminal acidosis in nonlactating dairy
- 645 cows. *American Journal of Animal and Veterinary Sciences* 7:84-91.
- 646 Dufner-Beattie J, Wang F, Kuo Y-M, Gitschier J, Eide D, and Andrews GK. 2003. The
- 647 Acrodermatitis Enteropathica Gene ZIP4 Encodes a Tissue-specific, Zinc-regulated Zinc

- Transporter in Mice. *Journal of Biological Chemistry* 278:33474-33481.
10.1074/jbc.M305000200
- Durbin BP, Hardin JS, Hawkins DM, and Rocke DM. 2002. A variance-stabilizing transformation for gene-expression microarray data. *Bioinformatics* 18:S105-S110.
10.1093/bioinformatics/18.suppl_1.S105
- Eckert RL, Sturniolo MT, Broome A-M, Ruse M, and Rorke EA. 2005. Transglutaminase Function in Epidermis. *J Invest Dermatol* 124:481-492.
- Eden E, Navon R, Steinfeld I, Lipson D, and Yakhini Z. 2009. GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* 10:48.
10.1186/1471-2105-10-48
- El-Kabbani O, Dhagat U, and Hara A. 2011. Inhibitors of human 20alpha-hydroxysteroid dehydrogenase (AKR1C1). *Journal of Steroid Biochemistry and Molecular Biology* 125:105-111. 10.1016/j.jsbmb.2010.10.006
- Ensembl. Sheep Genome v3.1 http://www.ensembl.org/Ovis_aries/Info/Index (accessed 03072015).
- Fox JT, Depenbusch BE, Drouillard JS, and Nagaraja TG. 2007. Dry-rolled or steam-flaked grain-based diets and fecal shedding of Escherichia coli O157 in feedlot cattle. *Journal of Animal Science* 85:1207-1212. 10.2527/jas.2006-079
- Freeman T, Ivens A, Baillie JK, Beraldi D, Barnett M, Dorward D, Downing A, Fairbairn L, Kapetanovic R, Raza S, Tomoiu A, Alberio R, Wu C, Su A, Summers K, Tuggle C, Archibald A, and Hume D. 2012. A gene expression atlas of the domestic pig. *BMC Biology* 10:90.
- Fujihara T, and Shem MN. 2011. Metabolism of microbial nitrogen in ruminants with special reference to nucleic acids. *Animal Science Journal* 82:198-208. 10.1111/j.1740-0929.2010.00871.x
- Funk S, Mark R, Bayo P, Flechtenmacher C, Grabe N, Angel P, Plinkert PK, and Hess J. 2015. High S100A8 and S100A12 protein expression is a favorable prognostic factor for survival of oropharyngeal squamous cell carcinoma. *International Journal of Cancer* 136:2037-2046. 10.1002/ijc.29262
- Georgi MI, Rosendahl J, Ernst F, Gunzel D, Aschenbach JR, Martens H, and Stumpff F. 2014. Epithelia of the ovine and bovine forestomach express basolateral maxi-anion channels permeable to the anions of short-chain fatty acids. *Pflügers Archiv European Journal of Physiology* 466:1689-1712. 10.1007/s00424-013-1386-x
- Georgi FM, Del Fabbro C, and Licausi F. 2013. Comparative study of RNA-seq- and Microarray-derived coexpression networks in Arabidopsis thaliana. *Bioinformatics* 29:717-724.
10.1093/bioinformatics/btt053
- Gould J. <http://www.broadinstitute.org/cancer/software/GENE-E/2015>.
- Gressley TF, Hall MB, and Armentano LE. 2011. Ruminant Nutrition Symposium: Productivity, digestion, and health responses to hindgut acidosis in ruminants. *Journal of Animal Science* 89:1120-1130. 10.2527/jas.2010-3460
- Groenen MAM, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, Rothschild MF, Rogel-Gaillard C, Park C, Milan D, Megens H-J, Li S, Larkin DM, Kim H, Frantz LAF, Caccamo M, Ahn H, Aken BL, Anselmo A, Anthon C, Auvin L, Badaoui B, Beattie CW, Bendixen C, Berman D, Blecha F, Blomberg J, Bolund L, Bosse M, Botti S, Bujie Z, Bystrom M, Capitanu B, Carvalho-Silva D, Chardon P, Chen C, Cheng R, Choi S-H, Chow W, Clark RC, Clee C, Crooijmans RPMA, Dawson HD, Dehais P, De Sapio F, Dibbitts B, Drou N, Du

- 694 Z-Q, Eversole K, Fadista J, Fairley S, Faraut T, Faulkner GJ, Fowler KE, Fredholm M, Fritz
695 E, Gilbert JGR, Giuffra E, Gorodkin J, Griffin DK, Harrow JL, Hayward A, Howe K, Hu Z-
696 L, Humphray SJ, Hunt T, Hornshoj H, Jeon J-T, Jern P, Jones M, Jurka J, Kanamori H,
697 Kapetanovic R, Kim J, Kim J-H, Kim K-W, Kim T-H, Larson G, Lee K, Lee K-T, Leggett
698 R, Lewin HA, Li Y, Liu W, Loveland JE, Lu Y, Lunney JK, Ma J, Madsen O, Mann K,
699 Matthews L, McLaren S, Morozumi T, Murtaugh MP, Narayan J, Truong Nguyen D, Ni P,
700 Oh S-J, Onteru S, Panitz F, Park E-W, Park H-S, Pascal G, Paudel Y, Perez-Enciso M,
701 Ramirez-Gonzalez R, Reecy JM, Rodriguez-Zas S, Rohrer GA, Rund L, Sang Y,
702 Schachtschneider K, Schraiber JG, Schwartz J, Scobie L, Scott C, Searle S, Servin B,
703 Southey BR, Sperber G, Stadler P, Sweedler JV, Tafer H, Thomsen B, Wali R, Wang J,
704 Wang J, White S, Xu X, Yerle M, Zhang G, Zhang J, Zhang J, Zhao S, Rogers J, Churcher
705 C, and Schook LB. 2012. Analyses of pig genomes provide insight into porcine demography
706 and evolution. *Nature* 491:393-398.
707 [http://www.nature.com/nature/journal/v491/n7424/abs/nature11622.html#supplementary-](http://www.nature.com/nature/journal/v491/n7424/abs/nature11622.html#supplementary-information)
708 [information](http://www.nature.com/nature/journal/v491/n7424/abs/nature11622.html#supplementary-information)
709 Handlogten ME, Hong SP, Zhang L, Vander AW, Steinbaum ML, Campbell-Thompson M, and
710 Weiner ID. 2005. Expression of the ammonia transporter proteins Rh B glycoprotein and Rh
711 C glycoprotein in the intestinal tract. *American Journal of Physiology Gastrointestinal and*
712 *Liver Physiology* 288:G1036-1047. 10.1152/ajpgi.00418.2004
713 Hassan MI, Waheed A, Yadav S, Singh TP, and Ahmad F. 2009. Prolactin inducible protein in
714 cancer, fertility and immunoregulation: structure, function and its clinical implications.
715 *Cellular and Molecular Life Sciences* 66:447-459. 10.1007/s00018-008-8463-x
716 Heger A, and Holm L. 2000. Rapid automatic detection and alignment of repeats in protein
717 sequences. *Proteins* 41:224-237.
718 Hofmann R. 1989. Evolutionary steps of ecophysiological adaptation and diversification of
719 ruminants: a comparative view of their digestive system. *Oecologia* 78:443-457.
720 Hoover WH. 1978. Digestion and absorption in the hindgut of ruminants. *Journal of Animal*
721 *Science* 46:1789-1799.
722 Hovnanian A. 2013. Netherton syndrome: skin inflammation and allergy by loss of protease
723 inhibition. *Cell & Tissue Research* 351:289-300. 10.1007/s00441-013-1558-1
724 Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, and
725 Zimmermann P. 2008. Genevestigator v3: a reference expression database for the meta-
726 analysis of transcriptomes. *Adv Bioinformatics* 2008:420747. 10.1155/2008/420747
727 Huber-Ruano I, Pinilla-Macua I, Torres G, Casado FJ, and Pastor-Anglada M. 2010. Link
728 between high-affinity adenosine concentrative nucleoside transporter-2 (CNT2) and energy
729 metabolism in intestinal and liver parenchymal cells. *Journal of Cellular Physiology*
730 225:620-630. 10.1002/jcp.22254
731 Hudson N, Dalrymple B, and Reverter A. 2012. Beyond differential expression: the quest for
732 causal mutations and effector molecules. *BMC Genomics* 13:356.
733 Huttenhower C, Haley EM, Hibbs MA, Dumeaux V, Barrett DR, Collier HA, and Troyanskaya
734 OG. 2009. Exploring the human genome with functional maps. *Genome Research* 19:1093-
735 1106. 10.1101/gr.082214.108
736 Ingle DL, Bauman DE, and Garrigus US. 1972. Lipogenesis in the Ruminant: in vivo Site of
737 Fatty Acid Synthesis in Sheep. *Journal of Nutrition* 102:617-623.

- Jenkins TC, Wallace RJ, Moate PJ, and Mosley EE. 2008. Recent advances in biohydrogenation of unsaturated fatty acids within the rumen microbial ecosystem. *Journal of Animal Science* 86:397-412. 10.2527/jas.2007-0588
- Jiang Y, Xie M, Chen W, Talbot R, Maddox JF, Faraut T, Wu C, Muzny DM, Li Y, Zhang W, Stanton J-A, Brauning R, Barris WC, Hourlier T, Aken BL, Searle SMJ, Adelson DL, Bian C, Cam GR, Chen Y, Cheng S, DeSilva U, Dixon K, Dong Y, Fan G, Franklin IR, Fu S, Fuentes-Utrilla P, Guan R, Highland MA, Holder ME, Huang G, Ingham AB, Jhangiani SN, Kalra D, Kovar CL, Lee SL, Liu W, Liu X, Lu C, Lv T, Mathew T, McWilliam S, Menzies M, Pan S, Robelin D, Servin B, Townley D, Wang W, Wei B, White SN, Yang X, Ye C, Yue Y, Zeng P, Zhou Q, Hansen JB, Kristiansen K, Gibbs RA, Flicek P, Warkup CC, Jones HE, Oddy VH, Nicholas FW, McEwan JC, Kijas JW, Wang J, Worley KC, Archibald AL, Cockett N, Xu X, Wang W, and Dalrymple BP. 2014. The sheep genome illuminates biology of the rumen and lipid metabolism. *Science* 344:1168-1173. 10.1126/science.1252806
- Josefsson M, Evilevitch L, Westrom B, Grunditz T, and Ekblad E. 2006. Sodium-iodide symporter mediates iodide secretion in rat gastric mucosa in vitro. *Experimental Biology and Medicine (Maywood, NJ)* 231:277-281.
- Kato D, Suzuki Y, Haga S, So K, Yamauchi E, Nakano M, Ishizaki H, Choi K, Katoh K, and Roh S-G. 2015. Utilization of digital differential display to identify differentially expressed genes related to rumen development. *Animal Science Journal*:n/a-n/a. 10.1111/asj.12448
- Kivel AJ, Kivel J, Saarnio J, and Parkkila S. 2005. Carbonic anhydrases in normal gastrointestinal tract and gastrointestinal tumours. *World Journal of Gastroenterology* 11:155-163.
- Kotani T, Ogata Y, Yamamoto I, Aratake Y, Kawano JI, Suganuma T, and Ohtaki S. 1998. Characterization of gastric Na⁺/I⁻ symporter of the rat. *Clinical Immunology and Immunopathology* 89:271-278.
- Kypriotou M, Huber M, and Hohl D. 2012. The human epidermal differentiation complex: cornified envelope precursors, S100 proteins and the 'fused genes' family. *Experimental Dermatology* 21:643-649. 10.1111/j.1600-0625.2012.01472.x
- Lane MA, Baldwin RL, and Jesse BW. 2002. Developmental changes in ketogenic enzyme gene expression during sheep rumen development. *Journal of Animal Science* 80:1538-1544.
- Langer P. 1988. *The mammalian herbivore stomach: comparative anatomy, function and evolution*: Gustav Fischer.
- Lapierre H, and Lobley GE. 2001. Nitrogen Recycling in the Ruminant: A Review. *Journal of Dairy Science* 84:E223-E236. 10.3168/jds.S0022-0302(01)70222-6
- Lees MS, H. Nagaraj S, Piedrafita DM, Kotze AC, and Ingham AB. 2012. Molecular cloning and characterisation of ovine dual oxidase 2. *Gene* 500:40-46.
- Li Y, Carrillo JA, Ding Y, He Y, Zhao C, Zan L, and Song J. 2015. Ruminant Transcriptomic Analysis of Grass-Fed and Grain-Fed Angus Beef Cattle. *PLoS ONE* 10:e0116437. 10.1371/journal.pone.0116437
- Love M, Anders S, and Huber W. 2013. Differential analysis of count data—the DESeq2 package.
- Martin AB, Aydemir TB, Guthrie GJ, Samuelson DA, Chang SM, and Cousins RJ. 2013. Gastric and colonic zinc transporter ZIP11 (SLC39A11) in mice responds to dietary zinc and exhibits nuclear localization. *Journal of Nutrition* 143:1882-1888. 10.3945/jn.113.184457

- 782 Mbanzamihigo L, van Nevel CJ, and Demeyer DI. 1996. Lasting effects of monensin on rumen
783 and caecal fermentation in sheep fed a high grain diet. *Animal Feed Science and Technology*
784 62:215-228.
- 785 McGrath JA, Bolling MC, and Jonkman MF. 2010. Lethal Acantholytic Epidermolysis Bullosa.
786 *Dermatologic Clinics* 28:131-135.
- 787 Miller JK, Swanson EW, and Spalding GE. 1975. Iodine Absorption, Excretion, Recycling, and
788 Tissue Distribution in the Dairy Cow. *Journal of Dairy Science* 58:1578-1593.
789 10.3168/jds.S0022-0302(75)84753-9
- 790 Milo R, Jorgensen P, Moran U, Weber G, and Springer M. 2010. BioNumbers--the database of
791 key numbers in molecular and cell biology. *Nucleic Acids Research* 38:D750-753.
792 10.1093/nar/gkp889
- 793 Naderi A, and Vanneste M. 2014. Prolactin-induced protein is required for cell cycle progression
794 in breast cancer. *Neoplasia* 16:329-342 e321-314. 10.1016/j.neo.2014.04.001
- 795 Nichols BL, Avery S, Sen P, Swallow DM, Hahn D, and Sterchi E. 2003. The maltase-
796 glucoamylase gene: common ancestry to sucrase-isomaltase with complementary starch
797 digestion activities. *Proc Natl Acad Sci U S A* 100:1432-1437. 10.1073/pnas.0237170100
- 798 Nichols BL, Eldering J, Avery S, Hahn D, Quaroni A, and Sterchi E. 1998. Human small
799 intestinal maltase-glucoamylase cDNA cloning. Homology to sucrase-isomaltase. *Journal of*
800 *Biological Chemistry* 273:3076-3081.
- 801 Ohana E, Shcheynikov N, Park M, and Muallem S. 2012. Solute carrier family 26 member a2
802 (SLC26A2) protein functions as an electroneutral SOFormula/OH-/Cl- exchanger regulated
803 by extracellular Cl. *Journal of Biological Chemistry* 287:5122-5132.
804 10.1074/jbc.M111.297192
- 805 Parkkila S, Parkkila AK, Juvonen T, and Rajaniemi H. 1994. Distribution of the carbonic
806 anhydrase isoenzymes I, II, and VI in the human alimentary tract. *Gut* 35:646-650.
807 10.1136/gut.35.5.646
- 808 Penner GB, Steele MA, Aschenbach JR, and McBride BW. 2011. Ruminant Nutrition
809 Symposium: Molecular adaptation of ruminal epithelia to highly fermentable diets. *Journal*
810 *of Animal Science* 89:1108-1119. 10.2527/jas.2010-3378
- 811 Penning TM. 1997. Molecular endocrinology of hydroxysteroid dehydrogenases. *Endocrine*
812 *Reviews* 18:281-305. 10.1210/edrv.18.3.0302
- 813 Reverter A, and Chan EKF. 2008. Combining partial correlation and an information theory
814 approach to the reversed engineering of gene co-expression networks. *Bioinformatics*
815 24:2491-2497. 10.1093/bioinformatics/btn482
- 816 Robinson MD, McCarthy DJ, and Smyth GK. 2010. edgeR: a Bioconductor package for
817 differential expression analysis of digital gene expression data. *Bioinformatics* 26:139-140.
818 10.1093/bioinformatics/btp616
- 819 Roder PV, Geillinger KE, Zietek TS, Thorens B, Koepsell H, and Daniel H. 2014. The role of
820 SGLT1 and GLUT2 in intestinal glucose transport and sensing. *PLoS ONE* 9:e89977.
821 10.1371/journal.pone.0089977
- 822 Rojen BA, Poulsen SB, Theil PK, Fenton RA, and Kristensen NB. 2011. Short communication:
823 Effects of dietary nitrogen concentration on messenger RNA expression and protein
824 abundance of urea transporter-B and aquaporins in ruminal papillae from lactating Holstein
825 cows. *Journal of Dairy Science* 94:2587-2591. 10.3168/jds.2010-4073

- 826 Rost D, Mahner S, Sugiyama Y, and Stremmel W. 2002. Expression and localization of the
827 multidrug resistance-associated protein 3 in rat small and large intestine. *Am J Physiol*
828 *Gastrointest Liver Physiol* 282:G720-G726. 10.1152/ajpgi.00318.2001
- 829 Scocco P, Mercati F, Brusafferro A, Ceccarelli P, Belardinelli C, and Malfatti A. 2013.
830 Keratinisation degree of rumen epithelium and body condition score in sheep grazing on
831 *Brachypodium rupestre*. *Veterinaria Italiana* 49:211-217.
- 832 Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, and
833 Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular
834 interaction networks. *Genome Research* 13:2498-2504. 10.1101/gr.1239303
- 835 Sivaprasad U, Askew DJ, Ericksen MB, Gibson AM, Stier MT, Brandt EB, Bass SA, Daines
836 MO, Chakir J, Stringer KF, Wert SE, Whitsett JA, Le Cras TD, Wills-Karp M, Silverman
837 GA, and Khurana Hershey GK. 2011. A nonredundant role for mouse SERPINB3A in the
838 induction of mucus production in asthma. *Journal of Allergy and Clinical Immunology*
839 127:254-261, 261.e251-256. 10.1016/j.jaci.2010.10.009
- 840 Spitzweg C, Joba W, Schriever K, Goellner JR, Morris JC, and Heufelder AE. 1999. Analysis of
841 human sodium iodide symporter immunoreactivity in human exocrine glands. *Journal of*
842 *Clinical Endocrinology and Metabolism* 84:4178-4184. 10.1210/jcem.84.11.6117
- 843 Steele MA, Croom J, Kahler M, AlZahal O, Hook SE, Plaizier K, and McBride BW. 2011a.
844 Bovine rumen epithelium undergoes rapid structural adaptations during grain-induced
845 subacute ruminal acidosis. *American Journal of Physiology Regulatory Integrative and*
846 *Comparative Physiology* 300:R1515-1523. 10.1152/ajpregu.00120.2010
- 847 Steele MA, Dionissopoulos L, AlZahal O, Doelman J, and McBride BW. 2012. Rumen epithelial
848 adaptation to ruminal acidosis in lactating cattle involves the coordinated expression of
849 insulin-like growth factor-binding proteins and a cholesterolgenic enzyme. *Journal of Dairy*
850 *Science* 95:318-327. 10.3168/jds.2011-4465
- 851 Steele MA, Vandervoort G, AlZahal O, Hook SE, Matthews JC, and McBride BW. 2011b.
852 Rumen epithelial adaptation to high-grain diets involves the coordinated regulation of genes
853 involved in cholesterol homeostasis. *Physiological Genomics* 43:308-316.
854 10.1152/physiolgenomics.00117.2010
- 855 Steentoft C, Vakhrushev SY, Joshi HJ, Kong Y, Vester-Christensen MB, Schjoldager KT,
856 Lavrsen K, Dabelsteen S, Pedersen NB, Marcos-Silva L, Gupta R, Bennett EP, Mandel U,
857 Brunak S, Wandall HH, Levery SB, and Clausen H. 2013. Precision mapping of the human
858 O-GalNAc glycoproteome through SimpleCell technology. *EMBO Journal* 32:1478-1488.
859 10.1038/emboj.2013.79
- 860 Su G, Kuchinsky A, Morris JH, States DJ, and Meng F. 2010. GLay: community structure
861 analysis of biological networks. *Bioinformatics* 26:3135-3137.
862 10.1093/bioinformatics/btq596
- 863 Tashian RE. 1989. The carbonic anhydrases: Widening perspectives on their evolution,
864 expression and function. *Bioessays* 10:186-192. 10.1002/bies.950100603
- 865 Tong L, Lan W, Lim RR, and Chaurasia SS. 2014. S100A proteins as molecular targets in the
866 ocular surface inflammatory diseases. *Ocul Surf* 12:23-31. 10.1016/j.jtos.2013.10.001
- 867 van Hasselt PM, Ferdinandusse S, Monroe GR, Ruiter JP, Turkenburg M, Geerlings MJ, Duran
868 K, Harakalova M, van der Zwaag B, Monavari AA, Okur I, Sharrard MJ, Cleary M,
869 O'Connell N, Walker V, Rubio-Gozalbo ME, de Vries MC, Visser G, Houwen RH, van der
870 Smagt JJ, Verhoeven-Duif NM, Wanders RJ, and van Haaften G. 2014. Monocarboxylate

- transporter 1 deficiency and ketone utilization. *New England Journal of Medicine* 371:1900-1907. 10.1056/NEJMoa1407778
- van Miert AS, and van Duin CT. 1991. Feed intake and rumen motility in dwarf goats. Effects of some alpha 2-adrenergic agonists, prostaglandins and posterior pituitary hormones. *Veterinary Research Communications* 15:57-67.
- Van Nevel CJ, and Demeyer DI. 1996. Influence of pH on lipolysis and biohydrogenation of soybean oil by rumen contents in vitro. *Reproduction Nutrition Development* 36:53-63.
- Veenendaal GH, Nijnanten FMAW-V, Duin CTMVAN, and Miert ASJPAMV. 1980. Role of circulating prostaglandins in the genesis of pyrogen (endotoxin)-induced ruminal stasis in conscious goats. *Journal of Veterinary Pharmacology and Therapeutics* 3:59-68. 10.1111/j.1365-2885.1980.tb00409.x
- Venturi S, and Venturi M. 1999. Iodide, thyroid and stomach carcinogenesis: evolutionary story of a primitive antioxidant? *European Journal of Endocrinology* 140:371-372. 10.1530/eje.0.1400371
- Wang G, Xu Z, Wang R, Al-Hijji M, Salit J, Strulovici-Barel Y, Tilley A, Mezey J, and Crystal R. 2012. Genes associated with MUC5AC expression in small airway epithelium of human smokers and non-smokers. *BMC Medical Genomics* 5:21.
- Wang J-H, Zhao L-F, Lin P, Su X-R, Chen S-J, Huang L-Q, Wang H-F, Zhang H, Hu Z-F, Yao K-T, and Huang Z-X. 2014. GenCLiP 2.0: a web server for functional clustering of genes and construction of molecular networks based on free terms. *Bioinformatics* 30:2534-2536. 10.1093/bioinformatics/btu241
- Wang LQ, Baldwin RL, and Jesse BW. 1996. Isolation and characterization of a cDNA clone encoding ovine type I carbonic anhydrase. *Journal of Animal Science* 74:345-353.
- Wang TY, Liu M, Portincasa P, and Wang DQ. 2013. New insights into the molecular mechanism of intestinal fatty acid absorption. *European Journal of Clinical Investigation* 43:1203-1223. 10.1111/eci.12161
- Watson-Haigh NS, Kadarmideen HN, and Reverter A. 2010. PCIT: an R package for weighted gene co-expression networks based on partial correlation and information theory approaches. *Bioinformatics* 26:411-413. 10.1093/bioinformatics/btp674
- Weaver BP, Dufner-Beattie J, Kambe T, and Andrews GK. 2007. Novel zinc-responsive post-transcriptional mechanisms reciprocally regulate expression of the mouse SLC39A4 and SLC39A5 zinc transporters (Zip4 and Zip5). *Biological Chemistry* 388:1301-1312. 10.1515/bc.2007.149
- Wrong OM, and Vince A. 1984. Urea and ammonia metabolism in the human large intestine. *Proceedings of the Nutrition Society* 43:77-86.
- Young JW. 1977. Gluconeogenesis in Cattle: Significance and Methodology. *Journal of Dairy Science* 60:1-15. 10.3168/jds.S0022-0302(77)83821-6
- Yu Z, Wilderboth JE, Wallace OAM, Gordon SW, Maqbool NJ, Maclean PH, Nixon AJ, and Pearson AJ. 2011. Annotation of sheep keratin intermediate filament genes and their patterns of expression. *Experimental Dermatology* 20:582-588. 10.1111/j.1600-0625.2011.01274.x

Table 1(on next page)

Gene Ontology enrichments of clusters

1 Table 1 Gene Ontology enrichments of clusters

Cluster	GO-term	FDR corrected <i>P</i> -value ¹
Rumen	EDC locus ²	7.1E-13 ³
Epithelia-rumen-tonsil	EDC locus ²	8.6E-15 ³
	Defense response to fungus	8.6E-03
Epithelia-rumen bias	Keratinization	2.4E-04
Epithelia-all	-	-
Epithelia-large intestine	Desmosome organization	4.7E-03
Epithelia-GI-liver	Cell junction organization	6.3E-03
Abomasum-intestine	-	-
Intestine-low in rectum	-	-
Large intestine	Regulation of chloride transport	4.5E-05
Intestine	-	-
Epithelia-intestine	Cell cycle process	1.4E-46
Abomasum	Digestion	3.8E-02
Small intestine	-	-
Rumen-abomasum	Platelet aggregation	2.2E-04
Rumen-intestine-liver	Flavonoid biosynthetic process	5.5E-10
Intestine-spleen	Humoral immune response	4.5E-02

¹Top significantly enriched pathway selected from GORILLA analysis (see methods) for each input gene cluster

²Genes in the EDC locus of the sheep genome.

³Enrichment of EDC locus genes was calculated using the hypergeometric distribution.

Table 2 (on next page)

Representation of the pig GIT gene clusters in the sheep GIT network

1 **Table 2 Representation of the pig GIT gene clusters in the sheep GIT network**

Pig cluster ¹	Pig tissues ¹	Pig cell type of origin ²	Overlap	<i>P</i> -value ²	Representation	Sheep tissues	Go term enrichment	<i>P</i> -value ³
Overall			179	8.1E-31	Over		Cell cycle process	2.0E-13
1, 7	Intestine -	Immune cells/cell cycle	58	2.4E-11	Over	Epithelia, intestine	Cell cycle process	1.5E-33
3, 8	Tongue-oesophagus	Stratified squamous epithelia	73	1.3E-34	Over	Rumen, epithelia, abomasum, large intestine	Epidermis development	2.9E-05
2, 4, 9	Oesophagus-stomach	Muscle	9	0.0002 ⁴	Under ⁴	Rumen, abomasum	na	
6, 13, 15	Salivary gland	Stratified columnar epithelia	4	0.1777	None		na	
5, 12, 14, 16	Stomach-intestine	Ciliate/glandular epithelia	35	5.4E-09	Over	Stomach intestine	na	
10	Stomach	Neuronal	0	na	na		na	

2 ¹Numbers, names and grouping of pig gene clusters by cell type of origin are according to (Freeman et al. 2012).

3 ²Calculated hypogeometric *P* values, representing the significance of representation of pig genes in sheep gene network.

4 ³FDR corrected GO term enrichment *P* values.

5 ⁴If overlap with just the rumen and rumen-abomasum clusters, significant (*P*=8E-05) over representation

6

Figure 1(on next page)

Transcriptomic sample clustering.

Each dot represents one tissue sample from a single animal. Circles indicate significant clusters (confidence interval = 95%). Raw PCA plots are available (Additional file 2: Figure S1).





Figure 2(on next page)

Gene co-expression network.

(A) Each dot represents a sheep transcript and different colors represent the tissue(s) where the transcript showed high expression, compared to the other tissues. Rectum↓: low in rectum. (B) The same gene co-expression network with only the orthologous genes present in specific pig GIT clusters (Freeman et al. 2012) highlighted (Additional file 1). The names and colors of pig cluster were determined according to the tissues where genes showed the highest and the second highest expression level in the pig GI gene network (Freeman et al. 2012) .



B




Figure 3 (on next page)

Expression profiles of innate immunity and epithelial development genes in sheep.

Data are presented with $\log_2$ Fragments Per Kilobase of exon per Million fragments mapped (FPKM) values along with the subcellular locations and/or tissues of pig (Freeman et al. 2012) and human (Genevestigator (Hruz et al. 2008) analysis) where these genes showed high expression. Cellular location information were derived from GENATLAS database (Frezal 1998) .



PeerJ PrePrints | <https://doi.org/10.7287/peerj.preprints.1766v1> | CC-BY 4.0 Open Access | rec: 21 Feb 2016, publ: 21 Feb 2016

Figure 4(on next page)

Gene expression profiles of metabolic processes discussed in the text.

Data are presented with $\log_2$ Fragments Per Kilobase of exon per Million fragments mapped (FPKM) values along with the subcellular locations and/or tissues of pig (Freeman et al. 2012) and human (Genevestigator (Hruz et al. 2008) analysis) where these genes showed high expression. Texts and bars on the left side of the heatmap indicate involved pathways for covered genes described in the article. Cellular location information were derived from GENATLAS database (Frezal 1998).




Figure 5(on next page)

Ruminant ketone body metabolism pathways.

Key enzyme encoding genes (red text) and pathways (black arrow) are highlighted.





Figure 6(on next page)

Organization of the MGAM2 carboxy-terminus.

Consensus motifs of the serine/threonine rich 40 amino acid repeats at the carboxy-terminus of predicted MGAM-like proteins. **(A)** sheep. **(B)** cattle. **(C)** Pig. **(D)** human.


A Sheep MGAM-like protein amino acid repeats



B Cattle MGAM-like protein amino acid repeats



C Pig MGAM-like protein amino acid repeats



D Human MGAM-like protein amino acid repeats

