

A peer-reviewed version of this preprint was published in PeerJ on 27 March 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.328) (peerj.com/articles/328), which is the preferred citable publication unless you specifically need to cite this preprint.

Bartomeus I, Potts SG, Steffan-Dewenter I, Vaissière BE, Woyciechowski M, Krewenka KM, Tscheulin T, Roberts SPM, Szentgyörgyi H, Westphal C, Bommarco R. 2014. Contribution of insect pollinators to crop yield and quality varies with agricultural intensification. PeerJ 2:e328
<https://doi.org/10.7717/peerj.328>

Contribution of insect pollinators to crop yield and quality varies with agricultural intensification

I. Bartomeus^{a,1}, S.G. Potts^b, I. Steffan-Dewenter^c, B. E. Vaissière^d, M. Woyciechowski^e, K. M. Krewenka^f, T. Tscheulin^{b,g}, S. P. M. Roberts^b, H. Szentgyörgyi^e, C. Westphal^f, R. Bommarco^a.

Affiliations:

^a Swedish University of Agricultural Sciences, Department of Ecology, Uppsala, Sweden.

^b University of Reading, School of Agriculture, Policy and Development, Reading, RG6 6AR, UK

^c University of Würzburg, Department of Animal Ecology and Tropical Biology, Biocentre, 97074 Würzburg, Germany

^d INRA, UR406 Abeilles & Environnement, 84914 Avignon, France

^e Jagiellonian University, Institute of Environmental Sciences, Gronostajowa 7, 30-387 Krakow, Poland.

^f Georg-August-University, Department of Crop Science, Agroecology, Grisebachstr. 6, 37077 Göttingen, Germany

^g University of the Aegean, Department of Geography, 81100 Mytilene, Greece

¹ Corresponding author: Ignasi Bartomeus; nacho.bartomeus@gmail.com; @ibartomeus

Running title: Pollinator contribution to crop yield

Keywords: agroecosystems, biodiversity, ecosystem services, honeybees, wild bees, pollination.

Data deposition: species richness and abundance of flower visiting insects, and mean crop yield values per site will be deposited in figshare:

<http://figshare.com/authors/Ignasi%20bartomeus/97352>

25 **Funding:** This work was funded by the European Union to the projects “ALARM - Assessing
26 large-scale environmental risks for biodiversity with tested methods” (2003-506675), and
27 “STEP – Status and trends of European pollinators” (244090) in the 6th and 7th Framework
28 Programme, respectively, and by the Swedish research council FORMAS.

29

Abstract

Background. Up to 75 % of crop species benefit at least to some degree from animal pollination for fruit or seed set and yield. However, basic information on the level of pollinator dependence and pollinator contribution to yield is lacking for many crops. Even less is known about how insect pollination affects crop quality. Given that habitat loss and agricultural intensification are known to decrease pollinator richness and abundance, there is a need to assess the consequences for different components of crop production.

Methods. We used pollination exclusion on flowers or inflorescences on a whole plant basis to assess the contribution of insect pollination to crop yield and quality in four flowering crops (spring oilseed rape, field bean, strawberry, and buckwheat) located in four regions of Europe. For each crop, we recorded abundance and species richness of flower visiting insects in ten fields located along a gradient from simple to heterogeneous landscapes.

Results. Insect pollination enhanced average crop yield between 18 and 71% depending on the crop. Yield quality was also enhanced in most crops. For instance, oilseed rape had higher oil and lower chlorophyll contents when adequately pollinated, the proportion of empty seeds decreased in buckwheat, and strawberries' commercial grade improved; however, we did not find higher nitrogen content in open pollinated field beans. Complex landscapes had a higher overall species richness of wild pollinators across crops, but visitation rates were only higher in complex landscapes for some crops. On the contrary, the overall yield was consistently enhanced by higher visitation rates, but not by higher pollinator richness.

Discussion. For the four crops in this study, there is clear benefit delivered by pollinators on yield quantity and/or quality, but it is not maximized under current agricultural intensification. Honeybees, the most abundant pollinator, might partially compensate the

54 loss of wild pollinators in some areas, but our results suggest the need of landscape-scale
55 actions to enhance wild pollinator populations.

Introduction

There is growing evidence that ecosystem services, such as biological pest control and crop pollination, benefit food production (Bommarco et al. 2013). Indeed, 75% of the crop species used for food depend on insect pollination to some degree (Klein et al. 2007). More than a decade of active pollination research has led to a greatly improved general understanding on animal pollination benefits to crop yields worldwide (e.g., Klein et al. 2007; Garibaldi et al 2011, 2013). However, major knowledge gaps remain.

First, we have surprisingly little information on the actual degree of pollinator dependence for some major crops. While some crops depend entirely on insect pollinator visits to set fruit, many others are only partly dependent on animal pollination and can produce more than 90% of the maximum seed or fruit yield without pollinators (Klein et al. 2007). The role of pollinators for crop production has mainly been examined in observational studies, relying primarily on natural variation in visitation rates among observed sites. Experiments directly manipulating insect flower visitation (e.g., excluded pollinators vs. open access of pollinators) are less common for most crops (but see Klein et al. 2003, Höhn et al. 2008). Assessing pollination dependence with proper controls is needed to correctly estimate the contribution that insect pollinators can provide to crop yields.

Second, most available studies quantify the number of fruits per plant. Fruit number can be a good proxy for yield (Garibaldi et al. 2013), which is the amount of produce harvested per unit area. However, the correlation between the number of fruit produced and yield is likely to be low in some crops. For example, interspecific plant competition can lead to high variability in plant size and thereby fruit production among plants. This is especially critical

for crops with indeterminate flowering and a high compensation capacity such as soybean (*Glycine max*) and oilseed rape (*Brassica napus*). For these, fruit set measured on a limited number of isolated plants is unlikely to be representative of the real production in a crop stand (Stivers & Swearingin 1980, Angadi et al. 2003). Moreover, plants can allocate resources for producing fruits of variable size based on the number of fruits per plant and the level of pollination received (e.g., Gonzalez et al. 1998 in kiwifruit *Actinidia deliciosa*), such that similar levels of fruit set can differ in total crop yield because of difference in fruit size (Bos et al. 2007). Again, the use of proper control plants from which pollinators are excluded is a way to better estimate the actual contribution of pollinators to yield in such crops.

Quality is also important in crop production, especially from an economic standpoint. Fruit quality can be negatively correlated with quantity when the fruit load on a tree or a vine is too high (e.g., Ferguson & Watkins 1992 in apple *Malus x domestica*), but it is not so otherwise, especially in crops with indeterminate flowering such as oilseed rape (Bommarco et al. 2012). Indeed, adequate pollination often leads to produce with enhanced quality in entomophilous crops such as orchard fruit production (e.g., in apple – Sheffield et al. 2005, Garratt et al. 2014), as well as in field crops (oilseed rape – Bommarco et al. 2012b) and small fruits and vegetables (e.g., strawberry *Fragaria x ananassa* – Andersson et al. 2012, Chagnon et al. 1993, Roselino et al. 2009; tomato *Solanum lycopersicum* – Hogendoorn et al. 2010; bell peppers *Capsicum annuum* – Roldan Serrano and Guerra-Sanz 2006 ; highbush blueberry *Vaccinium corymbosum* – Isaacs and Kirk 2010).

Given the drastic shifts in community composition of insects that visit flowering crops

104 (Winfree et al. 2011, Bommarco et al. 2011, Bartomeus et al. 2013a), and declines in
 105 pollinator species numbers observed in certain regions (Potts et al. 2010, Carvalheiro et al.
 106 2013), it is increasingly important to gather information on the extent to which different
 107 crops depend on insect pollination for yield, and if current pollinator communities fulfill the
 108 demand for pollination services such that both crop quality and yields are maximized
 109 (Breeze et al. 2011). Relationships between land use intensity, pollinator visitation, and fruit
 110 set have been well studied. While pollinator species richness consistently and drastically
 111 decays as agricultural landscapes are deprived of natural habitat and are more intensively
 112 cultivated (Kennedy et al. 2013), this relationship is much weaker for fruit set (Garibaldi et
 113 al. 2011, Chacoff et al. 2008, Ricketts et al. 2008). One explanation for this difference is that
 114 the remaining pollinators provide sufficient visitation even in monotonous, intensively
 115 cultivated landscapes, especially if the crop has a large degree of autonomous self-
 116 pollination. Moreover, intensive landscapes are characterized by harboring just a few
 117 generalist pollinator species (Bartomeus & Winfree 2013), but these might be in sufficient
 118 numbers to deliver enough crop pollination services. In fact, not all pollinator species
 119 respond equally to land use change (Williams et al. 2010, Winfree et al. 2011), and some
 120 even increase in abundance with agricultural intensification (Westphal et al 2003, Carré et
 121 al. 2009). This diversity of responses can in some cropping systems buffer a loss of
 122 pollination function (Cariveau et al. 2013), especially if the pollinators who are the main
 123 ecosystem service providers are adapted to the ephemeral and spatially disassociated
 124 resource distribution that is typical for agricultural landscapes. Moreover, although wild
 125 insects increase fruit set independently of honeybee visits (Garibaldi et al. 2013), honeybees
 126 are less dependent on landscape characteristics because they are mainly managed,
 127 particularly in North America and Europe, and can be moved around in the landscape.

Hence, honeybees can also help mitigate against wild pollinator loss in more intensively used landscapes where pollination services are degraded. In any case, the composition of the landscape in which the flowering crop field is embedded emerges as an important driver for pollinator community composition, and the landscape context needs to be considered when linking land use to pollination provisioning and benefits in field crops.

Here we used pollinator exclusion on the flowers or inflorescence on a whole plant basis in a set of crops under standard field conditions, to obtain information on pollinator dependency for four economically important annual crops in Europe. We assessed pollinator contribution to both yield quantity and quality. By replicating this experiment along a landscape gradient for each crop, we were able to assess changes in pollinator community composition and visitation rate following landscape level land use change and its consequences for crop pollination services and production.

Material and Methods

Study sites

The fieldwork was conducted in four European countries during May–August 2005 (Table 1). Spring oilseed rape was assessed in the region around the city of Uppsala, Sweden (see Bommarco et al. 2012 for details); field bean (*Vicia faba*) in around Reading, UK, strawberry around Göttingen, Germany; and buckwheat (*Fagopyrum esculentum*) near Krakow, Poland. For each crop, we selected ten fields that were separated by a minimum distance of 3 km, corresponding to the maximum foraging range of most bees (Greenleaf et al. 2007). Within each field, we established a 50 * 25 m study area (5*150 m for buckwheat as the fields were

long and narrow) with a homogeneous and continuous crop cover. For fields up to two ha in size, this study site was located in the middle of the field. For larger fields, it was located between the geometric center of the field and one of its margins (Vaissière et al. 2011).

Insect sampling

In each field, we assessed the abundance and species richness of the major groups of flower-visiting insects, including bees (Hymenoptera: Apoidea: Apiformes), hoverflies (Diptera: Syrphidae), and butterflies (Lepidoptera). We used standardized transect walks with an aerial net (Westphal et al. 2008). In each study site, a 150 m transect line was established in the field near the experimental plots. An observer walked this line for 30 min identifying visiting insects at species level and catching unidentified species within a corridor 4 m wide. We performed the transect walks between 0900 and 1700 hours only on days with temperatures at or above 15°C, with no precipitation, dry vegetation, and low windspeed ($<40 \text{ km.h}^{-1}$; Westphal et al. 2008). Specimens were pinned, labeled, and subsequently identified to species level. We returned four times to each study site during the main flowering period of each study crop.

Experimental design and yield analysis

In each of the ten fields, we established a block experiment with four blocks. Each block had two treatments with one plot per treatment and five to ten tagged contiguous plants monitored per plot. The first treatment (Open) was open pollinated with all the flowers of each plant accessible to autonomous self-, wind- and insect-pollination. In the second treatment (Net), all flowers were enclosed in nylon tulle bags with 1 * 1 mm openings (Diatex F510; <http://www.diatex.fr/-Agriculture-.html>) of an appropriate size to cover an

176 inflorescence (buckwheat, field bean & oilseed rape), or an individual flower (strawberry).
 177 Thus, in the Net-treatment all flowers were exposed to wind- and self-pollination, but not to
 178 insect pollination. Because such nets do not hinder the airborne pollen flow (Sacchi and
 179 Price 1988, Wragg and Johnson 2011), the difference between these treatments represents
 180 the contribution from insect pollination. We put the nets over the flower buds before the
 181 onset of flowering. Leaves and plant parts with no flowers were left as much as possible
 182 outside the net bag to minimize any effects of the bag on the photosynthesis (Howpage et
 183 al. 2001). As soon as flowers had wilted, we removed the nets, and the tagged plants were
 184 left to ripen in the field until harvest.

185
 186 For buckwheat, field bean, and oilseed rape, we cut all experimental plants from each plot
 187 and stored them individually in a linen bag just before commercial harvest. For strawberry,
 188 we followed the commercial harvest procedure and harvested the ripe strawberries twice a
 189 week. In each plot, we recorded the number of fruits per plant (field bean, oilseed rape, and
 190 strawberry), and the number and weight of seeds per plant (buckwheat, field bean, and
 191 oilseed rape) were measured using a precision scale and a seed counter. For each crop, we
 192 also measured the specific attributes of quality that affect its marketing value. For oilseed
 193 rape, we analyzed the oil content and chlorophyll contents of the seeds (performed by
 194 Svalof Weibull Lab AB, Svalov, Sweden). High chlorophyll contents decrease the durability
 195 and alter the color of the extracted oil. For field beans, we measured the nitrogen content
 196 of the seed as a proxy of their protein content. The nitrogen content was measured using
 197 oxidative combustion in an automated Dumas type combustion analyzer. For strawberry, we
 198 classified commercial quality as grade 1 (fully developed fruits of good quality), grade 2
 199 (marketable fruits with some changes in colour and shape) and grade 3 (non-marketable

200 fruits) according to guidelines of the German board of trade. For buckwheat, we measured
201 the proportion of filled seeds since high proportion of empty seeds leads to a penalty in the
202 market price. For buckwheat, six fields were destroyed due to a hailstorm, and hence we
203 do not have yield measures for those.

204

205 **Landscape context**

206 The ten fields for each crop were located along a gradient of surrounding landscape
207 complexity. The gradient ranged from intensive agricultural landscapes dominated by large
208 arable fields with few boundary features, to complex landscapes with smaller average
209 arable field sizes and more than 40% coverage of semi-natural habitats, such as pastures
210 and forest patches over 0.5, 1, 2, and 3 km radius around each study field. When selecting
211 the field sites, the proportion of arable land in the surrounding landscape was measured
212 around each experimental field and used as a proxy for landscape complexity (Steffan-
213 Dewenter et al. 2002, Fahrig 2013). The proportion of arable land in the landscape
214 surrounding each of the ten experimental fields varied depending on the region, with some
215 regions presenting more intense landscapes (e.g., range of 48 to 97 % of agricultural land for
216 oilseed rape fields at 1000 m radius), and other regions presenting more complex
217 landscapes (range of 4 to 45 % of agricultural land for field bean at 1000 m radius).

218

219 For oilseed rape, we used the Swedish digitized land cover terrain map database to
220 characterize the landscape surrounding each field (Lantmateriet 2008). For buckwheat and
221 strawberry, we used CORINE data from 2006 (European Environment Agency:
222 <http://www.eea.europa.eu/data-and-maps/data/corine-land-cover-2006-raster-2>). For field
223 beans we used the CORINE 2000 Land Cover Map

(<http://www.ceh.ac.uk/landcovermap2000.html>).

Data analysis and statistics

Landscape effects on bee richness and visitation

Because different organisms act in and react to the landscape at different spatial scales, it is necessary to find a suitable scale at which to measure the surrounding landscape (Steffan-Dewenter et al. 2002; Henry et al. 2012). Before exploring any significances, we ran models for each variable with each of the different radii (0.5–3 km) at which the landscapes had been measured. Hence, for each crop we regressed percentage of agricultural area against pollinator richness and abundance at different radii, and identified the radius that explained the highest proportion of variance (highest R^2). For species richness, all crops showed the highest R^2 at a radius of 0.5 km, while abundance was best explained at a 1 km radius with the exception of field bean bee communities, which also responded to a larger scale (1500 m). We performed joint models for all crops at 0.5 and 1 km radius for richness and visitation abundance, respectively. Bee species richness showed a similar relation to landscape complexity for all crops, and this permitted us to include 'crop' as random factor in the model to investigate the general influence of landscape on richness. Visitation abundance, however, followed contrasting trajectories in relation to landscape depending on the crop. We therefore included in the model crop and its interaction with landscape as fixed effects. Pollinator abundances were centered and scaled to a mean of zero and a deviation of one within each crop.

Yield quantity and quality

248 We first correlated fruit set with yield (i.e., total fruit or seed weight/plant) for each crop.
 249 Block was nested within site and included as random factor in all models. Second, we
 250 constructed two mixed effect model with yield as the response variable. In order to analyze
 251 all crops in the same model, yield and pollinator visitation abundance were centered and
 252 scaled to a mean of zero and a deviation of one within each crop. The first model had
 253 pollination treatment, species richness, total visitation abundance, and the interactions of
 254 treatment with the other two variables as predictors. The second model had treatment,
 255 landscape, and its interaction as predictors. Landscape was investigated at 0.5 and 1 km
 256 radius with similar results and so only models at 1 km are shown. Block, nested within site,
 257 nested within crop was included as a random factor in all models. In both models, a
 258 significant interaction with treatment would indicate that the factor had an effect on yield
 259 only in the open treatment. To account for heteroscedasticity, we added a constant
 260 variance structure (*varIdent* function in package nlme, R) in which the variance was
 261 independently specified for each crop (Cleasby et al. 2011).

262
 263 We also checked if quality was affected by the pollination treatment. Each crop was
 264 analyzed independently due to different measurement units and also because there was no
 265 homogeneous response among the crops. Block nested within site was included as a
 266 random factor in all models. In this case, we tested only for the effect of the pollination
 267 treatment, without including the interactions with species richness, visitation abundance, or
 268 landscape context due to sample size limitations. For buckwheat, we used block as a
 269 constant variance function to control for the different heteroscedasticity among blocks. The
 270 package *nlme* in R was used to fit all models (Pinheiro et al. 2013). Residual plots were
 271 used to check for normality and standardized residuals for heteroscedasticity.

Results

Landscape effects on bee richness and visitation

Pollinator species richness ranged from 2 to 26 species per site (Table 1). The flower visitors of all crops were highly dominated by one or two species of pollinators, in most cases managed honeybees. In field beans, the dominant species were bumblebees; *Bombus terrestris/lucorum* complex, followed by *B. hortorum* and *B. lapidarius* (Fig. 1). Simple landscapes had consistently lower species richness in all crops (GLMM: $F_{1,35} = 5.39$, $P = 0.02$; Fig. 2A). However, the pollinator abundance trend depended on the crop (Table 2; Fig. 2B). Visitation patterns were driven by managed honeybee densities in all crops except for field beans. While in most regions honeybee visits were also higher in complex landscapes (Table 2), in buckwheat there were higher honeybee visits in simple landscapes. For field beans, this positive relationship between number of visits recorded and landscape was even more pronounced at larger scales when we analyze the primary pollinators, the bumblebees, alone ($F_{1,8} = 6.44$, $P = 0.03$ at 1.5 km radius).

Yield quantity and quality

Seed numbers were in all cases positively correlated with yield. However, the correlation was stronger in some crops than others (oilseed rape: $R^2 = 0.95$, $P < 0.0001$; field bean: $R^2 = 0.90$, $P < 0.0001$; strawberry: $R^2 = 0.61$, $P < 0.0001$; buckwheat: $R^2 = 0.67$, $P < 0.0001$).

Open pollination increased fruit set and yield for all crops (Table 3; Fig. 3). We did not detect an interaction between treatment and species richness, which indicates that higher richness does not increase yield in any of the treatments. However, total visitation rate increased

yield in both treatments (Fig 4A). Interestingly, landscape complexity (both at 0.5 or at 1 km) also showed a significant interaction with treatment, indicating that simpler landscapes had lower yields only in the open pollinated plants (Table 3; Fig. 4B). However, landscape did not affect the yield of net-bagged plants.

In addition to quantity, the quality of oilseed rape, buckwheat and strawberry increased in the open pollination treatments (oilseed rape: oil content estimate = 1.28 ± 0.31 %, df = 39, $t = 4.18$ $P < 0.001$; Chlorophyll content estimate = -4.15 ± 1.76 ppm, df = 39, $t = -2.37$, $P = 0.02$; buckwheat: percentage of filled seeds estimate = 0.08 ± 0.01 %, df = 12, $t = 6.35$, $p < 0.001$; strawberry: commercial grade estimate = -0.32 ± 0.06 , df= 67, $t = -5.36$, $p < 0.001$). On the other hand, the nitrogen content of field beans did not increase on open-pollinated plants (estimate = -0.10 ± 0.08 %, df = 37, $t = -1.16$, $p = 0.25$; Fig 3).

Discussion

Four economically important entomophilous annual crops in Europe demonstrated highly different degrees of insect pollination dependence. When open pollinated, mean yield increases ranged from 18 to 71% depending on the crop. Three of these crops are listed as having a “modest” positive impact by animal pollination in the comprehensive review by Klein et al. (2007). However, despite being in the same category, oilseed rape and strawberry increased around 20 %, while field bean reached a 40 % increase in yield from insect pollination. The fourth crop, buckwheat is listed as having a large positive impact by animal pollination, in line of our reported 71 % increase. The Klein et al. (2007) review is currently the best available, most up to date source of animal pollination dependence on crops, but our data highlight a disparity of results among crops listed under the same

category. Our quantitative data on animal pollination dependence provides a first step to depart from the uncertainty embedded in a categorical approach. For example, animal pollination dependence can change by variety and region. Recent reports show variability in pollinator dependence between 0 to 30% among varieties of oilseed rape (Stanley et al. 2013, Garratt et al. 2013). While we were able to standardize variety for most studied crops, strawberry fields were planted with four different varieties and the presented data should be seen as an average across those varieties (but see Klatt et al. 2013).

As expected, we found that fruit set per plant was positively correlated with yield. However, this correlation was rather weak ($r^2 \sim 0.60$) for both strawberry and buckwheat. This indicates that for these crops, the total fruit weight was quite variable among plants with similar fruit or seed numbers. Indeed, for strawberry, the size of the receptacle is directly related to the number of fertilized achenes while for buckwheat, it is the proportion of filled seeds that can vary considerably and is a major component of yield besides fruit set. While previous research has focused mainly on exploring the effects of pollinators on fruit or seed set (e.g. Garibaldi et al. 2011, 2013), which is a more direct measure of plant reproduction, yield has the potential to better reflect economic value (Bommarco et al. 2012, Klatt et al. 2013), and hence, farmers' interest. For example, while less than 20% in mean yield increase may seem as a modest advantage from the plant perspective, for the farmers it can translate into a substantial difference in revenue.

Similarly, we report that the yield quality component is enhanced to different extents by open pollination in three out of four crops. For buckwheat, strawberry, and oilseed rape, quality is directly linked to the pollinating activity of insects. We find this despite the fact that the measure of quality and underlying mechanisms is specific for each crop, and largely

unrelated among crops. Empty seeds in buckwheat accumulate little or no starch (Björkman 1995). The shape of strawberries is also directly related to a complete pollination of all ovules, resulting in a homogeneously pollinated fruit (Zebrowska 1998). For oilseed rape, the plant allocate more oil resources to well pollinated seeds. In contrast, for field beans, the nitrogen content in the seeds was not affected by insect pollination. Other factors such as soil fertility and availability of the appropriate N-fixing bacteria (*Rhizobium* spp.) may play a more important role for field beans (Köpke and Nemecek 2010). Is interesting to note that we do not detect any trade-off between yield and nitrogen content of the seeds, as plants with more seeds do not present lower nitrogen content. Hence, the overall protein yield (i.e. nitrogen content at the plant level) was increased with open pollination. On a general note, it is interesting to observe that many different aspects of a plant's investments in its seed or fruit are affected by pollination.

The treatment with netted flowers gives us estimates for the extreme cases where pollinators are completely absent, and we show that the current levels of pollination do not suffice to maximize yield in the open pollinated treatment in all landscapes. As previously reported, we confirm that agricultural intensification has a drastic effect on bee species richness (Rickets et al. 2008, Garibaldi et al. 2011). However, total visitation does not always follow the same pattern as richness. This is the case for buckwheat and field bean, where fields presenting higher total visits were located in simple landscapes. For buckwheat, most of the visits in complex landscapes were due to increased honeybee densities managed for pollination. In field beans we found that bumblebees respond positively to agricultural simplification, noting, however, that even the more simple field beans landscapes contain a fair amount of semi-natural habitats. Overall, we find a general positive relationship

between total visitation rates and yield, but not with species richness. If the remaining species that thrive in intensively cultivated agricultural areas, including the managed honeybee, are effective pollinators, yield losses can be partly decoupled from losses of species (Bartomeus and Winfree 2013).

A recent global meta-analysis highlights the role of wild species in crop systems (Garibaldi et al. 2013). The flower visitors of three out of four crops were clearly dominated by honeybees (Fig. 1) and hence, are likely to be key pollinators for those crops. Garibaldi et al. (2013) show that an increase in wild insect visitation enhanced fruit set by twice as much as an equivalent increase in honeybee visitation. While this is generally the case in our target crops (three of which were included as part of Garibaldi's synthesis), the numerical advantage of honeybees in European agricultural landscapes needs to be acknowledged when calculating their total contribution to pollinated plants (e.g., as done in Winfree et al. 2007 and Rader et al. 2009). However, increasing or maintaining high pollinator diversity can enhance yield quantity and stability by improving the pollination efficiency of honeybees (Greenleaf & Kremen 2006) and reduce the risk of pollination failure due to climate change (Rader et al. 2013, Bartomeus et al. 2013b), or environmental disturbances such as extreme weather events (Brittain et al. 2012).

Overall, we also found a weak negative effect of land use intensity on yield (Garibaldi et al. 2011, but see Ricketts et al. 2008), but this was not directly mediated by increased pollinator visitation by itself, because the correlation between pollinator total visits and the proportion of agricultural land in the landscape was weak. The yield of experimental plots with net bagged flowers also increased in sites with more pollinators (Fig 4A). This suggests

that other environmental or biotic factors correlated with insect visitation may have been operating simultaneously. The release of airborne pollen by foraging bees could be such a factor (Pierre et al. 2009) (Fig 4B).

In order to make efficient management decisions and increase our power to predict the actual benefit from pollinators in a certain farming situation, we need to estimate the combined contribution of multiple ecosystem services and agricultural inputs (Boreux et al. 2013), as they may be influenced differently by landscape characteristics or have non-additive interactions among them (e.g., Lundin et al. 2013, Martin et al. 2013). Information on the benefit delivered by pollinators to yield quantity and quality in relation to landscape context provides an important baseline for this work.

Acknowledgements

We thank H. Dathe, G. Else, R. Fonfria, S. Iserbyt, M. Kuhlmann, G. Le Goff, D. Michez, H. Mouret, A. Müller, S. Patiny, A. Pauly, P. Rasmont, S. Risch, M. Schwarz, R. Theunert, Waldemar C., and P. Williams for bee identifications to species.

References

- Andersson GKS, Rundlöf M, Smith HG. 2012. Organic farming improves pollination success in strawberries. *PLoS ONE* 7(2):e31599 (doi:10.1371/journal.pone.0031599)
- Angadi SV, Cutforth HW, McConkey BG, Gan Y. 2003. Yield adjustment by canola grown at different plant populations under semiarid conditions. *Crop Sciences*. 43 :1358-1366 (doi:10.2135/cropsci2003.1358).

417
418 Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R. 2013.
419 Historical changes in northeastern US bee pollinators related to shared ecological traits.
420 *Proceedings of the National Academy of Sciences of the United States of America*, 110(12),
421 4656–60. doi:10.1073/pnas.1218503110
422
423 Bartomeus I, Park MG, Gibbs J, Danforth BN, Lakso AN, Winfree R. 2013. Biodiversity
424 ensures plant-pollinator phenological synchrony against climate change. *Ecology letters*,
425 16(11), 1331–8. doi:10.1111/ele.12170
426
427 Bartomeus I, Winfree R. 2013. Pollinator declines: reconciling scales and implications for
428 ecosystem services. *F1000Research*, 146, 2–4. doi:10.12688/f1000research.2-146.v1
429
430 Björkman T. 1995. Role of honey bees (Hymenoptera: Apidae) in the pollination
431 of buckwheat in Eastern North America. *Journal of Economic Entomology* 88(6): 1739-1745
432
433 Bommarco R, Kleijn D, Potts SG. 2013. Ecological intensification: harnessing ecosystem
434 services for food security. *Trends in ecology & evolution* 28(4) 230-238,
435 doi:10.1016/j.tree.2012.10.012.
436
437 Bommarco R, Lundin O, Smith HG, Rundlof M. 2011. Drastic historic shifts in bumble-bee
438 community composition in Sweden. *Proceedings of the Royal Society B*. 279(1727), 309-315.
439 doi:10.1098/rspb.2011.0647
440

- 441 Bommarco R, Marini L, Vaissière BE. 2012. Insect pollination enhances seed yield, quality,
442 and market value in oilseed rape. *Oecologia*, 169(4), 1025-1032.
- 443
- 444 Bos MM, Veddeler D, Bogdanski AK, Klein AM, Tschardt T, Steffan-Dewenter I, Tylianakis J
445 M. 2007. Caveats to quantifying ecosystem services: fruit abortion blurs benefits from crop
446 pollination. *Ecological Applications*, 17(6), 1841-1849.
- 447
- 448 Boreux V, Kushalappa CG, Vaast P, Ghazoul J. 2013. Interactive effects among ecosystem
449 services and management practices on crop production: pollination in coffee agroforestry
450 systems. *Proceedings of the National Academy of Sciences of the United States of America*,
451 110(21), 8387–92. doi:10.1073/pnas.1210590110
- 452
- 453 Breeze TD, Bailey AP, Balcombe KG, SG Potts. 2011. Pollination services in the UK: How
454 important are honeybees? *Agriculture, Ecosystems and Environment*, 142(3), 137-143.
- 455
- 456 Brittain C, Kremen C, Klein AM. 2012. Biodiversity buffers pollination from changes in
457 environmental conditions. *Global Change Biology*, 19(2), 540-547. doi:10.1111/gcb.12043
- 458
- 459 Cariveau DP, Williams NM, Benjamin FE, Winfree R. 2013. Response diversity to land use
460 occurs but does not consistently stabilise ecosystem services provided by native pollinators.
461 *Ecology letters*, 16(7), 903–11. doi:10.1111/ele.12126
- 462
- 463 Carré G, Roche P, Chifflet R, Morison N, Bommarco R, Harrison-Cripps J, Krewenka K, Potts
464 SG, Roberts SPM, Rodet G, Settele J, Steffan-Dewenter I, Szentgyörgyi H, Tscheulin T,
465 Westphal C, Woyciechowski M, Vaissière BE. 2009. Landscape context and habitat type as

466 drivers of bee biodiversity in European annual crops. *Agriculture, Ecosystems and*
 467 *Environment*, 133 :40-47.

468

469 Carvalho LG, Kunin WE, Keil P, Aguirre-Gutiérrez J, Ellis WN, Fox R, ... Biesmeijer JC. 2013.
 470 Species richness declines and biotic homogenisation have slowed down for NW-European
 471 pollinators and plants. *Ecology letters*, 16(7), 870–8. doi:10.1111/ele.12121

472

473 Chacoff NP, Aizen MA, Aschero V. 2008. Proximity to forest edge does not affect crop
 474 production despite pollen limitation. *Proceedings of the Royal Society B*, 275 :907-913 (doi:
 475 10.1098/rspb.2007.1547).

476

477 Chagnon M, Gingras J, Deoliveira D. 1993. Complementary aspects of strawberry pollination
 478 by honey and indigenous bees (Hymenoptera). *Journal of Economic Entomology*, 86, 416-
 479 420.

480

481 Cleasby IR, Nakagawa S. 2011. Neglected biological patterns in the residuals. *Behavioral*
 482 *Ecology and Sociobiology*, 65(12), 2361-2372.

483

484 Fahrig L. 2013. Rethinking patch size and isolation effects: the habitat amount hypothesis.
 485 *Journal of Biogeography*, 40(9), 1649–1663. doi:10.1111/jbi.12130

486

487 Ferguson LB, Watkins CB. 1992. Crop load affects mineral concentrations and incidence of
 488 bitter pit in 'Cox's Orange Pippin' apple fruit. *Journal of the American Society of Horticultural*
 489 *Sciences*, 117 :373-376.

490

- 491 Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA, ...
- 492 Klein AM. 2013. Wild pollinators enhance fruit set of crops regardless of honey bee
- 493 abundance. *Science*, 339(6127), 1608–11. doi:10.1126/science.1230200
- 494
- 495 Garibaldi LA, Steffan-Dewenter I, Kremen C, Morales JM, Bommarco R, Cunningham SA, ...
- 496 Klein AM. 2011. Stability of pollination services decreases with isolation from natural areas
- 497 despite honey bee visits. *Ecology Letters*, 14(10), 1062-1072.
- 498
- 499 Garratt MPD, Breeze T, Jenner N, Polce C, Biesmeijer JC, Potts SG. 2013. Avoiding a bad
- 500 apple: insect pollination enhances fruit quality and economic value. *Agriculture, Ecosystem*
- 501 *and Environment*, In press
- 502
- 503 Gomez NV, Miralles DJ. 2011. Factors that modify early and late reproductive phases in
- 504 oilseed rape (*Brassica napus* L.): its impact on seed yield and oil content. *Industrial Crops*
- 505 *and Products*, 34 :1277-1285.
- 506
- 507 Gonzalez MV, Coque M, Herrero M. 1998. Influence of pollination systems on fruit set and
- 508 fruit quality in kiwifruit (*Actinidia deliciosa*). *Annals of Applied Biology*, 132 :349-355.
- 509
- 510 Greenleaf SS, Kremen C. 2006. Wild bee species increase tomato production but respond
- 511 differently to surrounding land use in Northern California. *Biological Conservation*, 133, 81–
- 512 87.
- 513
- 514 Greenleaf SS, Williams NM, Winfree R, Kremen C. 2007. Bee foraging ranges and their
- 515 relationship to body size. *Oecologia*, 153(3), 589-596.

516

517 Henry M, Fröchen M, Maillet-Mezeray J, Breyne E, Allier F, Odoux J-F, Decourtye A. 2012.

518 Spatial autocorrelation in honeybee foraging activity reveals optimal focus scale for

519 predicting agro-environmental scheme efficiency. *Ecological Modeling*, 225 :103-114.

520 (doi.org/10.1016/j.ecolmodel.2011.11.015,)

521

522 Hogendoorn K, Bartholomaeus F, Keller MA. 2010. Chemical and sensory comparison of

523 tomatoes pollinated by bees and by a pollination wand. *Journal of Economic Entomology*,

524 103 :1286-1292.

525

526 Howpage D, Spooner Hart RN, Vithanage V. 2001. Influence of honey bee (*Apis mellifera*) on

527 kiwifruit pollination and fruit quality under Australian conditions. *New Zealand Journal of*

528 *Crop Horticultural Sciences*, 29 :51-59.

529

530 Höhn P, Tscharntke T, Tylianakis JM, Steffan-Dewenter I. 2008. Functional group diversity of

531 bee pollinators increases crop yield. *Proceedings of the Royal Society of London, Series B*,

532 275, 2283-2291

533

534 Isaacs R, Kirk AK. 2010. Pollination services provided to small and large highbush blueberry

535 fields by wild and managed bees. *Journal of Applied Ecology*, 47, 841-849.

536

537 Lundin O, Smith HG, Rundlöf M, Bommarco R. 2013. When ecosystem services interact: crop

538 pollination benefits depend on the level of pest control. *Proceedings of the Royal Society B:*

539 *Biological Sciences*, 280(1753).

540

541 Kennedy CM, Lonsdorf E, Neel MC, Williams NM, Ricketts TH, Winfree R, ... Kremen C. 2013.
 542 A global quantitative synthesis of local and landscape effects on wild bee pollinators in
 543 agroecosystems. *Ecology letters*, 16(5), 584–99. doi:10.1111/ele.12082
 544
 545 Kevan PG, Eisikowitch D. 1990. The effects of insect pollination on canola (*Brassica napus* L.
 546 cv. O.A.C. Triton) seed germination. *Euphytica*, 45 :39-41.
 547
 548 Klatt BK, Holzschuh A, Westphal C, Clough Y, Smit I, Pawelzik E, Tscharntke T. 2014. Bee
 549 pollination improves crop quality, shelf life and commercial value. *Proceedings of the Royal*
 550 *Society B: Biological Sciences*, 281(1775), 20132440.
 551
 552 Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke
 553 TD. 2007. Importance of pollinators in changing landscapes for world crops. *Proceedings of*
 554 *the Royal Society B-Biological Sciences*, 303–313.
 555
 556 Klein AM, Steffan-Dewenter I, Tscharntke T. 2003. Pollination of *Coffea canephora* in
 557 relation to local and regional agroforestry management. *Journal of Applied Ecology*, 40,
 558 837–845.
 559
 560 Köpke U, Nemecek T. 2010. Ecological services of faba bean. *Field Crops Research*, 115(3),
 561 217-233.
 562
 563 Le Guen J, Mesquida J, Pierre JS, Morin G, Taséi JN, Carré S. 1993. Efficacité pollinisatrice de
 564 différents traitements sur two lignées de féverole de printemps (*Vicia faba* L. var. equina

Steudel), à des niveaux d'autofertilité différents, avec utilisation de diverses espèces de
Bombus Latr. (Hymenoptera: Apidae). *Apidologie*, 24 :129-145.

Martin EA, Reineking B, Seo B, Steffan-Dewenter I. 2013. Natural enemy interactions
constrain pest control in complex agricultural landscapes. *Proceedings of the National
Academy of Sciences of the United States of America*, 110(14), 5534–9.
doi:10.1073/pnas.1215725110

Pierre J, Vaissière B, Vallée P, Renard M. 2010. Efficiency of airborne pollen released by
honeybee foraging on pollination in oilseed rape: a wind insect-assisted pollination.
Apidologie, 41(1), 109-115. (DOI: 10.1051/apido/2009056)

Pinheiro J, Bates D, DebRoy S, Sarkar D. 2011. nlme: linear and nonlinear mixed effects
models. R package version 3.1–98. *R Foundation for Statistical Computing, Vienna*.

Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010. Global
pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution*, 25(6), 345–
353.

Rader R, Howlett BG, Cunningham SA, Westcott DA, Newstrom-Lloyd LE, Walker MK, Teulon
DAJ, Edwards W. 2009. Alternative pollinator taxa are equally efficient but not as effective
as the honeybee in a mass flowering crop. *Journal of Applied Ecology*, 46(5), :1080-1087
(DOI: 10.1111/j.1365-2664.2009.01700.x).

- 589 Rader R, Reilly J, Bartomeus I, Winfree R. 2013. Native bees buffer the negative impact of
590 climate warming on honey bee pollination of watermelon crops. *Global change biology*,
591 19(10), 3103–10. doi:10.1111/gcb.12264
592
- 593 Ricketts TH, Regetz J, Steffan-Dewenter I, Cunningham SA, Kremen C, Bogdanski A, ... Viana
594 BF. 2008. Landscape effects on crop pollination services: are there general patterns? *Ecology*
595 *Letters*, 11(5), 499-515.
596
- 597 Roldan Serrano A, Guerra-Sanz JM. 2006. Quality fruit improvement in sweet pepper culture
598 by bumblebee pollination. *Scientia Horticulturae*, 110, 160-166.
599
- 600 Roselino AC, Santos SB, Hrncir M, Bego LR. 2009. Differences between the quality of
601 strawberries (*Fragaria x ananassa*) pollinated by the stingless bees *Scaptotrigona aff. depilis*
602 and *Nannotrigona testaceicornis*. *Genetics and Molecular Research*, 8, 539-545.
603
- 604 Sacchi CF, Price PW. 1988. Pollination of the arroyo willow, *Salix lasiolepis*: role of insects
605 and wind. *American Journal of Botany*, 1387-1393.
606
- 607 Stanley DA, Gunning D, Stout JC. 2013. Pollinators and pollination of oilseed rape crops
608 (*Brassica napus* L.) in Ireland: ecological and economic incentives for pollinator
609 conservation. *Journal of Insect Conservation*. doi:10.1007/s10841-013-9599-z
610
- 611 Steffan-Dewenter I, Munzenberg U, Burger C, Thies C, Tschardt TD. 2002. Scale-
612 dependent effects of landscape context on three pollinator guilds. *Ecology*, 1421–1432.
613

614 Stivers RK, Swearingin ML. 1980. Soybean yield compensation with different populations
 615 and missing plant patterns. *Agronomy Journal*, 72 :98-102
 616 (doi:10.2134/agronj1980.00021962007200010019x).
 617
 618 Vaissière BE, Freitas BM, Gemill-Herren B. 2011. Protocol to detect and assess pollination
 619 deficits in crops: a handbook for its use. FAO, Rome, Italy. 81 pp.
 620
 621 Westphal C, Bommarco R, Carré G, Lamborn E, Morison N, Petanidou T, ... Steffan-Dewenter
 622 I. 2008. Measuring bee diversity in different european habitats and biogeographical regions.
 623 *Ecological Monographs*, 78(4), 653–671. doi:10.1890/07-1292.1
 624
 625 Westphal C, Steffan-Dewenter I, Tscharntke T. 2003. Mass flowering crops enhance
 626 pollinator densities at a landscape scale. *Ecology Letters*, 6(11), 961–965.
 627 doi:10.1046/j.1461-0248.2003.00523.x
 628
 629 Williams NM, Crone EE, Roulston TH, Minckley RL, Packer L, Potts SG. 2010. Ecological and
 630 life-history traits predict bee species responses to environmental disturbances. *Biological*
 631 *Conservation*, 143, 2280–2291.
 632
 633 Williams IH, Martin AP, White RP. 1986. The pollination requirements of oilseed rape
 634 (*Brassica napus* L.). *Journal of Agricultural Science*, 106 :27-30.
 635
 636 Winfree R, Bartomeus I, Cariveau DP. 2011. Native pollinators in anthropogenic habitats.
 637 *Annual Review of Ecology Evolution and Systematics*, 42, 1–22. doi:10.1146/annurev-
 638 ecolsys-102710-145042

639

640 Winfree R, Williams NM, Dushoff J, Kremen C. 2007. Native bees provide insurance against
641 ongoing honey bee losses. *Ecology letters*, 10(11), 1105–13. doi:10.1111/j.1461-
642 0248.2007.01110.x

643

644 Wragg PD, Johnson SD. 2011. Transition from wind pollination to insect pollination in
645 sedges: experimental evidence and functional traits. *New Phytologist*, 191(4), 1128-1140.

646

647 Zebrowska, J. 1998. Influence of pollination modes on yield components in strawberry
648 (*Fragaria x ananassa* Duch). *Plant Breeding*, 117 :255-260 (DOI: 10.1111/j.1439-
649 0523.1998.tb01935.x).

650 Table 1. Characteristics of the four study systems.

	Variety	Distance between sites (range in km)	Field sizes (range in ha)	Mean richness of pollinators
Oil seed rape	Stratos	3 - 7	1.0 – 40.4	11.3
Field bean	Clipper	3 - 18	5.0 – 47.0	3.1
Strawberry	Honeoye, Korona, Darselect, Symphonie	3 - 26	0.3 – 1.3	12.9
Buckwheat	Kora	4 - 7	0.3 – 4.0	11.4

651

652 Table 2. Effects of land use complexity on total visitation and honeybee visitation (field
 653 beans excluded from the honeybee model). Visitation is scaled within each crop. Both
 654 models include block nested in site as random factors. Agriculture is the proportion of
 655 arable land in the surrounding landscape of each field.

Total visitation	F-value	D.f.	P-value
Crop	3.13	3	0.04
Agriculture 1km	0.05	1	0.81
Agriculture*crop	3.08	3	0.04
Residuals		32	
Honeybee visitation			
Crop	4.35	2	0.02
Agriculture 1km	0.10	1	0.75
Agriculture*crop	3.87	2	0.03
Residuals		32	


656
 657

658 Table 3. Effects of open pollination vs pollinator exclusion treatments, visitation and
 659 landscape context on yield in four entomophilous crops grown over 10 fields in Europe
 660 (buckwheat, field bean, spring oilseed rape and strawberry). Yield and visitation are scaled
 661 within each crop. Block, nested in site, nested in crop are included as a random factor.
 662 Agriculture is the proportion of arable land in the surrounding landscape of each field.

	F-value	Df	P-value
Pollination treatment	67.05	128	< 0.001
Pollinator richness	0.51	28	0.482
Total number of visits	7.25	28	0.012
Treatment*Pollinator richness	0.34	128	0.557
Treatment*Total number of visits	0.23	128	0.634
	F-value	Df	P-value
Pollination treatment	10.05	128	0.001
Agriculture 1 km radius	0.18	29	0.671
Treatment*Agriculture	9.73	128	0.002

663 **Figures:**

664 Fig 1: Total number of visits recorded per pollinator guild in each crop. All crops received the
 665 same sampling effort (i.e., four 30 minutes visits to 150 m transects). Note the strong
 666 dominance of honeybees in most crops.



667

668 Fig 2: Relationship of A) Pollinator richness per field and B) Total number of visits per field
 669 with landscape complexity at the appropriate radii. Each crop individual slope is plotted in a
 670 different color and the overall response, when appropriate, in black. Total visits are scaled
 671 within each crop.






Fig 3: Overall yield per plant (A, C, E, G) and quality (B, D, F, H) with pollinator exclusion (Net) and open pollination (Open) for each crop. Black dots are the mean values reported in the text, and the boxplots reflects the distribution of the data. Yield is measured in seed weight per plant (g) for all crops except strawberry, which was measured as fruit weight per plant (g). Commercial Grades of 1 and 2 are marketable, while grade 3 is considered non marketable.





681

Fig 4: Interaction plots showing the relationships of A) Yield per plant and total visitation
and B) Yield and landscape complexity for pollinator exclusion (open circles, dotted line) and
open pollination (black circles, solid line). Total visitation and yield are scaled to a mean of
zero within each crop.

