

A peer-reviewed version of this preprint was published in PeerJ on 13 September 2016.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.2376) (peerj.com/articles/2376), which is the preferred citable publication unless you specifically need to cite this preprint.

Savage AM, Hills J, Driscoll K, Fergus DJ, Grunden AM, Dunn RR. 2016. Microbial diversity of extreme habitats in human homes. PeerJ 4:e2376 <https://doi.org/10.7717/peerj.2376>

Microbial diversity of extreme habitats in human homes

Amy M Savage ^{Corresp., 1}, **Justin Hills** ², **Katherine Driscoll** ³, **Daniel J Fergus** ⁴, **Amy M Grunden** ⁵, **Robert R Dunn** ⁶

¹ Rutgers, The State University of New Jersey - Camden

² Laboratory of Cellular and Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, United States

³ Animal Management Department, The Wilds, Cumberland, OH, United States

⁴ Genomics and Microbiology, North Carolina Museum of Natural Sciences, Raleigh, NC, United States

⁵ Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, United States

⁶ Department of Applied Ecology and Keck Center for Behavioral Biology, North Carolina State University, Raleigh, NC, United States

Corresponding Author: Amy M Savage

Email address: amy.savage@rutgers.edu

High-throughput sequencing techniques have opened up the world of microbial diversity to scientists, and a flurry of studies in the most remote and extreme habitats on earth have begun to elucidate the key roles of microbes in ecosystems with extreme conditions. These same environmental extremes can also be found closer to humans, even in our homes. Here, we used high-throughput sequencing techniques to assess bacterial and archaeal diversity in the extreme environments inside human homes (e.g., dishwashers, hot water heaters, washing machine bleach reservoirs, etc.). We focused on habitats in the home with extreme temperature, pH, and chemical environmental conditions. We found a lower diversity of microbes in these extreme home environments compared to less extreme habitats in the home. However, we were nonetheless able to detect sequences from a relatively diverse array of bacteria and archaea. Habitats with extreme temperatures alone appeared to be able to support a greater diversity of microbes than habitats with extreme pH or extreme chemical environments alone. Microbial diversity was lowest when habitats had both extreme temperature and one of these other extremes. In habitats with both extreme temperatures and extreme pH, taxa with known associations with extreme conditions dominated. Our findings highlight the importance of examining interactive effects of multiple environmental extremes on microbial communities. Inasmuch as taxa from extreme environments can be both beneficial and harmful to humans, our findings also suggest future work to understand both the threats and opportunities posed by the life in these habitats.

1 Microbial diversity of extreme habitats in human homes

2 Amy M. Savage* ¹, Justin L. Hills², Katherine Driscoll³, Daniel J. Fergus⁴, Amy M. Grunden⁵,
3 Robert R. Dunn⁶

4 * Corresponding author: Amy.Savage@rutgers.edu

5 Other authors: JLHills14@gmail.com, Katherine.Driscoll14@gmail.com,
6 DanielJFergus@gmail.com, amgrunde@ncsu.edu, RRDunn@ncsu.edu

7 1- Department of Biology & Center for Computational and Integrative Biology, Rutgers
8 University, Waterfront Technology Center, 200 Federal Street, Camden, NJ 08102

9
10 2- Laboratory of Cellular and Molecular Biology, National Institute of Diabetes and
11 Digestive and Kidney Diseases, 8 Center Drive, MSC 0830, Bethesda, MD 20892-
12 08308322

13
14 3- Animal Management Department, the Wilds, 14000 International Road, Cumberland OH

15
16
17 4- Genomics and Microbiology, North Carolina Museum of Natural Sciences, 11 West
18 Jones St, Raleigh, NC 27601

19
20 5- Department of Plant and Microbial Biology, North Carolina State University, 4550A
21 Thomas Hall, Campus Box 7612, Raleigh, NC, 27695-7612

22
23 6- Department of Applied Ecology and Keck Center for Behavioral Biology, North Carolina
24 State University, Campus Box 7617, Raleigh, NC 27695-7617

25

26

Abstract:

28 High throughput sequencing techniques have opened up the world of microbial diversity to
 29 scientists, and a flurry of studies in the most remote and extreme habitats on earth have begun to
 30 elucidate the key roles of microbes in ecosystems with extreme conditions. These same
 31 environmental extremes can also be found closer to humans, even in our homes. Here, we used
 32 high throughput sequencing techniques to assess bacterial and archaeal diversity in the extreme
 33 environments inside human homes (e.g., dishwashers, hot water heaters, washing machine bleach
 34 reservoirs, etc.). We focused on habitats in the home with extreme temperature, pH and chemical
 35 environmental conditions. We found a lower diversity of microbes in these extreme home
 36 environments compared to less extreme habitats in the home. However, we were nonetheless
 37 able to detect sequences from a relatively diverse array of bacteria and archaea. Habitats with
 38 extreme temperatures alone appeared to be able to support a greater diversity of microbes than
 39 habitats with extreme pH or extreme chemical environments alone. Microbial diversity was
 40 lowest when habitats had both extreme temperature and one of these other extremes. In habitats
 41 with both extreme temperatures and extreme pH, taxa with known associations with extreme
 42 conditions dominated. Our findings highlight the importance of examining interactive effects of
 43 multiple environmental extremes on microbial communities. Inasmuch as taxa from extreme
 44 environments can be both beneficial and harmful to humans, our findings also suggest future
 45 work to understand both the threats and opportunities posed by the life in these habitats.

Keywords: Community Structure, Extreme environments, Human Homes, Interactive effects, Microbial Diversity

Introduction:

The innovation of culture-independent, high-throughput sequencing techniques has facilitated the discovery of high microbial diversity in many habitats once considered inhospitable to life (Rothschild and Mancinelli 2001). The species in these environments are frequent targets for the discovery of useful enzymes (Niehaus et al. 1999, van den Burg 2003, Elleuche et al. 2014), and studies of microbes living in extreme environments have provided key insights into the evolution of microbial metabolism (Valentine 2007, Hoehler and Jorgensen 2013). Often overlooked, however, is that the attributes that define many of the most extreme habitats on Earth, such as extremes of temperature, pH, water activity, or low nutrient levels, can also be found more immediate to everyday experience. Human homes, for example, contain microhabitats as hot, acidic, basic or salty as any encountered elsewhere on Earth (Martin et al. 2015).

We know of only two extreme habitats within homes where microbial diversity has been studied to date, and in both cases culture-dependent techniques were used. In 1973, Brock and Boylen discovered a species of the genus *Thermus* (*T. aquaticus*) living in hot water heaters. Species of this genus had previously been known only from hot springs (Brock and Boylen 1973). In addition, studies have considered the biology of tap water. Tap water is hospitable in terms of its abiotic conditions (e.g. temperature, pH, toxicity) but is very low in nutrients and so was long

assumed to be relatively devoid of life; until, that is, it was studied. Tap water has now been shown to contain many species of bacteria capable of surviving in low nutrient environments (Kalmbach et al. 1997, Szewzyk et al. 2000, Boe-Hansen et al. 2002). If life exists in hot water heaters and tap water, it seems possible and even likely that many extreme habitats in homes sustain life. That the environmental extremes imposed by these conditions in homes (cold, hot, acidic, alkaline, wet or dry) delineate which species are present seems inevitable. That they are lifeless is unlikely.

Here, we used culture-independent, high-throughput sequencing to address the following questions: (1) What is the relative diversity of microbes (specifically, Bacteria and Archaea) under extremes of temperature, pH and chemical environments of southeast US homes and how does it compare to habitats without each extreme conditions? Harrison et al. (2013) recently argued that because many extreme environments include simultaneous extremes in multiple environmental factors, interactive effects of these multiple sources of extreme conditions are likely to be important determinants of microbial diversity in extreme environments. Therefore, we additionally asked (2) how do multiple, simultaneous extreme conditions influence microbial diversity in human homes? Finally, we asked (3) which bacterial and archaeal genera from the broader home (Dunn et al. 2013) fail to persist in extreme home habitats, and which microbial genera persist only in these extreme habitats?

Methods:

Sampling extreme home environments

We sampled extreme environments in six houses in the Raleigh-Durham metropolitan area (Supp. Fig 1). In each house, we used dual-tipped sterile BBL™ CultureSwabs™ or 50 ml

conical tubes to collect water from each of 10 standardized extreme locations in homes. The sites sampled in all six houses included environments that were extreme in terms of their temperature, pH and/or chemical environments (Supp. Table 1). Our assumptions concerning these sampling locations are based upon publicly available consumer resources regarding certain commercial and industrial requirements (e.g. <http://www.nsf.org/consumer-resources/health-and-safety-tips/home-product-appliance-tips/sanitizing-dishwasher>, <http://energy.gov/energysaver/projects/savings-project-lower-water-heating-temperature>). For example, our sampling of dishwashers was influenced by the NSF/ANSI 184 standard for residential dishwashers to provide a final rinse at a temperature of at least 150 °F (65.6 °C). Additionally, temperature ranges for residential water heaters are 90 to 150 F (32 – 65.6 °C), depending on the manufacturer. Bleach receptacles in clothes washing machines would also be assumed to have a pH of 12 when bleach is present. Although the pH and chemical composition of laundry detergent and dishwasher detergent can be quite variable, manufacturing standards are generally within the 7-10 pH range. While measurements, opposed to assumptions, would be very useful, taking measurements of all the potential extreme axes under various sample sites in multiple homes was not feasible. All samples were preserved at -20°C immediately after collection.

Isolating and identifying microbes in extreme home environments

Genomic DNA was extracted from all samples using the MoBio Power Soil DNA extraction kit (MoBio, Carlsbad, CA) as described previously (Fierer et al., 2008; Lauber et al., 2009). For swabs, the tips were placed in PowerBead tubes containing solution C1 and swirled vigorously for approximately 10 seconds to release contents and removed. Water samples were thawed and filtered using Corning 50 ml 0.22µm cellulose acetate filters after which the filters were added to

the PowerBead tubes. The extractions were subsequently performed as directed by the manufacturer, except that the final elution was performed in 50µl of 70° C C6 elution buffer. Because the water samples were frozen prior to filtering and extraction, the results reported for the water samples likely under-represents the true diversity of taxa in those environments.

We used methods described in Bates et al (2011) to amplify bacterial and archaeal DNA from the samples collected from homes and six negative controls. Briefly, amplicons were produced by PCR with universal bacterial/archaeal 515F and 806R primers to which Roche 454 B pyrosequencing adapters had been added, as described in Hulcr et al. (2012). The 515F primer contained an additional 12-bp barcode sequence for individual sample identification. All the samples were amplified by triplicate PCR reactions, cleaned using the UltraClean-htp 96-well PCR Clean-up kit (MoBio), and quantified with a Quant-iT PicoGreen dsDNA Assay kit (Invitrogen). Equimolar amounts of each sample were pooled into a single sample to sequence. DNA pyrosequencing was performed at Selah Clinical Genomics Center at Innovista (University of South Carolina, USA) using a Roche Genome Sequencer 454 FLX system to facilitate comparison to previous related work that utilized this platform (Dunn et al, 2013). Though these methods here do not distinguish living from recently dead cells, with the comparative approach used here, we presume that taxa frequently identified in one habitat but rare or absent in most others are likely surviving in the habitat from which they are frequently identified. The sequences were submitted to NCBI (SRA accession number SRP071677).

The QIIME analysis package (Caporaso et al, 2010a) was used to process and analyze the barcoded microbial (bacterial and archaeal) amplicon sequences. Sequences were quality filtered to a minimum quality score of 25 with no unambiguous bases and sorted to each sample by the 12 bp barcodes. The 454 pyrosequencing produced 197,305 reads that passed the quality

screening. The sequences were grouped into Operational Taxonomic Units (OTUs) that shared at least 97% sequence similarity. A representative sequence was taken for each OTU group and PyNASt (Caporaso et al, 2010b) was employed to align these representative sequences to the Greengenes database (DeSantis et al, 2006) and the taxonomic identity of each OTU was determined using the RDP Classifier (Wang et al, 2007). Phylotypes were considered to be contaminants if they were seen in at least two of the six negative control samples. There were 152 OTUs at the genus level present in more than one negative sample, representing 9% of the total OTUs at this level. After removing contaminant sequences and singletons, the number of quality-filtered reads per sample was between 6 and 5861 (median=2306). Finally, we removed any OTU's represented by 20 or fewer reads to reduce the possibility of spurious results from the sequencing process. For among samples comparisons we rarefied to each to a depth of 1000 sequences. Thus, our assessments of microbial diversity are conservative.

Analysis of the relative diversity of microbes in extreme temperature, pH and chemical environments of homes and how it compares to habitats without each extreme condition

We compared microbial species accumulation among three extreme variables in homes: temperature, pH, and chemical extremes. Temperature was classified on a scale of 1-5, with 1 representing the coldest environments and 5 representing the hottest environments. We then binned 1 and 5 into an extreme temperature category and 2-4 into an intermediate temperature category. Similarly, environments were classified as acidic, basic or neutral and then binned into extreme pH (acidic or basic environments) *versus* neutral environments. Finally, chemical extremes were those environments characterized by the presence of detergent, bleach, metals, ammonia, or natural gas (Supp. Table 2).

We used EstimateS v. 9.1.0 (Colwell 2013) to construct individual-based species accumulations for all three extreme environments and their non-extreme counterparts. For these curves, reads were used as individuals and the curves were constructed using 1000 iterations. To formally assess differences in accumulated species by read, we used \pm 95% confidence intervals for each curve. Non-overlapping 95% confidence intervals are considered formal evidence of significance (Knezevic 2008).

Assessing how multiple, simultaneous extreme conditions influence microbial diversity in human homes

We were interested in testing the hypothesis that interactive effects of multiple, simultaneously extreme environmental conditions are important determinants of microbial diversity in extreme home environments (Harrison et al. 2013). Our study included multiple samples with more than one environmental extreme (Supp. Table 1); however, we only had sufficient replication to assess this hypothesis for 2-way interactions between extremely high temperatures and extreme pH as well as high temperature and chemical environments. Because number of reads varied significantly among different environmental extremes, we could not use a standard 2-way ANOVA. Instead, we assessed these effects using an ordination framework.

We visualized the composition of bacteria and archaea from extreme habitats in homes using non-metric multidimensional scaling ordination (NMDS) in Primer-E v.7.0.9 with PERMANOVA +1 (Clarke & Gorley, 2015). To do this, we first constructed NMDS plots with 100 restarts and a Type I Kruskal fit scheme based on a Dissimilarity matrix of Bray-Curtis distances. To assess the relationship between temperature (extreme vs. intermediate) and the other extremes (pH: extreme vs. neutral; chemicals: extreme vs. none) for α -diversity of microbes, we conducted a permuted multivariate analysis of variance (PERMANOVA) test with

temperature class and either pH or chemical class and their interaction as factors, 9,999 iterations and Type III sums of squares. When interactions were significant (Anderson et al. 2008), we conducted pairwise PERMANOVA to determine which treatment combinations significantly differed from one another. Similarly, we assessed these relationships in terms of β -diversity using a permuted dispersion (PermDisp) test of a presence/absence matrix of OTU occurrences. When these tests were significant, we conducted pairwise tests of extreme vs. non-extreme chemical and pH environments in habitats with intermediate and extreme temperatures (thus 2 tests per treatment combination). Finally, we conducted SIMPER analyses for each significant treatment combination to determine the OTUs that contributed the most to pairwise between-group differences in ordination space. Because we conducted two separate analyses for each level of diversity, we accounted for the additional error associated with multiple tests, using a revised $\alpha=0.05/2=0.025$ as our cut-off for statistical significance for the results of each test. This conservative α is particularly important because we did not have equal sample sizes in all groups for these analyses, which can increase the risk of Type I error (Anderson & Walsh 2013).

Determining which microbial genera differentiate extreme home habitats from the rest of the home

We compared the occurrences of microbes in our samples to those reported in less extreme home environments (Dunn et al. 2013). Human-associated microbes were common to both datasets, and we were particularly interested in those taxa unique to our dataset, relative to the broader home (Dunn et al. 2013). Therefore, we removed human-associated OTU's from our dataset. We identified these human-associated OTU's using databases that identified human gut (Flores et al. 2014) and skin (Urban et al. 2016) microbiomes. OTU's that occurred in at least 80% of the

samples in those databases were considered human-associates and excluded from our analyses of the microbial diversity of extreme habitats in human homes. We then determined the identity of microbes that were absent from the broader homes dataset, but present in extreme environments and then tabulated the extreme habitat(s) in which they were present. Likewise, we identified the non-human associated microbes that were present in the broader home environment, but absent from all extreme environments in our samples.

Results and Discussion:

What is the relative diversity of microbes in extreme temperature, pH and chemical environments of southeast US homes and how does it compare to habitats without each extreme condition?

The cumulative diversity (OTU richness) in habitats with extreme temperatures was more than twice as high as in habitats with extreme pH (maximum of 73 vs. 33, Fig. 1) and almost three times as high as habitats with extreme chemical environments (27.6; Fig. 1). Habitats with extreme temperatures also had higher OTU richness than habitats with intermediate temperatures (Fig. 2a). Conversely, previous research indicates that the diversity in habitats with either extremely high or extremely low temperatures is generally low, and dominated by a small number of abundant bacterial species (Lewin et al. 2013). For example, Sharp et al. (2014) recently found that OTU richness in hydrothermal vents peaked at intermediate temperatures (24°C), with reduced OTU richness in extremely hot or cold environments (Sharp et al. 2014). We did not detect significant differences in the rarefied species richness of bacterial and archaeal microbes in extreme vs. neutral pH conditions; however, the marginally non-significant trend suggests that extreme pH environments also had higher microbial diversity than neutral habitats

(Fig. 2b). Recent studies have demonstrated that pH is a key predictor of microbial diversity in both extreme environments, such as acid mine drainage sites (Kuang et al. 2013), and less extreme environments, such as tropical soils (Tripathi et al. 2012). In both cases, habitats with neutral pH had higher microbial diversity than those with a pH higher or lower than neutral. Thus, we again found different patterns in extreme home environments compared to other studies comparing extreme and non-extreme habitats. One possible explanation for the difference between our findings and these recent studies is that human-associated microbes are present in home environments with intermediate temperatures. Perhaps these species are able to dominate habitats with intermediate, but not extreme, conditions. Alternatively, the lower diversity in habitats with intermediate temperatures and neutral pH in our study could be due to the occurrence of extreme conditions along different axes (e.g. intermediate temperature, but extreme pH or chemical habitats). We examine potential interactive effects of these polyextreme habitats in the next section.

In contrast, habitats with extreme chemicals had significantly lower accumulated OTU richness than did habitats without these extreme conditions (Fig. 2c). Extreme chemical environments are poorly studied and understood (Rothschild and Mancinelli 2001). Our data suggest that they could act as strong filters in extreme environments.

How do multiple, simultaneous extreme conditions influence microbial diversity in extreme home environments?

Many of the habitats in this study were characterized by more than one extreme environmental condition. Therefore, we also examined the potential for interactive effects of multiple, simultaneous extreme conditions on microbial diversity. Due to limited replication across all environmental extremes, we were only able to examine extreme pH and chemical

habitats with and without extreme temperatures. We used an ordination framework to examine these interactive effects (see methods).

We found significant interactions between extreme temperature and both extreme pH (PERMANOVA: Pseudo- $F_{1,82} = 2.53$, $P=0.0001$; Figure 3A) and extreme chemical (PERMANOVA: Pseudo- $F_{1,82} = 3.16$, $P=0.0001$; Figure 3C) environments for OTU composition. When temperatures were intermediate, there were no significant differences in microbial composition in extreme vs. neutral pH habitats (pairwise PERMANOVA: $t_{1,38} = 1.02$, $P = 0.40$). However, when temperatures were extreme, there was a very large difference between the composition of microbes in extreme pH habitats, compared to neutral habitats (pairwise PERMANOVA: $t_{1,38} = 1.70$, $P=0.0001$; Fig. 3A). The five genera that contributed the most to differences between these two habitat types (from SIMPER analysis) were *Parascardovia*, *Micrococcus*, *Rothia*, *Brachybacterium*, and an unknown genus from Sphingomonadaceae. Most of these genera are associated with humans (Oshima et al. 2015, Gueimonde et al. 2012, Kloos et al. 1975, Kocur et al. 2006, Vaccher et al. 2007, Uchibori et al. 2012). Sphingomonadaceae are widespread in aquatic habitats, including drinking water (Vaz-Moreira et al. 2011), but also other aquatic environments (e.g. tree holes-Xu et al. 2008). *Brachybacterium* is usually associated with marine environments (Ward and Boru 2006), including Antarctic sea ice (Junge et al. 1998). However, it was recently detected in an urban shopping center (Tringe et al. 2008). All of these genera were more common in habitats with extreme temperatures and neutral pH than they were in habitats with both extreme temperatures and extreme pH. While different houses had significantly different microbial composition (3-way PERMANOVA, $P=0.0001$), there were no significant 2-or 3-way interactions with house (Supp. Table 3).

The interaction between temperature and chemical extremes was similar. Microbial composition was indistinguishable between the habitats that only had one extreme condition-regardless of whether it was temperature or chemicals that were extreme. There were also no significant differences between habitats with neither extreme temperatures nor extreme chemical conditions and habitats that had a single extreme condition. However, habitats with both extreme temperatures and extreme chemicals had significantly different microbial composition compared to all other groups (pairwise PERMANOVA; $t_{1,38} = 1.75$, $P=0.0001$; Fig. 3C). The five genera that contributed the most to compositional difference between these two habitats (from SIMPER analysis) were *Methylobacterium*, an unknown genus of Moraxellaceae, *Sejonia*, an unknown genus of Sphingomonadaceae, and *Flavobacterium*. With the exception of the unknown genus of Moraxellaceae, which was more common in extreme chemical and temperature environments, all of these genera were more common in the habitats without temperature and chemical extremes. Moraxallaceae have been found in other extreme environments, including deep sea sediments (Maruyama et al. 1997). Although it was more common in our less extreme environments, *Sejonia* is better known from Antarctic ice (Yi et al. 2005). Sphingomonadaceae as described above are common to aquatic habitats. *Methylobacterium* is a widespread habitat generalist that is facultatively methyltrophic (Green 2006). Finally, *Flavobacterium* is common in freshwater and marine ecosystems but tends to flourish in cold environments with high salinity (Bernardet and Bowman 2006).

There were also significant differences in the β -diversity in home environments with more than one extreme condition. When temperatures were intermediate, there were no significant differences between neutral and extreme pH environments (Figure 4A; PermDisp: $P = 0.3864$). However when temperatures were also extreme, habitats with extreme pH conditions had

significantly higher β -diversity than those with neutral pH conditions (Figure 4A; PermDisp: $P=0.0014$). Similarly, at intermediate temperatures, there was a non-significant trend (Figure 4B; PermDisp: $P=0.03$, Bonferroni-corrected $\alpha = 0.025$) in which habitats without extreme chemicals present had higher β -diversity than those with extreme chemicals present. However, when temperatures were also extreme, habitats with extreme chemicals present had higher β -diversity than those without extreme chemicals (Figure 4B; PermDisp: $P = 0.0006$). This increase in β -diversity in extreme pH and chemical environments when temperatures were also extreme suggests that polyextreme conditions may support a higher diversity of extremophiles and/or reduced occurrences of numerically dominant genera compared to environments with a single extreme condition, at least among habitats (in contrast to within habitats). The 5 genera that contributed the most to differences in β -diversity between neutral and extreme pH conditions when temperatures were also extreme were: *Veillonella*, *Kocuria*, *Peptoniphilus*, *Parascardovia*, and *Anaerococcus*. Interestingly, these were also the top 5 genera contributing to differences between habitats with and without extreme chemicals that also had extreme temperatures. All of these genera were less common in habitats with 2 extremes than they were in habitats with only extreme temperatures. They are also genera that include human-associated species (Bhatti & Frank 2000, Fadda et al. 2001, Song et al. 2007, Gueimonde et al. 2012).

Which microbial genera differentiate extreme home habitats from the rest of the home?

After removing all human-associated microbes (see methods), there were a total of 241 unique genera in the broader homes dataset (Dunn et al. 2013). Our extreme samples contained 135 of the remaining broader homes genera, but ~44% of the genera found in the broader homes were absent from our extreme home samples (Supp. Table 4), the absence of which might simply be

318 due to the larger number of samples in Dunn et al. (2013). More interestingly, we found 20
 319 genera present among our samples that were absent from the broader homes dataset. Nine of
 320 these genera were found in all three categories of extreme environments (Table 1); one genus
 321 (*Solibacter*) was absent from habitats with extreme pH, but occurred in both extreme chemical
 322 and temperature environments. *Solibacter* is a common and abundant soil microbe, especially in
 323 tropical regions (Guan et al. 2013, Wang et al. 2015). There was also one genus
 324 (*Brevundimonas*) that was absent from extreme chemical environments, but present in both
 325 extreme temperature and extreme pH environments; *Brevundimonas* is one of the only genera
 326 thought to be able to survive the low temperatures and ionizing radiation on Mars (Dartnell et al.
 327 2010). There were three genera (*Azobacteroides*, *Elizabethkingia*, and *Xiphinematobacter*) that
 328 occurred in both extreme pH and chemical environments that were absent in extreme
 329 temperature environments. Both *Azobacteroides* and *Xiphinematobacter* are gut symbionts of
 330 invertebrates; *Azobacteroides* is commonly found inside the protozoan symbionts of termites
 331 (Noda et al. 2007), and *Xiphinematobacter* is an endosymbiont of nematodes (Vandekerckhove et
 332 al. 2000). In invertebrate guts these microbes likely experience extreme chemical and pH
 333 environments frequently, while being relatively protected from temperature stress.
 334 *Elizabethkingia* is a cosmopolitan genus, with species that are endosymbionts of mosquitoes
 335 (Kämpfer et al. 2011), and others that are pathogens of both humans (Ceyhan and Celik 2011)
 336 and frogs (Xie et al. 2009). There was one genus that was only found in extreme chemical
 337 environments (*Helcococcus*). Interestingly, members of the genus *Helcococcus* possess the
 338 ability to degrade detergents. In fact, the detergent Tween-80 can be added to media to enrich
 339 *Helcococcus* (Collins et al. 1993, Chagla et al. 1998). Finally, we found 5 genera (*Brochothrix*,
 340 *Buchnera*, *Polynucleobacter*, *Ralstonia*, and *Thermicanus*) unique to extreme temperature

environments. *Brochothrix* is a common spoilage bacterium in meat (Rattanasomboon et al. 1999). *Buchnera* is a widespread aphid endosymbiont (Shigenobu et al. 2000). Recently, a survey of homes in Raleigh, NC demonstrated that aphids could be quite common in human homes (Bertone et al. 2016), which could explain how this genus arrived in the homes in our study (via aphids in the home). The genus *Polynucleobacter* includes both free-living species and species that are endosymbionts of nematodes (Vannini et al. 2007). *Ralstonia metallidurans* is a bacterium specifically adapted to toxic metal environments (Mergeay et al. 2003). Other species of *Ralstonia* have been shown to be effectively controlled using high temperature treatments in commercial crops (Kongkiattikajorn and Thepa 2007). In our study, *Ralstonia* were collected in both high and low temperature environments. Finally, *Thermicanus* is, as its name suggest, a thermophilic bacterial genus (Wrighton et al. 2008).

Conclusions:

This study has provided a glimpse into the microbial diversity that lives in habitats of human homes similar in their extreme temperature, pH and chemical conditions to some of the most extreme habitats on Earth. We discovered that these conditions have lower diversity than the surrounding home environment; yet tens of bacterial lineages can be found in these extreme habitats of the human home, including many taxa with known associations with extreme conditions. Habitats with extreme temperatures alone appear to be able to support a greater diversity of microbes than habitats with extreme pH or extreme chemical environments alone. Microbial diversity is significantly lowest when habitats have both extreme temperature and one of these other extremes. Interestingly, environments in homes often alternate between periods of extreme and non-extreme conditions. For example, dishwashers are only likely to have extremely high temperatures while cleaning and drying dishes. This variability could lead to temporal shifts

in microbial composition, similar to those found for human vaginal microbes (Gajer et al. 2012). This variability may also explain the presence of human-associated generalist species in our samples. Future work, with samples taken before and after appliances (like many of those used in our study) are operated, could elucidate the importance of episodic extreme conditions for microbial communities in homes. Additionally, a key next step is understanding which of the relatively few species that are found in these poly-extreme environments in the home are metabolically active there and both whether these polyextreme taxa pose health threats (as was recently suggested by Gümral et al. 2015) and/or might be useful industrially.

Acknowledgements: We thank the homeowners for allowing us to sample habitats in their homes. Thanks also to Dr. Holly Menninger for her support and guidance in addressing logistical challenges presented by the current work. The Genomics Laboratory at the North Carolina Museum of Natural Sciences provided critical logistical support for this project. This work was funded by A.P. Sloan Microbiology of the Built Environment Program grant awarded to RRD.

References

- Anderson, MJ, Gorley RN, Clark, KR (2008) PERMANOVA+ for PRIMER: Guide to software and statistical methods. PRIMER-E: Plymouth, UK.
- Bates ST, Berg-Lyons D, Caporaso JG, Walters WA, Knight R, Fierer N (2011) Examining the global distribution of dominant archaeal populations in soil. ISME J 5: 908.
- Bertone MA, Leong M, Bayless KM, Malow TLF, Dunn RR, Trautwein MD (2016) Arthropods of the great indoors: characterizing diversity inside urban and suburban homes. *PeerJ* 4:e1582 <https://doi.org/10.7717/peerj.1582>.

- 386 Boe-Hansen R, Albechtsen H-J, Arvin E, Jorgensen Claus (2002) Bulk water phase and biofilm
387 growth in drinking water at low nutrient conditions. Water Research 36: 4477-4486.
- 388 Brock TD, Boylen KL (1973) Presence of thermophilic bacteria in laundry and domestic hot
389 water heaters. Applied Microbiology 25: 72-76.
- 390 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña
391 AG, Goodrich JK, Gordon J , Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,
392 Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh
393 PJ, Walters WA, Widmann J, Yatsunenko T,Zaneveld J, Knight R. (2010a) QIIME allows
394 analysis of high-throughput community sequencing data. Nature Methods 7: 335–336.
- 395 Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, and Knight R. (2010b)
396 PyNAST: a flexible tool for aligning sequences to a template alignment. Bioinformatics 26:
397 266–267.Ceyhan M and Celik M (2011) *Elizabethkingia meningosepticum*
398 (*Chryseobacterium meningosepticum*) infections in children. International Journal of
399 Pediatrics. doi:10.1155/2011/215237/
- 400 Chagla AH, Borczyk AA, Facklam RR, Lovgren M. (1998) Breast abscess associated with
401 *Helcococcus kunzii*. Journal of Clinical Microbiology 36:2377–2379.
- 402 Collins MD, Facklam RR, Rodrigues UM, Ruoff KL (1993) Phylogenetic analysis of some
403 Aerococcus-like organisms from clinical sources: description of *Helcococcus kunzii* gen.
404 nov., sp. nov. Int. J. Syst. Bacteriol. 43:425– 429.
- 405 Colwell RK (2013) *EstimateS*: Statistical estimation of species richness and shared species from
406 samples. Version 9. Persistent URL <purl.oclc.org/estimates>

- 407 Dartnell LR, Hunter SJ, Lovell KV, Coates AJ, Ward JM (2010) Low temperature ionizing
408 radiation resistance of *Deinococcus radiodurans* and Antarctic Dray Valley bacteria.
409 *Astrobiology* 7: 717-732.
- 410 DeSantis, TZ, Hugenholtz P, Larsen N, Rojas M , Brodie EL, Keller K, Huber T, Dalevi D, Hu
411 P, and Andersen GL. (2006) Greengenes, a Chimera-Checked 16S rRNA Gene Database
412 and Workbench Compatible with ARB. *Applied Environmental Microbiology* 72:5069-72.
- 413 Dunn RR, Fierer N, Henley JB, Leff JW, Menninger HL (2013) Home life: factors structuring
414 the bacterial diversity found within and between homes. *PLoS One*. DOI:
415 10.1371/journal.pone.0064133
- 416 Elleuche S, Schröder C, Sahm K, Antranikian G. (2014) Extremozymes-biocatalysts with unique
417 properties from extremophilic microorganisms. *Current Opinion in Biotechnology* 29: 116-
418 123.
- 419 Fierer N, Hamady M, Lauber CL, Knight R. (2008) The influence of sex, handedness, and
420 washing on the diversity of hand surface bacteria. *PNAS* 105:17994-17999.
- 421 Flores GE, Caporaso JG, Henley JB, Rideout JR, Domogala D, Chase J, Leff JW, Vásquez-
422 Baeza Y, Gonzalez A, Knight R, Dunn RR, Fierer N (2014) Temporal variability is a
423 personalized feature of the human microbiome. *Genome Biology* 15: art531
- 424 Gajer P, Brotman RM, Bai G, Sakamoto J, Schütte UME, Zhong X, Koenig SK, Fu L, Ma Z,
425 Zhou X, Abdo Z, Forney LJ, Ravel, J. (2012). Temporal Dynamics of the Human Vaginal
426 Microbiota. *Science Translational Medicine*, 4(132), 132ra52.
427 doi.org/10.1126/scitranslmed.3003605

- 428 Guan X, Wang J, Zhao H, Wang J, Luo X, Zhao F (2013) Soil bacterial communities shaped by
429 geochemical factors and land use in a less-explored area, Tibetan Plateau. BMC Genomics
430 14: 820 (13pp).
- 431 Gueimonde M, Bottacini F, van Sinderen D, Ventura M, Margolles A, Sanchez B (2012)
432 Genome sequence of *Parascardovia denticolens* IPLA 20019, isolated from human breast
433 milk. Journal of Bacteriology 194: 4776-4777.
- 434 Gümral R, Özhak-Baysan, Tümgör A, Saraçlı MA, Yidrian ŞT, Ilkit M, Zupančič J, Novak-
435 Babič M, Gunde-Cimerman N, Zalar P, de Hoog GS (2015) Dishwashers provide a
436 selective extreme environment for human-opportunistic yeast-like fungi. Fungal Diversity
437 10.1007/s13225-015-0327-8.
- 438 Harrison JP, Gheeraert N, Tsigelnitskiy D, Cockell CS (2013) The limits of life under multiple
439 extremes. Trends in Microbiology 21: 204-212.
- 440 Hoehler TM, Jorgensen BB. (2013) Microbial life under extreme energy limitation. Nature
441 Reviews Microbiology 11: 83-95.
- 442 Hulcr J, Latimer AM, Henley JB, Rountree NR, Fierer N, Lucky A, Lowman MD, Dunn RR
443 (2012) A jungle in there: Bacteria in belly buttons are highly diverse, but predictable. PLoS
444 One 7(11): e47712.
- 445 Junge K, Gosink JJ, Hoppe HG, Staley JT (1998) *Arthrobacter*, *Brachybacterium* and
446 *Planococcus* isolates identified from Antarctic sea ice brine. Systematic and Applied
447 Microbiology 21: 306-314.

- 448 Kalmbach S, Manz W, Szewzyk U (1997) Isolation of new bacterial species from drinking water
449 biofilms and proof of their in situ dominance with highly specific 16S rRNA probes.
450 Applied and Environmental Microbiology 63: 4164-4170.
- 451 Kämpfer P, Matthews H, Glaeser SP, Martin K, Lodders N, Faye I (2011) *Elizabethkingia*
452 *anopheles* sp. nov., isolated from the midgut of the mosquito, *Anopheles gambiae*.
453 International Journal of Evolutionary Microbiology. doi: 10.1099/ijes.o68147-0.
- 454 Kloos WE, Musselwhite MS (1975) Distribution and persistence of *Staphylococcus* and
455 *Micrococcus* species and other aerobic bacteria on human skin. Applied Microbiology 30:
456 381-395.
- 457 Knezevic A. (2008) Overlapping confidence intervals and statistical significance ([http://www.](http://www.cscu.cornell.edu/news/statnews/stnews73.pdf)
458 [cscu.cornell.edu/news/statnews/stnews73.pdf](http://www.cscu.cornell.edu/news/statnews/stnews73.pdf)). In: Cornell University, Cornell Statistical
459 Consulting Unit.
- 460 Kocur M, Koos WE, Schleifer KH (2006) The genus *Micrococcus*. Prokaryotes 3: 961-971.
- 461 Kongkiattikajorn J, Thepa S (2007) Increased tomato yields by heat treatment for controlling
462 *Ralstonia solanacearum* in soil. Proceedings of the 45th Kasetsart University Annual
463 Conference.
- 464 Kuang J-L, Huang L-N, Chen L-X, Hua Z-S, Li S-J, Shu, W-S (2013) Contemporary
465 environmental variation determines microbial diversity patterns in acid mine drainage.
466 International Society for Microbial Ecology 7: 1038-1050.

- 467 Lauber CL, Hamady M, Knight R, Fierer N (2009) Pyrosequencing-based assessment of soil pH
468 as a predictor of soil bacterial community structure at the continental scale. *Applied and*
469 *Environmental Microbiology* 75: 5111-5120.
- 470 Lewin A, Wentzel A, Valla, S (2013) Metagenomics of microbial life in extreme temperature
471 environments. *Current Opinion in Biotechnology* 24: 516-525.
- 472 Martin LJ, Adams RI, Bateman A, Bik HM, Hawks J, Hird SM, Hughes D, Kembel SW, Kinney
473 K, Kolokotronis SO, Levy G, McClain C, Meadow JF, Medina RF, Mhuireach G, Moreau
474 CS, Munshi-South J, Nichols LM, Palmer C, Popova L, Schal C, Taubel M, Trautwein M,
475 Ugalde JA, Dunn RR. (2015) Evolution of the indoor biome. *Trends in Ecology and*
476 *Evolution* 1909:1-10.
- 477 Maruyama A, Taniguchi R, Tanaka H, Ishiwata H, Higashihara T (1997) Low-temperature
478 adaptation of deep-sea bacteria isolated from the Japan Trench. *Marine biology* 128: 705-
479 711.
- 480 Mergeay M, Monchy S, Vallaeys T, Auquier V, Benotmane A, Bertin P, Taghavi S, Dunn J, van
481 der Lelie D, Wattiez R (2003) *Ralstonia metallidurans*, a bacterium specifically adapted to
482 toxic metals: towards a catalogue of metal responsive genes. *FEMS Microbiology Reviews*
483 27: 385-410.
- 484 Niehaus F, Bertolodo C, Kähler M, Antranikian G (1999) Extremophiles as a source of novel
485 enzymes for industrial application. *Applied Microbiology and Biotechnology* 51:711-729.
- 486 Noda S, Kitade O, Inoue T, Kawi M, Kanuka M, Hiroshima K, Hongoh Y, Constantino R,
487 UysV, Zhong J, Kudo T, Ohkuma M (2007) Cospeciation in the triplex symbiosis of

termite gut protists (*Pseudotrichonympha* spp.), their hosts, and their bacterial endosymbionts. *Molecular Ecology* 16:1257-1266.

Oshima K, Hayashi J, Toh H, Nakano A, Shindo C, Komiya K, Morita H, Honda K, Hattori M (2015) Complete genome sequence of *Parascardovia denticolens* JCM 12538^T, isolated from human dental caries. *Genome Announcements* 3(3): e00485-15.

Rattanasomboon N, Bellara SR, Harding CL, Fryer PJ, Thomas CR, Al-Rubeai M, McFarlane CM (1999) Growth and enumeration of the meat spoilage bacterium *Brochothrix thermosphacta*. *International Journal of Food Microbiology* 51: 145-158.

Rothschild LJ and Mancinelli RL (2001) Life in extreme environments. *Nature* 409: 1092-1101.

Sharp CE, Brady AL, Sharp GH, Grasby SE, Stott MB, Dunfield PF (2014) Humboldt's spa: microbial diversity is controlled by temperature in geothermal environments. *ISME Journal* 8: 1166-1174.

Szewzyk U, Szewzyk R, Manz W, Schleifer KH (2000) Microbiological safety of drinking water. *Annual Review of Microbiology* 54: 81-127.

Tringe SF, Zhang T, Liu X, Yu Y, Lee WH, Yap J, Yao F, Suan ST, Ing SK, Haynes M, Rohwer F, Wei CL, Tan P, Bristow J, Rubin EM, Ruan Y (2008) The airborne metagenome in an indoor urban environment. *PLoS One* 3(4): e1862.

Tripathi BM, Kim M, Singh D, Lee-Cruz L, Lai-Hoe A, Ainuddin, AN, Go R, Rahim RA, Husni MHA, Chun J, Adams JM (2013) Tropical soil bacterial communities in Malaysia: pH dominates in the equatorial tropics too. *Microbial Ecology* 64: 474-484.

- 508 Uchibori S, Tsuzukibashi O, Kobayashi T, Aida M (2012) Localization of the genus *Rothia* in
509 the oral cavity. International Journal of Oral-Medical Science 11: 207-210.
- 510 Urban JM, Fergus DJ, Savage, AM, Ehlers M, Menninger HL, Dunn RR, Horvath JE (2016) The
511 effect of habitual antiperspirant and deodorant product use on the armpit microbiome. PeerJ
512 4: e1605.
- 513 Valentine DL (2007) Adaptations to energy stress dictate the ecology and evolution of Archea.
514 Nature Reviews Microbiology 5:316-323.
- 515 Vandekerckhove, TTM, Willems A, Gillis M, Coomans A (2000) Occurrence of novel
516 verrucomicrobial species endosymbiotic and associated with parthenogenesis in *Xiphinema*
517 *americanum*-group species (Nematoda, Longidoridae). International Journal of Systematic
518 and Evolutionary Microbiology 50: 2197-2205.
- 519 van den Burg B (2003) Extremophiles as a source of novel enzymes. Current Opinion in
520 Microbiology 6: 213-218.
- 521 Vannini C, Pöckl M, Petroni G, Wu QL, Lang E, Stackebrandt E, Schrällhammer M, Richardson
522 PM, Hahn MW (2007) Endosymbiosis in *Statu nascendi*: close phylogenetic relationship
523 between obligately endosymbiotic and obligately free-living *Polynucleobacter* strains
524 (Betaproteobacteria). Environmental Microbiology 9: 347-359.
- 525 Vaz-Moreira I, Nunes OC, Manaia, CM (2011) Diversity and antibiotic resistance patterns of
526 *Sphingomonadaceae* isolates from drinking water. Applied and Environmental
527 Microbiology 77: 5697-5706.

- Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Applied And Environmental Microbiology* 73: 5261–5267.
- Wang N, Ding L-J, Xu H-J, Li H-B, Su J-Q, Zhu Y-G *In Press* Variability in responses of bacterial communities and Nitrogen Oxide emission to Urea fertilization among various flooded paddy soils. *FEMS Microbiology and Ecology*.
- Wrighton KC, Agbo P, Warnecke F, Weber KA, Brodie EL, DeSantis, TZ, Hugenholtz P, Andersen GL, Coates JD (2008) A novel ecological role of the Firmicutes identified in thermophilic microbial fuel cells. *ISME Journal* 2: 1146-1156.
- Yi H, Yoon HI, Chun J (2005) *Sejongia antarctica* gen. nov., sp. nov. and *Sejongia jeonii* sp. nov., isolated from the Antarctic. 55: 409-416.

Figure Captions:

Figure 1: OTU accumulation curves for each extreme environment, expressed as number of OTUs by number of reads from sequencing. Each curve was constructed using 1000 iterations, and the dotted lines represent 95% confidence intervals. Non-overlapping confidence intervals indicate that the accumulation curves are significantly different. Thus, habitats with extreme temperatures had significantly more accumulated species than habitats with either extreme pH or extreme chemical environments. However, the accumulated species in habitats pH and chemical extremes did not differ significantly.

Figure 2: Comparison of rarefaction curves between extreme and non-extreme habitats. (A) extreme vs. intermediate temperatures, (B) extreme vs. neutral pH environments, and (C) extreme chemicals present vs. absent. Rarefaction curves are expressed as number of OTU by

number of reads from sequencing. Each curve was constructed using 1000 iterations, and the dotted lines represent 95% confidence intervals. Significance tests were as described for Figure 1.

Figure 3: NMDS ordinations OTU occurrence by (A-B) Temperature & pH and (C-D) Temperature & chemical environments in the home. Large symbols represent centroids ± 1 SE (A, C), and small symbols represent each sample (B, D). The interaction between temperature and pH was significant (PERMANOVA: (pseudo)- $F_{1, 85} = 2.53$, $P(\text{perm}) = 0.0001$), as was the interaction between extreme temperature and chemical conditions (PERMANOVA: (pseudo)- $F_{1, 85} = 3.16$, $P(\text{perm}) = 0.0001$). The ordination was constructed with Bray-Curtis distances and 100 restarts; 2-D stress was 0.21. PERMANOVA analyses were conducted using type III sums of squares and 9,999 iterations.

Figure 4: Average distances between samples and centroids (β -diversity) across home environments that differ with respect to extreme temperatures and (A) extreme pH conditions & (B) extreme chemical conditions. Data were assessed using PermDisp; dispersion was significantly different across extreme temperatures and extreme pH conditions ($F_{3, 82} = 4.08$, $P=0.024$) and across extreme temperatures and extreme chemical conditions ($F_{3, 82} = 6.99$, $P=0.0017$). Post-hoc pairwise tests: * $P<0.025$, ** $P<0.01$, *** $P<0.001$.

Table Captions:

Table 1: Summary of occurrences of microbes that were present in samples from extreme home environments, but absent from the broader home samples. Numbers indicate the number of reads of each genus by extreme environment. The first group includes genera that were only present in one extreme environment, the second group includes genera that were present in two extreme

environments, and the last group includes genera that were present in all three extreme home environments.

Supplementary Tables and Figures:

Raw Data: Output file from QIIME at the genus level (L6), rarefied to 1000 reads.

Supp. Table 1: Description of sample locations. Standardized locations were sampled in all 6 houses, while special locations were only sampled in a subset of the houses (due to availability of samples across houses)

Supp. Table 2: Classifications of sampled extreme home environments based upon temperature, pH and chemical conditions.

Supp. Table 3: Results from PERMANOVAs testing (a) the effects of house ID on microbial composition; and (b) the effects of temperature class, pH class, and house ID on microbial composition.

Supp. Table 4: List of non-human associated microbes in extreme and non-extreme (Dunn et al. 2013) home habitats

Supp. Figure 1: Map of houses that were sampled for the study

Supp. Figure 2: NMDS plot with houses and sampling locations labeled.

Supp. Figure 3: Histograms depicting the % of reads from bacterial and archaeal classes in (a) habitats with extreme temperatures, (b) habitats with extreme pH, and (c) habitats with extreme chemicals.

Table 1 (on next page)

Table 1

Summary of occurrences of microbes that were present in samples from extreme home environments, but absent from the broader home samples. Each X indicates that the genus was found in a given extreme environment. t0h6\00>

Table 1: Summary of occurrences of microbes that were present in samples from extreme home environments, but absent from the broader home samples. Numbers indicate the number of reads of each genus by extreme environment. The first group includes genera that were only present in one extreme environment, the second group includes genera that were present in two extreme environments, and the last group includes genera that were present in all three extreme home environments.

Genus	Extreme Temperatures	Extreme pH	Extreme Chemical
<i>Brochothrix</i>	265	0	0
<i>Buchnera</i>	22	0	0
<i>Polynucleobacter</i>	33	0	0
<i>Ralstonia</i>	21	0	0
<i>Thermicanus</i>	34	0	0
<i>Helcococcus</i>	0	0	22
<i>Solibacter</i>	86	0	30
<i>Brevundimonas</i>	184	189	0
<i>Azobacteroides</i>	0	33	33
<i>Elizabethkingia</i>	0	25	24
<i>Xiphinematobacter</i>	0	19	21
<i>Azospira</i>	139	33	44
<i>Brachybacterium</i>	101	52	69
<i>Enhydrobacter</i>	452	387	408
<i>Gluconobacter</i>	23	21	22
<i>Oligella</i>	40	74	77
<i>Parascardovia</i>	141	46	107
<i>Photobacterium</i>	71	65	93
<i>Propionibacterium</i>	73	31	40
<i>Salinibacterium</i>	108	334	355

Figure 1

OTU accumulation curves for each extreme environment

OTU accumulation curves for each extreme environment , expressed as the number of OTUs by the number of reads from sequencing. Each curve was constructed using 1000 iterations, and the dotted lines represent 95% confidence intervals. Non-overlapping confidence intervals indicate that the accumulation curves are significantly different. Thus, habitats with extreme temperatures had significantly more accumulated species than habitats with either extreme pH or extreme chemical environments. However, the accumulated species in habitats pH and chemical extremes did not differ significantly.




Figure 2

Comparison of rarefaction curves between extreme and non-extreme habitats

Comparison of rarefaction curves between extreme and non-extreme habitats . (A) extreme vs. intermediate temperatures, (B) extreme vs. neutral pH environments, and (C) extreme chemicals present vs. absent. Rarefaction curves are expressed as number of OTU by number of reads from sequencing. Each curve was constructed using 1000 iterations, and the dotted lines represent 95% confidence intervals. Significance tests were as described for Figure 1.




Figure 3

NMDS ordinations OTU occurrence by (A-B) Temperature & pH and (C-D) Temperature & chemical environments in the home.

NMDS ordinations OTU occurrence by (A-B) Temperature & pH and (C-D) Temperature & chemical environments in the home. Large symbols represent centroids ± 1 SE (A, C), and small symbols represent each sample (B, D). The interaction between temperature and pH was significant (PERMANOVA: (pseudo)- $F_{1,85} = 2.53$, $P(\text{perm}) = 0.0001$), as was the interaction between extreme temperature and chemical conditions (PERMANOVA: (pseudo)- $F_{1,85} = 3.16$, $P(\text{perm}) = 0.0001$). The ordination was constructed with Bray-Curtis distances and 100 restarts; 2-D stress was 0.21. PERMANOVA analyses were conducted using type III sums of squares and 9,999 iterations.




Figure 4

Average distances between samples and centroids (β -diversity) across home environments

Average distances between samples and centroids (β -diversity) across home environments that differ with respect to extreme temperatures and (A) extreme pH conditions & (B) extreme chemical conditions. Data were assessed using PermDisp; dispersion was significantly different across extreme temperatures and extreme pH conditions ($F_{3,82} = 4.08$, $P=0.024$) and across extreme temperatures and extreme chemical conditions ($F_{3,82} = 6.99$, $P=0.0017$). Post-hoc pairwise tests: * $P<0.025$, ** $P<0.01$, *** $P<0.001$.

