

Variation in energy intake of female chimpanzees: comparing estimates based on food item versus energy ingestion rates

Moreen Uwimbabazi¹, Richard Wrangham^{2,3}, Zarin Machanda^{2,3}, Nancy Conklin- Brittain², Jessica M. Rothman^{4,5}, Gilbert Basuta¹

¹Department of Zoology, Makerere University, ²Department of Human Evolutionary Biology, Harvard University, ³Kibale Chimpanzee Project, ⁴Department of Anthropology, Hunter College of the City University of New York, ⁵New York Consortium in Evolutionary Primatology

Abstract

Different food items can provide calories at different rates, yet nutritional studies of primates are generally based on the proportion of time spent eating different foods. Accordingly, estimates of energy intake are potentially wrong. To assess the importance of this problem we observed 15 female chimpanzees from the Kanyawara chimpanzee community in Kibale National Park, Uganda for 15 months. Feeding rates were recorded and nutritional analyses assembled for 90 food types. Across all foods the mean estimated rate of energy ingestion was 8.8 ± 7.6 Kcal/minute. Energy ingestion rates varied among food types ($F_{8,82}=2.24$, $p<0.05$), being higher for ripe fruits (12.2 ± 8.7 Kcals/min) than for young leaves (7.8 ± 7.9 Kcals/minute), piths (6.0 ± 5.2 Kcal/min) or seeds (2.8 ± 1.9 Kcal/min). Energy ingestion rates estimated assuming high %NDF fermentation (54.3%) were significantly higher than the rates at zero (0%) fermentation ($t=6.14$, $p<0.01$). At 54.3% NDF fermentation, estimated energy ingestion rate rose by 20.5% for ripe fruits (9.7 to 12.2 kcal/min), and by 39.7% for young leaves (4.7 to 7.8 kcal/min). Chimpanzees spent a mean of 314 ± 89 minutes feeding daily, indicating that daily energy intake averaged to 2433 ± 1221 Kcal. When chimpanzees spent >60% of feeding time on ripe fruits ($N=74$ days), the average energy intake (2494 Kcal) was significantly higher than when they spent >60% of feeding time on herbaceous vegetation (1774 Kcal, $N=39$ days) ($F_{1,111}=9.20$, $p<0.01$). Our results highlight the value of using feeding rates to understand chimpanzee nutrition, and suggest that seasonal fluctuation in nutrient intake is more pronounced than indicated by variation in feeding time.

Key words: feeding rates, feeding time, food quality, energy intake