

A peer-reviewed version of this preprint was published in PeerJ on 22 June 2016.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.2042) (peerj.com/articles/2042), which is the preferred citable publication unless you specifically need to cite this preprint.

Shields MW, Tompkins J, Saville DJ, Meurk CD, Wratten S. 2016.
Potential ecosystem service delivery by endemic plants in New Zealand vineyards: successes and prospects. PeerJ 4:e2042
<https://doi.org/10.7717/peerj.2042>

Potential ecosystem service delivery by endemic plants in New Zealand vineyards: successes and prospects

Morgan Shields, Jean-Marie Tompkins, David J Saville, Colin D Meurk, Stephen Wratten

Vineyards worldwide occupy over 7 million hectares and are typically virtual monocultures, with high and costly inputs of water and agro-chemicals. Understanding and enhancing ecosystem services can reduce inputs and their costs and help satisfy market demands for evidence of more sustainable practices. In this New Zealand work, low-growing, endemic plant species were evaluated for their potential benefits as Service Providing Units (SPUs) or Ecosystem Service Providers (ESPs). The services provided were weed suppression, conservation of beneficial invertebrates, soil moisture retention and microbial activity. The potential Ecosystem Dis-services (EDS) from the selected plant species by hosting the larvae of a key vine moth pest, the light-brown apple moth (*Epiphyas postvittana*), was also quantified. Questionnaires were used to evaluate winegrowers' perceptions of the value of and problems associated with such endemic plant species in their vineyards. Growth and survival rates of the 14 plant species, in eight families, were evaluated, with *Leptinella dioica* (Asteraceae) and *Acaena inermis* 'purpurea' (Rosaceae) having the highest growth rates in terms of area covered and the highest survival rate after 12 months. All 14 plant species suppressed weeds, with *Leptinella squalida*, *Geranium sessiliflorum* (Geraniaceae), *Hebe chathamica* (Plantaginaceae), *Scleranthus uniflorus* (Caryophyllaceae) and *L. dioica*, each reducing weed cover by > 95%. Plant species also differed in the diversity of arthropod taxa that they supported, with the Shannon Wiener diversity index (H') for these arthropods ranging from 0 to 1.3. *G. sessiliflorum* and *Muehlenbeckia axillaris* (Polygonaceae) had the highest invertebrate diversity. Density of spiders was correlated with arthropod diversity and *G. sessiliflorum* and *H. chathamica* had the highest densities of these arthropods. Several plant species led to higher soil moisture content than in control plots. The best performing species in this context were *A. inermis* 'purpurea' and *Lobelia angulata* (Lobeliaceae). Soil beneath all plant species had a higher microbial activity than in control plots, with *L. dioica* being highest in this respect. Survival proportion to the adult stage of the moth pest, *E. postvittana*, on all plant species was poor (<0.3). When judged by a ranking combining multiple criteria, the most promising plant species were (in decreasing order) *G. sessiliflorum*, *A. inermis* 'purpurea', *H. chathamica*, *M. axillaris*, *L. dioica*, *L. angulata*, *L. squalida* and *S. uniflorus*. Winegrowers surveyed said

that they probably would deploy endemic plants around their vines. This research demonstrates that enhancing plant diversity in vineyards can deliver SPUs, harbour ESPs and therefore deliver ES. The data also shows that growers are willing to follow these protocols, with appropriate advice founded on sound research.

1 Potential ecosystem service delivery by endemic plants in New Zealand vineyards: successes
2 and prospects

3 Morgan W. Shields¹, Jean-Marie L. Tompkins², David J. Saville³, Colin Meurk⁴, Stephen D.
4 Wratten¹

5 ¹ Bio-Protection Research Centre, Lincoln University, Lincoln, Canterbury, New Zealand

6 ² Environment Canterbury, Lincoln, Canterbury, New Zealand

7 ³ Saville Statistical Consulting Limited, Lincoln, Canterbury, New Zealand

8 ⁴ Landcare Research, Lincoln, Canterbury, New Zealand

9 Corresponding author:

10 Morgan Shields

11 Corner of Springs Rd & Ellesmere Junction Rds, P O Box 85084, Lincoln, Canterbury, 7647,
12 New Zealand

13 Email address: morgan.shields@lincolnuni.ac.nz

14 Abstract

15 Vineyards worldwide occupy over 7 million hectares and are typically virtual monocultures, with
16 high and costly inputs of water and agro-chemicals. Understanding and enhancing ecosystem
17 services can reduce inputs and their costs and help satisfy market demands for evidence of more
18 sustainable practices. In this New Zealand work, low-growing, endemic plant species were
19 evaluated for their potential benefits as Service Providing Units (SPUs) or Ecosystem Service
20 Providers (ESPs). The services provided were weed suppression, conservation of beneficial
21 invertebrates, soil moisture retention and microbial activity. The potential Ecosystem Dis-services
22 (EDS) from the selected plant species by hosting the larvae of a key vine moth pest, the light-
23 brown apple moth (*Epiphyas postvittana*), was also quantified. Questionnaires were used to
24 evaluate winegrowers' perceptions of the value of and problems associated with such endemic
25 plant species in their vineyards. Growth and survival rates of the 14 plant species, in eight
26 families, were evaluated, with *Leptinella dioica* (Asteraceae) and *Acaena inermis* 'purpurea'
27 (Rosaceae) having the highest growth rates in terms of area covered and the highest survival rate
28 after 12 months. All 14 plant species suppressed weeds, with *Leptinella squalida*, *Geranium*
29 *sessiliflorum* (Geraniaceae), *Hebe chathamica* (Plantaginaceae), *Scleranthus uniflorus*
30 (Caryophyllaceae) and *L. dioica*, each reducing weed cover by > 95%. Plant species also differed
31 in the diversity of arthropod taxa that they supported, with the Shannon Wiener diversity index
32 (H') for these arthropods ranging from 0 to 1.3. *G. sessiliflorum* and *Muehlenbeckia axillaris*
33 (Polygonaceae) had the highest invertebrate diversity. Density of spiders was correlated with
34 arthropod diversity and *G. sessiliflorum* and *H. chathamica* had the highest densities of these
35 arthropods. Several plant species led to higher soil moisture content than in control plots. The
36 best performing species in this context were *A. inermis* 'purpurea' and *Lobelia angulata*
37 (Lobeliaceae). Soil beneath all plant species had a higher microbial activity than in control plots,
38 with *L. dioica* being highest in this respect. Survival proportion to the adult stage of the moth

39 pest, *E. postvittana*, on all plant species was poor (<0.3). When judged by a ranking combining
40 multiple criteria, the most promising plant species were (in decreasing order) *G. sessiliflorum*, *A.*
41 *inermis* ‘*purpurea*’, *H. chathamica*, *M. axillaris*, *L. dioica*, *L. angulata*, *L. squalida* and *S.*
42 *uniflorus*. Winegrowers surveyed said that they probably would deploy endemic plants around
43 their vines. This research demonstrates that enhancing plant diversity in vineyards can deliver
44 SPUs, harbour ESPs and therefore deliver ES. The data also shows that growers are willing to
45 follow these protocols, with appropriate advice founded on sound research.

46 1. Introduction

47 Biodiversity and ecosystem-function relationships are a key component of agroecology, and
48 agriculturalists need help to understand how to deploy and manage functional diversity in the
49 most appropriate ways. A key question in agroecology is the extent to which ecosystem services
50 (ES) can be quantified and enhanced (MEA, 2005; Mooney, 2010; Allan et al., 2015; Sandhu et
51 al., 2015; Sandhu et al., 2016). ES are defined as goods and services such as biological control
52 that provide the foundation for sustainaning human life on Earth (Wratten et al., 2013). The
53 pathway for ES delivery includes the Service Providing Unit (SPU), defined as a the smallest
54 unit, population or community that provides ES or will provide it in the future, within a given
55 area (Luck, Daily & Ehrlich, 2003). An Ecosystem Service Provider (ESP) is defined as the
56 species, foodweb, habitat or system that faciliates and supports the provision of ES by an SPU
57 (Kremen, 2005). For example, a strip of flowering buckwheat, *Fagopyrum esculentum* Moench.
58 and the natural enemies which it supports can deliver multiple ES, including enhanced biological
59 control of insect pests (Scarratt, Wratten & Shishehbor, 2008).

60 Enhancing ES, SPU and ESPs may be achieved by a better understanding of how biodiversity
61 and its functions may contribute to reduced variable costs, sustainable agricultural production,
62 agro-ecotourism and wellbeing, among others (Wratten et al., 2013). Biodiversity delivers
63 ecosystem functions (Mooney & Ehrlich 1997; Swift & Anderson 2012) and many of these
64 functions have value for humans, thus becoming ES (Cardinale et al., 2012; Mace, Norris &
65 Fitter, 2012). The value of ES is increasingly being used to justify the incorporation of
66 biodiversity into farming practices (Fiedler, Landis & Wratten, 2008; Tschardt et al., 2012;
67 Tuck et al., 2014; Barral et al., 2015). *In situ* plant conservation continues to have a key role
68 (Keesing & Wratten, 1997) but with accelerating global biodiversity loss, policies and practices
69 which enhance biodiversity in agricultural landscapes are increasingly important (Wratten et al.,
70 2013). In that context, the provision of benefits by non-crop, low-growing, endemic New Zealand
71 plants is quantified here and prospects for end-user adoption are assessed.

72 Worldwide, vineyards occupy over 7 million hectares (The Wine Institute, 2012). Typically they
73 are virtual monocultures of *Vitis vinifera* L. with bare earth or mown ryegrass (*Lolium perenne*
74 L.) between the rows and sometimes with other naturally occurring species as weeds (Nicholls,
75 Altieri & Ponti, 2008). Ryegrass and forb plants are also sometimes deliberately sown below
76 vines, as in some organic vineyards (Reeve et al., 2005). It is well established that deployment of
77 non-native biodiversity in vine inter-rows can enhance at least one ES, that of pest biocontrol
78 (Berndt, Wratten & Hassan, 2002; Scarratt, Wratten & Shishehbor, 2008) but vegetation endemic
79 to the country involved may provide a wider range of ecosystem derived benefits, including
80 reduced soil erosion from increased ground cover and soil moisture (Ramos, Benito & Martínez-
81 Casasnovas, 2015), conservation and eco-tourism, as well as cultural values (Fieldler et al.,
82 2008). Here, experimental field work investigated the potential of 13 endemic and one native

83 plant species to provide ES in vineyards. For the purposes of this study, all the selected plant
84 species are termed 'endemic'.

85 To evaluate the usefulness and benefits to growers of this approach, winegrowers were sent a
86 questionnaire to elicit their perceptions of the barriers they face to deploy low-growing plants in
87 vineyards. These data provided the study not only with future research directions but also
88 practical insights on how best to achieve grower uptake. This socio-ecological aspect is a crucial
89 step so that the pathway for agroecology research is complete so that it is more likely to be
90 accepted (Warner, 2006).

91 **2. Materials and methods**

92 *2.1 Field experiment*

93 The trial was located in the Waipara region, North Canterbury, New Zealand (E2489521:
94 N5782109, altitude: 76 m) within the rows of grapevines (cv. Pinot Noir; 2.3 m inter-row width).
95 Mean annual rainfall at the site was 684 mm, mean January (summer) temperature was 23 °C and
96 soil type was Glasnevin silty loam (Jackson & Schuster, 2002). The field work, begun in October
97 2007, was a randomised complete block design comprising ten blocks, each with one replicate of
98 15 treatments. Each treatment comprised of 14 selected plant species and a control. The latter was
99 maintained as bare earth by hand weeding. Such a control was used because in conventional
100 viticulture worldwide, normal weed management practice comprises prophylactic use of
101 herbicides under vines. The work carried out here was conducted in a conventional vineyard,
102 therefore the control treatment comprised regular weed removal. Each block consisted of four
103 rows, each with 12 individual vines. Each experimental plot had two individual plants of one
104 species of the selected plants (or no plants in the case of the control): one on either side of a vine,

about 30 cm from the trunk, arranged along the irrigation drip line. Replicates were separated by two vines in each row and vines were 1.5 m apart. Within-row management consisted of hand weeding in all plots every 2 weeks or when required prior to the weed suppression assessment. Inter-row management consisted of mowing the perennial ryegrass (*L. perenne*) every 2 weeks. The whole experiment comprised an area of the vineyard which was allocated by the company. No further space was available so plot size had to be restricted to the area around a single vine. Although this has implications for invertebrates moving between treatments, the latter were separated by two vines within a row and by an inter-row distance of 2.3 m, the latter comprising dense *L. perenne*. Table 1 lists plant species used in the trial and indicates the ecosystem services which were delivered or had potential for delivery.

Table 1 about here

2.2 New Zealand plant species tested

Plant species were selected based on their growth habit (1–15 cm in height) to minimise interference with vine management. Species were further selected based on their shallow roots, floral characteristics and tolerance to frost, exposure, sun, drought and disturbance as well as practicalities such as cost and availability. All selected plants apart from *Muehlenbeckia axillaris* (Hook.f.) Endl. (also native to Australia) were New Zealand endemic species and all were perennial. Successful growth and survival of the plants were seen as prerequisites for their ability to provide benefits to the vineyard operation. Consequently, these parameters were assessed 6, 12 and 24 months after planting.

2.3 Weed suppression

In September 2008, 11 months after planting, hand weed removal was stopped in five selected blocks where the weed suppression assay was occurring. Normal vineyard management

prevented the cessation of weeding in the other five blocks, so they were excluded from this part of the overall experiment. In December 2008, 14 months after planting, weed suppression by the plants was assessed visually by placing a 20 cm × 20 cm quadrat over them and over the corresponding area in the control plots (where none of the selected plants were planted) in the five selected blocks. Percentage cover of the study plants and weeds was recorded. *Disphyma australe* (subsp. Australe) Aiton, *Muehlenbeckia ephedroides* Hook.f. and *Raoulia subsericea* Hook.f. were not assessed due to their poor condition, growth and survival. Data were statistically analysed using a randomised block analysis of variance (ANOVA), followed by the unprotected Least Significant Difference (LSD) procedure at $P = 0.05$ (Saville, 1990).

2.4 Invertebrate biodiversity conservation

In August 2008 and January and March 2009 (10, 15 and 17 months, respectively, after establishment of the plants) under-vine treatments were assessed for invertebrate diversity and abundance using a suction sampler (Arnold, 1994). In August 2008, samples were taken from the 14 plant treatments, the control and from the mid-point of inter-row areas (predominantly *L. perenne*) adjacent to the experimental plots in each of the ten blocks. The sampler was set on maximum power for 10 seconds, within which time an area of 0.04 m² was sampled at each location. Collected invertebrates were stored in 70 % ethanol before being brought to the laboratory for sorting and identification. Due to taxonomic limitations, individuals were assigned to RTUs (recognisable taxonomic unit) for statistical analysis of diversity and abundance. For the second and third sampling dates, *R. subsericea*, *M. ephedroides* and *D. australe* were not sampled because of their poor growth and survival. The Shannon Wiener diversity index (H') was used because it takes into account evenness and species richness (Magurran, 1988). Spiders are key predators of vineyard pests (Thomson & Hoffman, 2007), therefore spider density was analysed

separately. Data were statistically analysed using a randomised block ANOVA, followed by the unprotected LSD procedure at $P = 0.05$.

2.5 Soil quality

The effect of the plant species on soil moisture and microbial activity was assessed. Due to resource constraints, only six plant species (those with the greatest growth and survival) were assessed. These were *Geranium sessiliflorum* Simpson et Thomson, *Hebe chathamica* Cockayne et allan, *Leptinella dioica* Hook.f., *M. axillaris* and *Lobelia angulata* G. Forst. Control plots (bare earth) were also assessed.

To assess the effects of the treatments on soil microbial activity, the TCC method (see Alef & Nannipieri, 1995) was employed. This measures the rate of reduction of triphenyltetrazolium chloride (TTC) to triphenyl formazan (TPF) (Alef & Nannipieri, 1995). It is a non-specific enzyme assay which determines the dehydrogenase activity in the soil and thereby indicates one aspect of soil microbial activity. In December 2008, soil samples were taken from below the five plant species listed above and the control plots in the five randomly selected blocks used in Section 2.3. Within each plot, three 50 g subsamples of soil were collected at a depth of approximately 12 cm from around the selected plants' roots, or within the corresponding area in the control plot, were combined to make a 150 g soil sample per plot. These 150 g soil samples were kept at 4 °C, before being assessed for microbial activity on the following day using the TTC method. The soil sampling method used above, was repeated in December 2008 and September and November 2009 for determination of soil moisture percentage. In the above six plant and control treatments, this was calculated using a gravimetric method and expressed on a dry weight basis (Topp, Parkin & Ferré, 1993). Data for both soil parameters were statistically analysed using a randomised block ANOVA, followed by the unprotected LSD procedure.

2.6 Pest development and longevity on candidate plants

The larval development of *E. postvittana* on the vegetative parts of the plant species was recorded in a laboratory bioassay. Species supporting high larval development rates could potentially exacerbate pest problems in the vineyard by acting as a suitable host. However, there is also the possibility that these species could act as trap plants (Khan et al., 2008). Ten treatments including nine of the selected under-vine plant species and an artificial diet (Shorey & Hale, 1965) were tested. Some plant species were not included in this bioassay as they had poor growth and/or survival in the field trial and were unlikely to be considered suitable for vineyard deployment; they were *M. ephedroides*, *R. subsericea* and *D. australe*. Others were excluded because another species or sub-species of the same genus was included in the bioassay; these were *Leptinella squalida* Hook.f. and *Acaena inermis* Hook.f. Six newly emerged (<24 h) first-instar larvae were placed in each of six Petri dishes (15 × 120 mm) in each of ten treatments. Treatments comprised freshly cut plant material with shoots inserted into an Eppendorf tube filled with water. Each tube was placed in a Petri dish which was sealed with plastic food wrap to prevent larval escape. After 7 days, plant material was examined and water changed or the plant replaced as necessary. The artificial diet treatment consisted of cut squares of the diet substrate on which first instar-larvae were placed. There were six replicates of each treatment (a total of 6 × 6 = 36 larvae per treatment), arranged in a randomised block design under a 16:8 L/D photoperiod at 20 °C ±3. The number of larvae surviving to each development stage (second instar, third instar, final instar, pupa and adult) was recorded. A generalised linear model with a binomial distribution was used to determine the effect of treatment and development stage on *E. postvittana* survival.

2.7 A questionnaire to winegrowers

Experimental work on ecosystem services enhancement in agriculture is of limited practical value unless agriculturalists are provided with ESPs (Kremen, 2005) or similar to facilitate growers' adopting the work. To assess the likelihood of the latter, a questionnaire was mailed to 56 Waipara vineyard operators. Growers were asked "Which of the following uses of endemic plants would you consider adopting?" (see Section 3.7). Growers were also asked "To what extent do the following factors lead you NOT to use endemic plants in or around your vineyard in the above ways?" (see Section 3.7). This information was used to ensure that recommendations to growers were feasible and to identify future research directions.

3. Results

3.1 Growth and survival of the selected plants

Significant differences in coverage (difference from that of the initially planted area) between plant treatments were found after 6 and 12 months (Table 2). *L. dioica* and *A. inermis* 'purpurea' showed greatest growth after 12 months while *Anaphalioides bellidioides* Glenny, *M. ephedroides*, *R. subsericea* and *D. australe* had little or no growth. After 24 months, survival remained high ($\geq 90\%$) for *M. axillaris*, *L. dioica*, *Raoulia hookeri* Allan var. *hookeri*, *A. inermis* 'purpurea' and *G. sessiliflorum* while that of other plants had begun to decline.

Table 2 about here

3.2 Weed suppression

There was significantly more weed growth in the control compared to all plant treatments ($P < 0.05$) (Figure 1). *L. squalida*, *G. sessiliflorum*, *H. chathamica*, *Scleranthus uniflorus* P.A. Will and *L. dioica* had the most weed suppression (Figure 1). Weeds consisted primarily of *Trifolium* spp. (Fabaceae) but also included Poaceae, Malvaceae and Asteraceae families.

218

Figure 1 about here

219 3.3 Invertebrate biodiversity conservation

220 At all sampling dates there was a significant effect of treatment on invertebrate diversity and
 221 there was greater overall abundance in the summer (January and March) than in winter (August)
 222 (Table 3). A total of 3133 invertebrate individuals from 16 taxa were collected over all the
 223 sampling dates. During summer (January and March 2009), Hemiptera (1936 individuals),
 224 Araneae (203) and Formicidae (175) were the most abundant taxa. In winter (August 2008),
 225 Araneae (72), Diplopoda (54) and Diptera (37) were the dominant taxa.

226 During early summer (January 2009), *M. axillaris*, *G. sessiliflorum*, *A. bellidioides*, *L. dioica*, *L.*
 227 *squalida*, *L. angulata*, *A. inermis* and *R. hookeri* had significantly higher diversity than either of
 228 the controls ($P < 0.05$) (Table 3). In late summer (March 2009), *M. axillaris*, *G. sessiliflorum*, *A.*
 229 *inermis* ‘purpurea’ and *L. angulata* had significantly greater diversity than the ryegrass inter-row
 230 control ($P < 0.05$), while these and *A. inermis*, *H. chathamica*, *A. bellidioides*, *L. squalida* and *L.*
 231 *dioica* had significantly higher diversity indices than the bare earth control ($P < 0.05$) (Table 3).
 232 In winter (August 2008), *G. sessiliflorum*, *H. chathamica*, *A. bellidioides*, *A. inermis* ‘purpurea’,
 233 *L. dioica*, *M. axillaris* and *L. squalida* had significantly higher invertebrate diversity than either
 234 of the controls (bare earth and ryegrass inter-row treatments) ($P < 0.05$) (Table 3).

235

Table 3 about here

236 A significant effect of treatment on spider density was found for all sampling dates, with highest
 237 spider abundance in March 2009 (Table 4). Spider density was significantly correlated with
 238 arthropod diversity on the August and March sampling dates.

239

Table 4 about here

240 *G. sessiliflorum* and *H. chathamica* consistently had the highest densities of spiders. *A. inermis*
241 ‘*purpurea*’, *A. bellidioides*, *L. angulata* and *M. axillaris* also had significantly higher spider
242 densities than did the bare earth control treatment on at least one of the sampling dates.

243 Spider families included web-building spiders including members of the Theridiidae
244 (Sundervall), Linyphiidae (Blackwall), Agelenidae (Koch) and Amaurobiidae (Thorell) families.
245 Wandering/hunting spider families included Oxyopidae (Thorell), Salticidae (Blackwall),
246 Gnaphosidae (Pocock), Clubionidae (Wagner) and Pisauridae (Simon).

247 3.4 Soil quality – moisture and microbial activity

248 3.4.1 Soil moisture

249 Soil moisture in the bare earth control treatment was low relative to the other treatments on all
250 three sampling dates (Table 5). In September and November 2009, it was also low under the *L.*
251 *dioica* treatment. In November 2009, it was significantly higher below *L. angulata* and *A.*
252 *inermis* ‘*purpurea*’ compared to all other treatments ($P < 0.05$) (Table 5).

253

Table 5 about here

254 3.4.2 Soil microbial activity

255 Microbial activity in December 2008 was higher in all the plant treatments compared to the bare
256 earth control, while it was significantly higher beneath *L. dioica* compared to that under the other
257 plant treatments ($P < 0.05$) (Table 5). Although soil moisture may influence microbial activity, it
258 was very low in all treatments at the time of microbial activity assessment.

3.5 Development of *E. postvittana* larvae on the selected plant species

There was a significant effect of plant species ($P < 0.001$) and the larval instar reached ($P < 0.001$) on survival of the pest *E. postvittana*, but there was no significant interaction between treatment and instar ($P = 0.99$) (Figure 2). Survival across all stages was significantly higher on the diet than on any of the plant species used, suggesting that the selected plants provided below-optimal nutrition to *E. postvittana*. *E. postvittana* larval survival was significantly higher on *A. inermis* 'purpurea' than on any of the other tested plants. The other species in order of decreasing pest survival were *G. sessiliflorum*, *L. angulata*, *R. hookeri*, *L. dioica*, *M. axillaris*, *S. uniflorus*, *A. bellidioides* and *H. chathamica*. In the case of *H. chathamica*, no pest larvae survived to the adult stage.

Figure 2 about here

3.6 Overall ranking of endemic plant species

In Table 6, the 14 plant species are ranked for each of the characteristics summarised in Tables 2 to 5 and Figures 1 to 2. For most characteristics, the plant species with the highest mean value is assigned the rank of 1. However, for weed suppression and leafroller (pest) survival, a rank of 1 is assigned to the species that had the fewest weeds or had the lowest pest survival.

Table 6 about here

Some plant species were not evaluated for all characteristics, often because they had already been judged unsuitable. Only six species were assessed in all respects (Table 6). None of these was consistently the best. For example, *L. dioica* ranked first for growth, survival and microbial activity, but ranked 10 out of 11 for spider density, and 7 out of 11 for invertebrate diversity. By

280 comparison, *G. sessiliflorum* ranked first for weed suppression, invertebrate diversity and spider
281 density, but ranked 8 out of 9 for leafroller (pest) survival.

282 When judged by an overall ranking, the most promising plant species were (in decreasing order)
283 *G. sessiliflorum*, *A. inermis* ‘*purpurea*’, *H. chathamica*, *M. axillaris*, *L. dioica* and *L. angulata*,
284 with average ranks ranging from 3.8 to 5.0, respectively (Table 6). None of the other eight plant
285 species averaged a rank of 5.0 or more, when their ranks were averaged over the characteristics
286 for which they had been assessed.

287 3.7 Winegrower questionnaires

288 The survey response rate was 30 out of 56 growers (Table 7). The majority of respondents (who
289 had not already adopted endemic plants for any purpose) indicated that they would ‘definitely’ or
290 ‘maybe’ deploy endemic plants around or within their vineyard properties for the various uses
291 presented to them. Currently, the conservation of flora and fauna are the primary uses of endemic
292 plants within respondents’ properties and they stated that such plants are also likely to be
293 established for erosion control, enhancement of pest biological control or for weed suppression.

294 *Table 7 about here*

295 Growers were also asked to indicate whether certain factors had led them not to deploy endemic
296 plants for the uses listed above (Table 8). These, which may be seen as barriers to establishing
297 such plants for the various uses, included a lack of knowledge, cost of initial investment, risk,
298 disruption to normal practices or having no interest in such practices (Table 8). For most endemic
299 plant uses, the primary concern of growers was the initial investment required. Notably, however,
300 a lack of knowledge surrounding the use of such plants to suppress weeds beneath vines was
301 cited by an almost equal number of growers as was the barrier of initial investment. Risk was a

barrier cited by nearly half the growers for establishing endemic vegetation for conservation of flora and fauna. Risk was also stated by a significant proportion of growers as cause for not utilising endemic plants for marketing purposes.

Table 8 about here

4. Discussion

Findings here suggest the selected endemic plants deployed beneath vines have the potential to improve pathways to ES provision (i.e., SPU, ESP and ES themselves) ultimately improving value to growers. Overall, certain endemic plant species may preserve biodiversity, enhance biological control of vineyard pests, provide weed suppression and improve soil health. Clearly further research is required, such as repeating the trial in different regions. In the first trial described in this paper, however, the most promising plant species were *G. sessiliflorum*, *A. inermis* ‘purpurea’, *H. chathamica*, *M. axillaris*, *L. dioica* and *L. angulata*.

4.1 Weed suppression

Management of weeds is a major concern of vineyard managers as these plants can compete with the vines’ surface ‘feeder’ roots for resources and can act as refuges for pests (Tesic, Keller & Hutton, 2007; Waipara Valley North Canterbury Winegrowers, pers. comm. 2009). In this study, all the plant species assessed significantly suppressed weeds when compared to unplanted treatments. Whether suppression was sufficient to remove the need for further weed management would depend on the plant species deployed and the weed cover tolerances of individual growers. Plant cover and weed suppression were not significantly correlated, so while some plants may cover a large area, their growth form may not be dense enough to reduce weed penetration. The extent of weed pressure within the trial vineyard may be considered low (bare earth had only 30% weed cover) compared to other vineyards with higher rainfall. Consequently, if endemic

plant species are to be established in regions with higher weed pressure, suppression by the endemic plants may be lower, although they would also be expected to grow more rapidly.

4.2 Invertebrate biodiversity conservation

On all sampling dates, invertebrate diversity was higher for *G. sessiliflorum* than in the bare earth control or the ryegrass inter-row, whereas *M. axillaris* had the highest invertebrate diversity in summer (Table 3). Overall, diversity was lower in winter, which is not surprising considering typical invertebrate phenology (Dent & Walton, 1997; Bowie et al., 2014). However, invertebrate diversity levels were maintained over the winter period by *G. sessiliflorum*, *H. chathamica*, *A. bellidioides* and *A. inermis* ‘*purpurea*’ (Table 3), indicating that they provided suitable overwintering sites for invertebrates. This has implications for early-season pest biological control because early pest control by overwintering invertebrates may prevent pest outbreaks later in the season (Ramsden et al., 2015). While there is debate over the extent to which species richness correlates positively to ecosystem functioning (Loreau, Naeem & Inchausti, 2002; Cardinale et al., 2006), it remains the case that the extent of ecosystem functions depends on the traits of the species examined and their sensitivity to environmental change.

4.3 Conservation biological control (CBC)

Increasing plant diversity such as adding beneficial plants has become a fundamental part of integrated pest management (IPM) theory and practice (Bugg & Waddington, 1994; Landis, Wratten & Gurr, 2000; Gurr, Wratten & Snyder, 2012; Ratnadass et al., 2012). Increased rates of biological control under these conditions have often been attributed to the more diverse system providing natural enemies with resource subsidies including alternative food and shelter (Landis, Wratten & Gurr, 2000; Altieri & Nicholls, 2004; Gurr, Wratten & Altieri, 2004; Zehnder et al., 2007; Helyer, Cattlin & Brown, 2014). Also, diverse assemblages of arthropod taxa associated

with some of the selected plant treatments (Table 3) included potential alternative prey such as Collembola, Diptera, Hemiptera etc. For example, spider densities were higher for several plant treatments than the controls. This is consistent with other research (Thomson & Hoffmann, 2007) and was probably due to the plants providing suitable (and permanent) shelter. Spiders can reduce insect pest populations (Marc, Canard & Ysnel, 1999; Midega et al., 2008) and in vineyards have been implicated as key predators of pests (Hogg & Daane, 2010) including *E. postvittana*, mealybugs (*Pseudococcus* spp.), scales (Hemiptera: Coccidae) and mites (Acari: Eriophyidae) (Thomson & Hoffmann, 2007). The most abundant spider families represented in this study included web-building Linyphiidae and Theridiidae and the wandering/hunting Salticidae and Oxyopidae (Paquin, Vink & Duperre, 2010). These all predate *E. postvittana* and feed on both larval and adult stages of this pest (MacLellan, 1973; Danthanarayana, 1983; Hogg et al., 2014).

4.4 Soil improvements

For all plant species, the estimated soil moisture was always similar to or higher than the control on all three sample dates. It is well established that competition for water between the crop and added plant biodiversity can be a major factor in farmers' agronomic decision making (Warner, 2007). However, there was no obvious competition for water between the added plants and the vines, which obtain most of their water from deep roots, rather than surface 'feeder' roots (Jackson, 2000). Soil biological activity increased beneath grapevines with endemic plant understoreys which may correspond to enhanced nutrient cycling (Mader et al., 2002) compared to the control. The identity of those organisms responsible for such increases could be addressed to some extent by the use of molecular methods (Hirsch, Mauchline & Clark, 2010). The influence of the plants on the above parameters may increase over time, especially after further leaf litter accumulation and root development, although the dry conditions of many vineyards in

summer (occupying largely ‘Mediterranean’ climates (Hannah et al., 2013)) may limit soil microbial activity (Labeda, Kang-Chien & Casida, 1976).

4.5 Potential of the selected plants to host the pest E. postvittana: An ecosystem dis-service (EDS).

Results suggested that some of the plant species could be suitable hosts to the larvae of this pest. However, of the three plant species identified (*L. dioica*, *A. inermis* ‘purpurea’ and *M. axillaris*) by their growth and floral resource to be most promising for vineyard deployment, *L. dioica* and *M. axillaris* supported the lowest mean larval survival and, along with the other plants tested (Figure 2), pose little threat of enhancing *E. postvittana* populations.

4.6 Winegrower attitudes

The majority of growers indicated they would consider incorporating endemic plants into their properties (Table 7). However, several potential barriers to such action were identified and these would need to be overcome to achieve widespread establishment of endemic plants. These barriers centred on lack of knowledge of the other potential effects of plant establishment and the initial investment required (Table 8). This is probably because at the time of the survey, this practice was still in the research phase with protocols yet to be made available to winegrowers. Perceived risk was a notable barrier to growers establishing endemic plants in their vineyards (Table 8). This response is probably due to concerns that such vegetation may exacerbate bird damage to grapes by providing resources (shelter, food etc.) which may support pest bird populations (Waipara Valley North Canterbury Winegrowers, pers. comm. 2009).

4.7 Evaluating the benefits provided by non-crop plants in vineyards

It is critical that the establishment of endemic plants in vines is financially viable. Market-based incentives may exist for provision of enhanced ES, such as marketing, weed suppression or pest control. However, other ES that such plants provide may be public goods and lack any direct financial incentive to the grower; conservation, cultural value or aesthetics are examples. This involves paying for ecosystem services (PES) which have value beyond the farm (Wratten et al., 2013). Compensation for ES that are public goods would probably entail government incentives such as subsidies or tax reductions (Kroegeer & Casey, 2007) and could be delivered via agri-environment schemes such as those in the USA, UK and Europe, although these have achieved mixed results (Kleijn & Sutherland, 2003; Kleijn et al., 2006).

5. Conclusions

Endemic New Zealand plants beneath grapevines can provide multiple potential ecosystem services, including weed suppression, biodiversity conservation, soil improvement and conservation biological control. In some cases in the current work, the plants constituted RTUs and harboured ESPs. For example, the added plant populations were SPUs for services such as biological control, they enhanced ESPs such as spiders and provided ES in the form of weed suppression and enhanced soil quality, expressed as higher moisture and microbial activity. Winegrowers are likely to establish endemic plants within vineyards if perceived and real barriers to such action are overcome. These include growers' lack of knowledge, initial investment, risk and disruption to normal practices. Also, farmers learn about and adopt new practices in a range of ways, and social learning (Warner, 2007) is one of these. Orthodox teaching/technology-transfer methods rarely work (Cullen et al., 2008). This New Zealand work is highly relevant to other regions as the traits of the plants in this study are likely to be similar to other plant species in vineyard ecosystems worldwide. Also, although *A. inermis* is endemic to New Zealand, it is now available commercially in the UK and USA and as seeds in New Zealand

(<http://www.nzseeds.co.nz/contact-us>). The work presented here addresses a key current challenge, which is to maintain or enhance productivity of agro-ecosystems in a sustainable way and to reduce external costs by increasing the role that ES can play on farmland. Meeting this challenge has been called ‘sustainable intensification’ (Garnett et al., 2013; Pretty & Bharucha, 2014) and the current work, although not concerning food, contributes to that. It illustrates how simple enhancements of agricultural biodiversity can help translate ecosystem science into action, thereby supporting the goals of the intergovernmental science-policy platform on Biodiversity and Ecosystem Services (www.ipbes.net).

Acknowledgements

Thanks also to Mud House New Zealand (now Accolade Wines) for the generous provision of a field site and to Jean-Luc Dufour at that site for mentoring.

References

- Allan E, Manning P, Alt F, Binkenstein J, Blaser S, Blüthgen N, Böhm S, Grassein F, Hölzel N, Klaus VH, Kleinebecker T, Morris EK, Oelmann Y, Prati D, Renner SC, Rillig MC, Schaefer M, Schloter M, Schmitt B, Schöning I, Schrumpf M, Solly E, Sorkau E, Steckel J, Steffen-Dewenter I, Stempfhuber B, Tschapka M, Weiner CN, Weisser WW, Werner M, Westphal C, Wilcke W, Fischer M. 2015. Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition. *Ecology Letters* 18:834–843. DOI: 10.1111/ele.12469.
- Alef K, Nannipieri P. 1995. *Methods in Applied Soil Microbiology and Biochemistry*. New York: Academic Press.

- 437 Altieri MA, Nicholls CI. 2004. *Biodiversity and Pest Management in Agroecosystems*.
438 Binghamton, NY: Food Products Press.
- 439 Arnold AJ. 1994. A lightweight portable insect suction sampling system. *Crop Protection* 13:73–
440 76.
- 441 Barral MP, Benayas JMR, Meli P, Maceira NO. 2015. Quantifying the impacts of ecological
442 restoration on biodiversity and ecosystem services in agroecosystems: a global meta-
443 analysis. *Agriculture, Ecosystem and Environment* 202:223–231.
- 444 Berndt LA, Wratten SD, Hassan PG. 2002. Effects of buckwheat flowers on leafroller
445 (Lepidoptera: Tortricidae) parasitoids in a New Zealand vineyard. *Agriculture and Forestry*
446 *Entomology* 4:30–45.
- 447 Bowie MH, Klimaszewski J, Vink CJ, Hodge S, Wratten SD. 2014. Effect of boundary type and
448 season on predatory arthropods associated with field margins on New Zealand farmland. *New*
449 *Zealand Journal of Zoology* 41:268–284.
- 450 Bugg RL, Waddington C. 1994. Using cover crops to manage arthropod pests of orchards: A
451 review. *Agriculture, Ecosystem and Environment* 50:11–28.
- 452 Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM,
453 Tilman D, Wardle DA, Kinzig AP, [Daily](#) GC, Loreau M, [Grace](#) JB, Larigauderie A, Srivastava
454 DS, [Naeem](#) S. 2012. Biodiversity loss and its impact on humanity. *Nature* 486:59–67.

- 455 Cardinale BJ, Srivastava DS, Duffy JE, Wright JP, Downing AL, Sankaran M, Jouseau C. 2006.
456 Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature* 443:989–992.
- 457 Cullen R, Warner KD, Jonsson M, Wratten SD. 2008. Economics and adoption of conservation
458 biological control. *Biological Control* 45:272–280.
- 459 Danthanarayana W. (1983). Population ecology of the light brown apple moth, *Epiphyas*
460 *postvittana* (Lepidoptera: Tortricidae). *The Journal of Animal Ecology* 52:1-33.
- 461 Dent DR, Walton MP. 1997. Methods in Ecological and Agricultural Entomology. Wallingford:
462 CABI Publishing.
- 463 Fiedler AK, Landis DA, Wratten SD. 2008. Maximising ecosystem services from conservation
464 biological control: The role of habitat management. *Biological Control* 45:254-271.
- 465 Garnett T, Appleby MC, Balmford A, Bateman IJ, Benton TG, Bloomer P. 2013. Sustainable
466 intensification in agriculture: Premises and policies. *Science* 341:33–34.
- 467 Gurr G, Wratten SD, Altieri MA. 2004. *Ecological engineering for pest management: Advances*
468 *in habitat manipulation for arthropods*. Wallingford: CABI Publishing.
- 469 Gurr GM, Wratten SD, Snyder WE. 2012. *Biodiversity and Insect Pests: Key issues for*
470 *Sustainable Management*. Hoboken, NJ: Wiley Blackwell Publishers.

- 471 Hannah L, Roehrdanz PR, Ikegami M, Shepard AV, Shaw M-R, Tabord G, Zhie L, Marquet PA,
472 Hijmansj RJ. 2013. Climate change, wine, and conservation. *Proceeding of the National*
473 *Academy of Science* 110:6907–6912.
- 474 Helyer N, Cattlin ND, Brown KC. 2014. *Biological control in plant protection: a colour*
475 *handbook*. Boca Raton, FL: CRC Press.
- 476 Hirsch PR, Mauchline TH, Clark IM. 2010. Culture-independent molecular techniques for soil
477 microbial ecology. *Soil Biology and Biochemistry* 42:878–887.
- 478 Hogg BN, Daane KM. 2010. The role of dispersal from natural habitat in determining spider
479 abundance and diversity in California vineyards. *Agriculture, Ecosystem and Environment*
480 135:260–267.
- 481 Hogg BN, Wang XG, Mills NJ, Daane KM. 2014. Resident spiders as predators of the recently
482 introduced light brown apple moth, *Epiphyas postvittana*. *Entomologia Experimentalis et*
483 *Applicata* 151:65-74.
- 484 Jackson SRon. 2000. *Wine Science: Principles, practice and perception*. London: Academic
485 Press.
- 486 Jackson D, Schuster, D. 2002. *Canterbury grapes and wine 1840-2002*. Christchurch, New
487 Zealand: Shoal Bay Press.

- 488 Keesing V, Wratten SD. 1997. Integrating plant and insect conservation. In: Maxted N, Ford-
489 Lloyd BV, Hawkes JG, ed. *Plant Genetic Conservation*. London: Chapman and Hall, 220–235.
- 490 Khan ZR, James DG, Midega CAO, Pickett JA. 2008. Chemical ecology and conservation
491 biological control. *Biological Control* 45:210-24.
- 492 Kleijn D, Baquero RA, Clough Y, Diaz M, De Esteban J, Fernandez F, Gabriel D, Herzog F,
493 Holzschuh A, Johl R, Knop E, Kruess A, Marshall EJP, Steffan-Dewenter I, Tscharnkte T,
494 Verhulst J, West TM, Yela JL. 2006. Mixed biodiversity benefits of agri-environment schemes in
495 five European countries. *Ecology Letters* 9:243–254.
- 496 Kleijn D, Sutherland WJ. 2003. How effective are European agri-environment schemes in
497 conserving and promoting biodiversity? *Journal of Applied Ecology* 40:947–969.
- 498 Kremen C. 2005. Managing ecosystem services: What do we need to know about their ecology?
499 *Ecology Letters* 8:468-479.
- 500 Kroeger T, Casey F. 2007. An assessment of market-based approaches to providing ecosystem
501 services on agricultural lands. *Ecological Economics* 64:321–332.
- 502 Labeda DP, Kang-Chien L, Casida LE. 1976. Colonisation of soil by *Arthrobacter* and
503 *Pseudomonas* under varying conditions of water and nutrient availability as studied by plate
504 counts and transmission electron microscopy. *Applied Environmental Microbiology* 31:551–561.

- 505 Landis DA, Wratten SD, Gurr GM. 2000. Habitat management to conserve natural enemies of
506 arthropod pests in agriculture. *Annual Review of Entomology* 45:175–201.
- 507 Loreau M, Naeem S, Inchausti P. 2002. *Biodiversity and Ecosystem Functioning: Synthesis and*
508 *Perspectives*. Oxford: Oxford University Press.
- 509 Luck GW, Daily GC, Ehrlich PR. 2003. Population diversity and ecosystem services. *Trends in*
510 *Ecology and Evolution* 18:331–336.
- 511 Mace GM, Norris K, Fitter AH. 2012. Biodiversity and ecosystem services: a multilayered
512 relationship. *Trends in Ecology and Evolution* 27:19–26.
- 513 MacLellan CR. 1973. Natural enemies of the light brown apple moth, *Epiphyas postvittana*, in
514 the Australian Capital Territory. *The Canadian Entomologist* 105:681-700.
- 515 Mader P, Fliebbach A, Dubois D, Gunst L, Fried P, Niggli U. 2002. Soil fertility and biodiversity
516 in organic farming. *Science* 296:1694-1697.
- 517 Magurran AE. 1988. *Ecological Diversity and its Measurement*. Princeton, NJ: Princeton
518 University Press.
- 519 Marc P, Canard A, Ysnel F. 1999. Spiders (Araneae) useful for pest limitation and bioindication.
520 *Agriculture, Ecosystem and Environment*. 74:229–273.

- 521 MEA. 2005. Millennium Ecosystem Assessment. Ecosystems and Human Well-being:
522 Biodiversity Synthesis. Washington, DC: World Resources Institute. Available at
523 <http://www.millenniumassessment.org/en/Synthesis.html> (accessed 10 February 2016)

- 524 Midega CAO, Khan ZR, van den Berg J, Ogot CKPO, Dippenaar-Schoeman AS, Pickett JA,
525 Wadhams LJ. 2008. Response of ground-dwelling arthropods to a ‘push–pull’ habitat
526 management system: spiders as an indicator group. *Journal of Applied Entomology* 132:248–254.

- 527 Mooney HA, Ehrlich PR. 1997. Ecosystem services: A fragmentary history. In: Daily GC, ed.
528 *Nature's services: Societal dependence on natural ecosystems*. Washington, DC: Island Press, 11–
529 19.

- 530 Mooney HA. 2010. The ecosystem-service chain and the biological diversity crisis.
531 *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 365:31–39.

- 532 Nicholls CI, Altieri MA, Ponti L. 2008. Enhancing plant diversity for improved insect pest
533 management in northern California organic vineyards. *Acta horticulturae* 785:263–278.

- 534 Paquin P, Vink CJ, Duperre N. 2010. Spiders of New Zealand: annotated family key and species
535 list. Lincoln, New Zealand: Manaaki Whenua Press.

- 536 Pretty J, Bharucha ZP. 2014. Sustainable intensification in agricultural systems. *Annals of botany*
537 114:1571–1596.

- 538 Ramsden MW, Menéndez R, Leather SR, Wäckers F. 2015. Optimizing field margins for
539 biocontrol services: The relative role of aphid abundance, annual floral resources, and overwinter
540 habitat in enhancing aphid natural enemies. *Agriculture, Ecosystem & Environment* 199:94-104.
- 541 Ramos MC, Benito C, Martínez-Casasnovas JA. 2015. Simulating soil conservation measures to
542 control soil and nutrient losses in a small, vineyard dominated, basin. *Agriculture, Ecosystem &
543 Environment* 213:194-208.
- 544 Ratnadass A, Fernandes P, Avelino J, Habib R. 2012. Plant species diversity for sustainable
545 management of crop pests and diseases in agroecosystems: a review. *Agronomy for sustainable
546 development* 32:273–303.
- 547 Reeve JR, Carpenter-Boggs L, Reganold JP, York AL, McGourty G, McCloskey LP. 2005. Soil
548 and winegrape quality in biodynamically and organically managed vineyards. *American Journal
549 of Enology and Viticulture* 56:367-376.
- 550 Sandhu H, Wratten SD, Costanza R, Pretty J, Porter JR, Reganold J. 2015. Significance and value
551 of non-traded ecosystem services on farmland. *PeerJ* 3:e762.
- 552 Sandhu S, Wratten SD, Porter JR, Costanza R, Pretty J, Reganold JP. 2016. Mainstreaming
553 ecosystem services into future farming. *Solutions* March-April:40-47.
- 554 Saville DJ. 1990. Multiple comparison procedures: The practical solution. *The American
555 Statistician* 44:174–180.

- 556 Scarratt SL, Wratten SD, Shishehbor P. 2008. Measuring parasitoid movement from floral
557 resources in a vineyard. *Biological Control* 46:107–113.

- 558 Shorey HH, Hale RL. 1965. Mass rearing of the larvae of nine noctuid species on a simple
559 artificial medium. *Journal of Economic Entomology* 58:522–524.

- 560 Swift MJ, Anderson JM. 2012. Biodiversity and ecosystem function in agricultural systems. In:
561 Schulze E-D, Mooney HA, ed. *Biodiversity and Ecosystem Function*. Berlin: Springer Science &
562 Business Media, 15–42.

- 563 Tesic D, Keller M, Hutton RJ. 2007. Influence of vineyard floor management practices on
564 grapevine vegetative growth, yield, and fruit composition *American Journal of Enology and*
565 *Viticulture* 58:1–11.

- 566 The Wine Institute 2012. World vineyard acre by country 2009–2012. *Available at*
567 *www.wineinstitute.org* (accessed 24 Jun 2015).

- 568 Thomson LJ, Hoffmann AA. 2007. Effects of ground cover (straw and compost) on the
569 abundance of natural enemies and soil macro invertebrates in vineyards. *Agricultural and Forest*
570 *Entomology* 9:173–179.

- 571 Topp GC, Parkin GW, Ferré TP. 1993. Soil water content. In: Carter MR. ed. *Soil Sampling and*
572 *Methods of Analysis*. Ottawa: Canadian Society of Soil Science, 541-544.

- 573 Tschardtke T, Clough Y, Wanger TC, Jackson L, Motzke I, Perfecto I, Vandermeer J, Whitbread
574 A. 2012. Global food security, biodiversity conservation and the future of agricultural
575 intensification. *Biological Conservation* 151: 53–59.

- 576 Tuck SL, Winqvist C, Mota F, Ahnström J, Turnbull LA, Bengtsson J. 2014. Land-use intensity
577 and the effects of organic farming on biodiversity: a hierarchical meta-analysis. *Journal of*
578 *Applied Ecology* 51:746–755.

- 579 Warner KD. 2007. *Agroecology in Action Extending Alternative Agriculture through Social*
580 *Networks*. Cambridge: MIT Press.

- 581 Wratten SD, Sandhu H, Cullen R, Costanza R. 2013. *Ecosystem Services in Agricultural and*
582 *Urban Landscapes*. Hoboken, NJ: Wiley Blackwell Publishers.

- 583 Zehnder G, Gurr GM, Kühne S, Wade MR, Wratten SD, Wyss E. 2007. Arthropod pest
584 management in organic crops. *Annual Review of Entomology* 52:57–80.

585 Endemic^a plant species used in the vineyard trial and the ecosystem associated benefits assessed.

Plant species	Family	Ecosystem associated benefits				
		ES	ES	ES	ESP	EDS
		Weed suppression	Invertebrate conservation ^c	Improving Soil quality	Enhancing Predator densities ^c	Pest development
<i>Acaena inermis</i>	Rosaceae	+	+		+	
<i>Acaena inermis</i> 'purpurea' ^b	Rosaceae	+	+	+	+	+
<i>Anaphalioides bellidioides</i>	Asteraceae	+	+		+	+
<i>Disphyma australe</i>	Mesembryanthemaceae		+		+	
<i>Geranium sessiliflorum</i>	Geraniaceae	+	+	+	+	+
<i>Hebe chathamica</i>	Plantaginaceae	+	+	+	+	+
<i>Leptinella dioica</i>	Asteraceae	+	+	+	+	+
<i>Leptinella squalida</i>	Asteraceae	+	+		+	
<i>Lobelia angulata</i>	Lobeliaceae	+	+	+	+	+
<i>Muehlenbeckia ephedroides</i>	Polygonaceae		+		+	
<i>Muehlenbeckia axillaris</i>	Polygonaceae	+	+	+	+	+
<i>Raoulia hookeri</i>	Asteraceae	+	+		+	+
<i>Raoulia subsericea</i>	Asteraceae		+		+	
<i>Scleranthus uniflorus</i>	Caryophyllaceae	+	+		+	+

586 ^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

587 ^b A natural variation of *A. inermis* which has purplish coloration.

588 ^c Three sampling dates occurred, with some plant species sampled only once (*D. australe*, *M.*
589 *ephedroides* and *R. subsericea*).

590 Mean change in cover (m^2) of endemic plant species from planting to 6 or 12 months,
591 respectively, and their survival beneath grapevines at 12 and 24 months, respectively (for full
592 species names see Table 1).

Endemic plant ^a	Change in Cover (m^2) after:		Survival (%) at:	
	6 months	12 months ^b	12 months	24 months
<i>L. dioica</i>	0.24	0.38	100	100
<i>A. inermis</i> 'purpurea'	0.28	0.34	100	90
<i>L. angulata</i>	0.30	0.22	100	70
<i>L. squalida</i>	0.10	0.20	95	50
<i>G. sessiliflorum</i>	0.10	0.16	100	90
<i>M. axillaris</i>	0.20	0.15	100	100
<i>H. chathamica</i>	0.19	0.14	100	80
<i>R. hookeri</i>	0.13	0.13	100	100
<i>S. uniflorus</i>	0.06	0.13	100	80
<i>A. inermis</i>	0.07	0.12	60	60
<i>A. bellidioides</i>	0.06	0.04	90	40
<i>M. ephedrioides</i>	0.03	0.00	80	0
<i>R. subsericea</i>	-0.03	-0.03	60	10
<i>D. australe</i>	0.44	-0.14	0	0
LSD(5%)^c	0.10	0.12	-	-

593 ^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

594 ^b The table has been sorted into the order of decreasing growth to 12 months.

595 ^c LSD = Least Significant Difference. Means which differ by more than the LSD(5%) are
596 significantly different at $P < 0.05$.

Mean Shannon-Wiener diversity indices for invertebrates in under-vine treatments at three sampling dates, ranked for 2008 results. Treatments with means of 0 have been omitted from the analysis of variance, as denoted by placing these means in brackets. The variability of such treatments is zero, so a LS Effect (5%) has been calculated to allow comparison between bracketted and unbracketted means (for full species names see Table 1).

Endemic plant ^a	Invertebrate diversity (Shannon-Weiner H')		
	Aug 2008 ^b	Jan 2009	Mar 2009
<i>G. sessiliflorum</i>	1.11	1.17	1.31
<i>H. chathamica</i>	0.95	0.24	0.77
<i>A. bellidioides</i>	0.71	1.10	0.57
<i>A. inermis</i> 'purpurea'	0.45	0.55	1.10
<i>L. dioica</i>	0.35	1.09	0.50
<i>M. axillaris</i>	0.28	1.30	1.31
<i>L. squalida</i>	0.26	0.98	0.52
<i>L. angulata</i>	0.17	0.94	1.01
<i>A. inermis</i>	0.15	0.92	0.79
<i>D. australe</i>	0.07	-	-
<i>M. ephedrioides</i>	0.07	-	-
<i>R. hookeri</i>	0.07	0.71	0.24
<i>R. subsericea</i>	0.07	-	-
<i>S. uniflorus</i>	(0)	(0)	0.07
Ryegrass inter-row	(0)	0.19	0.43
Bare earth	(0)	0.07	(0)
LSD(5%)^c	0.36	0.49	0.45
LSEffect(5%)^d	0.25	0.34	0.32

^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

^b The table has been sorted into the order of decreasing Shannon-Wiener H' mean values in August 2008.

^c LSD = Least Significant Difference. Unbracketted means which differ by more than the LSD (5%) are significantly different at $P < 0.05$.

^d LSEffect = Least Significant Effect. If a bracketted mean and an unbracketted mean differ by more than the LS Effect(5%), then the two means are significantly different at $P < 0.05$.

- means plant species was not sampled.

Mean density of spiders/m² for different under-vine endemic plant treatments in August 2008, January 2009 and March 2009. Treatments with means of 0 or 3 (one spider in one plot) have been omitted from the analysis of variance, as denoted by placing these means in brackets. The variability of such treatments is nil or very low, so assuming it is zero, an LS Effect (5%) has been calculated to allow comparison between bracketted and unbracketted means (for full species names see Table 1).

Endemic plant ^{a, b}	Density of spiders /m ² in:		
	Aug 2008	Jan 2009	Mar 2009
<i>L. dioica</i>	8	(0)	5
<i>A. inermis</i> 'purpurea'	15	10	45
<i>L. angulata</i>	(0)	33	20
<i>L. squalida</i>	(3)	10	(0)
<i>G. sessiliflorum</i>	60	38	83
<i>M. axillaris</i>	20	8	30
<i>H. chathamica</i>	38	45	70
<i>R. hookeri</i>	(0)	8	13
<i>S. uniflorus</i>	(3)	(3)	(0)
<i>A. inermis</i>	5	15	18
<i>A. bellidioides</i>	18	18	23
<i>M. ephedrioides</i>	(0)	-	-
<i>R. subsericea</i>	(0)	-	-
<i>D. australe</i>	10	-	-
Ryegrass inter-row	(3)	10	5
Bare earth (Control)	(0)	(3)	(0)
LSD(5%)^c	25	29	32
LSEffect(5%)^d	18	20	23

^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

^b This table has been sorted into the same order of endemic plants as Table 2.

^c LSD = Least Significant Difference. Unbracketted means which differ by more than the LSD(5%) are significantly different at $P < 0.05$.

^d LSEffect = Least Significant Effect. If a bracketted mean and an unbracketted mean differ by more than the LSEffect(5%), then the two means are significantly different at $P < 0.05$.

- means plant species was not sampled.

Mean soil moisture percentage for different under-vine treatments in December 2008, September 2009 and November 2009, and mean microbial activity as measured by the TTC method on the first date. Soil moisture is expressed on a dry weight basis (for full species names see Table 1).

Endemic plant ^{a, b}	Soil moisture (%) in:			Mean microbial activity (TTC method) [(rate of reduction of TTC, µg)/(g dry soil/hr)]
	Dec 2008	Sep 2009	Nov 2009	
<i>L. dioica</i>	6.5	11.6	8.3	20.0
<i>A. inermis</i> 'purpurea'	7.7	14.8	14.3	13.3
<i>L. angulata</i>	7.0	-	16.2	12.2
<i>G. sessiliflorum</i>	5.2	17.1	8.7	12.2
<i>M. axillaris</i>	6.4	17.6	8.9	11.6
<i>H. chathamica</i>	5.0	16.3	8.3	12.9
Bare earth	5.3	10.3	7.1	6.7
LSD(5%)^c	2.6	4.0	5.0	4.4

^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

^b This table has been sorted into the same order of endemic plants as Table 2.

^c LSD = Least Significant Difference. Means which differ by more than the LSD (5%) are significantly different at $P < 0.05$.

- means plant species was not adequately sampled on this date.

Ranking of endemic plant species by change in growth, survival beneath grapevines and ecosystem associated benefits; weed suppression, mean invertebrate diversity, mean spider density, mean soil moisture, leafroller survival and microbial activity on one date. A rank of 1 was the best in terms of desirability. A mean ranking was calculated for only those endemic plants for which all attributes had been assessed. Ties were replaced by mean ranks; e.g., three 1= values were replaced by 2s, and two 4= values by 4.5s (for full species names see Table 1).

Endemic plant ^a	Growth (m ²) ^b to 12 months	Survival (%) to 24 months	Ecosystem associated benefits						Mean ranking
			ES	ES	ESP	ES	ES	EDS	
			Weed suppression at 11 months	Invertebrate diversity (Shannon-Wiener H')	Density of spiders/m ²	Soil moisture (%)	Mean microbial activity (TTC method) [(rate of reduction of TTC, µg)/(g dry soil)/hr]	Leaf-roller (pest) survival	
<i>L. dioica</i>	1	1=	5	7	10	6	1	4=	4.6
<i>A. inermis</i> 'purpurea'	2	4=	6	5	3	2	2	9	4.2
<i>L. angulata</i>	3	8	7	4	6	1	4	7	5.0
<i>L. squalida</i>	4	10	1=	9	9	-	-	-	-
<i>G. sessiliflorum</i>	5	4=	1=	1	1	4	5	8	3.8
<i>M. axillaris</i>	6	1=	8	2	4=	3	6	4=	4.5
<i>H. chathamica</i>	7	6=	3=	6	2	5	3	1	4.3
<i>R. hookeri</i>	8	1=	9	10	8	-	-	6	-
<i>S. uniflorus</i>	9	6=	3=	-	11	-	-	3	-
<i>A. inermis</i>	10	9	10	8	7	-	-	-	-
<i>A. bellidioides</i>	11	11	11	3	4=	-	-	2	-
<i>M. ephedrioides</i>	12	13=	-	-	-	-	-	-	-
<i>R. subsericea</i>	13	12	-	-	-	-	-	-	-
<i>D. australe</i>	14	13=	-	-	-	-	-	-	-

^a All plant species in this work apart from *M. axillaris* are endemic to New Zealand.

^b The table has been sorted into the order of decreasing growth to 12 months.

- means plant species was not assessed.


640 Current and potential use of endemic plants within Waipara vineyards (survey responses from n =
641 30 growers).

Endemic plant ecosystem benefit use	Number of growers establishing endemic plant for ecosystem associated benefits listed on left ^a						
	N/A	Already do this	Definitely	Maybe	Probably not	Definitely not	Already + Definitely
As groundcover to suppress weeds beneath vines	0	2	3	20	4	1	5
To provide resources to beneficial vineyard insects	0	10	6	14	0	0	16
To reduce soil erosion in the vineyard	7	6	12	4	0	1	18
To conserve beneficial invertebrates	1	17	8	4	0	0	25
To contribute to endemic plant conservation	1	18	8	2	1	0	26
For eco-marketing purposes	9	7	6	6	2	0	13


642 ^a Number of growers who currently or potentially would use endemic plants in the manner
643 indicated.

644 Potential barriers to deploying endemic plants within vineyard properties. For each plant use, the
645 number of respondents for which the use was applicable is given in the right-hand column.

Endemic plant ecosystem benefit use	Number of growers citing barriers to establishing endemic plant for various uses						Number of respondents to whom applicable
	N/A	Lack of knowledge	Initial investment	Risk	Disruption to normal practices	No interest by grower	
As groundcover to suppress weeds beneath vines	0	12	11	4	4	2	30
To provide resources to beneficial vineyard insects	0	4	10	1	5	0	30
To reduce soil erosion in the vineyard	7	3	6	1	1	1	23
To conserve beneficial invertebrates	1	3	7	13	1	0	29
To contribute to endemic plant conservation	1	3	7	13	13	0	29
For eco-marketing purposes	9	3	5	14	14	1	21



646 Mean weed penetration of under-vine treatments within the 0.04 m² areas assessed. Treatments
 647 with a letter in common are not significantly different from one another at $P < 0.05$. Letters were
 648 assigned using the unprotected LSD procedure (Saville, 1990); LSD(5%) = 13.



649 Mean proportion of leafroller, *Epiphyas postvittana*, larvae surviving at each development stage.
 650 Treatment names which have a letter in common indicate the two treatments are not significantly
 651 different in overall survival (averaged over all development stages) at $P < 0.05$.