

A peer-reviewed version of this preprint was published in PeerJ on 19 July 2016.

[View the peer-reviewed version](https://peerj.com/articles/2128) (peerj.com/articles/2128), which is the preferred citable publication unless you specifically need to cite this preprint.

Jaskuła R, Rewicz T, Płóciennik M, Grabowski M. 2016. Pleistocene phylogeography and cryptic diversity of a tiger beetle, *Calomera littoralis*, in North-Eastern Mediterranean and Pontic regions inferred from mitochondrial COI gene sequences. PeerJ 4:e2128
<https://doi.org/10.7717/peerj.2128>

Pleistocene phylogeography of tiger beetle, *Calomera littoralis*, in North-Eastern Mediterranean and Pontic regions inferred from mitochondrial COI gene sequence

Radomir Jaskuła, Tomasz Rewicz, Mateusz Płóciennik, Michał Grabowski

Background. *Calomera littoralis* is a Palearctic species, widely distributed in Europe; inhabiting predominantly its Atlantic, Mediterranean and Black Sea coastlines.

Methods. Its phylogeography on the Balkan Peninsula and on the north-western Black Sea coast was inferred using a 697 bp long portion of the mitochondrial COI gene, amplified from 169 individuals collected on 43 localities.

Results. The results revealed two genetically divergent groups/lineages, the southern one inhabiting both the Balkan Peninsula and the Pontic Region and the northern one found exclusively in the Pontic Region. Species delimitation based on DNA barcoding gap suggested an interspecific level of divergence between these groups. Multivariate analysis of eight male and female morphometric traits detected no difference between the groups, implying they may represent cryptic species. The Bayesian time-calibrated reconstruction of phylogeny suggested that the lineages diverged ca. 2.3 Ma, in early Pleistocene.

Discussion. This might result from contemporary isolation of the Pontic basin from the Mediterranean that broke the continuous strip of coastal habitats inhabited by *C. littoralis*. Demographic analyses indicated that both lineages have been in demographic and spatial expansion since ca. 0.15 Ma. It coincides with the terminal stage of MIS-6, i.e. Wartanian/Saalian glaciation, and beginning of MIS-5e, i.e. Eemian interglacial, during which, due to eustatic sea level rise, a wide connection between Mediterranean and the Pontic basin was re-established. This, along with re-appearance of coastal habitats could initiate north-east expansion of the southern lineage and its secondary contact with the northern one. Isolation of Pontic basin from Mediterranean during the Weichselian glaciation did not have probably any effect on their phylogeography.

1 **Pleistocene phylogeography of tiger beetle, *Calomera littoralis*, in North-Eastern**
2 **Mediterranean and Pontic regions inferred from mitochondrial COI gene sequences**

3 Radomir Jaskuła^{1*}, Tomasz Rewicz², Mateusz Płóciennik¹, Michał Grabowski¹

4 ¹Department of Invertebrate Zoology and Hydrobiology, University of Lodz, Banacha 12/16,
5 90-237 Lodz, Poland

6 ²Laboratory of Microscopic Imaging and Specialized Biological Techniques, University of
7 Lodz, Banacha 12/16, Łódź 90-237, Poland

8

9

10 Corresponding author:

11 Radomir Jaskuła¹

12 Ul. Banacha 12/16, 90-237 Łódź, Łódź, Poland

13 Email address: radekj@biol.uni.lodz.pl

14

15

16

17 **Abstract**

18 **Background.** *Calomera littoralis* is a Palearctic species, widely distributed in Europe;
19 inhabiting predominantly its Atlantic, Mediterranean and Black Sea coastlines.

20 **Methods.** Its phylogeography on the Balkan Peninsula and on the north-western Black Sea
21 coast was inferred using a 697 bp long portion of the mitochondrial COI gene, amplified from
22 169 individuals collected on 43 localities.

23 **Results.** The results revealed two genetically divergent groups/lineages, the southern one
24 inhabiting both the Balkan Peninsula and the Pontic Region and the northern one found
25 exclusively in the Pontic Region. Species delimitation based on DNA barcoding gap
26 suggested an interspecific level of divergence between these groups. Multivariate analysis of
27 eight male and female morphometric traits detected no difference between the groups,
28 implying they may represent cryptic species. The Bayesian time-calibrated reconstruction of
29 phylogeny suggested that the lineages diverged ca. 2.3 Ma, in early Pleistocene.

30 **Discussion.** This might result from contemporary isolation of the Pontic basin from the
31 Mediterranean that broke the continuous strip of coastal habitats inhabited by *C. littoralis*.
32 Demographic analyses indicated that both lineages have been in demographic and spatial
33 expansion since ca. 0.15 Ma. It coincides with the terminal stage of MIS-6, i.e.
34 Wartanian/Saalian glaciation, and beginning of MIS-5e, i.e. Eemian interglacial, during
35 which, due to eustatic sea level rise, a wide connection between Mediterranean and the Pontic
36 basin was re-established. This, along with re-appearance of coastal habitats could initiate
37 north-east expansion of the southern lineage and its secondary contact with the northern one.
38 Isolation of Pontic basin from Mediterranean during the Weichselian glaciation did not have
39 probably any effect on their phylogeography.

40

41 **Key words:** Coleoptera – Cicindelidae – ice age – interglacial – sea level changes – population
42 expansion – southern Europe

43

44 **Introduction**

45 The Eastern Mediterranean including the Pontic area is recognised as one of the major
46 biodiversity and endemism hot spots on a global scale as well as a major glacial refugium in
47 Europe (eg. Myers et al., 2000; Kotlík, Bogutskaya & Ekmekçi, 2004; Blondel et al., 2010).
48 Among others, it is a consequence of complex geological history of the region that was an
49 archipelago and united with rest of the European continent only in Neogene (Pffifner, 2014).
50 On the other side, a shallow epicontinental sea, Paratethys, occupied vast areas of the continent
51 and regressed gradually leaving relics, such as Black, Azov and Caspian Sea (Nahavandi et al.,
52 2013). Local isostatic and eustatic changes of sea level were among superior phenomena
53 shaping local landscapes. For example, there were at least twelve saline water intrusions from
54 the Mediterranean Sea, and eight intrusions from the Caspian Lake to the Black Sea during the
55 last 0.67 million years (Ma) (Badertscher et al., 2011). Inevitably, they played an important role
56 in modelling diversity and distribution patterns for numerous organisms, particularly those
57 inhabiting coastal ecosystems both in Mediterranean and in Pontic area. However, the evidence
58 comes mostly from aquatic, predominantly marine or brackish water, taxa (e.g. Audzijonyte,
59 Daneliya & Vainola, 2006; Neilson & Stepien, 2011). There is a deficiency of studies focusing
60 upon coastal species inhabiting terrestrial habitats in this region (Akin et al., 2010).

61 Tiger beetles, Cicindelidae Latreille, 1806, seem to be ideal model organisms to test such
62 assumptions. The family, with more than 2600 species, has a worldwide distribution with
63 exception of polar regions and some oceanic islands (Pearson & Cassola 2005). Most species,
64 both in larval and adult stage, prefer various types of sandy areas and are habitat specialists;
65 often inhabiting coastal areas (Pearson & Vogler 2001). Several studies dealt with

66 phylogeography of tiger beetles in various regions of the world (eg. Vogler et al., 1993; Cardoso
67 & Vogler, 2005; Woodcock *et al.*, 2007), yet so far only few focused on the role of sea level
68 oscillations in their evolutionary history (Vogler & DeSalle, 1993; Sota et al., 2011) or
69 compared the diversity patterns on both, the molecular and morphological, levels (Cardoso,
70 Serrano & Vogler, 2009; Tsuji et al., 2016).

71 The tiger beetle, *Calomera littoralis* Fabricius 1787, is widely distributed in Palearctic,
72 from the Iberian Peninsula and Morocco in the west to the Middle Asia and Russian Far East
73 in the east (Putchkov & Matalin, 2003; Serrano, 2013; Jaskuła, 2011, 2015). Generally, it is
74 recognised as euryoecious (Jaskuła, 2011, 2013, 2015). However, in Europe it occupies
75 predominantly the very narrow stretch of Atlantic, Mediterranean and Black Sea coastal
76 habitats (Cassola & Jaskuła, 2004; Franzen, 2006; Jaskuła, 2007a-b; Jaskuła, Pešić & Pavicević,
77 2005; Serrano, 2013).

78 Our study aims at (1) exploring and comparing spatial patterns of molecular and
79 morphological diversity of this coastal species in the Mediterranean and Pontic region, (2)
80 interpreting the observed patterns in the context of local paleogeography.

81

82 **Material and methods**

83 ***Sample collection and identification***

84 In total, 169 imagines of *Calomera littoralis* were collected with entomological hand net on 43
85 sites on the Mediterranean coasts of the Balkan Peninsula, Crete and Turkey as well as on the
86 northern and western coast of the Black and Azov Seas, in the years 2009-2012 (Figure 1, Table
87 1). At a site the material was fixed in 96% ethanol for DNA preservation. Taxonomic
88 identification of the collected material followed Mandl (1981).

89 DNA extraction, amplification and sequencing

90 Following Hillis et al. (1996) the standard phenol-chloroform method was used to extract DNA
91 from all the collected individuals. Air-dried DNA pellets were eluted in 100 µl of TE buffer,
92 pH 8.00, stored at 4°C until amplification, and subsequently at -20°C for long-term storage.

93 Fragments of mitochondrial cytochrome oxidase subunit I gene (COI), ca. 700 bp long,
94 were amplified using the Jerry and Pat pair of primers (Simon *et al.*, 1994). Each PCR reaction
95 was conducted in a total volume of 10 µl and contained DreamTaq Master Mix (1x) Polymerase
96 (ThermoScientific), 200nM of each primer and 1 µl of DNA template. The thermal regime
97 consisted of initial denaturation at 94°C for 2min, followed by 34 cycles of denaturation at 94°C
98 for 30s, annealing at 44°C for 30s, and elongation at 72°C for 60s, completed by a final
99 extension at 72°C for 10min. The amplified products were visualized on 2.0% agarose gels
100 stained with MidoriGreen (Nippon Genetics) to verify the quality of the PCR reactions. Then,
101 the PCR products were chemically cleaned up of dNTPs and primer residues by adding 1U of
102 Exonuclease I (Thermo Scientific) and 2U of FastAP Alkaline Phosphatase (Thermo Scientific)
103 per sample. Sequencing was performed using BigDye sequencing protocol (Applied
104 Biosystems 3730xl) by Macrogen Inc., Korea.

105 *Molecular data analysis*

106 First, all the obtained sequences were positively verified as *Calomera* DNA using
107 GenBankBLASTn searches (Altschul et al., 1990). Then they were edited and assembled with
108 CLUSTALW algorithm (Chenna et al., 2003) using BIOEDIT© 7.2.5. The resulting alignment was
109 697 bp long with no gaps, and composed of 169 COI sequences. The sequence data and trace
110 files were uploaded to BOLD and subsequently also to GenBank (accession nos KU905171-
111 KU905339).

112 Pairwise Kimura 2-parameter (K2p) distances between sequences were estimated using
113 MEGA 6.2 (Tamura et al., 2013). Haplotypes were retrieved using DNASP v5 (Librado & Rozas,
114 2009). Phylogenetic relationships between the haplotypes were visualised with phylogenetic

115 network computed using the neighbour-net algorithm and uncorrected p-distances in SplitsTree
116 ver. 4.13.1 (Huson & Bryant 2006).

117 To test for presence of distinct operational taxonomic units (OTUs) that may represent
118 potential cryptic species/subspecies in the sequenced pool of individuals we used the Automatic
119 Barcode Gap Discovery (ABGD) procedure (Puillandre et al., 2012). The default value of 0.001
120 was used as the minimum allowed intraspecific distance. The maximum allowed intraspecific
121 distance was set to $P_{max} = 0.03$ and 0.06, as both threshold values have been already used in
122 literature to delimit insect species (Hebert et al., 2003; Hebert, Ratnasingham & de Waard,
123 2003). We applied the K2P model sequence correction, which is a standard for barcode analyses
124 (Hebert et al., 2003). We used primary partitions as a principal for group definition for they are
125 usually stable over a wider range of prior values, minimise the number of false positive (over
126 split species) and are usually close to the number of groups described by taxonomists
127 (Puillandre et al., 2012).

128 To reveal the temporal framework for the divergence of the two phylogenetic lineages
129 (potential cryptic species) revealed in *Calomera littoralis*, the time calibrated phylogeny was
130 reconstructed in BEAST, version 1.8.1 (Drummond et al., 2012). A COI sequence of *Calomera*
131 *lugens aphrodisia* Baudi di Selve 1864 from GenBank (acc. no. KC963733) was used as an
132 outgroup. This analysis was performed on a reduced dataset, containing only the most distant
133 haplotypes from each OTU. The HKY model of evolution and coalescent model were set for
134 priors. The strict clock with rate 0.0115, widely used for phylogenetic studies upon insects, was
135 applied for the analyses (Brower, 1994). Five runs of 20M iterations of Markov chain Monte
136 Carlo (MCMC) sampled each 2000 iterations were performed. The runs were examined using
137 Tracer v 1.6 and all sampled parameters achieve sufficient effective sample sizes (ESS>200).
138 Tree files were combined using Log-Combiner 1.8.1 (Drummond et al., 2012), with removal of

139 the non-stationary 20% burn-in phase. The maximum clade credibility tree was generated using
140 TreeAnnotator 1.8.1 (Drummond et al., 2012).

141 To provide insight into historical demography, i.e. the temporal changes of the effective
142 population size of *Calomera littoralis* in the studied region, we performed Bayesian Skyline
143 Plot (BSP) analysis (Drummond et al., 2005) in BEAST, version 1.8.1 (Drummond et al., 2012).
144 Separate analysis was performed for each of the two phylogenetic lineages revealed in our study
145 (see Results). The Northern Lineage was represented by 84 individuals from 22 localities, while
146 the Southern Lineage was represented by 85 individuals from 32 localities. The HKY+I model
147 of evolution was used as the best fitting model in case of the Eastern Lineage, while TN93+I
148 was used in case of the Western Lineage. Two runs of MCMC, 20M iterations long sampled
149 each 2000 iterations, were performed. In both cases the runs were examined using Tracer v 1.6
150 (Drummond et al., 2012) and all sampled parameters achieved sufficient effective sample sizes
151 (ESS>200).

152 Two models of population expansion, demographic and spatial, were examined using
153 mismatch distribution analysis (Slatkin & Hudson, 1991; Rogers & Harpending, 1992) and
154 Tajima's D neutrality test (Tajima, 1989). Analyses were performed for the COI groups, using
155 Arlequin 3.5.1.3 (Excoffier & Lischer, 2010) with 1000 replicates.

156 ***Morphometric data analysis***

157 To test whether variation of morphometric traits reflects presence of two genetically divergent
158 lineages (potential cryptic species), measurements of eight body parameters (Fig. 2) were taken
159 from all the 69 males and 100 females used previously for the molecular analyses: 1 – right
160 mandible length (RML), 2 – length of head (LH), 3 – width of head (WH), 4 – maximum
161 pronotum width (MPW), 5 – pronotum length (PL), 6 – elytra length (EL), 7 – maximum elytra
162 width (MEW), and 8 – total body length (TBL). The principal component analysis (PCA) was
163 performed separately for each sex (Figure 3, Table 1). To test for significance ($p < 0.01$) of

164 morphological differences (separately for males and females) between the two divergent
165 lineages one-way ANOSIM Pairwise Test was performed. All the above statistical analyses
166 were done with PRIMER 6 software (Clarke & Gorley, 2006).

167

168 **Results**

169 *Molecular data*

170 A total of 81 haplotypes were identified in the dataset composed of 169 individuals from 43
171 sites from the Mediterranean and the Pontic areas (Table 1). The phylogenetic network
172 illustrating phylogenetic relationships among haplotypes (Fig.4) uncovered presence of two
173 distinct haplotype groups (phylogenetic lineages). The first group, from now on defined as
174 southern lineage, includes 36 haplotypes present all over the studied range including the Balkan
175 Peninsula and the Pontic area. The other group, from now on defined as northern lineage, is
176 composed of 45 haplotypes present exclusively along the north-western coast of the Black Sea.
177 The mean K2p genetic distance between both groups of haplotypes is relatively high (0.039,
178 SD 0.007). Both variants of the ABGD analysis resulted in partitioning of the dataset into two
179 OTUs, that may represent distinct operational taxonomic units – potential cryptic species or
180 subspecies within *Calomera littoralis* in the studied area (Fig. 3A).

181 The Bayesian time-calibrated reconstruction of phylogeny shows that the two lineages
182 split at ca. 2Ma, i.e. in early Pleistocene (Fig. 5A). Results of the BSP analyses showing the
183 temporal changes of the effective population size suggests that both lineages experienced rapid
184 population growth that has started ca. 0.15Ma (Fig. 5B). In both cases, a small decline in
185 effective population size may be observed in most recent times (<0.05 Ma). Results of the
186 mismatch analysis shows that both lineages are currently in the stage of both demographic and
187 spatial expansion (Fig 5C). Interestingly, geographical distribution of both lineages shows that
188 the spatial expansion of southern lineage was efficient enough to spread eastwards into the

189 Black Sea and colonise effectively the north-western Black Sea coast. The northern lineage has
190 spread only in the Pontic region.

191 ***Morphometric data***

192 The results of PCA and ANOSIM revealed no differences in the analysed morphometric traits
193 between the southern and the northern lineages, neither in males nor in females (Fig. 3B). In
194 PCA (Fig. 3B), a very weak gradient ($R=0.03$) could be seen in case of female body length.
195 Females from the northern lineage clade seem to be slightly larger than those from the southern
196 one (body length; ANOSIM Pairwise Tests $p=0.03$).

197

198 **Discussion and conclusions**

199 ***Cryptic diversity of Calomera littoralis***

200 Known as very important hotspot of biodiversity, endemism and cryptic diversity (eg. Myers
201 et al., 2000; Kryštufek & Reed, 2004; Huemer & Timossi, 2014; Blondel et al., 2010; Previšić
202 et al., 2014; Čaković et al., 2015), the southern Europe holds also most diverse tiger beetle
203 fauna in the entire Palearctic realm (Jaskuła, 2011). Presence of cryptic diversity was already
204 pointed out for *Cicindela hybrida* in the Mediterranean (Cardoso, Serrano & Vogler, 2009) as
205 well as for several species of tiger beetles occurring in other part of the world (Vogler &
206 Pearson, 1996; López-López, Hudson & Galián, 2012). Thus, existence of well-defined OTUs
207 within *Calomera littoralis* is not surprising in the studied area. The level of divergence, 0.04
208 K2p distance, between the northern and the southern lineage is similar as those found between
209 species of tiger beetles in other studies (eg. Cardoso & Vogler, 2005; López-López, Abdul Aziz
210 & Galián, 2015). Interestingly, we could not detect any conclusive morphological differences
211 between the two lineages based on the multivariate analysis of eight morphometric traits. It
212 must be mentioned that three subspecies of *Calomera littoralis*, described on the basis of
213 morphology, were reported from the studied area: *C. l. nemoralis* from all the studied Balkan

214 countries, Crete, Moldova, western Ukraine and western Turkey; *C. l. conjunctaepustulata*
215 (Dokhtouroff, 1887) from the Azov Sea area; *C. l. winkleri* (Mandl, 1934) from Crete and the
216 coastal zone of southern Turkey (Werner, 1991; Putschkov & Matalin, 2003; Avgin &
217 Özdikmen, 2007). However, the morphological differences between the subspecies, such as
218 body size, maculation of elytra and shape of aedeagus, are poorly defined and did not allow the
219 identification of the studies material further than to the species level. Unfortunately, we had no
220 opportunity to study the topotypical material – Provence, France, is *locus typicus* for *C. l.*
221 *nemoralis*, Tibet for *C. l. conjunctaepustulata*, and Cyprus for *C. l. winkleri*. Thus, we cannot
222 exclude a possibility that the two lineages we found in our material overlap with any of the
223 above mentioned subspecies. However, only a further taxonomic revision combining more
224 phenotypic traits, including e.g. cuticle ultrastructure, with several, mitochondrial and nuclear
225 DNA data, could help to resolve this problem. Until such revision is done, we propose to use
226 the tentative name “*Calomera littoralis* complex” for populations from the studied area.

227 ***Phylogeography of Calomera littoralis***

228 Occurrence of *C. littoralis* in Europe is restricted predominantly to marine shorelines with
229 sandy beaches and salt marshes as main habitats (e.g. Franzen, 2006; Jaskuła, 2011; Serrano,
230 2013). In the eastern Mediterranean it is distributed continuously all along the Adriatic and
231 Aegean coasts, Turkish Straits and the Black Sea coastline (Cassola & Jaskuła, 2004; Jaskuła
232 Pešić & Pavicević 2005; Franzen, 2006; Jaskuła, 2007a; Jaskuła 2007b). However, pronounced
233 genetic structure with two divergent operational taxonomic units (OTUs) implies prolonged
234 spatial isolation in the evolutionary past of this species. The observed level of divergence
235 indicates that this isolation initiated an allopatric speciation. Their present distribution i.e.
236 sympatry in the Pontic region reveals secondary contact of the already divergent lineages in this
237 area. The Bayesian time-calibrated reconstruction of phylogeny shows that split between these
238 OTUs begun in early Pleistocene. This coincides with beginning of recurrent glaciations

239 resulting in eustatic sea level changes and climate aridisation that ever since dominated the
240 global climate and landscape/habitat distribution (Fagan, 2009). In the Mediterranean and in
241 the Pontic region such global effects overlaid and strengthen the local effects of tectonic plate
242 collision leading to Alpine orogeny, i.e. local land uplift and subsidence resulting in isostatic
243 sea level changes, salinity fluctuations from freshwater to fully marine and habitat mosaicism
244 (Stanley & Blanpied, 1980). For example, during that time the connections of Pontic basin to
245 Mediterranean Sea was lost and regained for more than a dozen times (Kerey et al., 2004;
246 Badertscher et al., 2011). A profound impact of these events on the evolution and, hence,
247 distribution of local both aquatic (Audzijonyte, Daneliya & Vainola, 2006; Nahavandi et al.,
248 2013) and terrestrial taxa (eg. Böhme et al., 2007; Ferchaud et al., 2012). We can assume that
249 in case of *C. littoralis*, a halophilic species bound to coastal habitats, sea level fluctuations
250 would significantly affect its distribution. The 2 Ma divergence time for *C. littoralis* OTUs
251 derived from our data coincides with one particular disconnection of the Mediterranean and
252 Pontic basins. At that time, from ca. 2 to ca. 1.5 Ma, the Meothic Sea, one of several
253 predecessors of the Black Sea, turned into the predominantly freshwater Pontos Sea/Lake
254 (Grinevetsky et al., 2015). This surely broke the formerly continuous stretch of coastal
255 habitats connecting the two basins and thus, could be an effective barrier leading to split of
256 *C. littoralis* population into the allopatric southern and northern lineages. Their detailed
257 history is impossible to unravel, yet results of BSP analyses reconstructing past changes in
258 effective population size indicate that both lineages started their demographic expansions at ca.
259 0.15 Ma. This coincides with the terminal stage of MIS-6, i.e. Wartanian/Saalian glaciation,
260 and beginning of MIS-5e, i.e. Eemian interglacial (Lisiecki & Raymo, 2005; Marks, 2011). The
261 latter was characterized by warmer climate and sea level higher by 6-9 m in comparison to
262 Holocene (Kopp et al., 2009; Dutton & Lambeck, 2012). In result, a wide connection between
263 Mediterranean and the Pontic basin was re-established and the coastal habitats extended again,

264 enabling exchange of faunas. Due to deficiency of local studies, it is hard to compare our results
265 to evolutionary history of any other terrestrial taxa in the area. However, a wealth of studies
266 showing very similar spatiotemporal scenario in animal taxa comes from the coastal regions
267 of the Gulf of Mexico and the adjacent Atlantic coast (summarised by Avise, 1992). During
268 Pleistocene, Cuba was connected with a land bridge to the Florida Peninsula what lead to
269 divergence of populations of several terrestrial and aquatic animals, including also a local
270 tiger beetle species *Cicindela dorsalis* Say, 1817 (Vogler & DeSalle, 1993). Interestingly
271 enough, however according to our results both lineages are until now in the stage of
272 demographic and spatial expansion, only the southern one has crossed the present Turkish
273 straits. This asymmetry is hard to explain. Another interesting fact is that the isolation of Pontic
274 basin from Mediterranean during the following Weichselian glaciation did not have probably
275 any effect on the demography and phylogeography of the species. Based on the mitochondrial
276 DNA marker only we cannot also conclude, whether the secondary contact of the divergent
277 lineages effected in hybridization and or introgression. Answering this question requires
278 employment of nuclear marker, what leaves a space for the future studies – much wider in terms
279 of geographic coverage and molecular markers used.

280 Concluding, we have demonstrated that Pleistocene glaciations and associated sea level changes
281 in the Mediterranean/Pontic region had a profound effect on the genetic diversity and
282 distribution of widely distributed coastal insect species, generating some level of cryptic
283 diversity. Our case study casts more light on the evolutionary relationships between populations
284 of terrestrial animals inhabiting both the Mediterranean and Black Sea shorelines – a
285 phenomenon that is still weakly explored in literature.

286

287 **Acknowledgements**

288 The first author would like to thank to Iwona Jaroszevska, Piotr Józwiak, Błażej Pawicki,
289 Maciej Podsiadło, Anna Stepień, and Bartosz Ukleja for their kind help during and material
290 collecting TB-Quest Expedition to the Balkans.

291

292 **Literature**

293 Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment Search
294 Tool. *Journal of Molecular Biology* 215: 403-410.

295 Akin C, Bilgin CC, Beerli P, Westaway R, Ohst T, Litvinchuk SN, Uzzell T, Bilgin M, Hotz
296 H, Guex GD, Plötner J. 2010. Phylogeographic patterns of genetic diversity in eastern
297 Mediterranean water frogs were determined by geological processes and climate change in the
298 Late Cenozoic. *Journal of Biogeography* 37 (11): 2111-2124.

299 Audzijonyte A, Daneliya ME, Vainola R. 2006. Comparative phylogeography of Ponto-
300 Caspian mysid crustacean: isolation and exchange among dynamic inland sea basins.
301 *Molecular Ecology* 15: 2969–2984.

302 Avgin S, Özdikmen H. 2007. Check-list of tiger beetles of Turkey with review of distribution
303 and biogeography (Coleoptera: Cicindelidae). *Munis Entomology and Zoology* 2 (1): 87-102.

304 Avise JC. 1992. Molecular population structure and the biogeographic history of a regional
305 fauna: a case history with lessons for conservation biology. *Oikos* 63: 62-76.

306 Badertscher S, Fleitmann D, Cheng H, Edwards RL, Göktürk OM., Zumbühl A, Leuenberger
307 M, Tüysüz O. 2011. Pleistocene water intrusions from the Mediterranean and Caspian seas
308 into the Black Sea. *Nature Geoscience* 4: 236– 239.

309 Blondel J, Aronson J, Bodiou J-Y, Boeuf G. 2010. *The Mediterranean Region. Biological*
310 *Diversity in Space and Time*. 2nd Edition. Oxford: Oxford University Press.

- 311 Böhme MU, Fritz U, Kotenko T, Džukić G, Ljubisavljević K, Tzankov N, Berendonk TU.
312 2007. Phylogeography and cryptic variation within the *Lacerta viridis* complex (Lacertidae,
313 Reptilia). *Zoologica Scripta* 36: 119–131.
- 314 Brower AVZ. 1994. Rapid morphological radiation and convergence among races of the
315 butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA Evolution.
316 *Proceedings of the National Academy of Sciences USA* 91: 6491-6495.
- 317 Caković D, Stešević D, Schönswetter P, Frajman B. 2015. How many taxa? Spatiotemporal
318 evolution and taxonomy of *Amphoricarpos* (Asteraceae, Carduoideae) on the Balkan
319 Peninsula. *Organisms Diversity & Evolution* 15(3): 429-445.
- 320 Cardoso A, Vogler AP. 2005. DNA taxonomy, phylogeny and Pleistocene diversification of
321 the *Cicindela hybrida* species group (Coleoptera: Cicindelidae). *Molecular Ecology* 14: 3531-
322 3546.
- 323 Cardoso A, Serrano A, Vogler AP. 2009. Morphological and molecular variation in tiger
324 beetles of the *Cicindela hybrida* complex: is an ‘integrative’ taxonomy possible? *Molecular*
325 *Ecology* 18: 648-664.
- 326 Cassola F, Jaskuła R. 2004. Material to the knowledge of the tiger beetles of Romania
327 (Coleoptera: Cicindelidae). *Polskie Pismo Entomologiczne* 73: 193–214.
- 328 Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. 2003.
329 Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Research* 31:
330 3497–3500.
- 331 Clarke KR, Gorley RN. 2006. *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth.
- 332 Drummond AJ, Rambaut A, Shapiro B, Pybus OG. 2005. Bayesian coalescent inference of
333 past population dynamics from molecular sequences. *Molecular Biology and Evolution* 22:
334 1185-1192.

- 335 Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012. Bayesian Phylogenetics with BEAUti
336 and the BEAST 1.7. *Molecular Biology and Evolution* 29: 1969–1973.
- 337 Dutton A, Lambeck K. 2012. Ice volume and sea level during the last interglacial. *Science*
338 337 (6091): 216–219.
- 339 Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: a new series of programs to perform
340 population genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10:
341 564-567.
- 342 Fagan B (ed.). 2009. *The complete Ice Age. How climate change shaped the world*. London:
343 Thames & Hudson Ltd.
- 344 Ferchaud A-L, Ursenbacher S, Cheylan M, Luiselli L, Jelić D, Halpern B, Major A, Kotenko
345 T, Keyan N, Behrooz R, Crnobrnja-Isailović J, Tomović L, Ghira I, Ioannidis Y, Arnal V,
346 Montgelard C. 2012. Phylogeography of the *Vipera ursinii* complex (Viperidae):
347 mitochondrial markers reveal an east–west disjunction in the Palaeartic region. *Journal of*
348 *Biogeography* 39: 1836–1847.
- 349 Franzen M. 2006. Verbreitung und Lebensräume der Sandlaufkäfer der Peloponnes-Halbinsel,
350 Griechenland (Coleoptera, Cicindelidae). *Nachrichtenblatt der Bayerischen Entomologen* 55
351 (3/4): 46–64.
- 352 Grinevetsky SR, Zonn IS, Zhiltsov SS, Kosarev AN, Kostianoy AG. 2015. *The Black Sea*
353 *Encyclopedia*. Berlin: Springer.
- 354 Hebert PDN, Cywinska A, Ball SL, deWaard JR. 2003a. Biological identification through
355 DNA barcodes. to the fast publication and worldwide dissemination of high-quality research.
356 *Proceedings of the Royal Society of London B: Biological Sciences* 270: 313–321.
- 357 Hebert PDN, Ratnasingham S, deWaard JR. 2003b. Barcoding animal life: cytochrome c
358 oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal*
359 *Society of London B: Biological Sciences* Suppl. 27: 96–99.


- 360 Hillis DM, Moritz C, Mable BK. 1996. *Molecular Systematics*, 2nd ed. Sunderland: Sinauer
361 Associates.
- 362 Huemer P., Timossi G. 2014. *Sattleria* revisited: unexpected cryptic diversity on the Balkan
363 Peninsula and in the south-eastern Alps (Lepidoptera: Gelechiidae). *Zootaxa* 3780 (2): 282-
364 296.
- 365 Huson DH, Bryant D. 2006. Application of Phylogenetic Networks in Evolutionary Studies,
366 *Molecular Biology and Evolution* 23(2): 254-267.
- 367 Jaskuła R. 2007a. Remarks on distribution and diversity of the tiger beetle fauna (Coleoptera:
368 Cicindelidae) of Albania. *Fragmenta Faunistica* 50 (2): 127–138.
- 369 Jaskuła R. 2007b. Further records of tiger beetles from Romania (Coleoptera: Cicindelidae).
370 *Cicindela* 39 (1–2): 27–34.
- 371 Jaskuła R. 2011. How unique is the tiger beetle fauna (Coleoptera, Cicindelidae) of the
372 Balkan Peninsula? *ZooKeys* 100: 487-502.
- 373 Jaskuła R. 2013. Unexpected vegetarian feeding behaviour of a predatory tiger beetle
374 *Calomera littoralis nemoralis* (Olivier, 1790) (Coleoptera: Cicindelidae). *Journal of the*
375 *Entomological Research Society* 15(1): 01–06.
- 376 Jaskuła R. 2015. The Maghreb – one more important biodiversity hot spot for tiger beetle
377 fauna (Coleoptera, Carabidae, Cicindelinae) in the Mediterranean region. *ZooKeys* 482: 35-
378 53.
- 379 Jaskuła R, Pešić V, Pavicević D. 2005. Remarks on distribution and diversity of the tiger
380 beetle fauna of Montenegro (Coleoptera: Cicindelidae). *Fragmenta Funistica* 4 (1): 15–25.
- 381 Kerey IE, Meric E, Tunoglu C, Kelling G, Brenner RL, Dogan AU. 2004. Black Sea–
382 Marmara Sea Quaternary connections: new data from the Bosphorus, Istanbul, Turkey.
383 *Palaeogeography, Palaeoclimatology, Palaeoecology* 204: 277–295.

- 384 Kopp RE, Simons FJ, Mitrovica JX, Maloof AC, Oppenheimer M. 2009. Probabilistic
385 assessment of sea level during the last interglacial stage. *Nature* 462 (7275): 863–867.
- 386 Kotlík P, Bogutskaya NG Ekmekçi FG. 2004. Circum Black Sea phylogeography of *Barbus*
387 freshwater fishes: divergence in the Pontic glacial refugium. *Molecular Ecology* 13: 87-95.
- 388 Kryštufek B, Reed M. 2004 Patterns and Process in Balkan Biodiversity – an overview. In:
389 Griffiths HI, Kryštufek B, Reed JM. (Eds) *Balkan Biodiversity: Pattern and Process in the*
390 *European Hotspot*. Dordrecht, Kluwer Academic Publishers, 203–217.
- 391 Librado P, Rozas J. 2009. Dna SPv5: a software for comprehensive analysis of DNA
392 polymorphism data. *Bioinformatics* 25: 1451–1452.
- 393 Lisiecki LE., Raymo ME. 2005. A Pliocene-Pleistocene stack of 57 globally distributed
394 benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20 (10): 1-17.
- 395 López-López A, Abdul Aziz A, Galián J. 2015. Molecular phylogeny and divergence time
396 estimation of *Cosmodela* (Coleoptera: Carabidae: Cicindelinae) tiger beetle species from
397 Southeast Asia. *Zoologica Scripta* 44: 437-445.
- 398 López-López A, Hudson P, Galián J. 2012. The blackburni/murchisona species complex in
399 Australian *Pseudotetracha* (Coleoptera: Carabidae: Cicindelinae: Megacephalini): evaluating
400 molecular and karyological evidence. *Journal of Zoological Systematics and Evolutionary*
401 *Research* 50 (3): 177-183.
- 402 Mandl K. 1981. Revision der unter *Cicindela lunulata* F. im Weltkatalog der Cicindelinae
403 zusammengefaßten Formen (Col., Cicindelidae). *Entomologische Arbeiten aus dem Museum*
404 *Gg. Frey* 29: 117-176.
- 405 Marks L. 2011. Quaternary glaciations in Poland. [in:]: Ehlers J., Gibbard P.L., Hughes P.D.
406 (eds.) Quaternary Glaciations - Extent and Chronology, a closer look. *Developments in*
407 *Quaternary Science* 15: 299-303.

- 408 Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Jennifer K. 2000. Biodiversity
409 hotspots for conservation priorities. *Nature* 403: 853–858.
- 410 Nahavandi N, Ketmaier V, Plath M, Tiedemann R. 2013. Diversification of Ponto-Caspian
411 aquatic fauna: Morphology and molecules retrieve congruent evolutionary relationships in
412 *Pontogammarus maeoticus* (Amphipoda: Pontogammaridae). *Molecular Phylogenetics and*
413 *Evolution* 69: 1063–1076.
- 414 Neilson ME, Stepien CA. 2011. Historic speciation and recent colonization of Eurasian
415 monkey gobies (*Neogobius fluviatilis* and *N. pallasii*) revealed by DNA sequences,
416 microsatellites, and morphology. *Diversity and Distributions* 17: 688–702.
- 417 Pearson DL, Cassola F. 2005. A quantitative analysis of species descriptions of tiger beetles
418 (Coleoptera: Cicindelidae), from 1758 to 2004, and notes about related developments in
419 biodiversity studies. *The Coleopterologist Bulletin* 59 (2): 184–193.
- 420 Pearson DL, Vogler AP. 2001. *Tiger beetles: the evolution, ecology, and diversity of the*
421 *cicindelids*. Ithaca and London: Cornell University Press.
- 422 Pfiffner OA. 2014. *Geology of the Alps*. New York: Wiley Blackwell.
- 423 Previšić A, Wolfram G, Vitecek S, Waringer J. 2014. Cryptic diversity of caddisflies in the
424 Balkans: The curious case of *Ecclisopteryx* species (Trichoptera: Limnephilidae). *Arthropod*
425 *Systematics and Phylogeny* 72(3): 309-329.
- 426 Puillandre N, Lambert A, Brouillet S, Achaz G. 2012. ABGD, Automatic Barcode Gap
427 Discovery for primary species delimitation. *Molecular Ecology* 21(8): 1864-77.
- 428 Putschkov AV, Matalin AV. 2003. Subfamily Cicindelinae Latreille, 1802. In: Löbl L,
429 Smetana A. (Eds) *Catalogue of Palaearctic Coleoptera. V.1. Archeostemata – Myxophaga –*
430 *Adephaga*. Strenstrup: Apollo Books, 99–118.
- 431 Rogers AR., Harpending H. 1992. Population growth makes waves in the distribution of
432 pairwise genetic differences. *Molecular Biology and Evolution* 9: 552–569.

- 433 Serrano J. 2013. *New catalogue of the family Carabidae of the Iberian Peninsula*
434 *(Coleoptera)*. Madrid: Editum - Ediciones de la Universidad de Murcia.
- 435 Simon C, Frati F, Beckenbach AT, Crespi B, Liu H, Flook P. 1994. Evolution, weighting, and
436 phylogenetic utility of mitochondrial gene sequences and a compilation of conserved
437 polymerase chain reaction primers. *Annals of Entomological Society of America* 87: 51–701.
- 438 Slatkin M, Hudson RR. 1991. Pairwise comparisons of mitochondrial DNA sequences in
439 stable and exponentially growing populations. *Genetics* 129: 555–562.
- 440 Sota T, Liang H, Enokido Y, Hori M. 2011. Phylogeny and divergence time of island tiger
441 beetles of the genus *Cylindera* (Coleoptera: Cicindelidae) in East Asia. *Biological Journal of*
442 *the Linnean Society* 102: 715-727.
- 443 Stanley DJ, Blanpied C. 1980. Late Quaternary water exchange between the eastern
444 Mediterranean and the Black Sea. *Nature* 285 (5766): 537-541.
- 445 Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA
446 polymorphism. *Genetics* 123: 585–595.
- 447 Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular
448 Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–
449 2729.
- 450 Tsuji K, Hori M, Phyu MH, Liang H, Sota T. 2016. Colorful patterns indicate common
451 ancestry in diverged tiger beetle taxa: molecular phylogeny, biogeography, and evolution of
452 elytral coloration of the genus *Cicindela* subgenus *Sophiodela* and its allies. *Molecular*
453 *Phylogenetics and Evolution* 95: 1-10.
- 454 Vogler AF, DeSalle R. 1993. Phyllogeographic patterns in coastal North American tiger
455 beetles (*Cicindela dorsalis* Say) inferred from mitochondrial DNA sequences. *Evolution* 47
456 (4): 1192-1202.

- 457 Vogler AP, Pearson DL. 1996. A molecular phylogeny of the tiger beetles (Cicindelidae):
458 congruence of mitochondrial and nuclear rDNA data sets. *Molecular Phylogenetics and*
459 *Evolution* 6 (3): 321–338.
- 460 Vogler AP, Knisley CB, Glueck SB, Hill JM, DeSalle R. 1993. Using molecular and
461 ecological data to diagnose endangered populations of the puritan tiger beetle *Cicindela*
462 *puritana*. *Molecular Ecology* 2: 375-383.
- 463 Werner K. 1991. *Cicindelidae Regionis Palaearcticae. Megacephalini: Megacephala,*
464 *Cicindelini 1: Cicindela – Lophyridia. The Beetles of the World 13.* Compiègne: Sciences
465 Nat..
- 466 Woodcock MR, Kippenhan MG, Knisley CB, Foster JA. 2007. Molecular genetics of
467 *Cicindela (Cylindera) terricola* and elevation of *Cicindela lunalonga* to species level, with
468 comments on its conservation status. *Conservation Genetics* 8: 865-877.
- 469




470

471 **Figure 1.** Distribution and sampling of *Calomera littoralis* in Europe. (A) General distribution472 of *Calomera littoralis* in Europe shown as red-shaded area. (B) Picture of *Calomera littoralis*

473 beetle. (C) Sampling sites in Balkan Peninsula, Black Sea region and Turkey shown as black

474 dots. Localities coded as in Table 1.


475



476


477 **Figure 2.** Body parameters measured in *Calomera littoralis*. 1 – RLM – right mandible length,
478 2 – WH – width of head, 3 – LH – length of head, 4 – MPW – maximum pronotum width, 5 –
479 PL – pronotum length, 6 – EL – elytra length, 7 – MEW – maximum elytra width, 8 – TBL –
480 total body length.

481



482


483 **Figure 3.** (A) Automatic Barcode Gap Discovery (ABGD) analysis of *Calomera littoralis* and
 484 (B) Results of Principal Component Analysis performed for investigated specimens on main
 485 body dimensions. S – southern lineage, N – northern lineage, RLM – right mandible length,
 486 WH – width of head, LH – length of head, MPW – maximum pronotum width, PL – pronotum
 487 length, EL – elytra length, MEW – maximum elytra width, TBL – total body length. Both in
 488 ABGD and PCA analyses 169 specimens from 43 sites from the Mediterranean and the Pontic
 489 areas were used.



490

491 **Figure 4.** (A) Geographic distribution of haplogroups from southern (blue circles) and northern (yellow circles) lineages (B).

492



493

494 **Figure 5.** Phylogeny and demography of *Calomera littoralis*. (A) Maximum clade credibility chronogram with a strict molecular clock model
 495 inferred from COI sequences. The numbers next to the respective node indicate Bayesian posterior probabilities higher than 0.5. (B) Mismatch
 496 plots for southern and northern lineage. Thin solid lines indicate expected frequency under model of population demographic expansion, thick solid
 497 lines represent observed frequency, and dashed lines indicate 95% confidence intervals for the observed mismatch. SSD, sum of squared deviation;

498 r, Harpending's raggedness index; D, Tajima's D. (C) Bayesian skyline plots for southern and northern lineages of *Calomera littoralis*. Solid lines
499 indicate the median posterior effective population size through time; dashed lines indicate the 95% highest posterior density interval for each
500 estimate.