

**A peer-reviewed version of this preprint was published in PeerJ on 2 November 2016.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.2640) (peerj.com/articles/2640), which is the preferred citable publication unless you specifically need to cite this preprint.

Ovejero Aguilar RJA, Jahn GA, Soto-Gamboa M, Novaro AJ, Carmanchahi P. 2016. The Ecology of Stress: linking life-history traits with physiological control mechanisms in free-living guanacos. PeerJ 4:e2640 <https://doi.org/10.7717/peerj.2640>

# The Ecology of Stress: linking life-history traits with physiological control mechanisms in free-living guanacos.

Ramiro J.A. Ovejero Aguilar<sup>Corresp., 1</sup>, Graciela A Jahn<sup>2</sup>, Mauricio Soto-Gamboa<sup>3</sup>, Andrés Novaro<sup>4</sup>, Pablo Carmanchahi<sup>5</sup>

<sup>1</sup> Laboratorio de Ecología Conductual, Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Laboratorio de Interacciones Ecológicas (LIE) IADIZA-CONICET, Mendoza-Argentina.

<sup>2</sup> Laboratorio de Reproducción y Lactancia, IMBECU-CONICET, Mendoza, Argentina.

<sup>3</sup> Laboratorio de Ecología Conductual, Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile

<sup>4</sup> Programa Estepa-Andino Patagonica-CONICET-PATAGONIA NORTE-INIBIOMA, Neuquén, Argentina

<sup>5</sup> Grupo de Investigación de Eco-fisiología de Fauna Silvestre (GIEFAS), Asentamiento Universitario de San Martín de los Andes (AUSMA), Universidad Nacional del Comahue (INIBIOMA-CONICET -AUSMA-UNCo), Neuquén, Argentina

Corresponding Author: Ramiro J.A. Ovejero Aguilar

Email address: rovejero@mendoza-conicet.gob.ar

**Background**-Providing the context for the evolution of life-history traits, habitat features constrain successful ecological and physiological strategies. In vertebrates, a key response to life's challenges is the activation of the Stress (HPA) and Gonadal (HPG) axes. Much of the interest in stress ecology is motivated by the desire to understand the physiological mechanisms in which the environment affects fitness. As reported in the literature, several intrinsic and extrinsic factors affect variability in hormone levels. In both social and non-social animals, the frequency and type of interaction with conspecifics, as well as the status in social species, can affect HPA axis activity, resulting in changes in the reproductive success of animals. We predicted that a social environment can affect both guanaco axes by increasing the secretion of testosterone (T) and Glucocorticoid (GCs) in response to individual social interactions and the energetic demands of breeding. Assuming that prolonged elevated levels of GCs over time can be harmful to individuals, it is predicted that the HPA axis suppresses the HPG axis and causes T levels to decrease, as GCs increase.

**Methods**-All of the data for individuals were collected by non-invasive methods (fecal samples) to address hormonal activities. This is a novel approach in physiological ecology because feces are easily obtained through non-invasive sampling in animal populations.

**Results**- As expected, there was a marked adrenal (p-value= .344e-12) and gonadal (p-value= 0.002656) response due to seasonal variation in *Lama guanicoe*. No significant differences were found in fecal GCs metabolites between males/females\*season for the entire study period (p-value= 0.2839). Despite the seasonal activity variation in the hormonal profiles, our results show a positive correlation (p-value= 1.952e-11, COR=0.50) between the adrenal and gonadal system. The marked endocrine ( $r^2 = 0.806$ ) and gonad ( $r^2 = 0.7231$ ) response due to seasonal variation in male guanaco individuals highlights the individual's energetic demands according to life-history strategies. This is a remarkable result because no inhibition was found between the axes as theory suggests. Finally, the dataset was used to build a reactive scope model for guanacos.

**Discussion**-Guanacos cope with the trade-off between sociability and reproductive benefits and costs, by regulating their GCs and T levels on a seasonal basis, suggesting an adaptive role of both axes to different habitat pressures. The results presented here highlight the functional role of stress and gonad axes on a critical phase of a male mammal's life—the mating period—when all of the resources are at the

disposal of the male and must be used to maximize the chances for reproductive success.

The Ecology of Stress: linking life-history traits with physiological control mechanisms in free-living guanacos

Ovejero Ramiro<sup>abe\*</sup>, Jahn Graciela A<sup>d</sup>, Soto Gamboa Mauricio<sup>e</sup>, Novaro Andrés and Carmanchahi Pablo<sup>b</sup>.

<sup>a</sup>Laboratorio de Interacciones Ecológicas, LIE-IADIZA-CONICET-CCT-MENDOZA, Mendoza-Argentina.

<sup>b</sup>Grupo de Investigación de Eco-fisiología de Fauna Silvestre (GIEFAS), Asentamiento Universitario de San Martín de los Andes (AUSMA), Universidad Nacional del Comahue (INIBIOMA-CONICET -AUSMA-UNCo), Neuquén, Argentina.

<sup>d</sup>Laboratorio de Reproducción y Lactancia, IMBECU- CCT-MENDOZA, Mendoza, Argentina.

<sup>e</sup>Laboratorio de Ecología Conductual, Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile.

\*Corresponding author: Ramiro Ovejero: e-mail:rovejero@mendoza-conicet.gov.ar

## Abstract

**Background**-Providing the context for the evolution of life-history traits, habitat features constrain successful ecological and physiological strategies. In vertebrates, a key response to life's challenges is the activation of the Stress (HPA) and Gonadal (HPG) axes. Much of the interest in stress ecology is motivated by the desire to understand the physiological mechanisms in which the environment affects fitness. As reported in the literature, several intrinsic and extrinsic factors affect variability in hormone levels. In both social and non-social animals, the frequency and type of interaction with conspecifics, as well as the status in social species, can affect HPA axis activity, resulting in changes in the reproductive success of animals. We predicted that a social environment can affect both guanaco axes by increasing the secretion of testosterone (T) and Glucocorticoid (GCs) in response to individual social interactions and the energetic demands of breeding. Assuming that prolonged elevated levels of GCs over time can be harmful to individuals, it is predicted that the HPA axis suppresses the HPG axis and causes T levels to decrease, as GCs increase.

**Methods**-All of the data for individuals were collected by non-invasive methods (fecal samples) to address hormonal activities. This is a novel approach in physiological ecology because feces are easily obtained through non-invasive sampling in animal populations.

**Results**- As expected, there was a marked adrenal ( $p\text{-value} = .344\text{e-}12$ ) and gonadal ( $p\text{-value} = 0.002656$ ) response due to seasonal variation in *Lama guanicoe*. No significant differences were found in fecal GCs metabolites between males/females\*season for the entire study period ( $p\text{-value} = 0.2839$ ). Despite the seasonal activity variation in the hormonal profiles, our results show a positive correlation ( $p\text{-value} = 1.952\text{e-}11$ ,  $\text{COR} = 0.50$ ) between the adrenal and gonadal system. The marked endocrine ( $r^2 = 0.806$ ) and gonad ( $r^2 = 0.7231$ ) response due to seasonal variation in male guanaco individuals highlights the individual's energetic demands according to life-history strategies. This is a remarkable result because no inhibition was found between the axes as theory suggests. Finally, the dataset was used to build a reactive scope model for guanacos.

**Discussion**-Guanacos cope with the trade-off between sociability and reproductive benefits and costs, by regulating their GCs and T levels on a seasonal basis, suggesting an adaptive role of both axes to different habitat pressures. The results presented here highlight the functional role of stress and gonad axes on a critical phase of a male mammal's life—the mating period—when all of the resources are at the disposal of the male and must be used to maximize the chances for reproductive success.

**Key-words:** Stress ecology, hormonal profiles in wildlife, *Lama guanicoe*, sociality, reproduction, non-invasive methods.

# Introduction

Natural selection shapes phenotypic traits that permit organisms to respond appropriately to intrinsic and extrinsic factors (Roff, 1992; Stearns, 1992; Bozinovic, 2002; Nespolo et al., 2003). Genetic variation among organisms determines the resource allocation trade-offs among life-history traits (e.g. age-specific growth, reproduction and survival schedules, and sociability) and are also determined by phylogeny. Environmental, ecological and indirect demographic processes (extrinsic factors) create selective pressure that impinge on an organism's lifetime fitness (Stearns, 1992; Zera & Harshman, 2001). Therefore, understanding life-history variation (from an evolutionary perspective) among taxa requires an understanding of the physiological mechanisms that link genes with phenotype and how extrinsic factors regulate trade-offs between survival and reproduction. Ecophysiological, and particularly knowledge about endocrine mechanisms, is key to understanding how life-history variation and trade-offs arise and are retained (Ketterson & Nolan 1999; Zera & Harshman, 2001; Crespi et al., 2013).

The stress response is a mechanism by which an individual of a given species copes with environmental changes (stressors) (Boosntra et al., 2001; Winfield, 2005; Romero et al., 2009; McEwen and Wingfield, 2010; Ovejero and Carmanchahi, 2012; Ovejero 2013). Perception of a stressful situation activates the sympathetic adrenomedullary system and the hypothalamic pituitary adrenocortical (HPA) axis (Mostl and Palme, 2002). Activation of the HPA axis stimulates the release of steroid hormones (glucocorticoids GCs, e.g., corticosterone, cortisol). GCs in turn activate the mobilization of energy necessary to cope with adverse environmental conditions (Dallman et al., 2007; Sapolsky, 2000). In the last 30 years, the word "Stress" has been strongly criticized due to the ambiguity of the term and several authors have been researching allostasis and attempting to connect biomedical and ecological data (McEwen & Wingfield 2003; Romero et al., 2009, 2011). Allostasis is the daily and seasonal process of maintaining physiological parameters (homeostasis) within life-sustaining ranges, via allostatic mediators

(McEwen & Wingfield, 2003). The allostatic model has generated criticism and Romero et al. (2009) have proposed the reactive scope model, to maintain the strengths but avoid the weaknesses of the allostatic model. The present work contributes to the debate about the applicability of the allostatic model to nature.

Extrinsic factors, like anthropogenic disturbance, biological invasions, or climate change (in terms of allostasis) can negatively affect behavior, reproduction, immune function and growth through prolonged activation of the HPA axis (Moberg, 1991; Boonstra and Singleton, 1993; Goymann and Wingfield, 2004; Pride, 2005; Pereira et al., 2006). Glucocorticoids (GCs) are often considered the allostatic mediators of the stress response, and they have been the focus of research for decades (Sapolsky, 2000; Ovejero 2013). Much less is known, however, about the role of GCs' in free-living animals (Bonier et al., 2009). The role of baseline or stress-induced GC in the modulation of an individual or populations performance, is still essentially unknown. Understanding, social factors and their endocrine correlates (baseline measures of GCs and T) in wild animals has potential implications for wildlife management, population ecology, reproductive biology and evolution. The social environment is therefore one of the primary sources of information that can induce allostatic responses (Boonstra et al., 2001; Rubenstein and Shen, 2009; Creel et al., 2013. In both social and non-social animals, the frequency and type of interaction with conspecifics, as well as status in social species, can affect HPA and HPG axis activity and ultimately the reproductive success of animals.

During the breeding season of many mammals, males commonly increase testosterone levels in response to social stimuli such as presence of male competitors or potential mates (Enstrom et al., 1997; Cavigelli and Parer 2000; Sinervo et al., 2000; Ostner and Kappeler, 2002; Soto-Gamboa et al., 2005; Schradin, 2008; Creel et al., 2013. This physiological mechanism (androgen response) modulated by social environment, is likely to be mainly important when considering their roles (testosterone-mediated traits) in the evolution of mating systems and life histories (Wingfield et al., 1990; Knapp and Moore, 1997; Faulkes and Abbott, 1997; Hirschenhauser and Oliveira, 2006; Young et al., 2006; Rubenstein et al., 2009; McGlothilin et al., 2010. In addition, testosterone levels can also affect female preference (Enstrom et al., 1997), thus influencing fitness through both intra and intersexual selection; (Sinervo *et al.*, 2000). Although the role of short-term androgen changes has not been completely described, these have long been associated with territoriality; (McGlothilin et al., 2010). The "Challenge Hypothesis" (first described in bird studies) predicts that androgen levels and agonistic interactions are positively related and are socially-context

dependent Wingfield et al. (1997). Among mammals, these predictions have recently been tested, and a positive correlation was found between social cues and testosterone levels associated with territorial defense, group formation, social dominance and hierarchy formation during the breeding season Muller and Wrangham (2004); Creel et al. (2013). Consequently, social stressors provide a context where the individual must respond to a series of repeated stressors over time, depending on the frequency and intensity of the stimulus, and this can lead to various physiological problems, Sapolsky (2005).

Much of the variation in life histories strategies reflects individuals' phenotypic responses to environmental challenges and perceived risks. The adrenal and gonadal system strongly influence behavior, control the annual cycle of development, modulate behavioral and physiological responses to the environment, and establishes important incompatibilities in life stages. It provides a model for studying the connections between mammal physiology and life history. Guanacos (*Lama guanicoe*) are the largest social artiodactyls of South America, and these organisms have a polygynous resource-defense mating system. This wild and endemic camelid has been successfully used as a study model of behavioral interactions (De Lamo et al., 1998; Franklin, 1982; Marino and Baldi, 2008, Taraborelli et al. 2014), ecological studies, (Puig and Videla, 1995; Bank et al., 2003; Puig et al., 2008; Acebes et al., 2010; Ovejero et al. 2011; Ovejero, 2013; Marino and Baldi, 2014; Radovani, 2004 and management (Montes et al., 2006; Zapata et al., 2002; Zapata et al., 2004), but information about the guanaco's physiological ecology remains scarce. More recently, guanacos have also been used as a model of stress response; a study was conducted to determine how management including handling, shearing, and release induce changes in circulating steroid hormones Carmanchahi et al., (2011), however this work was based on an invasive sampling method.

To understand how physiology mediates the relationship between life history and the environment, we predict that: (A)- Social environment positively will affect the HPG and HPA axes by raising the secretion of T and GCs due to individual social interactions and the energetic demands of the breeding season. (B)-Male guanacos will show seasonal variation in the activity of their adrenal and gonadal systems, which perform incompatible functions in different life stages (survival vs reproduction). (C)-Prolonged elevated baseline levels of GCs in male guanacos due to long periods of intense social interaction will be detrimental, costly, and decrease fitness, and we would expect that in this case HPA inhibits the activity of the HPG. Together, these predictions provide a description of the reactive scope model and an understanding of how physiological mechanisms might constrain patterns of variation



in wild South-American camelid life histories. These predictions might also provide the basis for future mammalian studies targeted at understanding phenotype-environment interactions.

## Methods

### Ethics Statement

The present study did not warrant capturing or handling protected or endangered animals. All of the data for individuals were collected by non-invasive methods (fecal samples) to address hormonal activities. This is a novel approach in physiological ecology because feces are easily obtained through non-invasive sampling in animal populations. This is important because the handling necessary for collection of plasma can confound estimates of baseline GC measurements. The described field studies were carried out in a protected area with a permit (files:NO 4350-000019112/Res 238) from the DRNR (Dirección de Recursos Naturales Renovables, Mendoza).

### Non-invasive methods to address hormonal activities

Fresh fecal samples were collected from 334 wild adult guanacos from La Payunia reserve (36.000S and 36.360S; 68.340W and 69.230W, South of Mendoza Province, Argentina). The samples were collected during 7 field surveys (15 days each) over the course of a year. Fresh feces samples were collected from focal individuals (one sample per each indiv.) and were labeled according to the sample location, time, sex, age, social structure and environment variables. To slow microbial activity and to reduce immunoreactivity problems, the samples were froze immediately in liquid nitrogen and stored at -20 C° until the time of analysis (LARLAC-IMBECU-CCT CONICET-MENDOZA). Fieldwork was conducted during the breeding and non-breeding months to assess seasonal variation in hormone levels. Furthermore, 100% of the sampling surveys were done in the same sites. Steroids were extracted from lyophilized fecal samples according to a protocol developed for fecal steroid metabolite extraction for free-range ungulates (for more detail see, Ovejero 2013). The commercial <sup>125</sup>I-cortisol and <sup>125</sup>I-testosterone RIA-KITS (BECKMAN, Coulter Company IM-1841/IM-1119-Immunotech-, Prague, Czech Republic) were used to quantify the levels of cortisol and testosterone in the samples (for more detail see, Ovejero and Carmanchahi, 2012; Ovejero 2013). The analytical sensitivity of the cortisol assay is 3.075 ng / mL. The antibody used in the immunoassay is highly specific for cortisol (100% cross-reaction). The cross-reactivity to other naturally occurring steroids is very

low, for example: aldosterone <0.1%; 8.4% corticosterone; cortisone 1.5%; 11-desoxy-cortisol 18%, etc. The analytical sensitivity of the testosterone assay is 0.025 ng / mL. The antibody used in the immunoassay is highly specific for testosterone (100% cross-reaction). The cross-reactivity to other naturally occurring steroids is very low, for example: 0.00014% aldosterone; estrone <0.3%; 0.08 androsterone; 5-Dihydrotestosterone 10%  $\alpha$ ; 11 $\beta$ -Hydroxytestosterone 2%; corticosterone 8.4%; 1.5% cortisone; cortisol 11-deoxy-18%, etc. In addition, all samples were assayed in triplicate (extraction efficiency test) and duplicate (binding and parallelism test) and reanalyzed whenever the resulting coefficient of variation exceeded 20%. Intra- and inter-assay coefficients of variation of GCs-immunoassay were 6.5% and 11.2% and for testosterone immunoassays were 9.5 and 13.4%, respectively.

## Statistical analyses

Welch two sample T-tests were used to evaluate the seasonal variation response of steroid hormones. Furthermore, two-way ANOVAs were used to evaluate the interaction between sex and season. Generalized linear models were developed (n polynomials function =  $f(x) = a_n X^n + a_{n-1} X^{n-1} + a_{n-2} X^{n-2} + \dots$ ) to describe the hormonal profiles and their seasonal variation in response to the environment. Moreover, a Pearson's correlation was used to test the activity association between the adrenal and gonadal system throughout the year. All statistical analyses were conducted using R statistical software version 3.1.0 (Development Core Team, 2012).

## Results

### Ecological process and fecal-hormone profiles in wild guanacos

As expected, there was a marked adrenal ( $t=7.4016$ ,  $df = 302.174$ ,  $p\text{-value} = 1.344\text{e-}12$ ,  $1-\alpha(0.90)=7.661734\text{-}12.057391$ , mean =  $25.75864/15.89907$  breeding and non-breeding season respectively) and gonadal ( $t=3.2873$ ,  $df = 28.948$ ,  $p\text{-value} = 0.002656$ ,  $1-\alpha(0.90)=0.6213\text{-}1.9508$ , mean =  $3.209810/1.923700$  breeding and non-breeding season respectively for male groups) GCs and T response due to seasonal variation in *Lama guanicoe* individuals. This highlights the individual's energetic demands according to life-history strategies. No significant differences were found in the amounts of fecal GC metabolites between males/females\*season for the entire study period ( $F=1.0757$ ,  $df = 143.529$ ,  $p\text{-value} = 0.2839$ ; Male:  $\text{ng/gr}=22.42$ ,  $1-\alpha(0.90)=1.6887$ ,  $\text{std-e}=0.86$ ,

n=251; Female:  $\text{ng/gr} = 20,71,1 - \alpha(0.90) = 2.6079$ ,  $n = 76$ ,  $\text{std-e} = 1.33$ ). This result was not expected as it was thought that there would be a difference in the allocation of resources to cope with the different roles that males and females have in a population. The second prediction in our study is partially accepted. Despite the seasonal activity variation in the hormonal profiles, our results show a positive correlation ( $t = 7.1524$ ,  $df = 184$ ,  $p\text{-value} = 1.952e-11$ ,  $CI = 0.3457400 \text{ } 0.5719168$ ,  $COR = 0.50$ ) between the adrenal and gonadal system; both peaks of activity concur with the reproductive season. (C)-Prolonged elevated baseline levels of GCs due to long periods of intense social interaction are detrimental, costly and decrease fitness; it is expected that in this context, HPA inhibits the activity of the HPG. The marked endocrine ( $y = 0.0743x^5 - 1.9677x^4 + 19.136x^3 - 82.442x^2 + 147.4x - 58.902$ ;  $r^2 = 0.806$ ) and gonad ( $y = 0.0041x^5 - 0.1102x^4 + 1.0889x^3 - 4.6819x^2 + 7.914x - 0.7892$ ;  $r^2 = 0.7231$ ) response due to seasonal variation in male guanaco individuals (Figure 1) highlights the individual's energetic demands according to life-history strategies. This is a remarkable result because no inhibition was found between the axes as theory suggests. Finally, the dataset was used to build a reactive scope model for guanacos, see figure 2. Most studies measure only one physiological mediator at a time; here the concentration of two mediators is placed on the y-axis for a given time point. This was done because these mediators are correlated in the functional role that each one plays in the normal reactive scope range. In other words, both mediators encompass responses to cope with predictable and unpredictable changes in the environment. The values of each mediator are presumed to exist in four general ranges. Our result shows the ranges of predictive/reactive homeostasis and homeostatic overload/failure. These ranges consist of the seasonal set of point ranges for the physiological mediators.

## Discussion

Our results confirm the prediction that "Social environment positively affects the HPG and HPA axes by raising the secretion of T and GCs due to individual social interactions and the energetic demands of the breeding season." Further to this, our results suggest that social environment and breeding season are stressful scenarios for male and female guanacos. These scenarios are seen to produce associated costs such as increased metabolic rate and energy expenditure due to: territorial/female defense (Male-reproduction strategies), male and female social dominance hierarchies, reproduction, and increased exposure to predators during the breeding season. If the data value observed in March is taken as a baseline cortisol level, cortisol is seen to increase by one and a half times during the non-breeding season and three times during the breeding season (Fig. 1). If the March testosterone measurement is taken as the basal testosterone level, the same pattern is seen; the hormonal profile increases during

the non-breeding and breeding seasons (Fig.1). This highlights the importance of the functional roles of the HPA and HPG axes during the reproductive period, as well as the adaptive roles they play in the seasonal pattern observed, allowing the species to meet environmental challenges as well as challenges throughout its life cycle.

From the results gathered in this study, different immunoreactive GCs and T secretion patterns were observed according to the mating system and reproduction strategies (territorial defense and female defense) that guanaco males (of a migratory population) adopt during the breeding season (Ovejero 2013). Our results highlight the functional role of the stress and gonad axes during the critical mating period when all of the resources at a male's disposal must be used to maximize the chances of reproductive success. Generally, it appears that T is immediately involved in aggression associated with reproduction, rather than with other forms of aggression (e.g., anti-predator aggression), see Moyer (1968); Wingfield and Marler (1988). The data here presented support the correlation between T patterns and reproductive strategies (establishment and maintenance of a breeding territory and mate-guarding) proposed by Wingfield *et al.*, (1990) for male birds. However, this study extends Wingfield's finding to mammals. In 1984, the same authors proposed that males of polygynous species may have higher levels of T for longer periods during the breeding season than do males of monogamous species. This correlation was strengthened by the finding that when T was implanted in normally monogamous males, high T levels were maintained for longer than average periods than when these males became polygynous. Our results highlight the functional role that T levels have in determining and maintaining the polygynous mating system in guanacos and south american camelid wildlife, but we predict that this pattern could be altered in guanaco populations that coexist with human activities that fragment the guanaco's normal distributional range. For example, the use of fences by rural settlers has a direct impact on the space used by guanacos, and it is possible that this could promote some groups (within a population) to adopt monogamous behavior.

On the other hand, our results highlight the notorious decrease in the activity pattern of the stress and gonadal axes during the non-breeding season (Figure 1). This result was not expected because individuals must cope with another critical phase of autumnal migration at this time, therefore we expected that all of the resources (in terms of energy) would be available to maximize the chances for survival success during the migration trip. These results confirm our prediction that "guanacos show a seasonal variation in the activity of their endocrine and gonadal systems and incompatible functions (reproduction and self-maintenance) are established during different life stages." In winter, food intake by birds and mammals needs to be geared towards preventing starvation while at the same

time maintaining an optimum body mass to permit rapid escape from predators. GC concentrations may be the critical factor determining the size of the fat reserves and either environmental severity or predation risk; Wingfield (1990). The GC hormone, corticosterone, is thought to be important in the physiological orchestration of avian migration because of the identified elevated level of activity of the avian adrenal gland either prior to or during migration; Naik and George (1963); John (1965); Peczey (1976). Preparation for migration and for winter requires a period of hyperphagia and a laying down of fat reserves (for a review, see Holberton et al. (1999). In birds, Wingfield and Silverin (1986); Astheimer et al. (1992) and mammals, Dallman et al. (1993), GCs play a vital role in foraging behavior and hyperphagia, with low to moderate concentrations stimulating feeding behavior. As is noted by O'Reilly and Wingfield (1995) migrating birds show increased baseline concentrations of GCs but a reduced stress response. An adaptive explanation for this pattern is the migration modulation hypothesis, Holberton et al. (1996, 1999). It proposes that higher than average concentrations of baseline GCs are expressed prior to and during migration to facilitate migratory fattening. However, the stress response is lowered when challenged so that the negative, catabolic effects of high GCs on skeletal muscle do not occur; thus, critical skeletal muscles are preserved for flight during migration. As such, this hypothesis may explain the non-breeding season hormonal profiles generated in this study, but more data is needed to test these assumptions. Another explanation may suggest that migration (in terrestrial system) could be regulated by other endocrine mechanisms (and other hormones) involved in generating fat reserves and foraging behavior such as those that involve the thyroid gland Schwabl et al. (1984). Mattocks (1976) shows that gonad hormones (at least in some species) have a functional role (fat deposition) before spring migration. This may suggest that the increased activities of the gonad axis compared with the stress axis in guanacos could have an important role in the "migration to and from the winter range", but to confirm this prediction more experimental and field studies would be needed.

The last prediction of this study, that "prolonged elevated baseline levels of GCs due to long periods of intense social interaction are detrimental, costly and decrease fitness" was not validated by the results gathered. Our interest in the relationships between social or environmental conditions and stress-gonadal hormone concentrations (for both social and non-social species) is motivated by a desire to understand the physiological mechanisms by which the environment affects fitness. However, understanding the relationship between hormonal profiles and fitness is not straightforward, therefore this study attempts to obtain proxies that would help us begin to resolve this relationship. We consider that taken together, the interplay between stress/gonad hormones across seasonal variation

could be an interesting proxy to “establish” causal relationships between hormonal levels and ecological correlates. For example, some authors have stated that prolonged elevated baseline levels of GCs are commonly viewed to be detrimental, costly and decrease fitness, Bonier et al. (2009); However, there is scarce evidence for this negative association. Part of the problem with this approach is the variation among species in terms of their hormones/fitness relationships, and this variation can be attributed to a variety of demographic, social (e.g. social rank, age, sex, group size) or environmental factors (e.g. resource availability). Additionally, for a variety of vertebrate taxa, seasonal variation in GC levels has been reported to be due to varying food availability, temperature, rainfall and tourist activity Romero (2002); Tempel and Gutierrez (2004), however it still remains unclear if this holds true for mammals. The other issue is that many of the studies of HPA-fitness relationships reviewed by Bonier et al. (2009) were conducted during the breeding season alone. The study herein presented provides a full set of data (hormone profiles) across seasonal variation throughout the course of an entire year. As expected, guanacos cope with this trade-off between sociability and reproductive benefits and costs by regulating their GCs and testosterone levels on a seasonal basis like other mammals Boosntra (2005), and they may also experience fine adjustments on a daily basis. We suggest that elevated levels of GCs in guanacos due to highly social interactions during the breeding season may play a fundamental role in the regulation of enhanced metabolic needs during reproduction. Guanaco males, like other iteroparous males, exhibit high concentrations of cortisol, a gonadal axis that is not inhibited by high GC concentrations, so the last prediction presented in this work must be rejected. Furthermore, during the breeding season, our results show two GC peaks of activity, one at the beginning and another at the end of the breeding season. In the former, guanaco males are exposed to intense social instability caused by agonistic interactions with other males due to territory establishment and hierarchies. In the latter, guanaco males are exposed to female defense interactions. This suggests that for guanaco males, female defense strategies demand higher energetic costs when compared to territory defense strategies. This is expected if one assumes that defending a female (rather than a territory) is the principal objective (and target) of a male that wants to maximize the probabilities of leaving an offspring. Moreover, birth rate data for the population was collected from 2007-2008 (hormonal measures)-2009, so if one expects a negative relationship between the hormonal profiles, one would also expect to see a negative value for the proportion of offspring per adult. The birth rate values for those years were 0.14, 0.28 and 0.29 respectively; thus with these results one can expect that there is no detrimental effect on individual-population fitness.

The lesser HPA axis activity measured during the non-breeding season confirms the prediction "that reproduction and sociability may be long-term and predictable." This is likely the case because this period is associated with fewer social interactions. The individual hierarchies are more relaxed because the animals are preparing to migrate (Novaro pers. com.) and the species strategy at this time is to survive and to not reproduce.

Finally from the data collected here, a reactive scope model for guanacos has been generated, see figure 2. The results show the ranges of predictive/reactive homeostasis and homeostatic overload/failure ranges; these consist of the seasonal set of point ranges for the physiological mediators. For guanacos, the normal reactive scope is described with two GCs and T set point ranges. The model was made to reflect the natural progression of life-history stages such as breeding (figure 2). Furthermore, the minimum concentration of the mediator was presumed to constitute a threshold. Subsequently, a GC and T threshold was proposed, and below this limit it is seen that the individual/population enters into homeostatic failure. This threshold is predicted to indicate the minimum concentration of GCs and T mediator necessary to sustain guanacos seasonal life changes (in natural conditions). However, more studies are needed to test this prediction. On the other hand, above the upper end of the normal reactive scope, when a physiological mediator exceeds the reactive homeostatic range, the individual/population enters into homeostatic overload (pathological states). This threshold is predicted to indicate the maximum concentration of GCs and T mediator necessary to sustain seasonal life changes in guanacos (in natural conditions). However, more studies are needed to test this prediction especially considering that pharmacological stimulation may allow individuals to reach a higher threshold though this would not be considered within the scope of natural changes.

## Conclusions

Knowledge of physiological and behavioral mechanisms is key to understanding how life-history variation and trade-offs might arise and are maintained to cope with environmental variation. Mechanistically, trade-offs result from the need to differentially allocate limited resources to traits like reproduction versus self-maintenance with selection favoring the evolution of optimal allocation mechanisms. Guanacos cope with the trade-off between sociability and reproductive benefits and costs by regulating their GCs and T levels on a seasonal basis. This suggests the adaptive and functional role of both axes in regulating energy allocation for "predictable" life-history events. During breeding and highly social interactions, elevated baseline levels of GCs and T may actually be predictive of high reproductive success. Our results support the assumption that GCs via the HPA axis did not



suppress the HPG axis and did not diminish T levels. From this, it is proposed that guanacos anticipate stressors in which the impact on the animal is acute and the duration is moderated by physiological consequences (i.e reproduction). During the non-breeding season, a period with less intense social interactions, lower baseline levels of GCs and T may actually be predictive of a high probability of survival. Our study provides a model for studying the link between physiology and life history.

Studying both the HPA axis and the HPG axis is one of the best ways to understand the functional mechanisms that an animal uses to cope with stress. Studying these two axes are of importance because the key circulating steroid hormones of the stress axis, GCs, influence the expression of approximately 10% of the genome. Some of the targets of GCs include genes controlling metabolism, growth, repair, reproduction and the management of resource allocation Lee et al. (2005). However, both axes play fundamental roles not just when the animal is experiencing stress, but during normal periods of survival and growth. These axes mediate the adaptation of the organism to its environment. At the individual level, the stress axis plays a key role in allowing animals to respond to the changes and challenges of both environmental certainty and uncertainty. At the species level, the stress axis plays a central role in evolutionary adaptation to particular ecological pressures.

**Acknowledgements:** We appreciate the comments and suggestions made by Dr. Francisco Bozinovic, Donald R.Strong and Tim Coulson on earlier versions of this manuscript. We thank Nelly Horak and Emily Giles for helping with the English language. This study has been funded by the Rufford Small Grant Foundation (RSGF #120608); The Scientific Research Society/Sigma-Xi and FONDECYT-CONICYT-PROGRAM (N°3140237). The park rangers and other staff of La Payunia Provincial Park helped us throughout different stages of this work.

Figure 1. HPA & HPG axis activity patterns throughout the study period. Blue line= the seasonal baseline fecal testosterone profile (defined by  $y = 0.0041x^5 - 0.1102x^4 + 1.0889x^3 - 4.6819x^2 + 7.914x - 0.7892$ ;  $r^2 = 0.7231$ ), Pink line= the seasonal baseline fecal cortisol profile (defined by  $y = 0.0743x^5 - 1.9677x^4 + 19.136x^3 - 82.442x^2 + 147.4x - 58.902$ ;  $r^2 = 0.806$ ). There was marked adrenal and gonadal activity during the breeding and non-breeding (CI = 7.661734-12.057391, mean =25.75864/15.89907 respectively) season respectively for the male groups, the GCs and T response was due to seasonal variation in the guanacos highlighting the individual's energetic demands according to life-history strategies.



Figure 2. Most studies measure only one physiological mediator at a time, here the concentration of two mediators (cortisol-C /testosterone.T) is placed on the y-axis at a given time point. The mediators are correlated in terms of the functional role that each one plays in the normal reactive scope range. In other words, both mediators encompass responses for coping with predictable and unpredictable changes in the environment. The values of each mediator are presumed to exist in four general ranges. The minimum (5-10 Ng.gr for C;1-2 Ng.gr for T) and maximum (20-35 Ng.gr for C;3-4 Ng.gr for T) concentration of the mediator was presumed to constitute a threshold. The results show the ranges (double arrows) of predictive/reactive homeostasis (PH/RH) and homeostatic overload/failure, the stress-gonadal and feedback response (gray circles) indicates the max-min activity patterns for HPA & HPG axis during one year. The C activity patterns in guanacos shows two peak of RH (dotted lines) in the reproductive season, between Ag./O. males adopts a territory-defense strategies at the beginning of the reprod. season; during Ja./F. males changes to a female-defense strategies at the ends of the reprod. season. These results consist of the seasonal set of point ranges for the physiological mediators.

## References

- Bank MS, Sarno RJ, Franklin WL. 2003. Spatial distribution of guanaco mating sites in southern Chile: conservation implications. *Biological conservation*. 112: 427-434.
- Blanchard RJ, McKittrick CR, Blanchard DC. 2001. Animal models of social stress: effects on behavior and brain neurochemical systems. *Physiol. Behav.* 73, 261–271.
- Becker JB. 2002. *Behavioral endocrinology* 2nd ed. Massachusetts Institute of technology. 1-749.
- Bonier F, Moore IT, Martin PR & Robertson RJ. 2009. The relationship between fitness and baseline glucocorticoids in a passerine bird. *General and Comparative Endocrinology*, 163, 208–213.
- Boonstra R, Singleton GR. 1993. Population declines in the snowshoe hare and the role of stress, *Gen. Comp. Endocrinol.* 91, 126–143.
- Boonstra, R. 2005. Equipped for life: the adaptive role of the stress axis in male mammals. *Journal of Mammalogy*, 86, 236–247.
- Boonstra R, Barker JM, Castillo J & Fletcher QE. 2007. The role of the stress axis in life-history adaptations of rodents. *Rodent Societies: An Ecological and Evolutionary Perspective* (eds J.O. Wolff & P.W. Sherman), pp. 139–149, University of Chicago Press, Chicago.
- Boonstra R, McColl CJ, Karels TJ. 2001. Reproduction at all costs: the adaptive stress response of male arctic ground squirrels. *Ecology* 82: 1930–1946.

- Bozinovic F..2002. Physiological ecology and evolution. Theory and study cases in vertebrates. Ediciones Universidad Catolica de Chile . Pp.531.
- Buchanan KL, Goldsmith AR.2004. Noninvasive endocrine data for behavioural studies: the importance of validation. *Anim Behav* 67:183–185.
- Busch DS and Hayward LS. 2009. Stress in a conservation context: a discussion of glucocorticoid actions and how levels change with conservation-relevant variables, *Biol. Conserv.* 142, 2844–2853.
- Carmanchahi P, Ovejero RJA, Marull C, López C, Schroeder N, Jahn G, Novaro A y Somoza G. 2011. Physiological response of wild guanacos to capture for live shearing. *Wildlife research: csiro publishing*. vol. N°. P61 - 68.
- Cavigelli SA. 1999. Behavioural patterns associated with faecal cortisol levels in free-ranging female ringtailed lemurs, *Lemur catta*. *Anim Behav* 57:935–944.
- Cavigelli S, Pareira M. 2000. Mating season aggression and fecal testosterone levels in male ring-tailed lemurs (*Lemur catta*). *Horm. Behav.* 37, 246–255.
- Clarke FM, Faulkes CG. 2001. Intracolony aggression in the eusocial naked mole-rat, *Heterocephalus glaber*. *Anim. Behav.* 61, 311 –324.
- Clarke FM, Bennett NC. 2001. Family values: group dynamics and social control of reproduction in African mole-rats. *Trends Ecol. Evol.* 16, 184– 190.
- Clutton-Brock TH, Brotherton PNM, Russell AF, O’Riain MJ, Gaynor D, Kansky R, Griffin A, Manser M, Sharpe L, McIlrath GM, Small T, Moss A, Monfort S. 2001. Cooperation, control, and concession in meerkat groups. *Science* 291, 478–481.
- Co’té SD. 2000. Dominance hierarchies in female mountain goats: stability, aggressiveness and determinants of rank. *Behaviour* 137,1541– 1566.
- Creel S, Creel N y Monfort S. 1996. Social stress and dominance. *Nature* 379:212.
- Creel S. 2001. Social dominance and stress hormones. *Trends Ecol. Evol.* 16, 491– 497.
- Creel S, Dantzer B, Goymann W & Rubenstein DR. 2013. The ecology of stress: effects of the social environment. *Functional Ecology*, 27, 66–80.
- Creel S & Creel NM. 2002. *The African Wild Dog: Behavior, Ecology and Conservation*, Princeton University Press, Princeton, NJ.
- Dallman MF, Warne JP, Foster MT, Pecoraro NC. 2007. Glucocorticoids and insulin both modulate caloric intake through actions on the brain, *J. Physiol.- London* 583, 431–436.
- Enstrom DA, Ketterson ED & Nolan Jr V. 1997. Testosterone and mate choice in the dark-eyed junco. *Anim. Behav.*54, 1135–1146.
- Faulkes CG, Abbott DH.1997. The physiology of a reproductive dictatorship: regulation of male and female reproduction by a single breeding female in colonies of naked mole-rats. In: Solomon, N.G., French, J.A. (Eds.), *Cooperative Breeding in Mammals*. Cambridge Univ. Press, Cambridge, pp. 302–334.

- Franklin W. 1983. Constranting socioecologies of South America's wild camelids: the vicuna and guanaco. Pp. 573-629 in J.F.Eisenberg and D.Kleiman, eds. Advances in study of mammalian behavior. Spec. Publ. No. 7, Am.Soc.Mammal, Shippensburg, Pa.
- Gottreich A, Zuri I, Barel S, Hammer I, Terkel J. 2000. Urinary testosterone levels in the male blind mole rat (*Spalax ehrenbergi*) affect female preference. *Physiol Behav.*;69:309–315.
- Goymann W, Wingfield JC. 2004. Allostatic load, social status and stress hormones: the costs of social status matter, *Anim. Behav.* 67, 591–602.
- Goymann W and Wingfield JC. 2004 Allostatic load, social status and stress hormones: the costs of social status matter. *Animal Behaviour* 67:591–602.
- Jacobs JD and Wingfield JC. 2000. Endocrine control of life- cycle stages: a constraint on response to the environment? *Condor* 102:35-51.
- John TM. 1965. A histochemical study of adrenal corticoids in pre and post migratory wagtails *Motacilla alba* and *M. flava*. *Pavo* 4, 9–14.
- Kenagy GJ and Place NJ. 2000 Seasonal changes in plasma glucocorticosteroids of free- living female yellow-pine chipmunks: effects of reproduction and capture and handling, *Gen. Comp. Endocrinol.* 117. 189–199.
- Knapp R, Moore MC. 1997. Male morphs in tree lizards have different testosterone responses to elevated levels of corticosterone. *Gen. Comp. Endocrinol.* 107, 273–279.
- Le PP, Friedman JR, Schug J, Brestelli JE, Parker JB, Bochkis IM & Kaestner KH. 2005. Glucocorticoid receptor-dependent gene regulatory networks. *PLoS Genetics*, 1, 159–170.
- S. Levine, Stress: an historical perspective, in: T. Steckler, N. Kalin, J.M.H.M. Reul (Eds.), *Handbook of Stress and the Brain*, Elsevier, Amsterdam, the Netherlands, 2005, pp. 3–23.
- Lofts B & Marshall AJ. 1960. The experimental regulation of the sexual cycle in the Brambling *Fringilla montifringilla*. *Ibis* 102: 209-214.
- Naik DV and George JC. 1963. Histochemical demonstration of increased corticoid level in the adrenal of *Sturnus roseus* (Linnaeus) towards the migratory phase. *Pavo* 1, 103–105.
- Nelson R. 2000. An introduction to behavioural endocrinology. Sinauer, Sunderland, Mass.
- Nespolo RF, Bacigalupe LD & Bozinovic F. 2003. Heritability of energetics in a wild mammal, the leaf eared – mouse (*Phyllotis darwini*). *Evolution* 57: 1679-1688.
- Marino A & Baldi R. 2008. Vigilance patterns of territorial guanacos (*Lama guanicoe*): the role of reproductive interests and predation risk. *Ethology* 114, 413–423.
- Mattocks PW. 1976 The role of gonadal hormones in the regulation of the premigratory fat deposition in the White-crowned Sparrow *Zonotrichia leucophrys gambelii*. Unpublished Master's thesis. Seattle, Washington, Univ. Washington.
- Mateo JM, Cavigelli SA. 2005. A validation of extraction methods for noninvasive sampling of glucocorticoids in free-living ground squirrels. *Physiol Biochem Zool* 78: 1069–1084.

- McEwen BS, Wingfield JC. 2010. What is in a name? Integrating homeostasis, allostasis and stress, *Horm. Behav.* 57: 105–111.
- Millspaugh JJ, Woods RJ, Hunt KE, Raedeke KJ, Brundige GC, Washburn BE, Wasser SK. 2001 Using fecal glucocorticoid assays to study the physiological stress response of elk. *Wildlife Society Bulletin* 29, 899–907.
- Millspaugh JJ, Washburn BE. 2004. Use of fecal glucocorticoid metabolite measures in conservation biology research: considerations for application and interpretation. *Gen Comp Endocrinol* 138:189–199.
- Moberg GP. 1991. How behavioral stress disrupts the endocrine control of reproduction in domestic animals, *Journal Dairy Science.* 74, 304–311.
- Monclús R, Tiulim J, Blumstein DT. 2011. Older mothers follow conservative strategies under predator pressure: the adaptive role of maternal glucocorticoids in yellow-bellied marmots, *Horm. Behav.* 60, 660–665.
- Morton ML & Mewaldt LR. 1962 Some effects of castration on a migratory sparrow (*Zonotrichia atricapilla*). *Physiol. Zool.* 35: 237–247.
- Mostl E and Palme R. 2002. Hormones as indicators of stress, *Domest. Anim. Endocrinol.* 23, 67–74
- Mostl E, Rettenbacher S, Palme R. 2005. Measurement of corticosterone metabolites in birds' droppings: an analytical approach. *Ann NY Acad Sci* 1046:17–34.
- Muller MN, Wrangham RW. 2004. Dominance, aggression and testosterone in wild chimpanzees: a test of the challenge hypothesis. *Anim. Behav.* 67, 113– 123.
- Ovejero R and Carmanchahi P. 2012. "Stress In Nature? Integrating Physiology, Ecology and Natural History of guanacos (*Lama guanicoe*)". Proceedings of 2nd Latin-American Mammalian Congress.
- Ovejero R. 2013. "Variación del nivel de cortisol en función de factores sociales y ambientales en guanacos (*Lama guanicoe*). Implicancias para la conservación y manejo de las poblaciones silvestres". 315pp, Tesis doctoral. Facultad de Cs. Naturales Univ Nac
- Ostner J, Kappeler PM. 2002. Seasonal variation and social correlates of androgen excretion in male redfronted lemurs (*Eulemur fulvus rufus*). *Behav. Ecol. Sociobiol.* 52, 485– 495.
- Ostner J, Kappeler PM. 2002. Seasonal variation and social correlates of androgen excretion in male redfronted lemurs (*Eulemur fulvus rufus*). *Behav. Ecol. Sociobiol.* 52, 485– 495.
- Pereira RJG, Duarte JMB, Negrão JA. 2006. Effects of environmental conditions, human activity, reproduction, antler cycle and grouping on fecal glucocorticoids of free-ranging Pampas deer stags (*Ozotoceros bezoarticus*), *Horm. Behav.* 49, 114–122.
- Peczely P. 1976. Étude circannuelle de la fonction corticosurrénalienne chez les espèces de passereaux migrant et non migrant. *Gen. Comp. Endocrinol.* 30, 1–11.
- Pride RE. 2005. High faecal glucocorticoid levels predict mortality in ring-tailed lemurs (*Lemur catta*), *Biological Letters.* 1, 60–63.
- Raedeke K. 1979. *Population dynamics and socioecology of the guanaco (Lama guanicoe) of Magallanes, Chile.* Doctoral Dissertation. College of Forest Resources University of Washington.
- Raouf SA, Smith LC, Brown MB, Wingfield JC, Brown CR. 2006 Glucocorticoid hormone levels increase with group size and parasite load in cliff swallows. *Anim Behav* 71:39–48.

- Romero LM, Remage-Healey L. 2000. Daily and seasonal variation in response to stress in captive starlings (*Sturnus vulgaris*): corticosterone. *General and Comparative Endocrinology* 119, 52–59.
- Romero LM. 2002 Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. *General and Comparative Endocrinology*, 128, 1–24.
- Romero LM. 2004 Physiological stress in ecology: lessons from biomedical research. *T.R.E.E.* 19, 249–255.
- L.M. Romero, M.J. Dickens, N.E. Cyr. 2009. The reactive scope model – A new model integrating homeostasis, allostasis, and stress, *Horm. Behav.* 55 375– 389.
- Rubenstein DR & Shen SF. 2009. Reproductive conflict and the costs of social status in cooperatively breeding vertebrates. *The American Naturalist*, 173, 650–662.
- Sapolsky RM, Romero LM, Munck MU. 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions, *Endocr. Rev.* 21, 55–89.
- Sapolsky RM. 2002. Endocrinology of the stress response, in: J.B. Becker, S.M. Breedlove, D. Crews, M.M. McCarthy (Eds.), *Behavioral Endocrinology*, 2nd ed, Cambridge, Massachusetts, pp. 409–450.
- Sheriff MJ, Bosson CO, Krebs CJ, Boonstra R. 2009. A noninvasive technique for measuring fecal cortisol metabolites in snowshoe hares (*Lepus americanus*). *J Comp Physiol B* 179:305–313
- Sheriff MJ, Dantzer B, Delehanty B, Palme R, Boonstra R. 2011. Measuring stress in wildlife: techniques for quantifying glucocorticoids, *Oecologia* 166, 869–887.
- Schradin C. 2008. Seasonal changes in testosterone and corticosterone levels in four social classes of a desert dwelling sociable rodent, *Horm. Behav.* 53, 573–579.
- Schwabl H, Wingfield JC and Farner DS. 1984. Endocrine Correlates of Autumnal Behavior in Sedentary and Migratory Individuals of a Partially Migratory Population of the European Blackbird (*Turdus merula*). *The Auk*, Vol. 101, No. 3, pp. 499–507
- Sinervo B, Miles DB, Frankino WA, Klukowski M, DeNardo DF. 2000. Testosterone, endurance, and Darwinian fitness: natural selection and sexual selection on the physiological bases of alternative male behaviors in side-Blotched lizards. *Horm. Behav.* 38, 222–233.
- Smith JE, Whelan CJ, Taylor SJ, Denight ML, Stake MM. 2007 Novel predator-prey interactions: is resistance futile?, *Evol Ecol. Res.* 9, 433–446.
- Smith JE, Monclús R, Wantuck D, Florant GL, Blumstein DT. 2012. Fecal glucocorticoid metabolites in wild yellow-bellied marmots: Experimental validation, individual differences and ecological correlates. *General and Comparative Endocrinology* 178: 417–426.
- Soto-Gamboa M, Villalon M, Bozinovic F. 2005. Social cues and hormone levels in male *Octodon degus* (Rodentia): a field test of the challenge hypothesis. *Horm Behav* 47:311–318.
- Soto-Gamboa M, Gonzalez S, Hayes LD, Ebensperger L. 2009. Validation of a radioimmunoassay for measuring fecal cortisol metabolites in the hystricomorph rodent *Octodon degus*. *J Exp Zool* 311:496–503.
- Stearns SC. 1992. *The evolution of life histories*. Oxford: Oxford University Press.
- Tarlow EM, Blumstein DT. 2007. Evaluating methods to quantify anthropogenic stressors on wild animals, *Appl. Anim. Behav. Sci.* 102, 429–451.

- Tempel DJ, Gutierrez RJ. 2004. Factors related to fecal corticosterone levels in California Spotted Owls: implications for assessing chronic stress. *Conservation Biology* 18, 538–547.
- Touma C, Palme R. 2005. Measuring fecal glucocorticoid metabolites in mammals and birds: The importance of validation. *Ann. NY Acad. Sci.* 1046, 54–74.
- Vera F, Zenuto RR, Antenucci CD. 2012. Differential responses of cortisol and corticosterone to adrenocorticotrophic hormone (ACTH) in a subterranean rodent (*Ctenomys talarum*). *Journal of Experimental Zoology A*. 317:173–184.
- Von Holst D. 1998. The concept of stress and its relevance for animal behavior, *Stress and Behavior* 27, 1–131.
- Wasser SK, Hunt KE, Brown JL, Cooper K, Crockett CM, Bechert U, Millspaugh JJ, Larson S, Monfort SL. 2000. A generalized fecal glucocorticoid assay for use in a diverse array of nondomestic mammalian and avian species. *Gen Comp Endocrinol* 120:260–275.
- Weise CM. 1967. Castration and spring migration in the White-crowned Sparrow. *Condor* 69: 49– 68.
- Wingfield JC, Hunt K, Breuner C, Dunlap K, Fowler GS, Freed L, Lepson J. 1997. Environmental stress, field endocrinology, and conservation biology. In: Clemmons, JR.; Buchholz, R., editors. *Behavioral Approaches to Conservation in the Wild*. Cambridge University Press; Cambridge: p. 95–131.
- Wingfield JC, Sapolsky RM. 2003. Reproduction and resistance to stress: when and how. *J Neuroendocrinol* 15:711–724.
- Wingfield JC. 2005. The concept of allostasis: coping with a capricious environment, *J. Mammal.* 86: 248–254.
- Young JK & Franklin WL. 2004 Activity budget patterns in family-group and solitary territorial male guanacos. *Revista Chilena de Historia Natural*, 77, 617–625.
- Young AJ, Carlson AA, Monfort SL, Russell AF, Bennett NC & Clutton-Brock T. 2006. Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 12005–12010.




730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742

# **Figure 1**(on next page)

HPA & HPG axis activity patterns throughout the study period





## Figure 2(on next page)

The reactive scope model for guanacos

