

A peer-reviewed version of this preprint was published in PeerJ on 24 April 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.366) (peerj.com/articles/366), which is the preferred citable publication unless you specifically need to cite this preprint.

Kong X. 2014. Association between in-scanner head motion with cerebral white matter microstructure: a multiband diffusion-weighted MRI study. PeerJ 2:e366 <https://doi.org/10.7717/peerj.366>

Association between in-scanner head motion with cerebral white matter microstructure: a multiband diffusion-weighted MRI study

Xiang-zhen Kong

Diffusion-weighted Magnetic Resonance Imaging (DW-MRI) has emerged as the most popular neuroimaging technique used to depict the biological microstructural properties of human brain white matter. However, like other MRI technique, traditional DW-MRI data remains subject to head motion artifacts during scanning. For example, previous studies have indicated that, with traditional DW-MRI data, head motion artifacts significantly affect the evaluation of diffusion metrics. Actually, DW-MRI data scanned with higher sampling rate are important for accurately evaluating diffusion metrics because it allows for full-brain coverage through the acquisition of multiple slices simultaneously and more gradient directions. Here, we employed a publicly available multiband DW-MRI dataset to investigate the association between motion and diffusion metrics with the standard pipeline, tract-based spatial statistics (TBSS). The diffusion metrics used in this study included not only the commonly used metrics (i.e., FA and MD) in DW-MRI studies, but also newly proposed inter-voxel metric, local diffusion homogeneity (LDH). We found that the motion effects in FA and MD seems to be mitigated to some extent, but the effect on MD still exists. Furthermore, the effect in LDH is much more pronounced. These results indicate that researchers shall be cautious when conducting data analysis and interpretation. Finally, the motion-diffusion association is discussed.

Association between in-scanner head motion with cerebral white matter microstructure: A multiband diffusion-weighted MRI study

Diffusion-weighted Magnetic Resonance Imaging (DW-MRI) has emerged as the most popular neuroimaging technique used to depict the biological microstructural properties of human brain white matter. However, like other MRI technique, traditional DW-MRI data remains subject to head motion artifacts during scanning. For example, previous studies have indicated that, with traditional DW-MRI data, head motion artifacts significantly affect the evaluation of diffusion metrics. Actually, DW-MRI data scanned with higher sampling rate are important for accurately evaluating diffusion metrics because it allows for full-brain coverage through the acquisition of multiple slices simultaneously and more gradient directions. Here, we employed a publicly available multiband DW-MRI dataset to investigate the association between motion and diffusion metrics with the standard pipeline, tract-based spatial statistics (TBSS). The diffusion metrics used in this study included not only the commonly used metrics (i.e., FA and MD) in DW-MRI studies, but also newly proposed inter-voxel metric, local diffusion homogeneity (LDH). We found that the motion effects in FA and MD seems to be mitigated to some extent, but the effect on MD still exists. Furthermore, the effect in LDH is much more pronounced. These results indicate that researchers shall be cautious when conducting data analysis and interpretation. Finally, the motion-diffusion association is discussed.

1 **Title:** Association between In-scanner Head Motion and Cerebral White Matter Microstructure: A Multiband
2 Diffusion-weighted MRI study

3 **Running head:** Head Motion and White Matter Microstructure

4 **Authors:** Xiang-zhen Kong^{1,2}

5 **Affiliation:** ¹ State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for
6 Brain Research, Beijing Normal University, Beijing, 100875, China; ² Center for Collaboration and Innovation
7 in Brain and Learning Sciences, Beijing Normal University, Beijing, 100875, China.

8 **Correspondence Author:**

9 Xiang-zhen Kong; 2nd Floor, Brain Imaging Center, 19 Xijiekouwai St, Haidian District, Beijing 100875,
10 China. E-mail: kongxiangzheng@gmail.com

11 Introduction

12 Diffusion-weighted MRI (DW-MRI) has become one of the most popular MRI techniques in brain research, as
 13 well as in clinical practice. One key application of DW-MRI is diffusion tractography which can be used for the
 14 visualization of white matter (WM) tracts (Golby et al. 2011) and construction of the brain neuroanatomical
 15 connectome (Gong et al. 2009). Also, it has become a convenient tool for deriving regional measures of
 16 diffusivity and anisotropy. These metrics are believed to reflect biological microstructural properties of the
 17 white matter, and have been extensively applied as biological markers for studying WM under normal and
 18 clinical conditions (Johansen-Berg 2010; Le Bihan 2003; Le Bihan et al. 2001; Travers et al. 2012).

19 However, like other MRI technique, DW-MRI remains subject to specific biological factors (e.g.,
 20 temperature), uncertainty from the scanner (e.g., machine SNR, field shim) and, in particular, motion artifacts.
 21 Thus, movement of the head during scanning is undesirable, since it not only displaces the brain matter in space
 22 but also interferes with the readout of MR signals. Indeed, recent studies have discovered that head motion may
 23 introduce unwanted biases. Ling and his colleagues have shown that head motion is associated with both
 24 fractional anisotropy (FA) and mean diffusivity (MD) (the effect is greater for MD) (Ling et al. 2012). A recent
 25 study have also found group differences in head motion can induce group differences in white matter tract-
 26 specific diffusion metrics, and such effects can be more prominent in some specific tracts than others (Yendiki
 27 et al. 2013). However, these studies on head-motion artifacts have employed traditional DW-MRI data with
 28 relatively low sampling rate (e.g., 9 s) and hence few gradient directions (e.g., $n = 30$). In fact, previous works
 29 have indicated that more unique sampling directions may decrease bias of diffusion metrics (e.g., FA and MD)
 30 (Landman et al. 2007; Tjissen et al. 2009). Recently, several promising imaging techniques have been
 31 proposed, including MR-encephalography (Zahneisen et al. 2011) and multiband echo planar imaging (Moeller
 32 et al. 2010). Using the multiband scanning protocol, sampling across the whole brain at any given time is
 33 allowed through the acquisition of multiple slices simultaneously. Hence, additional gradient directions can be
 34 acquired in the same scan duration without loss in spatial resolution. Both of the advantages appear to result in
 35 evaluating diffusion metrics more accurately, but little is known about the head motion effects on diffusion
 36 metrics from the multiband dataset.

37 Here, the primary aim was therefore to investigate the relationship between head motion and diffusion
 38 metrics estimated from the multiband dataset. In this study, we examined the two tensor-based metrics most
 39 typically reported (i.e., FA and MD). Given the fact that they only reflect diffusion properties solely within the
 40 voxel, we also examined a newly proposed model-free inter-voxel metric, referred to as local diffusion

homogeneity (LDH) (Gong 2013). We hypothesized that motion effects would be mitigated in the multiband DW-MRI data. It has been suggested that head motion alters the measure of diffusion metrics even after motion and eddy current correction (Ling et al. 2012; Tijssen et al. 2009; Yendiki et al. 2013), and that it may also provide information regarding neuronal processing (Yan et al. 2013a; Yan et al. 2013b). Moreover, LDH is a recently proposed metrics and has not been fully validated yet (Gong 2013). In addition, unlike FA and MD, LDH directly depends on the raw diffusivity series without assuming a prior diffusion model (Gong 2013). Therefore, we also hypothesized that the association between head motion and LDH would be quite different to the tensor-based metrics, and may be more sensitive to motion artifacts. We tested these hypotheses by (1) confirming the test-retest reliability of both diffusion metrics and head motion across scan sessions, and (2) by examining the relationship between the averaged diffusion metrics and head motion. We also examined the relationship in each scan session.

Materials and Methods

Dataset

The dataset used in this study was from the NKI-RS Multiband Imaging Test-Retest Pilot Dataset (Mennes et al. 2012). There were 20 participants (34.3 ± 14.0 years). For each participant, the DW-MRI scans were performed twice (session 1 and session 2), around one week apart. Diffusion weighted images were collected a standard pulse sequence with 2-mm-thick axial slices and 137 directions: TE 85 ms; TR 2400 ms; b value, 1500 s/mm²; flip angle, 90°.

Image Processing

DW-MRI images were processed with FMRIB's Software Library (FSL, <http://www.fmrib.ox.ac.uk/fsl>). Non-brain tissue was removed using the Brain Extraction Tool (BET) with a fractional intensity threshold of 0.2, and then raw DW images were affinely registered to the nonDW image, to partially correct for the effects of motion and eddy currents. Then, by fitting a tensor model at each voxel using DTIFit from the FSL (Smith et al. 2004), we obtained the fractional anisotropy (FA) and mean (MD) diffusivity, used in subsequent TBSS analysis (Smith et al. 2006; Smith et al. 2007).

To compare between subjects, the TBSS framework was used. In detail, first, we non-linearly aligned the individual FA maps to FSL's standard 1 mm isotropic FA template (FMRIB58_FA) and averaged them to generate a study specific mean FA map. Next, voxels with an FA > 0.2 in the mean FA map were masked out, and the remainder was thinned to create a white matter "skeleton". The resulting skeleton contained WM tracts common to all subjects. Individual FA maps were then projected onto the mean FA skeleton by filling the skeleton with FA values from the nearest tract center. The same non-linear transformations derived for the FA maps were applied to the MD maps.

In terms of the LDH metric, it is a novel model-free metric that defines the regional inter-voxel coherence of diffusion series (Gong 2013). Technologically, LDH is quantified within the neighbors ($n = 27$) via the Kendall's coefficient concordance (KCC), after the estimation of the diffusivity strengths along each gradient direction. To compare between subjects, the LDH maps were also projected onto the WM skeleton mask using the TBSS framework described above. In addition, we used the same approach with different neighbor size (i.e., $n = 7$ and $n = 19$) for quantifying the LDH, and also used another approach for quantifying the regional coherence with information theory (Kong et al., 2014). The results in these cases were all similar to those of the original LDH (data not shown).

The DW-MRI data preprocessing and TBSS analysis pipelines were both implemented using Nipype (Gorgolewski et al. 2011), a flexible, lightweight and extensible neuroimaging data processing framework in python. The pipeline for calculating both original and improved LDH was implemented in python.

Assessment of in-scanner head motion

To retrospectively estimate head motion during scanning, DW images were realigned to the non-DW image with FMRIB's Linear Image Registration Tool (FLIRT), and at the same time, an affine transformation matrix was obtained for each image. Then, for each image, the root mean squared (RMS) deviation (Jenkinson et al. 2002), a summary statistic of in-scanner head motion, was calculated from its transformation using the tool *rmsdiff* from FSL. Since it summarizes six translational and rotational parameters, the RMS has been widely used in the neuroimaging community. For instance, it has been used in fMRI and DTI data processing to check the extent of head motion and make decisions about cohort formation or matching (e.g., Ikuta et al., 2014; Kochunov et al., 2013; Kong, 2014). Technically, the RMS can be calculated directly from the affine matrices with the formula (1).

94

$$RMS = \sqrt{\frac{1}{5} R^2 \text{Trace}(A^T A) + t^T t} \quad (1)$$

95 In formula (1), R is a radius specifying the volume of interest (R = 80mm, approximately the mean
96 distance from the cerebral cortex to the center of the head), A is a 3x3 'rotation' matrix and t is a 3x1 column
97 vector of translation. One thing to note is that since the RMS uses all the information from the affine matrices
98 (including the shear and scaling, if present), it could include the electrical properties of each participant's head.
99 Nevertheless, the RMS does provide a sensitive index of in-scanner head motion.

100 Here, the RMS was calculated from 2 transformations of consecutive images (Jenkinson et al. 2002). That
101 is, in-scanner head motion was measured as the summary measure of head motion relative to the preceding
102 volume as the previous studies (Satterthwaite et al. 2012; Van Dijk et al. 2012). Finally, head motion was
103 calculated by averaging the RMS deviations for all volumes.

104 **Test-retest reliability of diffusion metrics and head motion estimate**

105 The voxel-wise test-retest reliability for each diffusion metric was calculated with the intra-class correlation
106 coefficient (ICC) (Shrout & Fleiss 1979).

$$ICC = \frac{BMS - EMS}{BMS + (k - 1) EMS} \quad (2)$$

108 The formula estimates the correlation of the subject signal intensities between sessions, modeled by a two-
109 way ANOVA, with random subject effects and fixed session effects. In this model, the total sum of squares is
110 split into subject (BMS), session (JMS) and error (EMS) sums of squares; the k is the number of repeated
111 sessions. The reliability measure for whole-brain analysis was implemented in python and can be accessed
112 from Nipype (Gorgolewski et al. 2011). The test-retest reliability for head motion estimate was also calculated
113 with ICC.

114 **Relationship between In-Scanner Head Motion and Diffusion Metrics**

115 To maximize the signal to noise ratio of head motion estimates, first we calculated the average head motion for
116 each participant across two sessions. Analogously, for accurate measures of microstructure estimates, the MD,
117 FA and LDH metrics finally used were also taken from the average of the TBSS results across the two sessions.

To examine the possible relationship between head motion and diffusion regional metrics, we conducted a statistical analysis using general linear models (GLMs), for the three metrics respectively, with head motion as the variable of interest. In these models, gender, age and handedness were controlled as confounding covariates. Handedness was included here because it is associated with brain structure and the neural processing of attention, while attention deficit may cause more head motion (e.g., Durston et al., 2003; Schmidt et al., 2013). Voxel-wise statistical analysis was performed with Threshold-Free Cluster Enhancement (TFCE) correction (Smith & Nichols 2009) for multiple comparisons, considering fully corrected p-value < 0.05 as significant. In addition, the same statistical procedure was conducted for both session 1 and session 2 respectively.

Results

Test-retest reliability of diffusion metrics and head motion estimate

All of the diffusion metrics in this study showed relatively high test-retest reliability: FA: Mean ICC = 0.71; MD: Mean ICC = 0.71; LDH: Mean ICC = 0.75. In addition, the magnitude of head motion seemed acceptable (Table 1) and showed medium reliability (ICC = 0.54), which is consistent with previous studies (Van Dijk et al. 2012). Although all of the diffusion metrics, as well as the head motion estimate, are relatively reliable across the two scans, they were not exactly the same due to some random artifacts, including motion artifacts and machine noises. Thus, for accurate measures of this microstructure and the head motion estimate, we first averaged the head motion and diffusion metrics across the two sessions and mainly examined the results with the averaged data.

---insert Table 1. ---

Relationship between the head motion estimate and diffusion metrics

In Table 1, we also show a basic summary of the main head motion results of diffusion metrics across all three analyses.

Among the two mostly commonly used regional diffusion metrics (i.e., FA and MD), these results indicated that head motion was mainly associated with the MD values. The degree of head motion was positively associated with increased MD mainly within white matter tracts in left hemisphere, including

anterior limb of internal capsule, posterior limb of internal capsule, genu of corpus callosum and body of corpus callosum (Fig. 1). In the current report, we focus on voxels that survived the TFCE correction ($p < 0.05$) (Smith & Nichols 2009) within the whole white matter skeleton. For the analyses examining FA, no voxel survived correction for multiple comparisons.

For the analyses examining the inter-voxel diffusion metric (i.e., LDH), we found that wide-spread white matter showed significant negative association with head motion ($p < 0.05$, TFCE corrected; Fig. 1). This association mainly involved the bilateral superior longitudinal fasciculus, body and genu of corpus callosum, cingulum, superior, anterior and posterior corona radiata, retrolenticular part of internal capsule, fornix, cerebral peduncle, middle cerebellar peduncle, right anterior and posterior limb of internal capsule, right external capsule and sagittal stratum.

---insert Fig. 1. ---

Overall, with the same criterion of significance ($p < 0.05$, TFCE corrected), we found the most number of motion-related voxels with LDH (34551 voxels, 32.25%) and then MD (2686 voxels, 2.51%). No voxel survived with FA.

In addition, we also examined the association between head motion and diffusion metrics with data from the two sessions respectively. Although no voxel survived statistical correction for multiple comparisons ($p < 0.05$) in most of the analyses (except LDH in Session 2), there were some voxels that showed a significant trend ($p < 0.10$, TFCE corrected; Fig. 2).

---insert Fig. 2. ---

Discussion

Like any other MRI technique, DW-MRI signal is subject to head motion artifacts, however, the relationship between diffusion metrics and head motion remains incompletely understood. Previous studies have shown a significant relationship between diffusion metrics and head motion (Ling et al. 2012) with conventional scanning protocol. The current study expands on previous work by exploring the relationship between motion and diffusion metrics (including the recently proposed inter-voxel metric, LDH) with a multiband dataset. We found that the motion effects in FA and MD seems to be mitigated to some extent, but the effect on MD still

existed. In addition, the effect is much more pronounced in LDH. Since these results are present following standard processing procedures, researchers shall be cautious when conducting data analysis and interpretation.

Previous studies suggested a positive association between motion and MD, with increased magnitude of MD as a result of increased total motion (Ling et al. 2012). The results of this study, with multiband dataset, replicate this finding, as a positive relationship between head motion and the magnitude of MD was present in the left hemisphere tracts. The significant association was mainly located in the deeper white matter (e.g., corpus callosum and the internal capsule). Interestingly, these tracts have often been reported in the literature to differ between a variety of clinical populations and healthy subjects (Carrasco et al. 2012; Travers et al. 2012). For examining FA, we found no significant relationship between head motion and FA after multiple comparison correction. On the one hand, our findings appear consistent with the previous finding (Ling et al., 2012) that the head motion's bias is more pronounced in MD than FA. But on the other hand, given the reduction of the number of motion-related voxels (multiband dataset: 0 voxel for FA, 2686 voxels for MD; Ling et al., (2012): 2422 voxels for FA, 22679 voxels for MD), the motion effect seems to be mitigated in the multiband dataset. This may be due to several advantages of the multiband dataset. First, the multiband dataset was acquired with much more gradient directions than traditional datasets, which would result in more accuracy when evaluating diffusion metrics. Second, the multiband scanning protocol allows full-brain coverage through the acquisition of multiple slices simultaneously. This could avoid displacements of brain within a TR and further mitigate the motion effects. Finally, given the fact that multiband protocol is designed for a relatively high sampling rate (i.e., a shorter TR), motion effects from a shorter duration would be expected to decrease. All these advantages could result in higher accuracy and less irrelevant effects (e.g., head motion) when evaluating diffusion metrics.

In addition, we also explored the relationship between head motion and LDH values and found that there were widespread voxels significantly associated to head motion. It's worth noting that with a smaller neighbor sizes ($n = 19$ or $n = 7$) when calculating the LDH, we observed the similar result (number of voxels that survived the TFCE correction: 35570 voxels for $n = 19$; 40425 voxels for $n = 7$). This appears to be quite likely caused by motion artifacts. Indeed, the significant voxels were rarely located in the occipital white matter tracts, where motion artifacts may be much weaker than that in prefrontal lobe when subjects are laying supine. These results suggest that LDH values might be more subject to head motion artifacts. The increased susceptibility to motion may be due to the fact that it is an inter-voxel metric which would be subject to shear in the displacement, and that it is directly calculated with raw diffusivity series. Though previous studies have shown LDH values change during aging, the newly proposed metric has not yet well validated (Gong 2013) and simulation and experimental work is required to confirm the motion-LDH association.

So, why does the association between motion and diffusion metrics exist? The dominant view at present is that head motion introduces artifacts into diffusion signals, similar to what has been noted in the fMRI literature (Bullmore et al. 1999; Friston et al. 1996; Hajnal et al. 1994), which influence the calculation of diffusion metrics and further results of cross-subject analysis. A common strategy for controlling motion effects in neuroimaging cross-subject analysis is to regress or match motion estimates (Zuo et al. 2010a; Zuo et al. 2012; Zuo et al. 2010b). Another strategy for mitigating head-motion artifacts is to remove time series of high motion, which is called ‘scrubbing’ (Power et al. 2012). However, these strategies have their limitations. On the one hand, scrubbing volumes with high motion could not fundamentally change the relationship between motion and values of diffusion metrics (Ling et al. 2012). On the other hand, they may also reduce the ability to detect a significant effect of interest, and/or introduce sampling bias (Satterthwaite et al. 2012; Wylie et al. 2014).

While researchers attempt to propose more sophisticated algorithms, there is growing perception in the field that head motion reflects individual differences in psychological traits and clinical conditions. For instance, previous studies showed that head motion was correlated with some psychological and clinical measures, such as the autism symptom severity score (Yendiki et al. 2013). In addition, previous fMRI studies suggest that the association may reflect the neural processing related to head motion (Yan et al. 2013a; Yan et al. 2013b). However, it is important to note that this problem in dMRI would not be as serious as it is in fMRI, since in dMRI the neural processing causing the motion does not directly affect the signal intensity. Nevertheless, these findings do suggest that head motion might not simply be an uncorrelated random variable.

Taken together, as articulated previously (Van Dijk et al. 2012; Wylie et al. 2014; Yendiki et al. 2013), these findings demonstrate the significance of developing motion-compensated acquisition methods for DW-MRI and incorporating them into neuroimaging studies in the future. Nevertheless, with current technologies, it appears impossible to perfectly eliminate the motion effects. As a temporary solution, examining both models with and without motion being regressed out will be expected. But in this case, researcher should include both results in the report, rather than just pick a ‘better’ one. Additionally, researcher shall keep in mind that motion does not only influence MRI signals, but also correlated with some meaningful individual differences. Alternatively, replication in an independent sample would be helpful, since the effects of head motion on diffusion metrics are usually random and not specific to some brain regions. Nevertheless, for now, researchers shall be cautious when doing MRI data analysis and interpretation.

In sum, the results of this study indicate that, in the multiband diffusion data, there are also significant associations between head motion and diffusion metrics, although the motion effects appear to be mitigated

232 compared to those with traditional dataset. Specifically, head motion was associated with both MD and LDH,
 233 and no significant effect was found for FA. Future studies should investigate the association between head
 234 motion and diffusion metrics with larger multiband datasets.

235 References

- 236 Bullmore ET, Brammer MJ, Rabe-Hesketh S, Curtis VA, Morris RG, Williams SC, Sharma T, and
237 McGuire PK. 1999. Methods for diagnosis and treatment of stimulus-correlated motion in generic brain
238 activation studies using fMRI. *Hum Brain Mapp* 7:38-48.
- 239 Carrasco JL, Tajima-Pozo K, Diaz-Marsa M, Casado A, Lopez-Ibor JJ, Arrazola J, and Yus M. 2012.
240 Microstructural white matter damage at orbitofrontal areas in borderline personality disorder. *J Affect*
241 *Disord* 139:149-153.
- 242 Durston S, Tottenham NT, Thomas KM, Davidson MC, Eigsti I-M, Yang Y, Ulug AM, and Casey BJ.
243 2003. Differential patterns of striatal activation in young children with and without ADHD. *Biol Psychiatry*
244 53:871-878.
- 245 Friston KJ, Williams S, Howard R, Frackowiak RS, and Turner R. 1996. Movement-related effects in
246 fMRI time-series. *Magn Reson Med* 35:346-355.
- 247 Golby AJ, Kindlmann G, Norton I, Yarmarkovich A, Pieper S, and Kikinis R. 2011. Interactive
248 diffusion tensor tractography visualization for neurosurgical planning. *Neurosurgery* 68:496-505.
- 249 Gong G. 2013. Local diffusion homogeneity (LDH): an inter-voxel diffusion MRI metric for assessing
250 inter-subject white matter variability. *PLoS One* 8:e66366.
- 251 Gong G, He Y, Concha L, Lebel C, Gross DW, Evans AC, and Beaulieu C. 2009. Mapping anatomical
252 connectivity patterns of human cerebral cortex using in vivo diffusion tensor imaging tractography. *Cereb*
253 *Cortex* 19:524-536.
- 254 Gorgolewski K, Burns CD, Madison C, Clark D, Halchenko YO, Waskom ML, and Ghosh SS. 2011.
255 Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Front*
256 *Neuroinform* 5:13.
- 257 Hajnal JV, Myers R, Oatridge A, Schwieso JE, Young IR, and Bydder GM. 1994. Artifacts due to
258 stimulus correlated motion in functional imaging of the brain. *Magn Reson Med* 31:283-291.
- 259 Ikuta T, Shafritz KM, Bregman J, Peters BD, Gruner P, Malhotra AK, and Szeszko PR. 2014.
260 Abnormal cingulum bundle development in autism: a probabilistic tractography study. *Psychiatry Res*
261 221:63-68.
- 262 Jenkinson M, Bannister P, Brady M, and Smith S. 2002. Improved optimization for the robust and
263 accurate linear registration and motion correction of brain images. *Neuroimage* 17:825-841.
- 264 Johansen-Berg H. 2010. Behavioural relevance of variation in white matter microstructure. *Curr Opin*
265 *Neurol* 23:351-358.
- 266 Kochunov P, Du X, Moran LV, Sampath H, Wijtenburg SA, Yang Y, Rowland LM, Stein EA, and Hong
267 LE. 2013. Acute nicotine administration effects on fractional anisotropy of cerebral white matter and
268 associated attention performance. *Front Pharmacol* 4:117.
- 269 Kong, X-Z. (2014). Head motion in children with ADHD during resting-state brain imaging. PeerJ
270 PrePrints, 2.
- 271 Kong X-Z, Zhen Z, Liu J. (2014, April): Measuring Regional Diffusivity Dependency via Mutual
272 Information. Presented at the IEEE International Symposium on Biomedical Imaging. April 29-May 2.
273 2014. Beijing. China.
- 274 Landman BA, Farrell JA, Jones CK, Smith SA, Prince JL, and Mori S. 2007. Effects of diffusion
275 weighting schemes on the reproducibility of DTI-derived fractional anisotropy, mean diffusivity, and
276 principal eigenvector measurements at 1.5T. *Neuroimage* 36:1123-1138.

- 277 Le Bihan D. 2003. Looking into the functional architecture of the brain with diffusion MRI. *Nat Rev*
278 *Neurosci* 4:469-480.
- 279 Le Bihan D, Mangin JF, Poupon C, Clark CA, Pappata S, Molko N, and Chabriat H. 2001. Diffusion
280 tensor imaging: concepts and applications. *J Magn Reson Imaging* 13:534-546.
- 281 Ling J, Merideth F, Caprihan A, Pena A, Teshiba T, and Mayer AR. 2012. Head injury or head motion?
282 Assessment and quantification of motion artifacts in diffusion tensor imaging studies. *Hum Brain Mapp*
283 33:50-62.
- 284 Mennes M, Biswal BB, Castellanos FX, and Milham MP. 2012. Making data sharing work: The
285 FCP/INDI experience. *Neuroimage*.
- 286 Moeller S, Yacoub E, Olman CA, Auerbach E, Strupp J, Harel N, and Ugurbil K. 2010. Multiband
287 multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to
288 high spatial and temporal whole-brain fMRI. *Magn Reson Med* 63:1144-1153.
- 289 Power JD, Barnes KA, Snyder AZ, Schlaggar BL, and Petersen SE. 2012. Spurious but systematic
290 correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 59:2142-2154.
- 291 Satterthwaite TD, Wolf DH, Loughhead J, Ruparel K, Elliott MA, Hakonarson H, Gur RC, and Gur RE.
292 2012. Impact of in-scanner head motion on multiple measures of functional connectivity: relevance for
293 studies of neurodevelopment in youth. *Neuroimage* 60:623-632.
- 294 Schmidt SL, Simões EN, and Schmidt GJ. 2013. The Effects of Hand Preference on Attention.
295 *Psychology* 4:29-33.
- 296 Shrout PE, and Fleiss JL. 1979. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*
297 86:420-428.
- 298 Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, Watkins KE,
299 Ciccarelli O, Cader MZ, Matthews PM, and Behrens TE. 2006. Tract-based spatial statistics: voxelwise
300 analysis of multi-subject diffusion data. *Neuroimage* 31:1487-1505.
- 301 Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, Bannister PR,
302 De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang YY, De Stefano N, Brady JM,
303 and Matthews PM. 2004. Advances in functional and structural MR image analysis and implementation as
304 FSL. *Neuroimage* 23:S208-S219.
- 305 Smith SM, Johansen-Berg H, Jenkinson M, Rueckert D, Nichols TE, Miller KL, Robson MD, Jones
306 DK, Klein JC, Bartsch AJ, and Behrens TE. 2007. Acquisition and voxelwise analysis of multi-subject
307 diffusion data with tract-based spatial statistics. *Nat Protoc* 2:499-503.
- 308 Smith SM, and Nichols TE. 2009. Threshold-free cluster enhancement: addressing problems of
309 smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* 44:83-98.
- 310 Tijssen RH, Jansen JF, and Backes WH. 2009. Assessing and minimizing the effects of noise and
311 motion in clinical DTI at 3 T. *Hum Brain Mapp* 30:2641-2655.
- 312 Travers BG, Adluru N, Ennis C, Tromp do PM, Destiche D, Doran S, Bigler ED, Lange N, Lainhart
313 JE, and Alexander AL. 2012. Diffusion tensor imaging in autism spectrum disorder: a review. *Autism Res*
314 5:289-313.
- 315 Van Dijk KR, Sabuncu MR, and Buckner RL. 2012. The influence of head motion on intrinsic
316 functional connectivity MRI. *Neuroimage* 59:431-438.
- 317 Wylie GR, Genova H, Deluca J, Chiaravalloti N, and Sumowski JF. 2014. Functional magnetic
318 resonance imaging movers and shakers: Does subject-movement cause sampling bias? *Hum Brain Mapp*
319 35:1-13.

- 320 Yan C-G, Craddock RC, He Y, and Milham MP. 2013a. Addressing Head Motion Dependencies for
- 321 Small-World Topologies in Functional Connectomics. *Frontiers in Human Neuroscience* 7:910.
- 322 Yan CG, Cheung B, Kelly C, Colcombe S, Craddock RC, Di Martino A, Li Q, Zuo XN, Castellanos
- 323 FX, and Milham MP. 2013b. A comprehensive assessment of regional variation in the impact of head
- 324 micromovements on functional connectomics. *Neuroimage*.
- 325 Yendiki A, Koldewyn K, Kakunoori S, Kanwisher N, and Fischl B. 2013. Spurious group differences
- 326 due to head motion in a diffusion MRI study. *Neuroimage*.
- 327 Zahneisen B, Grotz T, Lee KJ, Ohlendorf S, Reiser M, Zaitsev M, and Hennig J. 2011. Three-
- 328 dimensional MR-encephalography: fast volumetric brain imaging using rosette trajectories. *Magn Reson*
- 329 *Med* 65:1260-1268.
- 330 Zuo XN, Di Martino A, Kelly C, Shehzad ZE, Gee DG, Klein DF, Castellanos FX, Biswal BB, and
- 331 Milham MP. 2010a. The oscillating brain: complex and reliable. *Neuroimage* 49:1432-1445.
- 332 Zuo XN, Ehmke R, Mennes M, Imperati D, Castellanos FX, Sporns O, and Milham MP. 2012.
- 333 Network centrality in the human functional connectome. *Cereb Cortex* 22:1862-1875.
- 334 Zuo XN, Kelly C, Di Martino A, Mennes M, Margulies DS, Bangaru S, Grzadzinski R, Evans AC,
- 335 Zang YF, Castellanos FX, and Milham MP. 2010b. Growing together and growing apart: regional and sex
- 336 differences in the lifespan developmental trajectories of functional homotopy. *J Neurosci* 30:15034-15043.

Table 1 (on next page)

A basic summary of head motion and the motion effects in three diffusion metrics.

The column Motion includes the averaged motion in the sample. The column Motion-Brain Association includes the summary of motion effects in different diffusion metrics (i.e., FA, MD, and LDH). n.r. indicates null results; Plus sign (+) indicates a positive relationship, while minus (−) indicates a negative relationship. *: $p < 0.10$ TFCE corrected, **: $p < 0.05$ TFCE corrected.

Sample	Motion	Motion-Brain Association		
		FA	MD	LDH
Session 1	1.1(0.32)	n.r.	+	-
Session 2	1.26(0.38)	n.r.	+	-
Averaged	1.18(0.29)	n.r.	+	-

Figure 1

Results from the tract-based spatial statistics (TBSS) analyses depicting the voxels that showed a significant association between head motion diffusion metrics.

Data are presented for the analyses involving both Mean Diffusivity (MD; A) and local diffusion Local Diffusion Homogeneity (LDH; B) as the dependent measure. Participants with higher motion exhibited higher apparent values of MD, but lower LDH. Voxels survived the TFCE correction ($p < 0.05$) across the whole white matter skeleton are displayed.




Figure 2

Results from the tract-based spatial statistics (TBSS) analyses depicting the voxels that showed a significant association between head motion diffusion metrics.

Since no voxel that survived statistical correction for multiple comparisons ($p < 0.05$) in most of the analyses (except LDH in Session 2), they are displayed at a more tolerant threshold ($p < 0.10$, TFCE corrected).

