

A peer-reviewed version of this preprint was published in PeerJ on 18 August 2016.

[View the peer-reviewed version](https://peerj.com/articles/2275) (peerj.com/articles/2275), which is the preferred citable publication unless you specifically need to cite this preprint.

Raina J, Tapiolas D, Motti CA, Foret S, Seemann T, Tebben J, Willis BL, Bourne DG. 2016. Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals. PeerJ 4:e2275
<https://doi.org/10.7717/peerj.2275>

Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals

Jean-Baptiste Raina ^{Corresp., 1, 2, 3, 4, 5}, Dianne Tapiolas ², Cherie A Motti ², Sylvain Foret ^{3, 6}, Torsten Seemann ⁷, Jan Tebben ^{8, 9}, Bette L Willis ^{3, 4}, David G Bourne ^{2, 4}

¹ Climate Change Cluster (C3), University of Technology Sydney, Sydney, NSW, Australia

² Australian Institute of Marine Science, Townsville, QLD, Australia

³ James Cook University, Australian Research Council Centre of Excellence for Coral Reef Studies, Townsville, QLD, Australia

⁴ Marine Biology and Aquaculture, College of Science and Engineering, James Cook University of North Queensland, Townsville, QLD, Australia

⁵ James Cook University, AIMS@JCU, Townsville, QLD, Australia

⁶ Research School of Biology, Australian National University, Canberra, ACT, Australia

⁷ Victorian Life Sciences Computation Initiative, University of Melbourne, Melbourne, Victoria, Australia

⁸ Section Chemical Ecology, Alfred Wegener Institute, Bremerhaven, Germany

⁹ University of New South Wales, Sydney, NSW, Australia

Corresponding Author: Jean-Baptiste Raina

Email address: Jean-Baptiste.Raina@uts.edu.au

Bacterial communities associated with healthy corals produce antimicrobial compounds that inhibit the colonization and growth of invasive microbes and potential pathogens. To date, however, bacteria-derived antimicrobial molecules have not been identified in reef-building corals. Here we report the isolation of an antimicrobial compound produced by *Pseudovibrio* sp. P12, a common and abundant coral-associated bacterium. This strain was capable of metabolizing dimethylsulfoniopropionate (DMSP), a sulfur molecule produced in high concentrations by reef-building corals and playing a role in structuring their bacterial communities. Bioassay-guided fractionation coupled with nuclear magnetic resonance (NMR) and mass spectrometry (MS), identified the antimicrobial as tropodithietic acid (TDA), a sulfur-containing compound likely derived from DMSP catabolism. TDA was produced in large quantities by *Pseudovibrio* sp., and prevented the growth of two previously identified coral pathogens, *Vibrio coralliilyticus* and *V. owensii*, at very low concentrations (0.5 µg/mL) in agar diffusion assays. Genome sequencing of *Pseudovibrio* sp. P12 identified gene homologs likely involved in the metabolism of DMSP and production of TDA. These results provide additional evidence for the integral role of DMSP in structuring coral-associated bacterial communities and underline the potential of these DMSP-metabolizing microbes to contribute to coral disease prevention.

1 **Isolation of an antimicrobial compound produced by bacteria associated with reef-building**
2 **corals**

3
4 Jean-Baptiste Raina^{a,b,c,d,e*}, Dianne M. Tapiolas^b, Cherie A. Motti^b, Sylvain Forêt^{d,f}, Torsten
5 Seemann^g, Jan Tebben^{c,h,i}, Bette L. Willis^{a,d} & David G. Bourne^{a,b}

6
7 ^aJames Cook University, Marine Biology and Aquaculture, College of Science and Engineering,
8 Townsville 4810, QLD, Australia.

9 ^bAIMS@JCU, James Cook University, Townsville 4810, QLD, Australia

10 ^cAustralian Institute of Marine Science, Townsville 4810, QLD, Australia.

11 ^dAustralian Research Council Centre of Excellence for Coral Reef Studies, James Cook
12 University, Townsville 4810, QLD, Australia

13 ^eClimate Change Cluster (C3), University of Technology Sydney, Sydney 2007, NSW, Australia

14 ^fAustralian National University, Research School of Biology, Canberra 2601, ACT, Australia.

15 ^gVictorian Life Sciences Computation Initiative, University of Melbourne, Melbourne 3168,
16 VIC, Australia.

17 ^hUniversity of New South Wales, Sydney 2052, NSW, Australia.

18 ⁱAlfred Wegener Institute, Section Chemical Ecology, Bremerhaven 27570, Germany

19
20 *Corresponding author

21 Tel: +61 295 144 092

22 Email: Jean-Baptiste.Raina@uts.edu.au

23

24

25 **Abstract**

26 Bacterial communities associated with healthy corals produce antimicrobial compounds that
27 inhibit the colonization and growth of invasive microbes and potential pathogens. To date,
28 however, bacteria-derived antimicrobial molecules have not been identified in reef-building
29 corals. Here we report the isolation of an antimicrobial compound produced by *Pseudovibrio* sp.
30 P12, a common and abundant coral-associated bacterium. This strain was capable of
31 metabolizing dimethylsulfoniopropionate (DMSP), a sulfur molecule produced in high
32 concentrations by reef-building corals and playing a role in structuring their bacterial
33 communities. Bioassay-guided fractionation coupled with nuclear magnetic resonance (NMR)
34 and mass spectrometry (MS), identified the antimicrobial as tropodithietic acid (TDA), a sulfur-
35 containing compound likely derived from DMSP catabolism. TDA was produced in large
36 quantities by *Pseudovibrio* sp., and prevented the growth of two previously identified coral
37 pathogens, *Vibrio coralliilyticus* and *V. owensii*, at very low concentrations (0.5 µg/mL) in agar
38 diffusion assays. Genome sequencing of *Pseudovibrio* sp. P12 identified gene homologs likely
39 involved in the metabolism of DMSP and production of TDA. These results provide additional
40 evidence for the integral role of DMSP in structuring coral-associated bacterial communities and
41 underline the potential of these DMSP-metabolizing microbes to contribute to coral disease
42 prevention.

43

44

45

46

47

48

49

50 **Introduction**

51 Coral reefs are one of the most biologically diverse ecosystems on the planet (Bellwood
52 & Hughes 2001; Knowlton 2001; Pauley 1997). Each square centimeter of a coral's surface
53 harbors several thousand species of microbes (Sunagawa et al. 2010) and up to 10^7 bacterial cells
54 (Garren & Azam 2010; Koren & Rosenberg 2006), an abundance ten times greater, on average,
55 than the surrounding sea water. These microbial assemblages are often highly specific to their
56 coral host and include large numbers of rare and sometimes even unique taxa (Sunagawa et al.
57 2010). Although the phylogenetic diversity and dynamics of coral-associated bacterial
58 communities have been studied for more than a decade (Bourne & Munn 2005; Bourne et al.
59 2013; Littman et al. 2009; Rohwer et al. 2001; Sunagawa et al. 2010), their ecological and
60 functional roles in the biology and health of corals are still poorly understood.

61

62 Recent studies have started to unravel the roles that coral-associated bacteria and their
63 interactions with their coral hosts are likely to play within the coral holobiont. For example,
64 some members of the *Cyanobacteria*, *Rhizobiales* and *Vibrionaceae* taxa are likely to fix
65 dissolved nitrogen, a particularly important process in oligotrophic environments such as coral
66 reefs (Lema et al. 2012; Lesser et al. 2004; Olson et al. 2009). Others, like *Roseobacter*,
67 *Pseudomonas* and *Oceanospirillales*, can metabolize dimethylsulfoniopropionate (DMSP), an
68 organic sulfur compound produced in large quantities by corals and suspected to play a role in
69 structuring coral-associated bacterial communities (Raina et al. 2010; Raina et al. 2013). It has
70 also been hypothesized that bacteria act as a line of defense against invasive pathogens, either by
71 competing for space and occupying coral niches (Ritchie & Smith 2004), or by directly

72 producing antimicrobial compounds that inhibit the growth of invasive microbes in coral mucus
73 (Ritchie 2006).

74

75 In artificial culture conditions, approximately 25% of the cultivable coral bacteria
76 produce antimicrobial compounds that prevent the growth of pathogenic micro-organisms
77 (Ritchie 2006; Shnit-Orland & Kushmaro 2008). Several of these antimicrobial-producing taxa,
78 such as *Pseudoalteromonas*, *Pseudomonas*, and the *Roseobacter* clade are commonly found in
79 association with numerous coral species (Nissimov et al. 2009; Radjasa et al. 2008; Rypien et al.
80 2010; Shnit-Orland & Kushmaro 2009). Although the presence of antimicrobial defences in reef-
81 building corals has been reported (Geffen et al. 2009; Geffen & Rosenberg 2005; Gochfeld &
82 Aeby 2008; Koh 1997), only few active compounds – all produced by the coral animal itself –
83 have been isolated to date (Fusetani et al. 1996; Kodani et al. 2013; Vidal-Dupiol et al. 2011).
84 The aim of this study was to identify specific antimicrobial compounds and thereby enhance our
85 understanding of the functional roles played by coral-associated bacteria. Our specific objectives
86 were to: (i) isolate a common coral-associated bacterium with antimicrobial activity and identify
87 the compound(s) responsible for the activity; (ii) evaluate the susceptibility of the coral
88 pathogens *Vibrio coralliilyticus* and *Vibrio owensii* to the isolated compound; (iii) determine the
89 effect that thermal stress might have on its production; and (iv) investigate the natural abundance
90 of the antimicrobial compound in coral extracts.

91

92 **Materials and Methods**

93 *Bacterial isolation*

94 Healthy colonies of the corals *Pocillopora damicornis*, *Acropora millepora* and
95 *Montipora aequituberculata* (one colony per species) were collected in November 2011 from
96 Davies Reef, Great Barrier Reef, Australia (latitude, 18°51'S; longitude, 147°41'E, Great Barrier
97 Reef Marine Park Authority permit G12/35236.1) and maintained in aquaria for 6 days at the
98 Australian Institute of Marine Science (Townsville, Queensland, Australia). Five replicate coral
99 fragments (approximately 25 mm in length, containing 60 to 70 polyps) were collected from
100 each colony and washed in sterile artificial seawater (ASW) to remove loosely attached
101 microbes. Tissue slurries were produced by airbrushing (80 lb/in²) each coral fragment into 5 mL
102 of ASW to remove coral tissues and associated microbes. These tissue slurries were
103 homogenized to break down tissue clumps, and a dilution series was plated immediately on
104 bacteriological agar (1%) in 1 L ASW supplemented with 0.3% casamino acids and 0.4%
105 glucose (Hjelm et al. 2004). After 2 days of incubation at 28°C, single colonies were transferred
106 into Marine Broth (MB; Difco, BD, Franklin Lakes, NJ) and grown overnight. Liquid cultures
107 were re-plated on minimal marine agar and the procedure was repeated until pure cultures were
108 obtained.

109

110 *Well diffusion assay with bacterial isolates*

111 Fifty bacteria isolated from the coral tissue slurries of the 3 species (*A. millepora* = 16, *P.*
112 *damicornis* = 17, *M. aequituberculata* = 17) were tested for growth-inhibitory activity against the
113 known coral pathogens *Vibrio coralliilyticus* P1 (LMG23696) and *V. owensii* DY05
114 (LMG25443) in a well diffusion agar assay. In brief, the *Vibrio* strains were seeded into two
115 different batches of minimal marine agar (after the agar temperature cooled to 40°C). Following
116 solidification, wells (diameter 5 mm) were cut into the agar and loaded with 20 µL of overnight

117 cultures (10^8 cells/ml) of the test isolates grown in MB (28°C, 170 rpm). Plates were incubated at
118 28°C and monitored every 24 h for a period of 72 h for inhibition zones. *Phaeobacter* strain 27-4
119 was used as a positive antagonistic control on each plate because of its broad spectrum inhibitory
120 activity against *Vibrio* (Bruhn et al. 2007; Hjelm et al. 2004).

121

122 *DNA extraction, gene sequencing genomic analyses*

123 One strain, P12 isolated from *Pocillopora damicornis*, produced the strongest growth-
124 inhibitory activity against the two target *Vibrio* strains. High molecular weight genomic DNA
125 from P12 was extracted using a miniprep phenol-chloroform based extraction. Briefly, 5 mL of
126 overnight liquid culture of P12 (10^8 cells/ml) were spun in a micro-centrifuge (10,000 rcf) for 2
127 min. The pellet was then resuspended in 567 μ L of TE buffer, 30 μ L of 10% SDS and 3 μ L of 20
128 mg/mL proteinase K. The tube was shaken thoroughly and incubated for 1 hr at 37°C. One
129 hundred microliters of 5M NaCl was subsequently added and the sample thoroughly mixed
130 before adding 80 μ L of CTAB/NaCl (10% CTAB in 0.7 M NaCl). The solution was incubated
131 for 10 min at 65°C, extracted with an equal volume of phenol/chloroform/isoamyl alcohol and
132 centrifuged for 10 mins (10,000 rcf). The supernatant was then extracted with an equal volume
133 of chloroform/isoamyl alcohol and centrifuged again for 10 min. The aqueous phase was
134 transferred to a new tube, DNA precipitated with equal volume of ice-cold isopropanol, washed
135 with 70% ethanol and dried.

136 The near complete 16S rRNA gene of the strain was PCR amplified with bacterial
137 specific primers 63F and 1387R, as outlined in Marchesi et al. (Marchesi et al. 1998). Amplified
138 PCR products were visualized by electrophoresis on 1% agarose gel stained with ethidium
139 bromide. The amplified DNA was dried in a vacuum centrifuge (Savant DNA 120) and

140 sequenced (Macrogen, Inc., Seoul, Korea). The 16S rRNA gene sequence of isolate P12 was
141 used for phylogenetic comparisons and Maximum Likelihood trees were constructed using the
142 ARB software.

143 We produced a draft genome assembly of P12. A paired-end library was prepared using
144 the Illumina Truseq protocol (Illumina, San Diego, CA, USA), with an insert size of 169 bp and
145 a read size of 150 bp. The library was sequenced on an Illumina MiSeq instrument at Monash
146 University (Melbourne, Australia). The genome was assembled with the SPAdes assembler
147 (v2.4.0) (Bankevich et al. 2012) and annotated with the Prokka software (v1.5.2) (Seeman 2014).
148 The presence of the genes involved in DMSP metabolism (*dmdA*, *dddD*, *dddL*, *dddP*, *dddY*,
149 *dddQ*, *dddW*) and TDA production (*tdaA-tdaH*) was investigated by searching for homologs of
150 the corresponding genes using reciprocal best blast hits.

151

152 *DMSP metabolic capabilities of the isolate P12*

153 Two different minimal media were used to examine the DMSP metabolic capabilities of
154 P12: a modified marine ammonium salt medium (MAMS) (Raina et al. 2009) lacking a carbon
155 source, and a modified basal salt medium lacking a sulfur source (Fuse et al. 2000) (25 g of
156 NaCl, 0.7 g of KCl, 0.05 g of KH₂PO₄, 1 g of NH₄NO₃, 0.2 g of MgCl₂·H₂O, 0.02 g of
157 CaCl₂·2H₂O, 0.005 g of FeEDTA, 1 g of Tris, 5 g of sodium succinate, 1.35 g of glucose in 1 L
158 of distilled water). DMSP was added to both media (1 mM), acting either as the sole carbon or
159 sulfur source. Five milliliters of each culture media were inoculated in triplicate with single P12
160 colonies and incubated at 28°C for 6 days. Negative controls containing only the basal media and
161 DMSP were used to account for possible chemical breakdown of DMSP. Bacterial growth was
162 assessed *via* optical density measurement (NanoDrop, Thermo Fisher, Waltham, MA). DMSP

163 metabolism was assessed by ^1H Nuclear Magnetic Resonance spectroscopy (NMR). Methanol
164 (CH_3OH ; 40 mL) was added to each culture tube, the mixture shaken vigorously and sonicated
165 for 10 mins before being dried *in vacuo* using a rotary evaporator (Buchi, Flawil, Switzerland).
166 The dried extracts were resuspended in a mixture of deuterium oxide (D_2O ; 250 μL) and
167 deuterated methanol (CD_3OD ; 750 μL) (Cambridge Isotope Laboratories, Andover, MA, USA).
168 A 750-mL aliquot of the particulate-free extract was transferred into a 5-mm Norell tube (Norell
169 Inc., Landisville, NJ, USA) and analyzed immediately using quantitative NMR (Tapiolas et al.
170 2013).

171

172 *Preparation of crude extracts for antagonist assays*

173 An overnight culture of P12 (8 mL) was used to inoculate 4 \times 250 mL of MB (total culture
174 volume = 1L). Bacterial cells were incubated for two days at 28°C (120 rpm); the culture broth
175 was then acidified to pH 2 with sulphuric acid before being exhaustively extracted with ethyl
176 acetate (3 \times 1.5 L). The extract was washed three times with MilliQ H_2O and dried *in vacuo*
177 using a rotary evaporator (Buchi). The dried extract was then weighed and resuspended in
178 CH_3OH (which was chosen for its ability to solubilize a wide range of compounds, its volatility
179 and its innocuity in small volume towards both *V. coralliilyticus* and *V. owensii*) and tested in
180 well-diffusion assays to confirm the extraction of the antimicrobial compound(s).

181

182 *Purification and characterization of active compound*

183 Purification of the crude extract was carried out using solid phase extraction on a
184 reversed phase C_{18} flash vacuum column (Septra C_{18} -E, Phenomenex, Torrance, CA, USA).
185 Eleven fractions were eluted sequentially with 20%, 40%, 60%, 80%, 90% CH_3OH in H_2O and

186 100% CH₃OH, followed by 20%, 50% and 100% dichloromethane (CH₂Cl₂) in CH₃OH, 40%
187 hexane in CH₂Cl₂ and finally 100% hexane. Each fraction was dried and resuspended in CH₃OH
188 (1 mg mL⁻¹). Well diffusion assays were prepared as described above. On each plate, test wells
189 were inoculated with 20 μL of each chromatographic fraction, or 20 μL of CH₃OH as a control,
190 and *Vibrio* growth monitored. The most active fraction (80% CH₃OH) presented an intense
191 yellow color. Fine orange-red needles were crystallized from this active fraction to yield
192 compound **1** (2.1 mg, 1.7 % dry weight of organic extract).

193

194 *NMR and FTMS analysis*

195 Identification and structural elucidation of compound **1** was achieved using liquid
196 chromatography - mass spectrometry (LC-MS), NMR, and Fourier Transform mass spectrometry
197 (FTMS). Likewise these techniques were used to monitor for the presence of compound **1** in
198 extracts and fractions. LC-MS analyses were performed on a Thermo Fisher Scientific Ultra
199 High Performance Liquid Chromatography system connected to an LTQ Orbitrap XL mass
200 spectrometer (Thermo Fisher Scientific, San Jose, CA). Samples were separated on a ACQUITY
201 UPLC BEH RP-C₁₈ column (130Å, 1.7 μm, 2.1 mm x 100 mm, solvents A = aqueous 0.1%
202 formic acid and B = acetonitrile, gradient elution 80% A:20% B for 0.5 min ramped up to 100%
203 B over 10 min, then held for 4 min, 400 μL) and detected by positive mode electrospray
204 ionisation using two different *m/z* ranges: 150-1500 and 170-400. ¹H and ¹³C NMR spectra of
205 compound **1** were acquired in a 5 mm 509-UP Norell NMR tube on a Bruker Avance 600 MHz
206 NMR spectrometer (Bruker, Germany) with a TXI cryoprobe using standard Bruker pulse
207 sequences. NMR spectra were referenced to residual ¹H and ¹³C resonances in deuterated
208 chloroform (CDCl₃). High resolution mass spectra of compound **1** were measured with a Bruker

209 BioApex 47e Fourier Transform Mass Spectrometer (FTMS) fitted with an Analytica of
210 Branford ESI source; ions were detected in negative mode within a mass range m/z 200-1,000 via
211 direct infusion at $120 \mu\text{l h}^{-1}$.

212

213 *Temperature-dependent activity*

214 The antimicrobial activity of P12 grown at 32°C (upper limit of coral thermal tolerance)
215 was compared to that of the control incubated at 28°C . The two cultures were grown overnight in
216 MB at the two different temperatures, and their densities were determined by flow-cytometry
217 (BD Accuri C6, Beckman Coulter, Brea, CA). Cell numbers were normalized prior to
218 inoculation into agar wells, and their activities against the two pathogens were compared using
219 well-diffusion assays as described above. The same procedure was repeated with compound **1**:
220 two vials containing equal concentrations ($2 \mu\text{M}$ of **1** in CH_3OH) were incubated overnight at
221 28°C or 32°C and their antimicrobial activities compared using the well diffusion assay.

222

223 *Preparation of coral extracts*

224 The coral species *Montipora aequituberculata*, *M. turtlensis*, *Pocillopora damicornis*,
225 *Acropora millepora*, and *Porites cylindrica* (one colony each; 500 g of dry skeleton per species)
226 were collected in July 2012 from Orpheus Island, Great Barrier Reef, Australia (latitude,
227 $18^\circ35'S$; longitude, $146^\circ20'E$, Great Barrier Reef Marine Park Authority permit G12/35236.1).
228 Coral tissues were airbrushed (80 lb/in^2) into $1 \mu\text{m}$ filtered seawater (FSW) (total volume = 500
229 mL), acidified to pH 2 with sulphuric acid and the solution exhaustively extracted with equal
230 volumes of ethyl acetate ($3 \times 750 \text{ mL}$). The combined organic layers were partitioned with
231 MilliQ H_2O , dried and tested in well-diffusion assays, as previously described for the bacterial

232 isolate extracts. The extracts of those coral species that exhibited antimicrobial activity were
233 subsequently fractionated as described above for the crude extract from P12 and tested in well-
234 diffusion assays. The active fractions were analyzed using ^1H NMR, FTMS and LC-MS.

235

236 Results

237 *Isolate P12: antimicrobial production, taxonomy and metabolic capabilities*

238 A total of 50 coral-associated bacterial isolates were obtained from tissue slurry
239 homogenates of the three coral species. Twelve of the 50 strains tested against the two
240 pathogenic *Vibrios* (*V. coralliilyticus* and *V. owensii*) inhibited their growth in well diffusion
241 assays. The bioactive isolate that exhibited the strongest *in vitro* activity against both pathogens,
242 isolate P12, originated from *Pocillopora damicornis* and produced growth inhibition zones of 5
243 mm (± 0.07 mm, $n=20$) against *V. owensii* and 2 mm (± 0.09 mm, $n=20$) against *V. coralliilyticus*.
244 The activity of P12 was temperature-dependent (Figure 1 A-B) and was significantly reduced
245 when grown at 32°C compared to 28°C (Unpaired T-Test, $n=20$, $df=38$, $t=30.61$, $*p<0.001$ for *V.*
246 *owensii* and $n=20$, $df=38$, $t=10.49$, $*p<0.001$ for *V. coralliilyticus*; Figure 1C). Based on its
247 bioactivity, the isolate P12 was selected for bioassay-guided fractionation.

248

249 According to its 16S rRNA gene sequence (NCBI accession number: KX198136), isolate
250 P12 is an alphaproteobacterium belonging to the *Rhodobacteraceae* family and the *Pseudovibrio*
251 genus. Its most closely related species is *Pseudovibrio denitrificans* (100% identity to the type
252 strain; Figure 2). Like other *P. denitrificans* strains (Enticknap et al. 2006), P12 colonies formed
253 brown mucoid colonies when grown on Marine Agar. The brown color was absent when the
254 strain was grown on minimal marine agar, with colonies appearing white. This strain was able

255 use DMSP as either a sole carbon or sole sulfur source (Figure 3). The complete utilization of
256 DMSP from the liquid media after 2 to 3 days of incubation, as well as the presence of its
257 metabolic byproduct dimethylsulfide (DMS), were confirmed by ^1H NMR. However acrylate,
258 another possible byproduct of DMSP metabolism, was not observed.

259

260 Among the seven different DMSP degradation pathways currently identified (Moran et
261 al. 2012), the full DMSP cleavage pathway (*dddD*, *dddB*, *dddC*, *dddT*, *dddR*; Table 1), involved
262 in the conversion of DMSP into DMS without formation of acrylate (Todd et al. 2007) (Table 1),
263 was identified in P12. We also identified possible orthologs for the demethylation pathway
264 (*dmdA*, *dmdB*, *dmdC* and *dmdD*) used by marine bacteria to assimilate sulfur from DMSP,
265 though these gene have low sequence identity to the genes originally identified in *Ruegeria*
266 *pomeroyi* DSS-3 (Howard et al. 2006; Reisch et al. 2011) (Table 1). The presence of these two
267 gene pathways corroborates the ^1H NMR measurements: the observed production of DMS
268 without acrylate formation following DMSP metabolism (DddD pathway); and the ability to use
269 DMSP as sole sulfur source (DmdA pathway) (Table 1).

270

271 *Identification of antimicrobial compounds produced by P12*

272 Well diffusion assays revealed that the crude extract from P12 retained the antimicrobial
273 properties of the strain against both *Vibrio* species. Purification of the active fractions using
274 reverse phase liquid chromatography yielded compound **1**: optically inactive orange-red crystals;
275 2.1 mg (1.7% dry weight); IR (film) ν_{max} 3420, 1660, 1280 cm^{-1} ; UV (PDA, CH_3OH) λ_{max} 512
276 nm; ^1H NMR spectrum (600 MHz, CD_3Cl): δ 7.12, 7.44, 7.45 and 16.7; ^{13}C NMR (150 MHz,
277 CD_3Cl): δ 120.3, 132.0, 136.0, 138.7, 149.5, 168.7, 171.7, and 183.5; HRESIMS m/z found

278 210.9534 (calculated for $C_8H_3O_3S_2^-$ 210.9529, Δ 2 ppm). Combined spectroscopic techniques
279 revealed that compound **1** was tropodithietic acid (TDA) (Brinkhoff et al. 2004; Penesyan et al.
280 2011) (Figure 4A).

281

282 Orthologs for 11 genes involved in TDA biosynthesis (Geng et al. 2008) were present in
283 the *Pseudovibrio* sp. P12 genome (Table 1). The biosynthesis of TDA correlated with production
284 of the yellow-brown pigmentation in the culture medium and antimicrobial activity, similar to
285 that previously reported (Brinkhoff et al. 2004; Bruhn et al. 2005; Porsby 2010). Both coral
286 pathogens were highly sensitive to TDA, with the pure compound still visually inhibiting their
287 growth at 0.5 μ g/mL (2.35 μ M; Figure 4B). In contrast to the decrease in antimicrobial activity
288 exhibited by *Pseudovibrio* sp. P12 after incubation at 32°C, TDA activity was not affected by
289 exposure to this temperature (Unpaired T-Test, $n=20$, $df=38$, $t=-0.94$, $p=0.355$ for *V. owensii*
290 and $n=20$, $df=38$, $t=0.632$, $p=0.531$ for *V. coralliilyticus*; Figure 1B-C).

291

292 *Investigating the presence of TDA in coral samples*

293 All of the extracts derived from the coral species investigated exhibited antimicrobial
294 activity against the two pathogens, with the inhibition zones for *P. cylindrica*, *M.*
295 *aequituberculata*, *M. turtlensis* and *P. damicornis* ranging from 3 to 5 mm in radius whilst the
296 inhibition zones for *A. millepora* were much smaller (1 mm on average). 1H NMR, LC-MS and
297 FTMS analyses of the extracts and the active fractions of all coral species did not confirm the
298 presence of TDA. The purified TDA could be detected by LC-MS in femtomolar concentrations
299 when the coral fractions were artificially spiked, indicating that this lack of detection was not due

300 to preferential ionization. Thus, TDA was either not present in the coral fractions tested or in
301 concentrations below the LC-MS detection threshold.

302

303 **Discussion**

304 While previous studies have identified corals as a source of bacteria with antimicrobial
305 activity (Radjasa et al. 2008; Sulistiyani et al. 2010), this study represents the first isolation and
306 identification of an antimicrobial compound produced by a reef building coral-associated
307 bacterium with activity against coral-associated pathogens. Tropodithietic acid (TDA) was
308 characterized from a pure culture of *Pseudovibrio* sp. P12 previously isolated from corals and
309 strongly inhibited the growth of two coral pathogens. Although we could not confirm the
310 presence of this molecule in the corals tested, TDA has the capacity to provide protective
311 antimicrobial properties to the coral host and prevent colonization by invasive bacterial species.

312

313 The strain P12 strongly inhibited the growth of *Vibrio coralliilyticus* and *V. owensii*, two
314 coral pathogens causing white syndrome (a collective term describing rapidly progressing tissue
315 loss, exposing band-like areas of white skeleton) (Ben-Haim et al. 2003; Sussman et al. 2008;
316 Ushijima et al. 2012; Willis et al. 2004). *Vibrio coralliilyticus* exhibits antimicrobial resistance to
317 a wide range of commercial antibiotics and is also resistant to the activities of a large number of
318 coral-associated bacteria (Rypien et al. 2010; Shnit-Orland & Kushmaro 2009; Vizcaino et al.
319 2010). Its resistance to commercial antibiotics is considerably greater than that of other marine
320 pathogens such as *V. parahaemolyticus* or *V. vulnificus*, and may contribute to its competitive
321 advantage within the coral holobiont, as well as its ability to infect corals (Vizcaino et al. 2010).

322 However, whilst *V. coralliilyticus* is resistant to many coral-associated bacteria, its growth was
323 strongly inhibited by the strain P12, emphasizing its antimicrobial capabilities.

324

325 The isolate P12 belongs to the bacterial genus *Pseudovibrio* (Shieh et al. 2004), and is
326 ubiquitously found in association with healthy sponges (Enticknap et al. 2006; Thiel & Imhoff
327 2003; Webster & Hill 2001) and corals (see Table 2). *Pseudovibrio* are thought to be involved in
328 symbiotic relationships with various organisms; they are vertically transmitted in large densities
329 by adult sponges to their larvae (Enticknap et al. 2006) and their presence is required for the
330 growth of the sulfur-oxidizing bacteria *Beggiatoa* in culture (Bondarev et al. 2013). Furthermore,
331 their genome is organized similarly to that of *Rhizobia*, a well-characterized symbiotic bacterium
332 (Bondarev et al. 2013; Enticknap et al. 2006; Kennedy et al. 2009). The full genome sequences
333 of *Pseudovibrio* FO-BEG1 (KEGG genome T01669; isolated from a Caribbean coral, and
334 sharing 100% sequence similarity with P12 based on its 16S rRNA gene sequence) reveal the
335 presence of genes involved in host-cell adhesion, interactions with eukaryotic cell machinery,
336 and production of secondary metabolites (Bondarev et al. 2013), further suggesting that this
337 bacterium is involved in symbiotic relationships with its hosts.

338

339 The *Pseudovibrio* genus is also known for its antimicrobial properties, especially against
340 human pathogens such as *Mycobacterium tuberculosis*, *Bacillus cereus*, *Yersinia enterocolitica*,
341 *Listeria monocytogenes* or methicillin-resistant *Staphylococcus aureus* (O'Halloran et al. 2011;
342 Sulistiyani et al. 2010). To date, three active compounds have been isolated from different
343 *Pseudovibrio* strains: heptylprodigiocin in tunicate-associated *P. denitrificans* Z143-1 (Sertan-de
344 Guzman et al. 2007), pseudovibrocin in *P. denitrificans* strain PaH3.28 (Vizcaino 2011) and

345 tropodithietic acid (TDA) from red alga-associated *P. ascidiaceicola* D323 (Penesyan et al.
346 2011). In the present study, we isolated TDA from P12, a strain closely related to *P.*
347 *denitrificans*. The ability of P12 to produce TDA was further supported by the the presence of
348 the genes essential for TDA biosynthesis in the genome of strain P12 (Geng et al. 2008). TDA
349 inhibits the growth of a wide range of marine pathogens (Bruhn et al. 2007; Bruhn et al. 2005)
350 and is produced almost exclusively by bacteria from the *Roseobacter* clade, especially the genera
351 *Phaeobacter*, *Silicibacter*, and *Ruegeria* (Brinkhoff et al. 2004; Bruhn et al. 2005; Geng & Belas
352 2010; Geng et al. 2008; Wilson et al. 2016) that are commonly associated with DMSP-producing
353 dinoflagellates (Miller et al. 2004; Wagner-Dobler & Biebl 2006) and reef-building corals
354 (Bourne et al. 2013; Littman et al. 2009; Raina et al. 2009).

355

356 Many members of the *Roseobacter* clade, including coral-associated isolates, have been
357 implicated in sulfur cycling (Miller et al. 2004; Moran et al. 2003; Raina et al. 2010).
358 Interestingly, TDA contains two sulfur atoms ($C_8H_4O_3S_2$) and *Pseudovibrio* sp. P12 was able to
359 use DMSP either as sole carbon or sole sulfur source, a common trait among
360 *Alphaproteobacteria* and especially the *Roseobacter* clade (Bruhn et al. 2005; Wagner-Dobler &
361 Biebl 2006). Bacteria from this clade preferentially metabolize DMSP rather than sulphate (SO_4^{2-}
362), despite the latter being between 10^6 to 10^7 -fold more abundant in seawater (Geng & Belas
363 2010; Kiene et al. 1999). Based on genomic and chemical analyses, DMSP metabolism in P12
364 could occur *via* two likely routes: the cleavage pathway (encoded by *dddD*, (Todd et al. 2007))
365 that releases the climate-regulating molecule DMS, and the demethylation pathway (encoded by
366 *dmdA*, (Howard et al. 2006)) by which the bacterium can retain the sulfur contained in DMSP
367 molecules.

368

369 The biosynthetic pathway of TDA has not been fully elucidated (Brock et al. 2014). Both
370 labelling (Cane et al. 1992; Thiel et al. 2010) and genetic dissection (Geng & Belas 2010) studies
371 have shown that its aromatic skeleton is derived from phenylacetyl-CoA produced by the
372 shikimate pathway. However, the sulfur donor allowing the incorporation of the two sulfur atoms
373 into the TDA molecule has not been clearly identified. It has been proposed that sulfur
374 originating from DMSP metabolism might be used to synthesize TDA (Bruhn et al. 2007; Bruhn
375 et al. 2005; Geng & Belas 2010; Porsby 2010; Wagner-Dobler & Biebl 2006). For example,
376 DMSP increases TDA synthesis two-fold in comparison to other sulfur sources (Geng & Belas
377 2010), suggesting that DMSP is a preferred source of sulfur for TDA biosynthesis. Even though
378 other sources of sulfur, such as the amino-acids cysteine and methionine present in artificial
379 media like Marine Broth, might be used to synthesize TDA (Geng & Belas 2010), DMSP is by
380 far the most readily available reduced sulfur source in the marine environment (Simo 2001). It is
381 therefore likely that in DMSP-rich environments, such as reef-building corals, DMSP
382 metabolism provides the sulfur needed to produce TDA *via* the demethylation pathway (Howard
383 et al. 2006).

384

385 The presence of TDA in extracts derived from five coral species (*Montipora*
386 *aequituberculata*, *M. turtlensis*, *Pocillopora damicornis*, and *Porites cylindrica*) could not be
387 confirmed. Three possibilities can explain this lack of detection: (i) TDA is not synthesized in
388 corals; (ii) TDA is present in corals in concentrations below the detection limit of our
389 instruments (which would imply that this compound has a very limited role in coral defense);
390 (iii) our sampling effort was not sufficient. Indeed, we only sampled one colony per species,

391 from a location more than 100 km away from the site where the TDA-producing bacteria was
392 isolated and without prior characterization of the bacterial communities present in the colony
393 sampled. Given that TDA-producing *Roseobacters* are among the first bacteria to colonize the
394 surface of marine microalgae and corals (Apprill et al. 2009; Dang & Lovell 2000; Miller et al.
395 2004) and can be highly abundant in some coral species (Raina et al. 2009), it would be
396 premature to rule out possible TDA biosynthesis in corals.

397

398 The activity of P12 against *V. coralliilyticus* sharply decreased at elevated temperatures
399 (32°C), however, the activity of the purified TDA did not. This reveals that the loss of
400 antimicrobial activity observed for P12 at 32°C is not due to thermal sensitivity of TDA but
401 likely to a decrease in its production. Our results are in line with previously reported decline in
402 the antibacterial activity of other TDA-producing *Roseobacter* with temperature increase (Bruhn
403 et al. 2005). Clear links have previously been identified between warm thermal anomalies and
404 outbreaks of white syndromes (Bruno et al. 2007; Heron et al. 2010; Maynard et al. 2011). If
405 indeed TDA is synthesized *in vivo*, a decrease in its production during anomalously high
406 seawater temperatures could facilitate pathogen outbreaks in corals following thermal stress,
407 especially since the virulence of some disease-causing bacteria (i.e. *V. coralliilyticus*) increase at
408 32°C (Sussman et al. 2008).

409

410 This study demonstrates that a common coral-associated bacterium, *Pseudovibrio* sp.
411 P12, produces TDA, a potent antimicrobial compound that inhibits the growth of marine and
412 coral pathogens, including *V. coralliilyticus*. The bacterium can use DMSP as a sole sulfur or
413 carbon source and potentially as a precursor in the biosynthesis of TDA. The production of TDA

414 by *Pseudovibrio* sp. P12 is greatly reduced at temperatures causing thermal stress in corals,
415 potentially providing a window of opportunity for the growth of pathogens. These results provide
416 additional evidence for the integral role of DMSP in structuring healthy, coral-associated
417 bacterial communities and suggest that these DMSP-metabolizing communities may contribute
418 to the prevention of coral diseases.

419

420 **Acknowledgments**

421 The authors would like to thank E. Botté, C. Gao and M. Garren for their laboratory assistance.

422

423

424

425

426

427

428

429

430


431

432

433

434

435 **Figures**




436

437 **Figure 1:** Representative well diffusion assays of (A) *Pseudovibrio* sp. P12 and (B) pure TDA,
 438 incubated at two different temperatures (28°C and 32°C) and then inoculated onto agar plates
 439 with embedded *Vibrio owensii* [(-) Negative control]. (C) Comparison of the radius of inhibition
 440 zones between the two temperature treatments for both *Pseudovibrio* sp. P12 (Unpaired T-Test,
 441 $n=20$, $df=38$, $t=30.61$, $*p<0.001$ for *V. owensii* and $n=20$, $df=38$, $t=10.49$, $*p<0.001$ for *V.*
 442 *coralliilyticus*) and pure TDA (2 μ M, Unpaired T-Test, $n=20$, $df=38$, $t=-0.94$, $p=0.355$ for *V.*
 443 *owensii* and $n=20$, $df=38$, $t=0.632$, $p=0.531$ for *V. coralliilyticus*).

444


445



446
 447

448 **Figure 2:** Maximum likelihood phylogenetic tree based on 16S rRNA gene sequences showing
 449 the isolate used in this study (P12 in red) and closely associated *Pseudovibrio* spp. Note: the
 450 strain FO-BEG1 has been fully sequenced. *Phaeobacter* sp. 27-4 [AJ536669] was used as
 451 outgroup. Maximum parsimony bootstrap values (10,000 replicates) are given when different
 452 from 100. The scale bar indicates the number of substitution per nucleotide position.

453
 454




455

456

457 **Figure 3:** ^1H NMR spectra showing DMSP utilization as (A) sole carbon source and (B) sole
 458 sulfur source in minimal media at the end of a 6-day incubation. The “control” lines in all cases
 459 are the growth medium (with no bacterial inoculation). The black and green spectra show the
 460 results from inoculation with *Pseudovibrio sp.* P12 and *V. coralliilyticus* (negative control),
 461 respectively. In both cases, the DMSP signals (within the three boxes, see (Tapiolas et al. 2013))
 462 disappeared in the *Pseudovibrio* treatment and remain unchanged between the no-bacteria

463 control and the *V. coralliilyticus* treatment. In the case of DMSP as a sole sulfur source,
464 *Pseudovibrio* consumed the DMSP and other carbon sources present and produced secondary
465 metabolites (appearance of new signals). a: solvent peak (methanol).



466

467 **Figure 4:** (A) Tropodithietic acid (TDA). (B) Dilution series of pure TDA, showing zones of
468 growth inhibition against *Vibrio coralliilyticus*: [(1): 500 $\mu\text{g/mL}$, (2): 50 $\mu\text{g/mL}$, (3): 5 $\mu\text{g/mL}$,
469 (4): 500 ng/mL , (5): 50 ng/mL , (6): 5 ng/mL , (7): 500 pg/mL , (8): 50 pg/mL , (-) negative control
470 with solvent only].

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491
492
493
494
495
496
497
498
499
500
501
502
503

Table 1: Orthologous genes involved in DMSP degradation and TDA biosynthesis in *Pseudovibrio* sp. P12 genome. Accession numbers available in NCBI (<http://www.ncbi.nlm.nih.gov/genbank/>). Percent of similarity compared to ^a *Marinomonas* sp. MWY11 (Todd et al. 2007); ^b *Ruegeria pomeroyi* DSS-3 (Reisch et al. 2011); ^c *Pseudovibrio* sp. FO-BEG1 (Alcolombri et al. 2015); ^d *Phaeobacter inhibens* (Brock et al. 2014); ^e *Nesiotobacter exalbescens*; ^f *Rhodopseudomonas palustris*; ^g *Stappia stellulata*.

Gene	Function	Percent of identity	Accession number
<i>DddD</i>	L-carnitine dehydratase	70% ^a	KM819464
<i>DddT</i>	BCCT transporter	70% ^a	KM819465
<i>DddB</i>	Alcohol dehydrogenase	70% ^a	KM819466
<i>DddC</i>	Methylmalonate-semialdehyde dehydrogenase	56% ^a	KM819467
<i>DddR</i>	Transcriptional regulator, LysR family protein	53% ^a	KM819468
<i>DmdA</i>	Aminomethyl transferase family protein	24% ^b	KU521525
<i>DmdB</i>	Acyl-CoA synthetase	30% ^b	KU521526
<i>DmdC</i>	MMPA-CoA dehydrogenase	43% ^b	KU521527
<i>DmdD</i>	MTA-CoA hydratase	32% ^b	KU521528
<i>Alma1</i>	Hypothetical Alma1 ortholog	99% ^c	KU521524
<i>tdaA</i>	Transcriptional regulator, LysR family protein	66% ^d	KU760700
<i>tdaB</i>	Glutathione S-transferase domain protein	67% ^d	KU760701
<i>tdaC</i>	Prephenate dehydratase	67% ^d	KU760702
<i>tdaD</i>	Acyl-CoA thioester hydrolase	81% ^d	KU760703
<i>tdaE</i>	Acyl-CoA dehydrogenase	69% ^d	KU760704
<i>tdaF</i>	Phosphopantothenoylecysteine decarboxylase	72% ^d	KU760705
<i>cysI</i>	Sulfite reductase hemoprotein beta-component	76% ^e	KU760706
<i>malY</i>	Cystathione beta-lyase	68% ^e	KU760707
<i>paal</i>	Phenylacetate-CoA oxygenase	59% ^f	KU760708
<i>paaJ</i>	Phenylacetate-CoA oxygenase, PaaJ subunit	73% ^g	KU760709
<i>paaK</i>	Phenylacetic acid degradation oxidoreductase	57% ^g	KU760710

504
505
506
507
508

509
510
511
512
513
514
515
516
517
518
519
520

Table 2: Summary of *Pseudovibrio* isolated or sequenced from corals, accession numbers are displayed when available (NA: not available).

Host	Location	Method	Reference	Accession numbers
<i>Acropora palmata</i>	Panama	Amplicon	(Sunagawa et al. 2010)	GU118050, GU118108, GU119014
<i>Porites astreoides</i>	Panama	Amplicon	(Sunagawa et al. 2010)	GU118050, GU118108, GU119014
<i>Acropora cervicornis</i>	Panama	Amplicon	(Sunagawa et al. 2010)	GU118050, GU118108, GU119014
<i>Montastrea franksi</i>	Panama	Amplicon	(Sunagawa et al. 2010)	GU118050, GU118108, GU119014
<i>Tubastraea coccinea</i>	China	Amplicon	(Yang et al. 2013)	JF925014
<i>Pseudopterogorgia americana</i>	Puerto Rico	Isolated	(Vizcaino et al. 2010)	GQ406787, GQ406798, GQ391966, GQ406786
<i>Platygyra carnosus</i>	Hong Kong	Isolated	(Chiu et al. 2012)	JF411474, JF411466, JF411439, JF411464
<i>Oculina patagonica</i>	Israel	Isolated	(Koren & Rosenberg 2006)	DQ416557, AY654776
<i>Montastrea anularis</i>	Florida	Isolated	(Rypien et al. 2010)	FJ952798, FJ952774, FJ952804
<i>Sinularia sp.</i>	Indonesia	Isolated	(Sulistiyani et al. 2010)	NA
<i>Acropora almata</i>	Florida	Isolated	(Ritchie 2006)	DQ530540
<i>Sarcophyton sp.</i>	Java	Isolated	(Sabdono & Radjasa 2006)	NA
<i>Oculina patagonica</i>	Israel	Isolated	(Nissimov et al. 2009)	NA
<i>Lobophytum sp.</i>	Taiwan	Isolated	(Chen et al. 2012)	JQ342682, JQ342695, JQ342696, JQ342697
Hard coral	Florida	Isolated	(Bondarev et al. 2013)	CP003147

521

522 References

- 523 Alcolombri U, Ben-Dor S, Feldmesser E, Levin Y, Tawfik DS, and Vardi A. 2015. Identification
524 of the algal dimethyl sulfide-releasing enzyme: a missing link in the marine sulfur cycle.
525 *Science* 348:1466-1469.
- 526 Apprill A, Marlow HQ, Martindale MQ, and Rappe MS. 2009. The onset of microbial
527 associations in the coral *Pocillopora meandrina*. *ISME J* 3:685-699.
- 528 Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko
529 SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev
530 MA, and Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its
531 application to single-cell sequencing. *Journal of Computational Biology* 19:455-477. doi:
532 10.1089/cmb.2012.0021
- 533 Bellwood DR, and Hughes TP. 2001. Regional-scale assembly rules and biodiversity of coral
534 reefs. *Science* 292:1532-1535.

- 535 Ben-Haim Y, Thompson FL, Thompson CC, Cnockaert MC, Hoste B, Swings J, and Rosenberg
536 E. 2003. *Vibrio coralliilyticus* sp. nov., a temperature dependent pathogen of the coral
537 *Pocillopora damicornis*. *IJSEM* 53:309-315.
- 538 Bondarev V, Richter M, Romano S, Piel J, Schwedt A, and Schulz-Vogt HN. 2013. The genus
539 *Pseudovibrio* contains metabolically versatile bacteria adapted for symbiosis. *Environ*
540 *Microbiol* 15:2095-2113.
- 541 Bourne D, and Munn C. 2005. Diversity of bacteria associated with the coral *Pocillopora*
542 *damicornis* from the Great Barrier Reef. *Environ Microbiol* 7:1162-1174.
- 543 Bourne DG, Dennis PG, Uthicke S, Soo RM, Tyson GW, and Webster NS. 2013. Coral reef
544 invertebrate microbiomes correlate with the presence of photosymbionts. *ISME J* In
545 press. 10.1038/ismej.2012.172
- 546 Brinkhoff T, Bach G, Heidorn T, Liang L, Schlingloff A, and Simon M. 2004. Antibiotic
547 production by a *Roseobacter* clade-affiliated species from the German Wadden Sea and
548 its antagonistic effects on the indigenous isolates. *Appl Environ Microbiol* 70:2560-2565.
- 549 Brock NL, Nikolay A, and Dickschat JS. 2014. Biosynthesis of the antibiotic tropodithietic acid
550 by the marine bacterium *Phaeobacter inhibens*. *Chemical Communications* 50:5487-5489.
551 10.1039/C4CC01924E
- 552 Bruhn JB, Gram L, and Belas R. 2007. Production of antibacterial compounds and biofilm
553 formation by *Roseobacter* species are influenced by culture conditions. *Appl Environ*
554 *Microbiol* 73:442-450.
- 555 Bruhn JB, Nielsen KF, Hjelm M, Hansen M, Bresciani J, Schulz S, and Gram L. 2005. Ecology,
556 inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to
557 the *Roseobacter* clade. *Appl Environ Microbiol* 71:7263-7270.
- 558 Bruno JF, Selig ER, Casey KS, Page CA, Willis BL, Harvell DC, Sweatman H, and Melendy
559 AM. 2007. Thermal stress and coral cover as driver of coral disease outbreaks. *PLoS Biol*
560 5:e124. 10.1371/journal.pbio.0050124
- 561 Cane DE, Wu Z, and van Epp JE. 1992. Thiotropocin biosynthesis. Shikimate origin of a sulfur-
562 containing tropolone derivative. *J Am Chem Soc* 114:8479-8483.
- 563 Chen Y-H, Kuo J, Sung P-J, Chang Y-C, Lu M-C, Wong T-Y, Liu J-K, Weng C-F, Twan W-H,
564 and Kuo F-W. 2012. Isolation of marine bacteria with antimicrobial activities from
565 cultured and field-collected soft corals. *World Journal of Microbiology and*
566 *Biotechnology* 28:3269-3279. 10.1007/s11274-012-1138-7
- 567 Chiu JMY, Li S, Li A, Po B, Zhang R, Shin PKS, and Qiu J-W. 2012. Bacteria associated with
568 skeletal tissue growth anomalies in the coral *Platygyra carnosus*. *FEMS Microbiology*
569 *Ecology* 79:380-391. 10.1111/j.1574-6941.2011.01225.x
- 570 Dang HY, and Lovell CR. 2000. Bacterial primary colonization and early succession on
571 surfaces in marine waters as determined by amplified rRNA gene restriction analysis and
572 sequences analysis of 16S rRNA genes. *Appl Environ Microbiol* 66:467-475.
- 573 Enticknap JJ, Kelly M, Peraud O, and Hill RT. 2006. Characterization of a culturable
574 Alphaproteobacterial symbiont common to many marine sponges and evidence for
575 vertical transmission *via* sponge larvae. *Appl Environ Microbiol*.
- 576 Fuse H, Takimura O, Murakami K, Yamaoka Y, and Omori T. 2000. Utilization of dimethyl
577 sulfide as a sulfur source with the aid of light by *Marinobacterium* sp. strain DMS-S1.
578 *Appl Environ Microbiol* 66:5527-5532.

- 579 Fusetani N, Toyoda T, Asai N, Matsunaga S, and Maruyama T. 1996. Montiporic acids A and B,
580 cytotoxic and antimicrobial polyacetylene carboxylic acids from eggs of the scleractinian
581 coral *Montipora digitata*. *J Nat Prod* 59:796-797.
- 582 Garren M, and Azam F. 2010. New method for counting bacteria associated with coral mucus.
583 *Appl Environ Microbiol* 76:6128-6133.
- 584 Geffen Y, Ron EL, and Rosenberg E. 2009. Regulation of release of antibacterials from stressed
585 scleractinians corals. *FEMS Microbiology Letters* 295:103-109.
- 586 Geffen Y, and Rosenberg E. 2005. Stress-induced rapid release of antibacterials by scleractinian
587 corals. *Mar Biol* 146:931-935.
- 588 Geng H, and Belas R. 2010. Expression of tropodithietic acid biosynthesis is controlled by a
589 novel autoinducer. *J Bacteriol* 192:4377-4387.
- 590 Geng H, Bruhn JB, Nielsen KF, Gram L, and Belas R. 2008. Genetic dissection of tropodithietic
591 acid biosynthesis by marine roseobacters. *Appl Environ Microbiol* 74:1535-1545.
- 592 Gochfeld D, and Aeby GS. 2008. Antibacterial chemical defences in Hawaiian corals provide
593 possible protection from disease. *MEPS* 362:119-128.
- 594 Heron SF, Willis BL, Skirving WJ, Eakin M, Page CA, and Miller IR. 2010. Summer hot snaps
595 and winter conditions: modelling white syndrome outbreaks on Great Barrier Reef corals.
596 *PLoS One* 5:e12210. 10.1371/journal.pone.0012210
- 597 Hjelm M, Bergh O, Riaza A, Nielsen J, Melchiorson J, Jensen S, Duncan H, Ahrens P, Birkbeck
598 H, and Gram L. 2004. Selection and identification of autochthonous potential probiotic
599 bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst Appl Microbiol*
600 27:360-371.
- 601 Howard EC, Henriksen JR, Buchan A, Reisch CR, Burgmann H, Welsh R, Ye W, González JM,
602 Mace K, Joye SB, Kiene RP, Whitman WB, and Moran MA. 2006. Bacterial taxa that
603 limit sulfur flux from the ocean. *Science* 314:649-652.
- 604 Kennedy J, Baker P, Piper C, Cotter PD, Walsh M, Mooij MJ, Bourke MB, Rea MC, O'Connor
605 PM, Ross RP, Hill C, O'Gara F, Marchesi JR, and Dobson ADW. 2009. Isolation and
606 analysis of bacteria with antimicrobial activities from the marine sponge *Haliclona*
607 *simulans* collected from Irish waters. *Mar Biotechnol* 11:384-396.
- 608 Kiene RP, Linn LJ, Gonzalez J, Moran MA, and Bruton JA. 1999. Dimethylsulfoniopropionate
609 and methanethiol are important precursors of methionine and protein-sulfur in marine
610 bacterioplankton. *Appl Environ Microbiol* 65:4549-4558.
- 611 Knowlton N. 2001. Coral reef biodiversity - habitat size matters. *Science* 292:1493-1494.
- 612 Kodani S, Sato K, Higuchi T, Casareto BE, and Suzuki Y. 2013. Montiporic acid D, a new
613 polyacetylene carboxylic acid from scleractinian coral *Montipora digitata*. *Natural*
614 *Product Research* 27:1859-1862.
- 615 Koh EGL. 1997. Do scleractinian corals engage in chemical warfare against microbes. *Journal of*
616 *Chemical Ecology* 23:379-398.
- 617 Koren O, and Rosenberg E. 2006. Bacteria associated with mucus and tissues of the coral
618 *Oculina patagonica* in summer and winter. *Appl Environ Microbiol* 75:254-259.
- 619 Lema KA, Willis BL, and Bourne DG. 2012. Corals form characteristic associations with
620 symbiotic nitrogen-fixing bacteria. *Appl Environ Microbiol* 78:3136-3144.
- 621 Lesser MP, Mazel CH, Gorbunov MY, and Falkowski PG. 2004. Discovery of symbiotic
622 nitrogen-fixing cyanobacteria in corals. *Science* 305:997-1000.

- 623 Littman RA, Willis BL, Pfeffer C, and Bourne DG. 2009. Diversity of coral-associated bacteria
624 differ with location but not species for three Acroporids on the Great Barrier Reef. *FEMS*
625 *Microbiology Letters* 68:152-163.
- 626 Marchesi JR, Sato T, Weightman AJ, Martin AT, Fry JC, and Wade WG. 1998. Design and
627 evaluation of useful bacterium-specific primers that amplify genes coding for 16S rRNA.
628 *Appl Environ Microbiol* 64:795-799.
- 629 Maynard JA, Anthony KRN, Harvell DC, Burgman MA, Beeden R, Sweatman H, Heron SF,
630 Lamb JB, and Willis BL. 2011. Predicting outbreaks of a climate-driven coral disease in
631 the Great Barrier Reef. *Coral Reefs* 30:485-495.
- 632 Miller TR, Hnilicka K, Dziedzic A, Desplats P, and Belas R. 2004. Chemotaxis of *Silicibacter*
633 sp. strain TM1040 toward dinoflagellate products. *Appl Environ Microbiol* 70:4692-
634 4701.
- 635 Moran MA, González JM, and Kiene RP. 2003. Linking a bacterial taxon to sulfur cycling in the
636 sea: studies of the marine *Roseobacter* group. *Geomicrobiol J* 20:375-388.
- 637 Moran MA, Reisch CR, Kiene RP, and Whitman WB. 2012. Genomic insights into bacterial
638 DMSP transformations. *Annual Review of Marine Science* 4:523-542.
- 639 Nissimov J, Rosenberg E, and Munn CB. 2009. Antimicrobial properties of resident coral mucus
640 bacteria of *Oculina patagonica*. *FEMS Microbiol Lett* 292:210-215.
- 641 O'Halloran JA, Barbosa TM, Morrissey JP, Kennedy J, O'Gara F, and Dobson ADW. 2011.
642 Diversity and antimicrobial activity of *Pseudovibrio* spp. from Irish marine sponges. *J*
643 *Appl Microb* 110:1495-1508.
- 644 Olson ND, Ainsworth TD, Gates RD, and Takabayashi M. 2009. Diazotrophic bacteria
645 associated with Hawaiian *Montipora* corals: diversity and abundance in correlation with
646 symbiotic dinoflagellates. *J Exp Mar Biol and Ecol* 371:140-146.
- 647 Pauley G. 1997. Diversity and distribution of reef organisms. In: Birkeland C, ed. *Life and death*
648 *of coral reefs*. New York: Chapman and Hall, 298-345.
- 649 Penesyan A, Tebben J, Lee M, Thomas T, Kjelleberg S, Harder T, and Egan S. 2011.
650 Identification of the antibacterial compound produced by the marine epiphytic bacterium
651 *Pseudovibrio* sp. D323 and related sponge-associated bacteria. *Mar Drugs* 9:1391-1402.
- 652 Porsby CH. 2010. Antagonism of *Roseobacter* clade bacteria against pathogenic bacteria PhD
653 Thesis. Technical University of Denmark.
- 654 Radjasa OK, Wiese J, Sabdono A, and J.F. I. 2008. Coral as source of bacteria with antimicrobial
655 activity. *J Coastal Development* 11:121-130.
- 656 Raina JB, Dinsdale EA, Willis BL, and Bourne DG. 2010. Do the organic sulfur compounds
657 DMSP and DMS drive coral microbial associations? *Trends in Microbiology* 18:101-108.
- 658 Raina JB, Tapiolas D, Willis BL, and Bourne DG. 2009. Coral-associated bacteria and their role
659 in the biogeochemical cycling of sulfur. *Appl Environ Microbiol* 75:3492-3501.
- 660 Raina JB, Tapiolas DM, Forêt S, Lutz A, Abrego D, Ceh J, Seneca F, Clode PL, Bourne DG,
661 Willis BL, and Motti CA. 2013. DMSP biosynthesis by an animal and its role in coral
662 thermal stress response. *Nature* 502:677-680.
- 663 Reisch CR, Stoudemayer MJ, Varaljay VA, Amster IJ, Moran MA, and Whitman WB. 2011.
664 Novel pathway for assimilation of dimethylsulfoniopropionate widespread in marine
665 bacteria. *Nature* 473:208-211.
- 666 Ritchie KB. 2006. Regulation of microbial populations by coral surface mucus and mucus-
667 associated bacteria. *Marine Ecology Progress Series* 322:1-14.

- 668 Ritchie KB, and Smith GW. 2004. Microbial communities of coral surface mucopolysaccharide
669 layer. In: Rosenberg E, and Loya Y, eds. *Coral health and disease*. Berlin: Springer-
670 Verlag, 259-263.
- 671 Rohwer F, Breitbart M, Jara J, Azam F, and Knowlton N. 2001. Diversity of bacteria associated
672 with the Caribbean coral *Monastera franksi*. *Coral Reefs* 20:85-91.
- 673 Rypien KL, Ward JR, and Azam F. 2010. Antagonistic interactions among coral-associated
674 bacteria. *Environmental Microbiology* 12:28-39.
- 675 Sabdono A, and Radjasa OK. 2006. Antifouling activity of bacteria associated with soft coral
676 *Sarcophyton* sp. against marine biofilm-forming bacteria. *J Coastal Development* 10:55-
677 62.
- 678 Seeman T. 2014. Prokka: rapid prokariotic genome annotation. *Bioinformatics* 30:2068-2069.
- 679 Sertan-de Guzman AA, Predicala RZ, Bernardo EB, Neilan BA, Elardo SP, Mangalindan GC,
680 Tasdemir D, Ireland CM, Barraquio WL, and Concepcion GP. 2007. *Pseudovibrio*
681 *denitrificans* strain Z143-1, a heptylprodigiosin-producing bacterium isolated from a
682 Philippine tunicate. *FEMS Microbiol Lett* 277:188-196.
- 683 Shieh WY, Lin Y, and Jean WD. 2004. *Pseudovibrio denitrificans* gen. nov., sp. nov., a marine,
684 facultative anaerobic, fermentative bacterium capable of denitrification. *IJSEM* 54:2307-
685 2312.
- 686 Shnit-Orland M, and Kushmaro A. 2008. Coral mucus bacteria as a source of antibacterial
687 activity. *Proceedings of the 11th International Coral Reef Symposium*:257-259.
- 688 Shnit-Orland M, and Kushmaro A. 2009. Coral mucus-associated bacteria: a possible first line of
689 defence. *FEMS Microbiol Lett* 67:371-380.
- 690 Simo R. 2001. Production of atmospheric sulfur by oceanic plankton: biogeochemical, ecological
691 and evolutionary links. *Trends in Ecology and Evolution* 16:287-294.
- 692 Sulistiyani S, Nugraheni A, Radjasa OK, Sabdono A, and Khoeri MM. 2010. Antibacterial
693 activities of bacterial symbionts of soft coral *Simularia* sp. against tuberculosis bacteria. *J*
694 *Coastal Development* 14:45-50.
- 695 Sunagawa S, Woodley CM, and Medina M. 2010. Threatened corals provide underexplored
696 microbial habitats. *PLoS One* 5:e9554. 10.1371/journal.pone.0009554
- 697 Sussman M, Willis BL, Victor S, and Bourne DG. 2008. Coral pathogens identified for white
698 syndrome epizootic in the indo-pacific. *PLoS One* 3:e2393.
- 699 Tapiolas DM, Raina JB, Lutz A, Willis BL, and Motti CA. 2013. Direct measurement of
700 dimethylsulfoniopropionate (DMSP) in reef-building corals using quantitative nuclear
701 magnetic resonance (qNMR) spectroscopy. *JEMBE* 443:85-89.
- 702 Thiel V, Brinkhoff T, Dickschat JS, Wickel S, Grunenberg J, Wagner-Dobler I, Simon M, and
703 Schulz S. 2010. Identification and biosynthesis of tropone derivatives and sulfur volatiles
704 produced by bacteria of the marine *Roseobacter* clade. *Org Biomol Chem* 2010:234-246.
- 705 Thiel V, and Imhoff JF. 2003. Phylogenetic identification of bacteria with antimicrobial
706 activities isolated from Mediterranean sponges. *Biomolecular Engineering* 20:421-423.
- 707 Todd JD, Rogers R, Li YG, Wexler M, Bond PL, Sun L, Cuurson ARJ, Malin G, Steinke M, and
708 Johnston AWB. 2007. Structural and regulatory genes required to make the gas
709 dimethylsulfide in bacteria. *Science* 315:666-669.
- 710 Ushijima B, Smith A, Aeby GS, and Callahan SM. 2012. *Vibrio owensii* induces the tissue loss
711 disease *Montipora* white syndrome in the Hawaiiin reef coral *Montipora capitata*. *PLoS*
712 *One* 7:10.

- 713 Vidal-Dupiol J, Ladriere O, Destoumieux-Garzon D, Sautiere PE, Meistertzheim AL, Tambutte
714 E, Tambutte S, Duval D, Foure L, Adjeroud M, and Mitta G. 2011. Innate immune
715 responses of a scleractinian coral to vibriosis. *J Biol Chem* 286:22688-22698.
- 716 Vizcaino MI. 2011. The chemical defense of *Pseudopternogorgia americana*: a focus on the
717 antimicrobial potential of a *Pseudovibrio* sp. PhD Thesis. University of South Carolina.
- 718 Vizcaino MI, Johnson WR, Kimes NE, Williams K, Torralba M, Nelson KE, Smith GW, Weil E,
719 Moeller PDR, and Morris PJ. 2010. Antimicrobial resistance of the coral pathogen *Vibrio*
720 *coralliilyticus* and sister phylotypes isolated from a diseased Octocoral. *Microb Ecol*
721 59:646-657.
- 722 Wagner-Dobler I, and Biebl H. 2006. Environmental biology of the marine Roseobacter lineage.
723 *Annual Reviews Of Microbiology* 60:255-280.
- 724 Webster NS, and Hill RT. 2001. The culturable microbial community of the Great Barrier Reef
725 sponge *Rhopaloeides odorabile* is dominated by an Alphaproteobacterium. *Marine*
726 *Biology* 138:843-851.
- 727 Willis BL, Page CA, and Dinsdale EA. 2004. Coral disease on the Great Barrier Reef. In:
728 Rosenberg E, and Loya Y, eds. *Coral health and disease*: Springer, 69-104.
- 729 Wilson MZ, Wang R, Gitai Z, and Seyedsayamdost MR. 2016. Mode of action and resistance
730 studies unveil new roles for tropodithietic acid as an anticancer agent and the γ -glutamyl
731 cycle as a proton sink. *Proceedings of the National Academy of Sciences* 113:1630-1635.
732 10.1073/pnas.1518034113
- 733 Yang S, Sun W, Zhang F, and Li Z. 2013. Phylogenetically Diverse Denitrifying and Ammonia-
734 Oxidizing Bacteria in Corals *Alcyonium gracillimum* and *Tubastraea coccinea*. *Marine*
735 *Biotechnology* 15:540-551. 10.1007/s10126-013-9503-6

736

737