

A peer-reviewed version of this preprint was published in PeerJ on 13 September 2016.

[View the peer-reviewed version](https://peerj.com/articles/2444) (peerj.com/articles/2444), which is the preferred citable publication unless you specifically need to cite this preprint.

Kelly RP, O'Donnell JL, Lowell NC, Shelton AO, Samhoury JF, Hennessey SM, Feist BE, Williams GD. 2016. Genetic signatures of ecological diversity along an urbanization gradient. PeerJ 4:e2444
<https://doi.org/10.7717/peerj.2444>

Genetic signatures of ecological diversity along an urbanization gradient

Ryan P. Kelly^{Corresp., 1}, James L. O'Donnell¹, Natalie C. Lowell², Andrew O. Shelton³, Jameal F. Samhuri³, Shannon M. Hennessey⁴, Blake E. Feist³, Gregory D. Williams³

¹ School of Marine and Environmental Affairs, University of Washington, Seattle, Washington, United States of America

² School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, United States of America

³ Northwest Fisheries Science Center, NOAA Fisheries, Seattle, Washington, United States of America

⁴ Department of Integrative Biology, Oregon State University, Corvallis, Oregon, United States of America

Corresponding Author: Ryan P. Kelly

Email address: rpkelly@uw.edu

Despite decades of work in environmental science and ecology, estimating human influences on ecosystems remains challenging. This is partly due to complex chains of causation among ecosystem elements, exacerbated by the difficulty of collecting biological data at sufficient spatial, temporal, and taxonomic scales. Here, we demonstrate the utility of environmental DNA (eDNA) for quantifying associations between human land use and changes in an adjacent ecosystem. We analyze metazoan eDNA sequences from water sampled in nearshore marine eelgrass communities and assess the relationship between these ecological communities and the degree of urbanization in the surrounding watershed. Counter to conventional wisdom, we find strongly increasing richness and decreasing beta diversity with greater urbanization, and similar trends in the diversity of life histories with urbanization. We also find evidence that urbanization influences nearshore communities at local (hundreds of meters) rather than regional (tens of km) scales. Given that different survey methods sample different components of an ecosystem, we then discuss the advantages of eDNA—which we use here to detect hundreds of taxa simultaneously—as a complement to traditional ecological sampling, particularly in the context of broad ecological assessments where exhaustive manual sampling is impractical. Genetic data are a powerful means of uncovering human-ecosystem interactions that might otherwise remain hidden; nevertheless, no sampling method reveals the whole of a biological community.

1
2
3
4 **Genetic Signatures of Ecological Diversity Along an**
5 **Urbanization Gradient**
6

7 **Ryan P. Kelly*¹, James L. O'Donnell¹, Natalie C. Lowell², Andrew O.**
8 **Shelton³, Jameal F. Samhuri³, Shannon M. Hennessey⁴, Blake E. Feist³,**
9 **Gregory D. Williams³**

10 *¹School of Marine and Environmental Affairs, University of Washington, 3707 Brooklyn Ave NE, Seattle,*
11 *Washington 98105, USA*

12 *²School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle, Washington*
13 *98105, USA*

14 *³NOAA Fisheries, Northwest Fisheries Science Center, 2725 Montlake Blvd E, Seattle, Washington 98112, USA*

15 *⁴Department of Integrative Biology, Oregon State University, 3029 Cordley Hall, Corvallis, Oregon 97331,*
16 *USA*

17
18
19 **Corresponding author: rpkelley@uw.edu*
20
21

22 Introduction

23 An enduring question of environmental science and ecology is how to measure the effects of human
24 activities on nearby biological communities and ecosystems. While in some cases such impacts are so
25 obvious that in-depth sampling is unnecessary to reveal them—such as paving over a wetland or
26 clear-cutting a rainforest—many human activities are likely to have more subtle effects on the
27 surrounding system. More adequately measuring human impacts is a core challenge as human
28 demands on natural resources continue to grow; such measurement is a prerequisite for identifying
29 sustainable development pathways.

30
31 The difficulty of surveying ecological communities generally results in a depth-vs.-breadth (i.e.,
32 specificity, (Rice & Rochet, 2005)) tradeoff in sampling strategy. For example, one might
33 comprehensively survey indicator taxa with the idea that they reflect larger changes to the ecological
34 community (Niemi & McDonald, 2004), or instead build limited data from many taxa into
35 multimetric indices in an attempt to reflect some more holistic sense of ecosystem integrity (Karr,
36 1981; Weisberg et al., 1997). Environmental DNA (eDNA) could substantially improve upon
37 existing survey methods by mitigating this tradeoff (Karr, 1981; Weisberg et al., 1997) by providing
38 in-depth views of ecosystems at levels of effort comparable to traditional sampling. Indeed, microbial
39 ecology has used these same core techniques for a decade or more (Tyson et al., 2004; Venter et al.,
40 2004; Yutin et al., 2007). Sequencing the diagnostic traces of genetic material in environmental
41 samples makes it possible to detect hundreds or thousands of animals, plants, and other organisms
42 from target habitats on ecological time scales of hours to days (Thomsen et al., 2012; Turner et al.,
43 2014). Yet although the rapid rise of eDNA as a tool for ecological studies has featured
44 methodological leaps and assessments of performance (Thomsen et al., 2012; Ficetola et al., 2014;
45 Thomsen & Willerslev, 2015; Evans et al., 2016), the value that community-level eDNA methods
46 add to traditional ecological sampling is just beginning to be apparent.

47
48 Measuring the influence of urban development on surrounding ecosystems is one application for
49 which the broad scope of eDNA sampling may be particularly useful, in part because of the many
50 pathways through which correlates of urbanization are likely to influence nearby ecological
51 communities. Accordingly, it may be difficult to identify diffuse urban impacts using traditional
52 ecological sampling alone, a particularly pressing problem as coastal urbanization increases globally
53 (Neumann et al., 2015). For example, in Puget Sound, Washington, USA, as in many coastal areas,
54 homeowners modify or harden their shorelines with concrete or other materials to protect their
55 properties from erosion (Scyphers, Picou & Powers, 2015). Permitting for shoreline armoring can
56 create conflicts between individual property rights and the communal benefits that arise from
57 unarmored shoreline, which include storm- and flood mitigation, habitat, waterline access, and other
58 services. Laborious manual sampling has documented some shifts in ecology as a result of shoreline
59 armoring (Heerhartz et al., 2014), but the ability to detect the ecosystem effects of any stressor
60 depends strongly upon the choice of taxa sampled. Making such informed decisions about the scope
61 of sampling is a general problem in ecology and environmental sciences.

62
63 We assessed the effects of upland watershed urbanization on nearshore estuarine eelgrass (*Zostera*
64 *marina*) communities in Puget Sound, Washington, USA using eDNA sampling at four pairs of more-
65 and less-urban sites (Fig. 1). Puget Sound has experienced rapid urbanization over the past century,
66 its human population increasing nearly six-fold since 1920 (Minnesota Population Center, 2011), and
67 nearly 4 million people live within 20 km of its shore (Bright et al., 2012). Although preserving

68 biogenic eelgrass habitat is now a policy priority for state and federal agencies (Puget Sound
69 Partnership, 2011; US Army Corps of Engineers, 2012), the effect of such urbanization on eelgrass-
70 associated fauna has been difficult to characterize with traditional sampling techniques (e.g., (Blake,
71 Duffy & Richardson, 2014) in Chesapeake Bay). As such, the steep urbanization gradient of Puget
72 Sound makes a compelling setting for evaluating eDNA as a means of detecting ecological
73 differences associated with human development. Here, we report significant changes in community
74 composition, diversity, and life-history composition associated with upland urbanization, as
75 measured by the genetic signatures of animals detected in the water.

76 77 Methods

78
79 We selected 8 sites in nearshore eelgrass habitats adjacent to watersheds along a gradient of
80 urbanization in Puget Sound, Washington, USA (Fig. 1). We employed a paired study design, in
81 which each more-urbanized site had a companion less-urbanized site at approximately the same
82 latitude (Fig. 1), controlling for well-known geographic, oceanographic, and ecological gradients
83 within the Sound (Dethier, 2010). These were a subset of the sampling sites described in (Samhuri
84 et al.), and included Big Gulch Creek (BG), Clearwater Casino (CC), Clinton-Whidbey (CW),
85 Manchester (MA), Pipers Creek (PC), Redondo Beach Cold Creek (RB), Sinclair Inlet (SI), and
86 Shingle Mill Creek (SM). Further site details and coordinates are given in (Samhuri et al.).

87 88 **Environmental Setting**

89 We chose sites on the basis of watershed-scale patterns of urbanization as further described in
90 (Samhuri et al.). All watershed basins were less than 1,000 ha, and contained perennial streams
91 (Puget Sound Nearshore Ecosystem Restoration Project, 2010). We used three different geospatial
92 data layers that captured various aspects of terrestrial urbanization—imperviousness (Fry et al.,
93 2011), roadways (OpenStreetMap, 2013), and percent developed land cover (NOAA, 2013)—as well
94 as percent shoreline armoring (Puget Sound Nearshore Ecosystem Restoration Project, 2010), to
95 characterize urbanization at each site. Each of these individual metrics positively covaried and
96 ordination techniques did not result in an index that was significantly more useful than any one
97 urbanization variable alone. We therefore simply used imperviousness (the area-weighted mean
98 percent cover of impervious surface) here as a proxy for human population and other urbanization-
99 related parameters. This layer represents highly- to completely impermeable surfaces such as
100 building roofs, concrete or asphalt roads and parking lots, concrete, asphalt or brick sidewalks,
101 pedestrian walkways, and malls. We used Environmental Systems Research Institute's (Esri) ArcGIS
102 software suite (v. 10.1) for all spatial analyses. Within site pairs, more-urban sites had higher values
103 of imperviousness than their less-urban counterparts. Other environmental variables such as sea-
104 surface temperature (mean, max, SD) and salinity did not systematically vary with urbanization
105 across our sites.

106 107 **eDNA Collection, Extraction, and Sequencing**

108 In July 2014, we collected 1-liter water samples for eDNA analysis at each of three transects within
109 each site, and kept these on ice until they could be processed in the lab (within hours of collection).
110 We filtered samples onto cellulose acetate filters (47mm diameter; 0.45um pore size) under vacuum
111 pressure, and preserved the filter at room temperature in Longmire's buffer following Renshaw et al.
112 (Renshaw et al., 2015). Deionized water (1-liter) served as a negative control for filtering. We
113 extracted total DNA from the filters using the phenol:chloroform:isoamyl alcohol protocol in
114 (Renshaw et al., 2015), resuspended the eluate in 200uL water, and used 1uL of diluted DNA extract

115 (1:100, diluted to reduce amplification inhibition) as template for PCR. Total DNA recovered from
116 samples (quantified using a Qubit fluorometer) was uncorrelated with site urbanization, indicating
117 our results were not due to an accumulation of eDNA in environments near urban sites. See
118 Supplementary Methods for additional sampling details.

119
120 We designed a novel set of primers using ecoPrimers (Riaz et al., 2011) to amplify approximately
121 114-140bp of mitochondrial 16S DNA from metazoans exclusively. These primers effectively
122 amplify most major animal phyla—including representatives from Chordata, Arthropoda, Mollusca,
123 Echinodermata, Nemertea, and others—while excluding non-metazoans entirely. Their sequences are
124 as follows (5' to 3'): 16s_Metazoa_fwd AGTTACYYTAGGGATAACAGCG; 16s_Metazoa_rev
125 CCGGTCTGAACTCAGATCAYGT.

126
127 We generated amplicons using a two-step PCR procedure, described in (O'Donnell James L. et al.,
128 2016), to avoid the taxon-specific amplification bias that results from the use of differentially indexed
129 PCR primers (commonly used to include multiple samples onto the same high-throughput sequencing
130 run to minimize costs). The specific PCR protocol is included in Supplementary Methods.

131
132 Each of the 24 environmental samples (3 samples/site, 8 sites) was amplified in a total of four PCR
133 reactions, twice with each of two distinct indexed primer sets (see Supplementary Methods for
134 indexing details), for a total of $24 \times 4 = 96$ individual sets of amplicons for sequencing. All but one of
135 the environmental samples (from site CW) was sequenced successfully. We also sequenced four
136 positive (*Tilapia*; *Oreochromis niloticus* tissue) and three negative controls, treated the same way
137 (twice with each of two indexed primers, for a total of 16 replicates of positive controls and 12
138 replicates of negative controls). Using tissue-derived DNA as a positive control allowed us to assess
139 non-amplifications as deriving from sample-specific, rather than PCR-condition-specific causes, and
140 selecting a non-native species as the tissue source allowed us to identify putative cross-contamination
141 among samples (all *Tilapia* sequences should derive from the laboratory rather than the field). 150bp
142 paired-end sequencing was carried out on an Illumina Nextseq.

143 144 **Sequence Processing and Bioinformatics**

145 We processed the Nextseq reads with a custom Unix-based script (O'Donnell, 2015), which calls
146 existing third-party scripts to move from raw sequence data to a quality-controlled dataset of
147 operational taxonomic units (OTUs). See Supplementary Methods for further bioinformatics details.

148 149 **Contamination Removal and Sequencing-Depth Normalization**

150 We used a Bayesian site-occupancy modeling method to estimate the probability of the OTU
151 representing a true positive detection (Ficetola et al., 2014; Lahoz-Monfort, Guillera-Arroita &
152 Tingley, 2015), fitting a binomial distribution to OTU occurrences across replicates of each
153 environmental sample, and rarefied OTUs in each sample using the smallest number of reads we
154 observed in a single sample (124,041 reads; (Gotelli & Colwell, 2001)) to standardize estimates of
155 taxon richness across samples. We generated 1000 rarefied datasets, and unless otherwise specified
156 below, we report results from one representative rarefied dataset consisting of 11.8×10^6 reads
157 representing 1664 unique OTUs. The results do not depend significantly on the choice of rarefaction
158 replicates; for example, replicates differed only trivially in OTU richness (mean = 1662, sd = 9.5)
159 and did not show different spatial trends among replicates. For beta and gamma diversity measures,
160 in particular, OTU identity is of importance, and accordingly we show data derived from the entire
161 set of rarefaction replicates. Finally, for each water sample, we then averaged across the four PCR

162 replicates to estimate the abundance of each OTU. The complete eDNA dataset and analytical scripts
163 are publicly available on Dryad (Accession: doi:10.5061/dryad.04tq4). See Supplementary
164 Methods for further sequence processing details.

165
166 Our results do not depend strongly on decontamination or normalization procedures. Analyses of raw
167 OTU data (with no decontamination or normalization), of only the most common 100 OTUs, and of
168 only the least-common 500 OTUs, all produce the same trends in the quality-controlled and
169 normalized data (Suppl. Fig. 1). Similarly, rarefaction replicates retain the same strong trends
170 observed in our representative single replicate (Suppl. Fig. 2).

171 172 **Taxonomic Annotation of eDNA Sequences**

173 We annotated the final set of OTU sequences using the command-line BLAST+ software (Camacho
174 et al., 2009), searching against the complete NCBI nucleotide database (as of 12 October, 2015), with
175 word size = 7 and up to 1000 hits per query sequence retained. Those with no hits at $e = 10^{-13}$ (<
176 ca. 85% identity) or better were treated as unannotated. Conflicting sequence annotations were
177 resolved using the last common ancestor algorithm implemented in MEGAN (Huson et al., 2011).
178 Disagreement among hits for a given OTU (i.e., where a single OTU is an equally good match to >1
179 taxon) was generally resolved at the level of taxonomic Family (83.2% of reads; Supp. Table 1).

180 181 **Data Analysis, Community Composition, and Diversity**

182 Although amplicon sequencing produces read counts that may contain valuable information about
183 target species abundances (Evans et al., 2016; Port et al., 2016) it remains difficult to interpret the
184 results of amplicon studies in the context of quantitative ecology because the precise relationship
185 between amplicon abundance and taxon abundance remains unknown and likely varies among taxa
186 (Evans et al., 2016). Accordingly, our analyses used presence/absence information derived from
187 sequence count data.

188
189 To assess the appropriateness of the spatial scale of sampling, we apportioned the observed
190 variance in ecological distance (Jaccard) among sites, among transects (within sites), and among PCR
191 replicates using a PERMANOVA. We calculated alpha diversity (= richness, or “density”, sensu
192 (Gotelli & Colwell, 2001)) at both the OTU level and at the level of taxonomic family, treating
193 individual transects as replicates within a geographic site. We calculated beta diversity (sensu
194 Whittaker 1960, a measure of faunal change) both among transects within sites and among sites
195 (using transect means within sites to calculate the latter), focusing on OTUs because of the loss of
196 resolution associated with incomplete taxonomic annotation. We used Raup-Crick dissimilarity
197 (Chase et al., 2011) to ensure the observed beta diversity trends were not strictly dependent upon
198 changes in alpha diversity. We then evaluated gamma diversity (richness across sites within a region)
199 by generating an accumulation curve for three sets of sites: more-urban (N = 4 sites), less-urban (N =
200 4 sites), and all sites (N = 8). We sampled each set of sites (with replacement) 25 times at each step in
201 the accumulation curve to capture the distribution of site-specific richness.

202
203 We evaluated the relationships between diversity metrics and urbanization using linear and
204 generalized linear regression, as well as mixed-effects models. Our data were nested, with three
205 transect samples per site, and with each site having a single imperviousness value. To avoid
206 pseudoreplication among transects, we used site means for linear and generalized linear regressions.
207 For the mixed-effects models, we considered imperviousness as a fixed covariate and both site pair
208 and site identity as a random intercept terms.

209
210 To approximate life-history diversity, we organized all OTUs for which a Family-level annotation
211 was possible and classified each according to the following natural history attributes: Category
212 (epifauna, infauna, demersal, pelagic, terrestrial); Habitat (terrestrial, freshwater, intertidal, subtidal);
213 and Mobility (motile, sessile) using available reference materials such as (Kozloff, 1983). In some
214 cases, Families included species with a range of classifications (e.g., Cardiidae are a bivalve family
215 which includes infaunal and epifaunal cockles found both intertidal and subtidal habitats, with a
216 range of motility); in such cases the Family was listed as having both attributes. In all, there were 19
217 unique life-history niches that combinations of these attributes described (e.g., “Sessile Intertidal
218 Epifauna”, etc.; Suppl. Table 2). We used these classifications to assess trends in the richness of these
219 life-history groups with respect to imperviousness, and in a principal components analysis to assess
220 differences in faunas among sites.

221
222 Finally, we used logistic regression and binomial tests to identify particular taxa, OTUs, and life-
223 history characteristics significantly associated with imperviousness. We conducted all analyses in R
224 v3.2.2 (R Core Team, 2015).

225 226 Results

227 Our representative rarefied eDNA (16s mtDNA) dataset recovered 1664 operational taxonomic units
228 (OTUs; mean of 1000 rarefaction replicates = 1662 OTUs \pm 9.5) from a wide array of taxa
229 characteristic of the Puget Sound estuarine environment, with 10 animal phyla represented across 27
230 Classes, 65 Orders, and 135 Families (Table 1). Detections included iconic groups such as
231 *Metacarcinus* (i.e., *Cancer*) crabs, birds of prey (Accipitridae), and marine mammals (Delphinidae),
232 with the bulk of unique OTUs reflecting molluscs (45.1%), chordates (20.2%), and arthropods
233 (15.9%). 92% of reads (70% of OTUs) could be annotated with high confidence ($e < 10^{-32}$). These
234 annotations included many animal taxa common to Puget Sound or the surrounding environment
235 (Table 1; see Supplemental Table 1 for full Family-level annotations).

236
237 The total variance in community-level ecological distance was attributable to differences among sites
238 (38.6%), among transects within sites (45.4%), or among PCR replicates of the same water samples
239 (15.9%; PERMANOVA with Jaccard distance, $p < 0.001$, 999 permutations, using OTU presence-
240 absence data). These results are consistent with earlier work in nearshore habitats (Port et al., 2016),
241 reflecting differences in eDNA profiles at spatial scales on the order of tens to hundreds of meters
242 (here, between transects separated by ca. 50-100m) and limited variability due to PCR and
243 sequencing processes. Ordination of OTU data shows transect samples largely, but not exclusively,
244 clustering within geographic sites (Suppl. Fig. 3).

245 246 OTU Diversity and Urbanization

247
248 OTU richness increased significantly with upland imperviousness (Fig. 2). Family-level richness
249 reflected the overall richness trends (Fig. 2). The results were highly robust to different
250 decontamination or normalization procedures (Suppl. Figs. 1 and 2).

251
252 Our paired sampling design controlled for potentially confounding geographically associated
253 differences among sites. We observe the same strong positive OTU richness correlation with
254 imperviousness in all 4 site pairs (Fig. 2), evidence that some aspect of urbanization—rather than
255 confounding spatial differences among site pairs—explains the observed pattern. A mixed-effects

256 model showed that imperviousness had a positive effect on richness after accounting for pair and site
257 identity ($p = 0.018$).

258
259 We calculated beta diversity (faunal turnover) at two different hierarchical scales: between sites and
260 among transects within sites. Consistent with the high level of heterogeneity we observed among
261 transects within sites, between-site beta diversity was uniformly high and did not differ for more- or
262 less-urban sites (Whittaker's beta (1960); Wilcoxon test, $p = 0.58$). Focusing on the individual
263 transects, however, revealed a strong decrease in within-site beta diversity with urbanization across
264 all four site pairs: communities became more homogeneous (transects within sites became more
265 similar) as watershed imperviousness increased (Fig. 2). Whittaker's beta (Whittaker, 1960)
266 decreased from a mean of 0.816 when imperviousness was less than 10% to a mean of 0.546 when
267 imperviousness was greater than 25% ($R^2 = 0.93$, $p = 8.4 \times 10^{-5}$). Raup-Crick dissimilarity among
268 transects showed a similar trend, indicating that the urbanization-associated trend in transect-to-
269 transect variation in eDNA composition was greater than expected due to changes in alpha diversity
270 alone.

271
272 Consistent with the trend in richness, more-urban sites had consistently higher gamma diversity than
273 less-urban sites, as reflected in the completely non-overlapping OTU accumulation curves in those
274 sets of sites (Fig 2). In total, more-urban sites had 1295 unique OTUs in 116 Families, while less-
275 urban sites had 790 OTUs from 80 Families, respectively.

276 277 **Life-History Diversity and Urbanization**

278
279 Assessing individual characteristics of habitat and mobility, taxa with differing natural history
280 characteristics were differentially associated with urbanization. For example, OTU richness tripled
281 with greater urbanization among sessile taxa ($p = 1.7 \times 10^{-5}$), but motile taxa increased only
282 nonsignificantly ($p = 0.054$). Similarly, OTU richness in intertidal ($p = 7.5 \times 10^{-6}$) and subtidal ($p =$
283 3×10^{-5}) taxa increased with imperviousness, terrestrial taxa showed no such trend ($p = 0.16$).

284
285 Community shifts among natural history types reflected richness changes by taxonomic groups. At
286 both the OTU- and Family level, eDNA richness increased with urbanization, most notably among
287 bivalves and gastropods (Suppl. Fig. 4). Family-level bivalve richness rose, for example, from an
288 average of 5 Families (37 OTUs) at $<10\%$ imperviousness to 7.4 Families (111 OTUs) at $>25\%$
289 imperviousness (Poisson GLM with log-link, $p < 0.01$ at family level, 10^{-16} at OTU level). Other
290 taxa showed a more gradual increase in richness with imperviousness (Suppl. Fig. 5), resulting in an
291 overall increase in the number of taxonomic Families. No abundant Family declined with
292 imperviousness.

293
294 Combining ecological characteristics into tri-variate life-history categories (e.g., “intertidal sessile
295 epifauna”) revealed 19 unique Family categories present. Life-history richness increased with
296 urbanization (Suppl. Fig. 6; $R^2 = 0.74$, $p = 0.006$), from a mean of 12.5 life histories per site in less-
297 urban sites to a mean of 14.7 in more-urban sites, due to the concomitant increase in taxon richness at
298 more-urban sites. Normalizing by the number of Families present at each site reveals a strong
299 decrease in occupied life-histories-per-taxon with urbanization, from a mean of 0.66 in less-urban
300 sites to 0.47 in more-urban sites (although the trend is nonsignificant; Suppl. Fig. 6; $R^2 = 0.38$, $p =$
301 0.1). Ordination of the life histories results in identifiable sites and urbanization categories (Suppl.
302 Fig. 7), similar to the ordination plot for OTUs.

303
304 Beyond community measures, we identified 46 individual OTUs—again dominated by bivalves (33
305 OTUs from 5 families)—that were positively correlated ($p < 0.01$; logistic regression) with upland
306 imperviousness. Gastropods (5; limpets), urchins or sand dollars (7; not classifiable to family level),
307 and one fish OTU comprised the remaining 13 OTUs. Conversely, a single OTU was negatively
308 correlated with imperviousness (a mytilid mussel OTU). Providing some direct indication of human
309 influence on the nearshore Puget Sound, human OTU richness increased significantly with
310 imperviousness ($p = 0.01$; Poisson GLM), as did richness in selected taxa cultivated commercially
311 (*Panopea*, $p = 5 \times 10^{-4}$; *Bos*, $p = 0.005$) or introduced taxa (*Mya*, $p = 5.9 \times 10^{-6}$).
312

313 Discussion

314 All organisms leave behind residual genetic signatures in their environments, which provide the
315 opportunity to explore patterns of diversity and community structure that may not be possible
316 otherwise. Here, we recovered these signatures from nearshore estuarine habitats along an urban
317 gradient, revealing strong trends in the diversity of animals and ecological roles present. While alpha
318 (site richness) and gamma (regional richness) diversity strongly increased with upland urbanization,
319 more-urban sites were significantly more homogeneous (within sites) than less-urban sites. Life-
320 history diversity largely paralleled these same trends, with a greater richness of ecological life
321 histories among taxa found in more urban areas, but greater redundancy in life-history niches among
322 these taxa. Taken together, our results suggest that more urbanized upland areas support larger suites
323 of species, with less compositional variation, in and around downstream eelgrass habitats. Further,
324 we find evidence that the mechanisms of land-sea interaction act at watershed scales, rather than at
325 the larger scale of Puget Sound. These results also substantiate the idea that eDNA can be a powerful
326 addition to traditional means of assessing human-ecosystem interactions.
327

328 Trends in Diversity and Ecological Function with Urbanization

329 Although dense urban areas do not necessarily decrease biodiversity in general (Ives et al., 2016) and
330 the effects of urbanization on species richness appear to be taxon- and spatial-scale-specific (Shochat
331 et al., 2006), the positive richness trend we see in Puget Sound 16s eDNA is nevertheless striking.
332 Several plausible mechanisms could explain the increase in 16s eDNA richness, although our study
333 design prevents us from assessing causation explicitly.
334

335
336 One likely explanation for the trend is the interaction between fauna sampled with eDNA and the
337 kinds of habitat that are more common near urban settlements. Our study design attempted to sample
338 identical habitats across all sites, however, there may be unobserved differences in habitats. For
339 example, our results may reflect an increase in availability of muddy habitats associated with
340 urbanization, and a concomitant increase in richness within those habitat patches.
341

342 A second plausible mechanism is that greater anthropogenic nutrient inputs into urban areas yields
343 greater productivity. Urbanization greatly increases total nitrogen fluxes into rivers and estuaries
344 (Rabalais et al., 2009; Mohamedali et al., 2011), and increased primary productivity, which may
345 result from such fertilization, is generally—but not strictly—associated with increased secondary
346 productivity (Leslie et al., 2005) and taxonomic diversity (Mittelbach et al., 2001; Whittaker &
347 Heegaard, 2003). However, Puget Sound, like many coastal systems, is dominated by marine derived
348 nutrients (Mackas & Harrison, 1997; Mohamedali et al., 2011), suggesting that any fertilization effect
349 from small watersheds such as those we focus on here is unimportant. Each of the urban sites we

350 sampled also has a wastewater treatment facility in the vicinity. However, all outflows from
351 treatment facilities occur in deep water offshore, far from our sampling areas, making any effect of
352 fertilization indirect at best. Wastewater treatment facilities could also increase richness by
353 concentrating genetic material originating elsewhere. However, although the increase in human
354 OTUs we observe is consistent with this hypothesis, the great majority of DNA recovered stems from
355 Puget Sound species rather than taxa likely to be dominant in human waste streams and none of our
356 results is driven by exogenous eDNA.

357
358 Intriguingly, as eDNA communities increased in richness with urbanization, they also became more
359 homogeneous. Others have found that increased subtidal sedimentation—associated with the kind of
360 low-energy environments we sampled here—tended to make rocky reef communities more similar to
361 one another (Balata, Piazzini & Benedetti-Cecchi, 2007), and nutrient enrichment can have the same
362 effect in lakes (Donohue et al., 2009). Our results are consistent with the idea that urbanization tends
363 to homogenize communities even though the total number of unique taxa may increase (Urban et al.,
364 2006; Piazzini & Balata, 2008). A similar effect is also associated with non-indigenous species
365 introductions (Rosenzweig, 2001), but non-indigenous species do not drive the trends we observe
366 here. Although a comprehensive list of native taxa is not available against which to compare our
367 results, the annotated Families are nearly all familiar native taxa from Puget Sound; moreover, the
368 trends we report are consistent across even small subsets of the data (Suppl. Figs 1 & 2), indicating
369 our results do not depend upon a small set of potentially non-indigenous taxa.

370
371 More generally, beta diversity can help disentangle the ecological forces behind community
372 assembly (Condit et al., 2002; Tuomisto, Ruokolainen & Yli-Halla, 2003; Dornelas, Connolly &
373 Hughes, 2006; Chase, 2007, 2010; Chase & Myers, 2011), by distinguishing niche-related
374 deterministic processes from stochastic ones. Our observations are consistent with the idea that that
375 deterministic, possibly niche-related, processes significantly influence Puget Sound nearshore
376 communities: transect-to-transect beta-diversity declined steadily with an environmental gradient of
377 urbanization independent of geographic space, and per-taxon life-history richness similarly declined
378 (albeit nonsignificantly) across this same environmental gradient.

379
380 We expect different ecological patterns to be apparent at different spatial scales, and conversely, the
381 scales of ecological patterns provide hints about the mechanisms driving those patterns (Levin,
382 1992). Given the site- and transect-level differences we observed, it seems likely that the mechanisms
383 mediating the human-ecosystem interactions in Puget Sound occur at the watershed scale (~100s of
384 meters), rather than at larger scales of urbanization (e.g., Puget Sound scale, 10s of km). Urbanization
385 does not appear to homogenize communities across sites; more-urban sites were just as different from
386 one another as less-urban sites were, and the gamma diversity accumulation curve indicated that
387 additional urbanized sites continued to feature new OTUs. The real differences associated with
388 urbanization occurred within sites, with more-urban sites being more homogeneous (i.e., smaller
389 differences among transects) than less-urban sites. In sum, we did not observe a generalized “urban”
390 fauna at urban sites. Instead, each urbanized site had a distinct ecological community, exhibiting
391 greater richness, lower spatial variability, and greater life-history redundancy than a similar less-
392 urban site, but without a shared, characteristic community.

393
394 Regardless of the precise mechanism, the eDNA data reveal a strong signal of land-sea interaction
395 (Samhuri & Levin, 2012). Especially in light of ever-increasing human population density in coastal

396 areas worldwide (Neumann et al., 2015), our results suggest that eDNA can be a powerful tool for
397 uncovering human-ecosystem interactions that might otherwise remain hidden.

398

399 **eDNA as an Emerging Tool for Ecological Analysis: Scale and Selectivity**

400

401 Ecology and related disciplines depend upon techniques to sample and describe communities,
402 ecosystems, and their properties. However, any one set of samples yields a necessarily biased view of
403 the world; ten different sampling methods can yield ten different results even with small numbers of
404 target taxa (Valentini et al., 2015). This selectivity is usually intentional—e.g., settlement plates are
405 designed to sample bryozoans rather than seals—but where unintentional, such selectivity can bias
406 results in ways that often remain unexplored (Baker et al., 2016).

407

408 The rise of eDNA sampling has led to studies comparing molecular techniques either to traditional
409 methods or to known communities. Single-taxon qPCR studies have compared favorably with
410 traditional surveys in terms of detection rates (Jerde et al., 2011; Takahara et al., 2012; Eichmiller,
411 Bajer & Sorensen, 2014a; Laramie, Pilliod & Goldberg, 2015), with sequence-based (i.e.,
412 metabarcoding) analyses proving more difficult to interpret relative to traditional sampling, in part
413 because of difficulty of comparing detection rates across methods (Cowart et al., 2015). eDNA is an
414 in-depth sampling technique that yields interesting and repeatable results; however, the absence of
415 eDNA detection does not imply absence of taxon of interest (Roussel et al., 2015). One eDNA locus,
416 or even several loci, will not reveal all of the taxa present in an area. Indeed, eDNA sampling with a
417 different genetic locus—or even a different set of primers at the same locus—would have yielded a
418 different suite of taxa (e.g., (Cowart et al., 2015)).

419

420 Consistent with earlier observations from a study of *Zostera* communities (Cowart et al., 2015), our
421 single eDNA locus failed to detect epifauna known from the sampled sites. Hippolytid and crangonid
422 shrimp, littorinid snails, idoteid isopods, and others were common in the field but absent from the
423 eDNA (Samhuri et al.), likely due to amplification bias and primer mismatches. Such performance
424 does not make eDNA inappropriate for biodiversity monitoring, but rather put sequenced-based
425 sampling in the company of every other sampling technique (Shelton et al., 2016). Because the “true”
426 community remains unknown (Shelton et al., 2016), it is impossible to evaluate error rate in an
427 absolute sense for any field-based method. Given that nearly all (>99%) of the taxa we detect here
428 are known from local waters or the surrounding area, our false positive rate for eDNA appears to be
429 very low. We suggest that community-level eDNA surveys be viewed in a light appropriate to any
430 new sampling technique: biased relative to some unknown true value, but significantly
431 complementing existing imperfect sampling techniques such as tow nets and other manual
432 collections.

433

434 Finally, our results suggest that eDNA recovers fine-scale differences in ecological communities,
435 such that transects tens of meters apart can be as different as transects kilometres apart. Nearly half
436 (45%) of the variance in ecological distance was due to differences between transects at the same
437 sampling site, consistent with the fine-grained spatial resolution reported by (Port et al., 2016) in
438 another nearshore eDNA amplicon study. This observation supports a growing sense that eDNA may
439 travel only limited distances away from its sources, depending upon the environmental context
440 (Eichmiller, Bajer & Sorensen, 2014a; Deiner & Altermatt, 2014; Laramie, Pilliod & Goldberg,
441 2015), and provide further evidence that eDNA variation at small spatial scales is more likely signal
442 than noise. Nevertheless, it is not obvious why eDNA might exhibit such variability on the order of

443 tens of meters (in this study and in others; (Eichmiller, Bajer & Sorensen, 2014b; Port et al., 2016)),
444 but simultaneously feature the genetic signatures of species that are not in the immediate vicinity.
445 Examples here include terrestrial and aquatic taxa, whose DNA must have travelled at least some
446 distance into the intertidal habitats sampled. One explanation is that—if genetic material is detectable
447 as a steady-state balance of generation, degradation, advection, and diffusion away from a point
448 source—such transportation is to be expected at low levels, even when the bulk of genetic material
449 remains close to its source. Consistent with this model, the great majority of taxa in our data are
450 marine, with non-marine taxa only at low levels (6% of reads including human DNA; 3% not
451 including human DNA; see Suppl. Table 1).

452

453 **Conclusion**

454

455 Sampling using eDNA sequencing offers a breadth of taxonomic coverage valuable for both basic
456 and applied ecology. Our results demonstrate the power of this technique for assessing human-
457 ecosystem interactions in a nearshore environment, revealing significant trends in animal diversity
458 and life history likely linked to human alteration of upland habitats. Like all sampling methods,
459 eDNA offers a view of the world that is both biased and incomplete, in the sense that surveys using a
460 given gene will detect some taxa and not others. Traditional sampling has analogous drawbacks.
461 Here, data from a single genetic locus provided a reasonably holistic view of the Puget Sound
462 nearshore ecosystem—encompassing taxa as diverse as high-intertidal barnacles, birds of prey, and
463 subtidal bivalves, from a wide variety of ecologically-linked habitats—that strongly suggests
464 urbanization has generated unexpected consequences for a large number of nearshore taxa,
465 particularly those with sessile lifestyles. Consistent with (Samhuri et al.; Blake, Duffy &
466 Richardson, 2014; Ives et al., 2016), we see these results as a counterexample to the idea that humans
467 uniformly decrease biodiversity. Rather, the observation that more urbanized areas support larger, but
468 more homogeneous, suites of species indicates a more nuanced effect of human alteration on
469 nearshore communities.

470

471 **Additional Information**

472 **Acknowledgments**

473 We thank J.Port, L. Sassoubre, and A. Boehm; A. Stier, and P. Levin; M. Dethier, E. Heery, J. Toft, and J. Cordell; R.
474 Morris and V. Armbrust; J. Kralj; A. Wong, E. Garrison, J. Levy, M. Klein, and E. Buckner; coastal property owners for
475 access to field sites; and the Helen R. Whiteley Center at Friday Harbor Laboratorie, and two anonymous reviewers.

476

477 **Data Accessibility**

478 The article's supporting data, metadata, and analytical code can be accessed at Dryad (accession:
479 doi:10.5061/dryad.04tq4) and Genbank (accession number pending).

480

481

482

483

484

485

486

487

488 **Figure captions**

489

490 **FIGURE 1**

491 Study site sampling locations and associated stream basins. Matched site pairs share a stream basin color.
 492 More urban sites are open boxes, less urban are black boxes. Two-letter codes correspond to site names in
 493 the Methods. Brown shading indicates areas with greater than 50% imperviousness.

494 495 FIGURE 2

496 **Top Row:** Rarefied OTU richness and imperviousness—a proxy for urbanization—in Puget Sound. Analysis of
 497 a single focal rarefaction draw. **Left:** Rarefied 16s eDNA richness (solid trendline reflects OTUs; dashed
 498 trendline reflects taxonomic Families). Site means (larger circles) among transect-level data points (smaller
 499 circles). Family data shifted slightly for clarity. **Right:** The same data by site pair ($N = 4$ pairs of more- and
 500 less-urban sites), means plotted. Red lines indicate significant trends, grey lines indicate non-significant
 501 trends. Legends correspond to 2-letter site codes in Fig. 1. **Middle Row: Left:** Mean among-transect (within-
 502 site) Whittaker's beta diversity for each of 1000 rarefaction draws from the overall OTU dataset, rarefied to
 503 create comparable sample sizes ($N = 1.3 \times 10^5$ OTUs per transect). Linear regression on site means,
 504 $R^2 = 0.95$, $p = 3.38 \times 10^{-5}$. **Right:** Site means highlight the site-pair trends for single focal rarefaction draw.
 505 **Bottom Row:** Regional (gamma) diversity, in OTUs-per-site, as an accumulation curve. Boxplots show
 506 variance due to sampling each set of sites (with replacement) 1000 times from a pool of 1000 rarefaction
 507 draws from the overall OTU dataset, rarefied to create comparable sample sizes ($N = 1.3 \times 10^5$ OTUs per
 508 transect). Best-fit logarithmic curves shown for more-urban sites ($N = 4$), less-urban sites ($N = 4$), and all sites
 509 ($N = 8$).

510
 511
 512

513 References

- 515
 516 Baker DGL., Eddy TD., Mclver R., Schmidt AL., Thériault M-H., Boudreau M., Courtenay SC., Lotze HK.
 517 2016. Comparative analysis of different survey methods for monitoring fish assemblages in coastal
 518 habitats. *PeerJ* 4:e1832. DOI: 10.7717/peerj.1832.
 519 Balata D., Piazzini L., Benedetti-Cecchi L. 2007. Sediment disturbance and loss of beta diversity on
 520 subtidal rocky reefs. *Ecology* 88:2455–2461.
 521 Blake RE., Duffy JE., Richardson JP. 2014. Patterns of seagrass community response to local shoreline
 522 development. *Estuaries and Coasts* 37:1549–1561.
 523 Bright EA., Coleman PR., Rose AN., Urban ML. 2012. LandScan 2011. *Digital dataset, Oakridge National*
 524 *Laboratory, Oakridge, TN, USA, web. ornl.gov/sci/landscan/index.shtml.*
 525 Camacho C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., Madden TL. 2009. BLAST+:
 526 architecture and applications. *BMC Bioinformatics* 10:421.
 527 Chase JM. 2007. Drought mediates the importance of stochastic community assembly. *Proceedings of*
 528 *the National Academy of Sciences of the United States of America* 104:17430–17434. DOI:
 529 10.1073/pnas.0704350104.
 530 Chase JM. 2010. Stochastic community assembly causes higher biodiversity in more productive
 531 environments. *Science (New York, N.Y.)* 328:1388–1391. DOI: 10.1126/science.1187820.
 532 Chase JM., Kraft NJB., Smith KG., Vellend M., Inouye BD. 2011. Using null models to disentangle
 533 variation in community dissimilarity from variation in α -diversity. *Ecosphere* 2:art24.
 534 Chase JM., Myers JA. 2011. Disentangling the importance of ecological niches from stochastic processes
 535 across scales. *Philosophical transactions of the Royal Society of London. Series B, Biological*
 536 *sciences* 366:2351–2363. DOI: 10.1098/rstb.2011.0063.
 537 Condit R., Pitman N., Leigh EG., Chave J., Terborgh J., Foster RB., Núñez P., Aguilar S., Valencia R.,
 538 Villa G., Muller-Landau HC., Losos E., Hubbell SP. 2002. Beta-diversity in tropical forest trees.
 539 *Science (New York, N.Y.)* 295:666–669. DOI: 10.1126/science.1066854.
 540 Cowart DA., Pinheiro M., Mouchel O., Maguer M., Grail J., Miné J., Arnaud-Haond S. 2015.
 541 Metabarcoding Is Powerful yet Still Blind: A Comparative Analysis of Morphological and Molecular
 542 Surveys of Seagrass Communities. *PLoS One* 10:e0117562.
 543 Deiner K., Altermatt F. 2014. Transport Distance of Invertebrate Environmental DNA in a Natural River.
 544 *PLoS One* 9:e88786. DOI: 10.1371/journal.pone.0088786.
 545 Dethier MN. 2010. Variation in recruitment does not drive the cline in diversity along an estuarine

- 546 gradient. *Marine Ecology Progress Series* 410:43–54.
- 547 Donohue I., Jackson AL., Pusch MT., Irvine K. 2009. Nutrient enrichment homogenizes lake benthic
548 assemblages at local and regional scales. *Ecology* 90:3470–3477.
- 549 Dornelas M., Connolly SR., Hughes TP. 2006. Coral reef diversity refutes the neutral theory of
550 biodiversity. *Nature* 440:80–82. DOI: 10.1038/nature04534.
- 551 Eichmiller JJ., Bajer PG., Sorensen PW. 2014a. The Relationship between the Distribution of Common
552 Carp and Their Environmental DNA in a Small Lake. *PLoS One* 9:e112611.
- 553 Eichmiller JJ., Bajer PG., Sorensen PW. 2014b. The relationship between the distribution of common carp
554 and their environmental DNA in a small lake. *PLoS one* 9:e112611. DOI:
555 10.1371/journal.pone.0112611.
- 556 Evans NT., Olds BP., Turner CR., Renshaw MA., Li Y., Jerde CL., Mahon AR., Pfrender ME., Lamberti
557 GA., Lodge DM. 2016. Quantification of mesocosm fish and amphibian species diversity via eDNA
558 metabarcoding. *Molecular Ecology Resources* 16:25–41.
- 559 Ficetola GF., Pansu J., Bonin A., Coissac E., Giguët-Covex C., De Barba M., Gielly L., Lopes CM., Boyer
560 F., Pompanon F., Others. 2014. Replication levels, false presences and the estimation of the
561 presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources* 15:543–556.
- 562 Fry JA., Xian G., Jin S., Dewitz JA., Homer CG., LIMIN Y., Barnes CA., Herold ND., Wickham JD. 2011.
563 Completion of the 2006 national land cover database for the conterminous United States.
564 *Photogrammetric engineering and remote sensing* 77:858–864.
- 565 Gotelli NJ., Colwell RK. 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and
566 comparison of species richness. *Ecology Letters* 4:379–391.
- 567 Heerhartz SM., Dethier MN., Toft JD., Cordell JR., Ogston AS. 2014. Effects of shoreline armoring on
568 beach wrack subsidies to the nearshore ecotone in an estuarine fjord. *Estuaries and Coasts*
569 37:1256–1268.
- 570 Huson DH., Mitra S., Ruscheweyh H-J., Weber N., Schuster SC. 2011. Integrative analysis of
571 environmental sequences using MEGAN4. *Genome research* 21:1552–1560.
- 572 Ives CD., Lentini PE., Threlfall CG., Ikin K., Shanahan DF., Garrard GE., Bekessy SA., Fuller RA.,
573 Mumaw L., Rayner L., Others. 2016. Cities are hotspots for threatened species. *Global Ecology and*
574 *Biogeography* 25:117–126.
- 575 Jerde CL., Mahon AR., Chadderton WL., Lodge DM. 2011. “Sight-unseen” detection of rare aquatic
576 species using environmental DNA. *Conservation Letters* 4:150–157.
- 577 Karr JR. 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6:21–27.
- 578 Kozloff EN. 1983. *Seashore life of the northern Pacific coast: an illustrated guide to northern California,*
579 *Oregon, Washington, and British Columbia.* University of Washington Press Seattle.
- 580 Lahoz-Monfort JJ., Guillera-Arroita G., Tingley R. 2015. Statistical approaches to account for false
581 positive errors in environmental DNA samples. *Molecular Ecology Resources*.
- 582 Laramie MB., Pilliod DS., Goldberg CS. 2015. Characterizing the distribution of an endangered salmonid
583 using environmental DNA analysis. *Biological Conservation* 183:29–37.
- 584 Leslie HM., Breck EN., Chan F., Lubchenco J., Menge BA. 2005. Barnacle reproductive hotspots linked to
585 nearshore ocean conditions. *Proceedings of the National Academy of Sciences of the United States*
586 *of America* 102:10534–10539.
- 587 Levin SA. 1992. The problem of pattern and scale in ecology. *Ecology* 73.
- 588 Mackas DL., Harrison PJ. 1997. Nitrogenous Nutrient Sources and Sinks in the Juan de Fuca Strait/Strait
589 of Georgia/Puget Sound Estuarine System: Assessing the Potential for Eutrophication. *Estuarine,*
590 *Coastal and Shelf Science* 44:1–21. DOI: 10.1006/ecss.1996.0110.
- 591 Minnesota Population Center. 2011. National historical geographic information system: Version 2.0.
592 *Minneapolis, MN: University of Minnesota.*
- 593 Mittelbach GG., Steiner CF., Scheiner SM., Gross KL., Reynolds HL., Waide RB., Willig MR., Dodson SI.,
594 Gough L. 2001. What is the observed relationship between species richness and productivity?
595 *Ecology* 82:2381–2396.
- 596 Mohamedali T., Roberts M., Sackmann B., Kolosseus A. 2011. *Puget Sound Dissolved Oxygen Model*
597 *Nutrient Load Summary for 1999-2008.*
- 598 Neumann B., Vafeidis AT., Zimmermann J., Nicholls RJ. 2015. Future coastal population growth and
599 exposure to sea-level rise and coastal flooding—a global assessment. *PLoS one* 10:e0118571. DOI:
600 10.1371/journal.pone.0118571.
- 601 Niemi GJ., McDonald ME. 2004. Application of ecological indicators. *Annual Review of Ecology,*

- 602 *Evolution, and Systematics*:89–111.
- 603 NOAA. 2013. NOAA's Coastal Change Analysis Program (C-CAP) 2006 Regional Land Cover Data -
604 Coastal United States. *National Ocean Service (NOS), Office for Coastal Management (OCM)*.
- 605 O'Donnell JL. 2015.banzai. Available at <https://github.com/jimmyodonnell/banzai>
- 606 O'Donnell James L., Kelly Ryan P., Lowell Natalie., Port JA. 2016. Indexed PCR Primers Induce
607 Template-Specific Bias In Large-Scale DNA Sequencing Studies. *PLoSOne* in press.
- 608 OpenStreetMap. 2013. OpenStreetMap - Washington State Roads, OpenStreetMap contributors.
- 609 Piazzini L., Balata D. 2008. The spread of *Caulerpa racemosa* var. *cylindracea* in the Mediterranean Sea:
610 an example of how biological invasions can influence beta diversity. *Marine Environmental*
611 *Research* 65:50–61.
- 612 Port JA., O'Donnell JL., Lowell N., Romero-Maraccini O., Kelly RP. 2016. Assessing the Vertebrate
613 Community of a Kelp Forest Ecosystem Using Environmental DNA. *Molecular Ecology* 25:527–541.
- 614 Puget Sound Nearshore Ecosystem Restoration Project. 2010. Puget Sound Basin PSNERP Database.
- 615 Puget Sound Partnership. 2011. *Leadership Council Resolution 2011-01: Adopting an Ecosystem*
616 *Recovery Target for Eelgrass*.
- 617 R Core Team. 2015. R: A Language and Environment for Statistical Computing.
- 618 Rabalais NN., Turner RE., Diaz RJ., Justic D. 2009. Global change and eutrophication of coastal waters.
619 *ICES Journal of Marine Science* 66:1528–1537. DOI: 10.1093/icesjms/fsp047.
- 620 Renshaw MA., Olds BP., Jerde CL., McVeigh MM., Lodge DM. 2015. The room temperature preservation
621 of filtered environmental DNA samples and assimilation into a phenol--chloroform--isoamyl alcohol
622 DNA extraction. *Molecular Ecology Resources* 15:168–176.
- 623 Riaz T., Shehzad W., Viari A., Pompanon F., Taberlet P., Coissac E. 2011. ecoPrimers: inference of new
624 DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Research* 39:e145–
625 e145.
- 626 Rice JC., Rochet M-J. 2005. A framework for selecting a suite of indicators for fisheries management.
627 *ICES Journal of Marine Science* 62:516–527.
- 628 Rosenzweig ML. 2001. The four questions: What does the introduction of exotic species do to diversity?
629 *Evolutionary Ecology Research* 3:361–367.
- 630 Roussel J-M., Paillisson J-M., Treguier A., Petit E. 2015. The downside of eDNA as a survey tool in water
631 bodies. *Journal of Applied Ecology* 52:823–826.
- 632 Samhuri JF., Shelton AO., Williams GD., Feist B., Hennessey S., Bartz K., O'Donnell JL., Sheer M.,
633 Levin PS. How much city is too much city? Biodiversity and ecosystem functions along an urban
634 gradient at the land-sea interface in Puget Sound. *Journal of Applied Ecology* In Revision.
- 635 Samhuri JF., Levin PS. 2012. Linking land-and sea-based activities to risk in coastal ecosystems.
636 *Biological Conservation* 145:118–129.
- 637 Scyphers SB., Picou JS., Powers SP. 2015. Participatory Conservation of Coastal Habitats: The
638 Importance of Understanding Homeowner Decision Making to Mitigate Cascading Shoreline
639 Degradation. *Conservation Letters* 8:41–49. DOI: 10.1111/conl.12114.
- 640 Shelton AO., O'Donnell JL., Samhuri JF., Lowell N., Williams GD., Kelly RP. 2016. A framework for
641 inferring biological communities from environmental DNA. *Ecological Applications* in press.
- 642 Shochat E., Warren PS., Faeth SH., McIntyre NE., Hope D. 2006. From patterns to emerging processes
643 in mechanistic urban ecology. *Trends in ecology & evolution* 21:186–191.
- 644 Takahara T., Minamoto T., Yamanaka H., Doi H., Kawabata Z. 2012. Estimation of fish biomass using
645 environmental DNA. *PloS One* 7:e35868.
- 646 Thomsen PF., Kielgast J., Iversen LL., Møller PR., Rasmussen M., Willerslev E. 2012. Detection of a
647 diverse marine fish fauna using environmental DNA from seawater samples. *PLoS One* 7:e41732.
- 648 Thomsen PF., Willerslev E. 2015. Environmental DNA—an emerging tool in conservation for monitoring
649 past and present biodiversity. *Biological Conservation* 183:4–18.
- 650 Tuomisto H., Ruokolainen K., Yli-Halla M. 2003. Dispersal, environment, and floristic variation of western
651 Amazonian forests. *Science (New York, N.Y.)* 299:241–244. DOI: 10.1126/science.1078037.
- 652 Turner CR., Barnes MA., Xu CCY., Jones SE., Jerde CL., Lodge DM. 2014. Particle size distribution and
653 optimal capture of aqueous microbial eDNA. *Methods in Ecology and Evolution* 5:676–684.
- 654 Tyson GW., Chapman J., Hugenholtz P., Allen EE., Ram RJ., Richardson PM., Solovyev V V., Ruben
655 EM., Rokhsar DS., Banfield JF. 2004. Community structure and metabolism through reconstruction
656 of microbial genomes from the environment. *Nature* 428:37–43.
- 657 Urban MC., Skelly DK., Burchsted D., Price W., Lowry S. 2006. Stream communities across a rural--

- 658 urban landscape gradient. *Diversity and Distributions* 12:337–350.
- 659 US Army Corps of Engineers. 2012. *User's Guide For Nationwide Permits in Washington State*.
- 660 Valentini A., Taberlet P., Miaud C., Civade R., Herder J., Thomsen PF., Bellemain E., Besnard A.,
661 Coissac E., Boyer F., Others. 2015. Next-generation monitoring of aquatic biodiversity using
662 environmental DNA metabarcoding. *Molecular ecology*.
- 663 Venter JC., Remington K., Heidelberg JF., Halpern AL., Rusch D., Eisen JA., Wu D., Paulsen I., Nelson
664 KE., Nelson W. 2004. Environmental genome shotgun sequencing of the Sargasso Sea. *Science*
665 304:66–74.
- 666 Weisberg SB., Ranasinghe JA., Dauer DM., Schaffner LC., Diaz RJ., Frithsen JB. 1997. An estuarine
667 benthic index of biotic integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20:149–158.
- 668 Whittaker RH. 1960. Vegetation of the Siskiyou mountains, Oregon and California. *Ecological*
669 *Monographs* 30:279–338.
- 670 Whittaker RJ., Heegaard E. 2003. What is the observed relationship between species richness and
671 productivity? Comment. *Ecology* 84:3384–3390.
- 672 Yutin N., Suzuki MT., Teeling H., Weber M., Venter JC., Rusch DB., Béjà O. 2007. Assessing diversity
673 and biogeography of aerobic anoxygenic phototrophic bacteria in surface waters of the Atlantic and
674 Pacific Oceans using the Global Ocean Sampling expedition metagenomes. *Environmental*
675 *microbiology* 9:1464–1475.
- 676

Figure 1

Study site sampling locations and associated stream basins.

Study site sampling locations and associated stream basins. Matched site pairs share a stream basin color. More urban sites are open boxes, less urban are black boxes. Two-letter codes correspond to site names in the Methods. Brown shading indicates areas with greater than 50% imperviousness.




Figure 2 (on next page)

Alpha, beta, and gamma diversity recovered from water samples in Puget Sound along an urbanization gradient.

Top Row: Rarefied OTU richness and imperviousness—a proxy for urbanization—in Puget Sound. Analysis of a single focal rarefaction draw. **Left** : Rarefied 16s eDNA richness (solid trendline reflects OTUs; dashed trendline reflects taxonomic Families). Site means (larger circles) among transect-level data points (smaller circles). Family data shifted slightly for clarity. **Right**: The same data by site pair ($N = 4$ pairs of more- and less-urban sites), means plotted. Red lines indicate significant trends, grey lines indicate non-significant trends. Legends correspond to 2-letter site codes in Fig. 1. **Middle Row: Left**: Mean among-transect (within-site) Whittaker's beta diversity for each of 1000 rarefaction draws from the overall OTU dataset, rarefied to create comparable sample sizes ($N = 1.3 \times 10^5$ OTUs per transect). Linear regression on site means, $R^2 = 0.95$, $p = 3.38 \times 10^{-5}$. **Right**: Site means highlight the site-pair trends for single focal rarefaction draw. **Bottom Row**: Regional (gamma) diversity, in OTUs-per-site, as an accumulation curve. Boxplots show variance due to sampling each set of sites (with replacement) 1000 times from a pool of 1000 rarefaction draws from the overall OTU dataset, rarefied to create comparable sample sizes ($N = 1.3 \times 10^5$ OTUs per transect). Best-fit logarithmic curves shown for more-urban sites ($N = 4$), less-urban sites ($N = 4$), and all sites ($N = 8$).








Table 1 (on next page)

Summary of 16s Read Annotations

Summary of taxonomic annotations for 16S reads; for full annotations, see Supplementary Material.

1 Tables

PHYLUM	CLASSE S	ORDERS	FAMILIES	OTHER RANK S
MOLLUSCA	3	6	34	9
ARTHROPODA	6	13	29	10
CHORDATA	5	28	37	28
BRYOZOA	1	2	10	2
ECHINODERMATA	5	8	13	4
NEMERTEA	3	2	8	0
HEMICHORDATA	1	1	1	0
ENTOPROCTA	1	1	1	0
PORIFERA	1	2	2	0

2

3 TABLE1

4 Summary of taxonomic annotations for 16S reads; for full annotations, see Supplementary Material.

5