

**A peer-reviewed version of this preprint was published in PeerJ on 13 October 2016.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.2552) (peerj.com/articles/2552), which is the preferred citable publication unless you specifically need to cite this preprint.

Ong OTW, Young LJ, Old JM. 2016. Evaluation of reference genes for gene expression in red-tailed phascogale (*Phascogale calura*) liver, lung, small intestine and spleen. PeerJ 4:e2552  
<https://doi.org/10.7717/peerj.2552>

# Evaluation of reference genes for gene expression in red-tailed phascogale (*Phascogale calura*) liver, lung, small intestine and spleen

Oselyne TW Ong<sup>1</sup>, Lauren J Young<sup>1</sup>, Julie M Old<sup>Corresp.</sup><sup>1</sup>

<sup>1</sup> School of Science and Health, Western Sydney University, Richmond, New South Wales, Australia

Corresponding Author: Julie M Old

Email address: j.old@westernsydney.edu.au

**Background:** Reference genes serve an important role as an endogenous control/standard for data normalisation in gene expression studies. Although reference genes have recently been suggested for marsupials, independent analysis of reference genes on different immune tissues is yet to be tested. Therefore, an assessment of reference genes is needed for the selection of stable, expressed genes across different marsupial tissues.

**Methods:** The study was conducted on red-tailed phascogales (*Phascogale calura*) using five juvenile and five adult males. The stability of five reference genes (glyceraldehyde-3-phosphate dehydrogenase, *GAPDH*;  $\beta$ -Actin, *ACTB*; 18S rRNA, *18S*; 28S rRNA, *28S*; and ribosomal protein L13A, *RPL13A*) was investigated using SYBR Green and analysed with the geNorm application available in qBase<sup>PLUS</sup> software.

**Results:** Gene stability for juvenile and adult tissue samples combined show that *GAPDH* was most stable in liver and lung tissue, and *18S* in small intestine and spleen. While all reference genes were suitable for small intestine and spleen tissues, all reference genes except *28S* were stable for lung and only *18S* and *28S* were stable for liver tissue. Separating the two age groups, we found that two different reference genes were considered stable in juveniles (*ACTB* and *GAPDH*) and adults (*18S* and *28S*), and *RPL13A* was not stable for juvenile small intestine tissue. Except for *28S*, all reference genes were stable in juvenile and adult lungs, and all five reference genes were stable in spleen tissue.

**Discussion:** Based on expression stability, *ACTB* and *GAPDH* are suitable for all tissues when studying the expression of marsupials in two age groups, except for adult liver tissues. The expression stability between juvenile and adult liver tissue was most unstable, as the stable reference genes for juveniles and adults were different. Juvenile and adult lung, small intestine and spleen share similar stable reference genes, except for small intestine tissues where all reference genes were stable in adults but *RPL13A* was not suitable in juveniles.

Evaluation of reference genes for gene expression in red-tailed phascogale (*Phascogale calura*) liver,  
lung, small intestine and spleen

Oselyne T. W. Ong<sup>1</sup>, Lauren J. Young<sup>1</sup>, Julie M. Old<sup>1\*</sup>

<sup>1</sup>School of Science and Health, Western Sydney University, Penrith, NSW 2751, Australia

Ms Oselyne T. W. Ong

E: [o.ong@westernsydney.edu.au](mailto:o.ong@westernsydney.edu.au)

Dr Lauren Young

E: [ljmbyoung@gmail.com](mailto:ljmbyoung@gmail.com)

\*Corresponding author:

Associate Professor Julie Old

School of Science and Health

Western Sydney University,

Hawkesbury Campus

Locked Bag 1797,

Penrith NSW 2751

P: +61 2 4570 1283

E: [j.old@westernsydney.edu.au](mailto:j.old@westernsydney.edu.au)

*Running title:* Red-tailed phascogale qPCR expression

## Abstract

**Background:** Reference genes serve an important role as an endogenous control/standard for data normalisation in gene expression studies. Although reference genes have recently been suggested for marsupials, independent analysis of reference genes on different immune tissues is yet to be tested. Therefore, an assessment of reference genes is needed for the selection of stable, expressed genes across different marsupial tissues.

**Methods:** The study was conducted on red-tailed phascogales (*Phascogale calura*) using five juvenile and five adult males. The stability of five reference genes (glyceraldehyde-3-phosphate dehydrogenase, *GAPDH*;  $\beta$ -Actin, *ACTB*; 18S rRNA, 18S; 28S rRNA, 28S; and ribosomal protein L13A, *RPL13A*) was investigated using SYBR Green and analysed with the geNorm application available in qBase<sup>PLUS</sup> software.

**Results:** Gene stability for juvenile and adult tissue samples combined show that *GAPDH* was most stable in liver and lung tissue, and 18S in small intestine and spleen. While all reference genes were suitable for small intestine and spleen tissues, all reference genes except 28S were stable for lung and only 18S and 28S were stable for liver tissue. Separating the two age groups, we found that two different reference genes were considered stable in juveniles (*ACTB* and *GAPDH*) and adults (18S and 28S), and *RPL13A* was not stable for juvenile small intestine tissue. Except for 28S, all reference genes were stable in juvenile and adult lungs, and all five reference genes were stable in spleen tissue.

**Discussion:** Based on expression stability, *ACTB* and *GAPDH* are suitable for all tissues when studying the expression of marsupials in two age groups, except for adult liver tissues. The expression stability between juvenile and adult liver tissue was most unstable, as the stable reference genes for juveniles and adults were different. Juvenile and adult lung, small intestine and spleen share similar stable reference genes, except for small intestine tissues where all reference genes were stable in adults but *RPL13A* was not suitable in juveniles.

## INTRODUCTION

Fluorescence-based quantitative real-time polymerase chain reaction (qPCR) has the capacity to monitor the amplification of cDNA during thermocycling, starting with the use of ethidium bromide for the detection of fluorescence (Higuchi et al. 1992). Over the past 13 years, qPCR has developed into the most accurate and sensitive method to study gene expression with low concentrations of mRNA (Bustin

2000; Schmittgen & Livak 2008). When studying the expression of a target gene, it is important to have a stable reference gene for the normalisation of gene expression (Pierzchała et al. 2011). Although it may seem scientifically sound to use reference genes that have been used in other similar studies, it has to be reported that common reference genes such as *GAPDH* and *ACTB* have various stabilities depending on the type of tissue or experimental conditions (Glare et al. 2002; Selvey et al. 2001; Zhong & Simons 1999). Commonly used rRNA genes, 18S and 28S are also affected by various biological factors (Warner 1999). At present, there has not been a study on the stability of reference genes in a marsupial. In most cases, reference genes found stable in other mammalian groups (particularly eutherians) have been used in studies without considering the stability of the reference genes, which may differ between mammal groups and tissues or cells.

Reference genes should not only be stable enough for normalisation, but the use of more than one reference gene is encouraged in expression studies as the use of only one reference gene has led to expression study errors (Bustin 2000; Vandesompele et al. 2002). By selecting the most stable reference genes for the type of tissue/cells or experimental condition, an accurate normalisation of qPCR can be performed, avoiding variation in reference gene expression in tissues/cells investigated (Vandesompele et al. 2002). Since the stability of reference genes have not been reported in a marsupial, gene expression studies using similar tissues/cells or marsupial species can use this study as a baseline, and therefore be used in future marsupial gene expression studies.

The aim of this study was to identify stable reference genes across a range of tissues of a marsupial species – the Red-tailed phascogale (*Phascogale calura*). Red-tailed phascogales are a model species as they are small and relatively easy to maintain in captivity (Foster et al. 2006; Russell 1982; Stannard et al. 2013). In the wild, the red-tailed phascogale inhabits a small corner of the south west of western Australia (Bradley et al. 2008) and are distinguished from other small marsupials by their long brush-like hairs on the end of their red tail (Kennedy & Williams 1990).

This study evaluates the expression stability of five different reference genes in four different tissues associated with immunity in the red-tailed phascogale; liver, lung, small intestine and spleen in two different age groups. Optimal reference genes should be considered stable and expressed at constant levels in various tissues and age groups. Liver and lung tissues were primarily chosen because

they contain large populations of macrophages and are in regular contact with pathogens (Laskin et al. 2001). In addition, unlike eutherian livers that cease haematopoiesis prior to birth, the marsupial liver is the main site of haematopoiesis during early postnatal life (reviewed by (Borthwick et al. 2014; Old & Deane 2000).

Lung and small intestine were chosen as representative mucosal-associated lymphoid tissue (MALT). The MALT in lungs is responsible for protection of the respiratory system (Mak et al. 2013), whilst the small intestine is an important gut-associated lymphoid tissue (GALT). In marsupials, GALT can be localised in Peyer's patches or follicular aggregations, or appear as scattered cells distributed throughout the gut (Old & Deane 2002). The last of the tissues chosen for use in this study was spleen. The spleen is an important haematopoietic site and actively involved in the adaptive immune response (reviewed in Borthwick et al. 2014; Old & Deane 2000). The four red-tailed phascogale tissues (liver, lung, small intestine and spleen) were therefore chosen based on their immunological capacity and function. The expression stability of five reference genes in these tissues was investigated using the geNorm application in the qBase<sup>PLUS</sup> software.

## **MATERIALS AND METHODS**

### **Animal and tissue collection**

Ten male red-tailed phascogales from two age groups (juveniles: 3.5 – 5 months, and adults: 1.2 – 1.5 years) were utilised in this study. Tissue samples were opportunistically obtained from a the Small Native Mammal Teaching and Research Facility, a captive colony housed at the Western Sydney University (WSU) (Richmond, NSW) as per standard operating procedures approved by the UWS Animal Ethics Committee (A9694) during population maintenance. Samples of liver, lung, small intestine and spleen were dissected, and immediately stored at -80°C until total RNA extraction.

### **RNA extraction and cDNA synthesis**

Total RNA was extracted using the SV Total RNA Isolation System (Promega, Wisconsin, USA) according to the manufacturer's protocol. The quantity and quality of total RNA was estimated using a Nanodrop 2000 Spectrophotometer (Thermo Scientific, Delaware, USA) with the OD260nm/OD280nm ratio expected to be between 1.8 and 2. One µg of total RNA was reverse transcribed with SuperScript®

III First-Strand Synthesis SuperMix (Invitrogen, California, USA) according to the manufacturer's protocol. The quantity of the final cDNA was assessed using a Nanodrop 2000 Spectrophotometer, and final cDNA products were aliquoted and stored at -20°C until use.

#### Primers and real-time PCR

We selected five reference genes used previously in marsupial gene expression studies (Maher et al. 2014; Markey et al. 2007; Yu et al. 2006): glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -actin (*ACTB*), 18S rRNA (18S), 28S rRNA (28S) and ribosomal protein L13a (*RPL13A*). Real-time PCR primers, *GAPDH* and *ACTB*, were designed using consensus sequences based on marsupial species obtained from GenBank (Tasmanian devil *Sarcophilus harrisii*) *GAPDH*: XM\_012550750.1; gray short-tailed opossum (*Monodelphis domestica*) *GAPDH*: XM\_007503905.1; Tasmanian devil *ACTB*: XM\_003761554.2). Primers for 18S and 28S were obtained from Daly et al. (2009), and *RPL13A* from Siddle et al. (2013). Specific information for each primer is listed in Table 1.

PCR amplification was performed using a Rotor Gene Q (Qiagen, Hilden, Germany) and the Rotor-Gene SYBR® Green PCR kit (Qiagen). A PCR mix (25  $\mu$ L) was prepared: 7.5  $\mu$ L water, 2.5  $\mu$ L primers (forward and reverse; 10  $\mu$ M), 1  $\mu$ L (100 ng) cDNA, and 12.5  $\mu$ L of Rotor-Gene SYBR Green PCR Master Mix. The following amplification program was used: 5 min denaturation at 95°C, 35 cycles of amplification with 5 s at 95°C (denaturation), 10 s at 60°C (annealing), and 15 s at 72°C (elongation). Annealing temperatures were optimised according to individual genes and primers by testing several annealing temperatures ranging from 55°C to 65°C around the respective primer  $T_m$ , and the annealing temperature with the best efficiency was chosen. A melting step was performed to confirm a single gene-specific peak by a stepwise temperature increase ranging from 60°C to 95°C at ramp rate 1°C/s with continuous monitoring of fluorescence. Further analysis of amplicon specificity and size were also evaluated running qPCR products in a standard 2% agarose gel electrophoresis. Standard curves were made to calculate the amplification efficiency during real-time PCR using five-fold serial dilutions of cDNA for each tissue and each reference gene in one adult male red-tailed phascogale. The quantification cycle ( $C_q$ ) was automatically determined for each reaction by the Rotor Gene Software (v. 1.7.94).

#### Data analysis

Gene expression variation was calculated for individual reference genes based on cycle threshold ( $C_q$ ) values and real-time PCR efficiencies ( $E$ ). The real-time  $E$  value was calculated from the given slopes in the qBase<sup>PLUS</sup> software (Hellemans et al. 2007) according to the equation: ( $E = 10^{(-1/\text{slope})-1}$ ). Only  $C_q$  values <40 were used for calculation of  $E$  values.  $C_q$  and  $E$  values were then analysed in geNorm on the qBase<sup>PLUS</sup> software, which ranks the reference genes based on the  $M$  values (reference genes with the lowest  $M$  value is considered most stable). A one-way ANOVA was also performed on  $C_q$  values obtained from the expression in juveniles and adults of the four reference genes.

## RESULTS

Five reference genes were amplified in four tissues, and all real-time PCR assays produced a single peak on the melting curve (*GAPDH*, Supplementary Figure 1; *ACTB*, Supplementary Figure 2; *18S*, Supplementary Figure 3; *28S*, Supplementary Figure 4 and *RPL13A*, Supplementary Figure 5) and have been submitted into GenBank. GenBank accession numbers are as follows: *GAPDH*, KX 88916; *ACTB*, KX788917; *18S*, KX788914; *28S*, KX788915 and *RPL13A*, KX788918. The linear correlation coefficient ( $R^2$ ) of all genes ranged from 0.97-1.00.  $C_q$  values for all genes in all samples were within 10.41-33.19 cycles, and were covered by the range of their respective standard curves.  $E$  value of reference genes, mean  $C_q$  values and range of  $C_q$  values for each tissue are depicted in Table 2. All reference genes in all tissues had a  $C_q$  value below 29, indicating an abundance of target nucleic acid in cDNA samples (Fig. 1).  $C_q$  values observed for all reference genes in this study were insignificant in juvenile and adult tissues ( $p \geq 0.05$ ).

$C_q$  and  $E$  values were used in qBase<sup>PLUS</sup> to calculate expression stability ( $M$  value).  $M$  values are used to rank reference genes based on the stability, depending on the type of tissue (Fig. 2).  $M$  values below 1.5 indicate a stable expression (Nygard et al., 2007). According to the results obtained, the most stable genes in liver are *ACTB* and *GAPDH*, lungs *RPL13A* and *GAPDH*, small intestine *ACTB* and *18S*, and spleen *ACTB* and *18S*.

As per Nygard et al. (2007),  $M$  values lower than 1.5 were considered stable reference genes. Upon analysis using the geNorm application,  $M$  values for *ACTB* and *GAPDH* in all liver tissue samples were below 1.5, indicating their stability for use in the developing red-tailed phascogale studies. In



addition, all reference genes except 28S were stable for lung tissue, and all reference genes were stable in both small intestine and spleen tissue samples (Fig. 2; Table 3).

When the two age groups were analysed individually, all reference genes had  $M$  values below 1.5 for adult small intestine, juvenile and adult spleen tissue samples. The two most stable reference genes differed in juvenile and adult liver tissues:  $M$  values for 18S and 28S for adult liver samples were considered stable, whereas *ACTB* and *GAPDH* were stable in juvenile liver samples. All reference genes except 28S were considered stable in both juvenile and adult lung samples, and all reference genes except *RPL13A* were considered stable in juvenile small intestine tissue samples (Fig. 3).

## DISCUSSION

The ideal reference gene should constantly be transcribed in the type of tissue being examined (Nygard et al. 2007). Studies looking at the expression of reference genes in multiple tissues have however demonstrated that the regulation of reference genes are tissue specific (Lisowski et al. 2008). In this study, we provided a detailed analysis of the stability and expression levels of five different reference genes previously used in marsupial expression studies (Daly et al. 2009; Maher et al. 2014; Menzies et al. 2009; Siddle et al. 2013), in four different red-tailed phascogale tissues. We found at least two stable reference genes with  $M$  values  $>1.5$  for liver, lung, small intestine and spleen, and that all reference genes were suitable for expression studies of red-tailed phascogale small intestine and spleen tissues, which is useful as the normalisation of gene expression requires at least two reference genes (Bustin 2000). The average  $C_q$  values for all reference genes in all tissues were below 29 cycles, indicating strong positive reactions of the target tissue to the reference genes (Fig. 1).

Not unexpectedly, the results of this study showed that expression stability differs between different tissues, and confirm that reference genes are expressed in every cell but are regulated differently in different tissues (Lisowski et al. 2008). *GAPDH* is one of the most commonly used reference genes for normalisation in mammalian tissues. Studies have found *GAPDH* expression to be unstable as its expression differs, for example, according to age and sex of individuals (Barber et al. 2005). Therefore, it was not surprising that there were significant differences in *GAPDH* expression across tissues used in this study; *GAPDH* expression was most stable for liver and lung, and least stable for spleen tissues (Fig.

2; Table 3). When separated into two age groups, *GAPDH* expression had the highest stability in juvenile liver tissues, and a combination of the two ages. *GAPDH*, along with *ACTB*, were stable for all tissue samples, except for adult liver tissues.

*ACTB* is another reference gene commonly used for normalisation in mammalian tissues (Menzies et al. 2012; Nygard et al. 2007). Foss et al. (1998) found that the levels of *ACTB* were more variable than *GAPDH*, and that high levels of *ACTB* were found in the porcine small intestine and spleen. However, we found that *ACTB* was more stable than *GAPDH* in small intestine and spleen tissue. In particular, juvenile small intestine and spleen tissues had the highest *ACTB* stability. Selvey et al. (2001) found that *ACTB* is an unstable reference gene in mouse sarcoma cells (matrigel), and found *18S* to be more stable. When combining both age groups, our study agrees with Selvey et al. (2001), as *ACTB* was suitable for normalisation in small intestine and spleen tissue, however it was still less stable than *18S* in the same tissues. The same results were also observed using adult small intestine and spleen tissue samples.

Both *18S* and *28S* are often recommended as reference genes because ribosomal RNA has little variation among mammalian tissues and is often used as a successful internal standard (Goidin et al. 2001; Thellin et al. 1999). In addition, *18S* and *28S* were found to work effectively for normalisation in marsupial tissues (Janke et al. 2002; Maher et al. 2014). In this study, *18S* and *28S* were suitable for normalisation in small intestine and spleen tissue, with *18S* being the most stable. *18S* and *28S* were also the only two stable reference genes for adult liver tissue.

The last reference gene used in this study was *RPL13A*, a gene that encodes a protein in the 60S subunit of ribosomes (Vandesompele et al. 2002). Szabo et al. (2004) found *RPL13A* to be the best universal reference gene in various human tissues, including lung and small intestine. In addition, *RPL13A* had been selected by Ahn et al. (2008) as one of two ideal reference genes in rhesus macaques (*Macaca mulatta*), which included the comparison of eight reference genes in six tissues, including liver and lung. In this study, *RPL13A* was found to be least stable (highest *M* value) in liver and small intestine tissue, but suitable for normalisation in lung, small intestine and spleen tissue when both ages are combined. When divided into different age groups, *RPL13A* was stable for both age groups for lung and

spleen tissue and adult small intestine, showing its suitability for normalisation in most tissue samples, with the exception of liver.

The findings of this study confirmed previous research that demonstrated tissue specific regulation of some reference genes in eutherian mammals (Lisowski et al., 2008; Nygard et al., 2007; Uddin et al., 2011) also apply to a marsupial. Pierzchala et al. (2011) and Uddin et al. (2011) did not identify any of the reference genes used in this study as a stable reference gene in porcine liver, which shows that the regulation of certain reference genes may be different in marsupials. This study also demonstrated the stability of reference genes in some, but not all, marsupial tissues that were tested. This will aid in the selection of reference genes for normalisation in future expression studies in marsupials, particularly where studies of immune-related whole tissue preparations are performed. As in eutherian mammals, marsupial expression studies are increasing because of the ability of qPCR to detect and quantify nucleic acids (Bustin 2000). For example, expression studies have been conducted in koala (*Phascolarctos cinereus*) to test the up- or down-regulation of specific immune genes in stimulated cells (Maher et al. 2014), and whether viral RNA levels increased or decreased in association with age (Tarlinton et al. 2005). Expression studies are also useful for comparing gene expression in eutherians and marsupials (Hübler et al. 2013).

## CONCLUSIONS

We have successfully found stable reference genes in lung, small intestine and spleen tissue preparations from a dasyurid marsupial. It is possible to apply this study to whole tissue gene expression studies, especially when it is associated with immunity. While gene expression may occur at the single-cell level, whole tissue studies show the mean expression of several cell types available in the tissue (Kahlem et al., 2004), which is relevant especially when more than one type of cell relates to immunity in an immune system study. Future studies that focus on isolated cell preparations from these tissues will shed further light on reference gene expression and whether or not whole tissue preparations can be directly compared with cell culture studies. Results from the present study enable recommendations on reference genes suitable for use in various marsupial tissues and for normalisation in gene expression experiments in developing marsupials.

249

## 250 Additional Information and Declarations

### 251 List of abbreviations

252 GAPDH Glyceraldehyde-3-phosphate dehydrogenase

253 ACTB b-actin

254 28S 28S Ribosomal RNA

255 18S 18S Ribosomal protein L13a

256 Cq Cycle Threshold

257 E Reaction efficiency

258 M Expression stability value

259 qPCR real-time polymerase chain reaction

260 MALT mucosal-associated lymphoid tissue

261 GALT gut-associated lymphoid tissue

262 WSU Western Sydney University

263 PCR polymerase chain reaction

264

### 265 Availability of data and materials

266 Nucleic acid sequences have been deposited in the GenBank database and the accession

267 numbers have been included in the manuscript.

268

### 269 REFERENCES

270 Ahn K, Huh J-W, Park S-J, Kim D-S, Ha H-S, Kim Y-J, Lee J-R, Chang K-T, and Kim H-S. 2008.

271 Selection of internal reference genes for SYBR green qRT-PCR studies of rhesus monkey  
272 (*Macaca mulatta*) tissues. *BMC Molecular Biology* 9:78.

273 Barber RD, Harmer DW, Coleman RA, and Clark BJ. 2005. GAPDH as a housekeeping gene: analysis of  
274 GAPDH mRNA expression in a panel of 72 human tissues. *Physiological Genomics* 21:389-395.

275 Borthwick CR, Young LJ, and Old JM. 2014. The development of the immune tissues in marsupial pouch  
276 young. *Journal of Morphology* 275:822-839.

277 Bradley A, Foster W, Taggart D. 2008. Red-tailed phascogale. In: Van Dyck S, Strahan R, eds. *The*  
278 *Mammals of Australia*. Sydney: Reed New Holland, 101-102.

279 Bustin SA. 2000. Absolute quantification of mRNA using real-time reverse transcription polymerase chain  
280 reaction assays. *Journal of Molecular Endocrinology* 25:169-193.

- 281 Daly KA, Mailer SL, Digby MR, Lefèvre C, Thomson P, Deane E, Nicholas KR, and Williamson P. 2009.
- 282 Molecular analysis of tammar (*Macropus eugenii*) mammary epithelial cells stimulated with
- 283 lipopolysaccharide and lipoteichoic acid. *Veterinary Immunology Immunopathology* 129:36-48.
- 284 Foss DL, Baarsch MJ, and Murtaugh MP. 1998. Regulation of hypoxanthine phosphoribosyltransferase,
- 285 glyceraldehyde-3-phosphate dehydrogenase and  $\beta$ -actin mRNA expression in porcine immune
- 286 cells and tissues. *Animal Biotechnology* 9:67-78.
- 287 Foster W, Bradley A, Caton W, and Taggart D. 2006. Comparison of growth and development of the red-
- 288 tailed phascogale (*Phascogale calura*) in three captive colonies. *Australian Journal of Zoology*
- 289 54:343-352.
- 290 Glare E, Divjak M, Bailey M, and Walters E. 2002.  $\beta$ -Actin and GAPDH housekeeping gene expression in
- 291 asthmatic airways is variable and not suitable for normalising mRNA levels. *Thorax* 57:765-770.
- 292 Goidin D, Mamessier A, Staquet M-J, Schmitt D, and Berthier-Vergnes O. 2001. Ribosomal 18S RNA
- 293 prevails over glyceraldehyde-3-phosphate dehydrogenase and  $\beta$ -actin genes as internal standard
- 294 for quantitative comparison of mRNA levels in invasive and noninvasive human melanoma cell
- 295 subpopulations. *Analytical Biochemistry* 295:17-21.
- 296 Hellemans J, Mortier G, De Paepe A, Speleman F, and Vandesompele J. 2007. qBase relative
- 297 quantification framework and software for management and automated analysis of real-time
- 298 quantitative PCR data. *Genome Biology* 8:R19.
- 299 Higuchi R, Dollinger G, Walsh PS, and Griffith R. 1992. Simultaneous amplification and detection of
- 300 specific DNA sequences. *Nature Biotechnology* 10:413-417.
- 301 Hübler M, Molineaux AC, Keyte A, Schecker T, and Sears KE. 2013. Development of the marsupial
- 302 shoulder girdle complex: a case study in *Monodelphis domestica*. *Evolution & Development*
- 303 15:18-27.
- 304 Janke A, Magnell O, Wieczorek G, Westerman M, and Arnason U. 2002. Phylogenetic analysis of 18S
- 305 rRNA and the mitochondrial genomes of the wombat, *Vombatus ursinus*, and the spiny anteater,
- 306 *Tachyglossus aculeatus*: increased support for the *Marsupionta* hypothesis. *Journal of Molecular*
- 307 *Evolution* 54:71-80.
- 308 Kennedy M, Williams G. 1990. *Australia's endangered species: The extinction dilemma*. New York, USA:
- 309 Prentis Hall Press.
- 310 Laskin DL, Weinberger B, and Laskin JD. 2001. Functional heterogeneity in liver and lung macrophages.
- 311 *J Leukocyte Biol* 70:163-170.
- 312 Lisowski P, Pierzchala M, Goscik J, Pareek CS, and Zwierzchowski L. 2008. Evaluation of reference
- 313 genes for studies of gene expression in the bovine liver, kidney, pituitary and thyroid. *Journal of*
- 314 *Applied Genetics* 49:367-372.
- 315 Maher IE, Griffith JE, Lau Q, Reeves T, and Higgins DP. 2014. Expression profiles of the immune genes
- 316 CD4, CD8 $\beta$ , IFN $\gamma$ , IL-4, IL-6 and IL-10 in mitogen-stimulated koala lymphocytes (*Phascolarctos*
- 317 *cinereus*) by qRT-PCR. *PeerJ* 2:e280.
- 318 Mak TW, Saunders ME, and Jett BD. 2013. *Primer to the immune response*: Newnes.
- 319 Markey B, Wan C, Hanger J, Phillips C, and Timms P. 2007. Use of quantitative real-time PCR to monitor
- 320 the shedding and treatment of chlamydiae in the koala (*Phascolarctos cinereus*). *Veterinary*
- 321 *Microbiology* 120:334-342.
- 322 Menzies BR, Shaw G, Fletcher TP, Pask AJ, and Renfree MB. 2012. Maturation of the growth axis in
- 323 marsupials occurs gradually during post-natal life and over an equivalent developmental stage
- 324 relative to eutherian species. *Molecular and Cellular Endocrinology* 349:189-194.
- 325 Menzies BR, Shaw G, Fletcher TP, and Renfree MB. 2009. Early onset of ghrelin production in a
- 326 marsupial. *Molecular and Cellular Endocrinology* 299:266-273.
- 327 Nygard A-B, Jørgensen CB, Cirera S, and Fredholm M. 2007. Selection of reference genes for gene
- 328 expression studies in pig tissues using SYBR green qPCR. *BMC Mol Biol* 8:67.
- 329 Old J, and Deane E. 2002. The gut-associated lymphoid tissues of the northern brown bandicoot
- 330 (*Isodon macrourus*). *Developmental & Comparative Immunology* 26:841-848.
- 331 Old JM, and Deane EM. 2000. Developmental of the immune system and immunological protection in
- 332 marsupial pouch young. *Developmental & Comparative Immunology* 24:445-454.
- 333 Pierzchala M, Lisowski P, Urbanski P, Pareek CS, Cooper RG, Kuryl J. 2011. Evaluation based selection
- 334 of housekeeping genes for studies of gene expression in the porcine muscle and liver tissues.
- 335 *Journal of Animal and Veterinary Advances* 10:401-405.

- 336 Pierzchała M, Pareek PL, Urbański P, Goluch UC, Kamyczek MR, Cooper JK. 2011. Expression profile of  
337 MYF5 and MYF6 genes in skeletal muscles of young growing gilts of five breeds at different ages,  
338 based on the most stable reference genes. *Animal Science Papers and Reports* 29:231-246.
- 339 Russell EM. 1982. Patterns of parental care and parental investment in marsupials. *Biological Reviews*  
340 57:423-486.
- 341 Schmittgen TD, and Livak KJ. 2008. Analysing real-time PCR data by the comparative Ct method. *Nature*  
342 *Protocols* 3:1101-1108.
- 343 Selvey S, Thompson EW, Matthaei K, Lea RA, Irving MG, and Griffiths LR. 2001.  $\beta$ -Actin—an unsuitable  
344 internal control for RT-PCR. *Molecular and Cellular Probes* 15:307-311.
- 345 Siddle HV, Kreiss A, Tovar C, Yuen CK, Cheng Y, Belov K, Swift K, Pearce A-M, Hamede R, and Jones  
346 ME. 2013. Reversible epigenetic down-regulation of MHC molecules by devil facial tumour  
347 disease illustrates immune escape by a contagious cancer. *PNAS* 110:5103-5108.
- 348 Stannard HJ, Borthwick CR, Ong O, and Old JM. 2013. Longevity and breeding in captive red-tailed  
349 phascogales (*Phascogale calura*). *Australian Mammalogy* 35:217-219.
- 350 Szabo A, Perou CM, Karaca M, Perreard L, Quackenbush JF, and Bernard PS. 2004. Statistical modeling  
351 for selecting housekeeper genes. *Genome Biol* 5:R59.
- 352 Tarlinton R, Meers J, Hanger J, and Young P. 2005. Real-time reverse transcriptase PCR for the  
353 endogenous koala retrovirus reveals an association between plasma viral load and neoplastic  
354 disease in koalas. *Journal of General Virology* 86:783-787.
- 355 Thellin O, Zorzi W, Lakaye B, Borman BD, Coumans B, Hennen G, Grisar T, Igout A, and Heinen E.  
356 1999. Housekeeping genes as internal standards: use and limits. *Journal of Biotechnology*  
357 75:291-295.
- 358 Uddin MJ, Cinar MU, Tesfaye D, Looft C, Tholen E, and Schellander K. 2011. Age-related changes in  
359 relative expression stability of commonly used housekeeping genes in selected porcine tissues.  
360 *BMC Res Notes* 4:441.
- 361 Vandesompele J, Preter KD, Pattyn F, Poppe B, Roy NV, Paepe AD, and Speleman F. 2002. Accurate  
362 normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal  
363 control genes. *Genome Biol* 3:research0034.
- 364 Warner JR. 1999. The economics of ribosome biosynthesis in yeast. *Trends in Biochemical Sciences*  
365 24:437-440.
- 366 Yu H, Pask AJ, Shaw G, and Renfree MB. 2006. Differential expression of WNT4 in testicular and ovarian  
367 development in a marsupial. *BMC Developmental Biology* 6:44.
- 368 Zhong H, and Simons JW. 1999. Direct comparison of GAPDH,  $\beta$ -actin, cyclophilin, and 28S rRNA as  
369 internal standards for quantifying RNA levels under hypoxia. *Biochemical and Biophysical*  
370 *Research Communications* 259:523-526.



372 Table 1

373 **Candidate reference genes evaluated in this study.**

| Gene symbol   | Gene name                                | Oligo sequence (5' → 3')                                                                                                                                 | Amplicon size (bp) | Annealing temp. (°C) |
|---------------|------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|----------------------|
| <i>GAPDH</i>  | Glyceraldehyde-3-phosphate dehydrogenase | Forward: CAGGCGGAGTAGACATTG<br>Reverse: CCTTGAAGTTGCCATGGG                                                                                               | 63                 | 60                   |
| <i>ACTB</i>   | β-Actin                                  | Forward: TTGCTGACAGGATGCAGAAG<br>Reverse: GAGCCTCCAATCCAGACAGA<br>Rindfleisch et al. (2010).                                                             | 66                 | 60                   |
| <i>28S</i>    | 28S Ribosomal RNA                        | Forward: CGATGTCGGCTCTTCCTATC<br>Reverse: TCCTCAGCCAAGCACATACA*<br>Daly et al. (2009).<br>*Reverse primer was modified according to marsupial sequences. | 165                | 60                   |
| <i>18S</i>    | 18S Ribosomal RNA                        | Forward: CCAACACGGGAAACCTCA<br>Reverse: AACCAGAAATCGCTCCAC<br>(Daly et al. 2009).                                                                        | 83                 | 60                   |
| <i>RPL13A</i> | Ribosomal protein L13a                   | Forward: CCCCACAAGACCAAGCGAGGC<br>Reverse: ACAGCCTGGTATTTCCAGCCAACC<br>(Siddle et al. 2013).                                                             | 146                | 60                   |

374

375

376 Table 2

377 **Cycle threshold ( $C_q$ ) and reaction efficiency ( $E$ ) values for individual genes in examined tissues of**  
 378 **juveniles and adults.**

|                        | <i>GAPDH</i>      | <i>ACTB</i>       | <i>18S</i>        | <i>28S</i>        | <i>RPL13A</i>     |
|------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| <b>Liver</b>           |                   |                   |                   |                   |                   |
| Mean $C_q$             | 26.242            | 27.226            | 14.526            | 15.382            | 27.907            |
| Range of $C_q$         | 22.55-29.98       | 22.07-30.09       | 9.66-18.48        | 9.83-21.28        | 22.27-33.19       |
| $E$ (out of 2)         | $1.855 \pm 0.047$ | $1.716 \pm 0.08$  | $1.953 \pm 0.035$ | $1.89 \pm 0.011$  | $1.851 \pm 0.029$ |
| <b>Lung</b>            |                   |                   |                   |                   |                   |
| Mean $C_q$             | 24.6              | 21.746            | 12.995            | 14.476            | 21.924            |
| Range of $C_q$         | 22.74-28.06       | 19.17-24.84       | 10.41-16.64       | 10.8-20.14        | 19.72-24.43       |
| $E$ (out of 2)         | $1.94 \pm 0.058$  | $1.732 \pm 0.04$  | $1.912 \pm 0.026$ | $1.909 \pm 0.035$ | $1.763 \pm 0.032$ |
| <b>Small Intestine</b> |                   |                   |                   |                   |                   |
| Mean $C_q$             | 23.981            | 23.849            | 13.867            | 15.04             | 22.618            |
| Range of $C_q$         | 21.75-25.50       | 20.67-27.47       | 11.56-17.67       | 12.32-17.50       | 18.89-28.53       |
| $E$ (out of 2)         | $1.919 \pm 0.043$ | $1.753 \pm 0.058$ | $1.833 \pm 0.033$ | $1.703 \pm 0.013$ | $1.811 \pm 0.067$ |
| <b>Spleen</b>          |                   |                   |                   |                   |                   |
| Mean $C_q$             | 23.541            | 23.566            | 15.361            | 16.673            | 20.661            |
| Range of $C_q$         | 21.24-26.89       | 21.64-26.56       | 13.46-17.81       | 14.8-17.59        | 18.92-23.40       |
| $E$ (out of 2)         | $1.879 \pm 0.033$ | $1.693 \pm 0.043$ | $1.768 \pm 0.036$ | $1.978 \pm 0.029$ | $1.801 \pm 0.028$ |

379

380



381 Table 3

382 Control genes ranked in order of their expression stability\*. Reference genes with *M* values below

383 1.5 are considered as suitable reference genes for a particular immune tissue and are bold.

| Liver               | Lung                | Small Intestine    | Spleen             |
|---------------------|---------------------|--------------------|--------------------|
| <i>RPL13A</i>       | <i>28S</i>          | <i>RPL13A</i>      | <i>GAPDH</i>       |
| <i>28S</i>          | <b><i>18S</i></b>   | <i>GAPDH</i>       | <i>RPL13A</i>      |
| <i>18S</i>          | <b><i>ACTB</i></b>  | <i>28S</i>         | <i>28S</i>         |
| <b><i>ACTB</i></b>  | <i>RPL13A</i>       | <b><i>ACTB</i></b> | <b><i>ACTB</i></b> |
| <b><i>GAPDH</i></b> | <b><i>GAPDH</i></b> | <b><i>18S</i></b>  | <b><i>18S</i></b>  |

384 \*Increasing expression stability from top to bottom.

385

386 Figure 1

387 **Average RNA transcription levels of putative reference genes ( $\pm$ SE; n=10) presented as absolute**  
 388 **Cq values.** The reference genes are: Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -Actin  
 389 (*ACTB*), 18S rRNA (*18S*), 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*) in liver, lung, small  
 390 intestine and spleen tissue samples.

391 Figure 2

392 **Overall expression stability values (*M*) of reference genes in tissues of different ages (n=10).** Gene  
 393 expression stability of reference genes in juvenile and adult red-tailed phascogale tissues **(A)** liver, **(B)**  
 394 lung, **(C)** small intestine, and **(D)** spleen, analysed by geNorm application (qBase<sup>PLUS</sup>). The reference  
 395 genes are: Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -Actin (*ACTB*), 18S rRNA (*18S*),  
 396 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*). Genes with the lowest *M* values have the most  
 397 stable expression. Reference gene points below *M* value = 1.5 indicate that the reference gene is stable  
 398 and can be used for normalisation in the respective tissue. *M* value threshold (=1.5) is indicated by  
 399 dashed lines.

400 Figure 3

401 **Expression stability values (*M*) of reference genes in tissues of juvenile (n=5) and adults (n=5).**  
 402 Gene expression stability of reference genes in **(A)** juvenile and adult liver, **(B)** juvenile and adult lung,  
 403 **(C)** juvenile and adult small intestine, **(D)** juvenile and adult spleen tissues, analysed by geNorm  
 404 application (qBase<sup>PLUS</sup>). The reference genes are: Glyceraldehyde 3-phosphate dehydrogenase  
 405 (*GAPDH*),  $\beta$ -Actin (*ACTB*), 18S rRNA (*18S*), 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*).  
 406 Genes with the lowest *M* values have the most stable expression. Reference gene points below *M* value  
 407 = 1.5 indicate that the reference gene is stable and can be used for normalisation in the respective tissue.  
 408 *M* value threshold (=1.5) is indicated by dashed lines.

409

410 Supplementary Figure 1

411 **Melting curves for glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) expression.** The melting  
412 curves for *GAPDH* expression in (A) liver, (B) lung, (C) small intestine and (D) spleen of red-tailed  
413 phascogales (n=10), performed in triplicate.

414 Supplementary Figure 2

415 **Melting curves for  $\beta$ -Actin (*ACTB*) expression.** The melting curves for *ACTB* expression in (A) liver,  
416 (B) lung, (C) small intestine and (D) spleen of red-tailed phascogales (n=10, performed in triplicate.

417 Supplementary Figure 3

418 **Melting curves for 18S rRNA (*18S*).** The melting curves for *18S* expression in (A) liver, (B) lung, (C)  
419 small intestine and (D) spleen of red-tailed phascogales (n=10), performed in triplicate.

420 Supplementary Figure 4

421 **Melting curves for 28S rRNA (*28S*).** The melting curves for *28S* expression in (A) liver, (B) lung, (C)  
422 small intestine and (D) spleen of red-tailed phascogales (n=10), performed in triplicate.


423 Supplementary Figure 5

424 **Melting curves for ribosomal protein L13a (*RPL13A*).** The melting curves for *RPL13A* expression in  
425 (A) liver, (B) lung, (C) small intestine and (D) spleen of red-tailed phascogales (n=10), performed in  
426 triplicate.x

# Figure 1

Average RNA transcription levels of putative reference genes presented as absolute C<sub>q</sub> values.


The reference genes are: Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -Actin (*ACTB*), 18S rRNA (*18S*), 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*) in liver, lung, small intestine and spleen tissue samples.



# Figure 2

Overall expression stability values ( $M$ ) of reference genes in tissues of different ages.


Gene expression stability of reference genes in juvenile and adult red-tailed phascogale tissues (A) liver, (B) lung, (C) small intestine and (D) spleen analysed by geNorm application (qBase<sup>PLUS</sup>). The reference genes are: Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -Actin (*ACTB*), 18S rRNA (*18S*), 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*). Genes with the lowest  $M$  values have the most stable expression.



# Figure 3

Expression stability values ( $M$ ) of reference genes in tissues of juvenile and adults.

Gene expression stability of reference genes in (A) juvenile liver, (B) adult liver, (C) juvenile lung, (D) adult lung, (E) juvenile small intestine, (F) adult small intestine, (G) juvenile spleen and (H) adult spleen tissues, analysed by geNorm application (qBase<sup>PLUS</sup>). The reference genes are: Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*),  $\beta$ -Actin (*ACTB*), 18S rRNA (*18S*), 28S rRNA (*28S*) and ribosomal protein L13a (*RPL13A*). Genes with the lowest  $M$  values have the most stable expression.



# **Table 1**(on next page)

Candidate reference genes evaluated in this study.



| Gene symbol   | Gene name                                | Oligo sequence (5' → 3')                                                                                                                                 | Amplicon size (bp) | Annealing temp. (°C) |
|---------------|------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|----------------------|
| <i>GAPDH</i>  | Glyceraldehyde-3-phosphate dehydrogenase | Forward: CAGGCGGAGTAGACATTG<br>Reverse: CCTTGAACCTGCCATGGG                                                                                               | 90                 | 60                   |
| <i>ACTB</i>   | β-Actin                                  | Forward: CTCTTCCAGCCATCTTTCTT<br>Reverse: GACATCCGTAAGGATCTGTA                                                                                           | 100                | 60                   |
| <i>28S</i>    | 28S Ribosomal RNA                        | Forward: CGATGTCGGCTCTTCCTATC<br>Reverse: TCCTCAGCCAAGCACATACA*<br>Daly et al. (2009).<br>*Reverse primer was modified according to marsupial sequences. | 203                | 60                   |
| <i>18S</i>    | 18S Ribosomal RNA                        | Forward: CCAACACGGGAAACCTCA<br>Reverse: AACCAGAAATCGCTCCAC<br>Daly <i>et al</i> (Daly et al. 2009).                                                      | 121                | 60                   |
| <i>RPL13A</i> | Ribosomal protein L13a                   | Forward: CCCCACAAGACCAAGCGAGGC<br>Reverse: ACAGCCTGGTATTTCCAGCCAACC<br>Siddle (Siddle et al. 2013).                                                      | 145                | 60                   |

## Table 2 (on next page)

Cycle threshold ( $C_q$ ) and reaction efficiency ( $E$ ) values for individual genes in examined tissues of juveniles and adults.

|                         | <i>GAPDH</i>  | <i>ACTB</i>   | <i>18S</i>    | <i>28S</i>    | <i>RPL13A</i> |
|-------------------------|---------------|---------------|---------------|---------------|---------------|
| <b>Liver</b>            |               |               |               |               |               |
| Mean C <sub>q</sub>     | 26.242        | 27.226        | 14.526        | 15.382        | 27.907        |
| Range of C <sub>q</sub> | 22.55-29.98   | 22.07-30.09   | 9.66-18.48    | 9.83-21.28    | 22.27-33.19   |
| <i>E</i> (out of 2)     | 1.855 ± 0.047 | 1.716 ± 0.08  | 1.953 ± 0.035 | 1.89 ± 0.011  | 1.851 ± 0.029 |
| <b>Lung</b>             |               |               |               |               |               |
| Mean C <sub>q</sub>     | 24.6          | 21.746        | 12.995        | 14.476        | 21.924        |
| Range of C <sub>q</sub> | 22.74-28.06   | 19.17-24.84   | 10.41-16.64   | 10.8-20.14    | 19.72-24.43   |
| <i>E</i> (out of 2)     | 1.94 ± 0.058  | 1.732 ± 0.04  | 1.912 ± 0.026 | 1.909 ± 0.035 | 1.763 ± 0.032 |
| <b>Small Intestine</b>  |               |               |               |               |               |
| Mean C <sub>q</sub>     | 23.981        | 23.849        | 13.867        | 15.04         | 22.618        |
| Range of C <sub>q</sub> | 21.75-25.50   | 20.67-27.47   | 11.56-17.67   | 12.32-17.50   | 18.89-28.53   |
| <i>E</i> (out of 2)     | 1.919 ± 0.043 | 1.753 ± 0.058 | 1.833 ± 0.033 | 1.703 ± 0.013 | 1.811 ± 0.067 |
| <b>Spleen</b>           |               |               |               |               |               |
| Mean C <sub>q</sub>     | 23.541        | 23.566        | 15.361        | 16.673        | 20.661        |
| Range of C <sub>q</sub> | 21.24-26.89   | 21.64-26.56   | 13.46-17.81   | 14.8-17.59    | 18.92-23.40   |
| <i>E</i> (out of 2)     | 1.879 ± 0.033 | 1.693 ± 0.043 | 1.768 ± 0.036 | 1.978 ± 0.029 | 1.801 ± 0.028 |

1  
2

### Table 3 (on next page)

Control genes ranked in order of their expression stability\*. Reference genes with  $M$  values below 1.5 are considered as suitable reference genes for a particular immune tissue and are bold.

| Liver         | Lung          | Small Intestine | Spleen        |
|---------------|---------------|-----------------|---------------|
| <i>RPL13A</i> | <i>28S</i>    | <i>RPL13A</i>   | <i>GAPDH</i>  |
| <i>28S</i>    | <i>18S</i>    | <i>GAPDH</i>    | <i>RPL13A</i> |
| <i>18S</i>    | <i>ACTB</i>   | <i>28S</i>      | <i>28S</i>    |
| <i>ACTB</i>   | <i>RPL13A</i> | <i>ACTB</i>     | <i>ACTB</i>   |
| <i>GAPDH</i>  | <i>GAPDH</i>  | <i>18S</i>      | <i>18S</i>    |

\*Increasing expression stability from top to bottom.