

A peer-reviewed version of this preprint was published in PeerJ on 3 November 2016.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.2656) (peerj.com/articles/2656), which is the preferred citable publication unless you specifically need to cite this preprint.

Fort T, Robin C, Capdevielle X, Delière L, Vacher C. 2016. Foliar fungal communities strongly differ between habitat patches in a landscape mosaic. PeerJ 4:e2656 <https://doi.org/10.7717/peerj.2656>

Foliar fungal communities strongly differ between habitat patches in a landscape mosaic

Thomas Fort ^{Corresp., 1}, **Cécile Robin** ¹, **Xavier Capdevielle** ¹, **Laurent Delière** ², **Corinne Vacher** ³

¹ BIOGECO, UMR 1202, INRA, Université de Bordeaux, Cestas, France

² SAVE, UMR 1065, INRA, ISVV, Université de Bordeaux, INRA, Villenave d'Ornon, France

³ BIOGECO, UMR 1202, INRA, Université de Bordeaux, Pessac, France

Corresponding Author: Thomas Fort

Email address: thomaslc.fort@gmail.com

Dispersal events between habitat patches in a landscape mosaic can structure ecological communities and influence the functioning of agrosystems. Here we investigated whether short-distance dispersal events between vineyard and forest patches shape foliar fungal communities. We hypothesized that these communities homogenize between habitats over the course of the growing season, particularly along habitat edges, because of aerial dispersal of spores.

We monitored the richness and composition of foliar and airborne fungal communities over the season, along transects perpendicular to edges between vineyard and forest patches, using Illumina sequencing of the ITS2 region.

In contrast to our expectation, foliar fungal communities in vineyards and forest patches increasingly differentiate over the growing season, even along habitat edges. Moreover, the richness of foliar fungal communities in grapevine drastically decreased over the growing season, in contrast to that of forest trees. The composition of airborne communities did not differ between habitats. The composition of oak foliar fungal communities change between forest edge and centre.

These results suggest that dispersal events between habitat patches are not major drivers of foliar fungal communities at the landscape scale. Selective pressures exerted in each habitat by the host plant, the microclimate and the agricultural practices play a greater role, and might account for the differentiation of foliar fungal communities between habitats.

Foliar fungal communities strongly differ between habitat patches in a landscape mosaic

Thomas Fort¹, Cécile Robin¹, Xavier Capdevielle¹, Laurent Delière², Corinne Vacher³

¹ BIOGECO, UMR 1202, INRA, Université de Bordeaux, F-33610 Cestas, France

² SAVE, UMR1065, INRA, ISVV, Université de Bordeaux, F-33883 Villenave d'Ornon, France

³ BIOGECO, UMR 1202, INRA, Université de Bordeaux, F-33615 Pessac, France

Corresponding Author :

Thomas Fort¹

69 route d'Arcachon, 33612 Cestas, France

Email address: thomaslc.fort@gmail.com

13 Summary

14 **Background.** Dispersal events between habitat patches in a landscape mosaic can structure
15 ecological communities and influence the functioning of agrosystems. Here we investigated
16 whether short-distance dispersal events between vineyard and forest patches shape foliar fungal
17 communities. We hypothesized that these communities homogenize between habitats over the
18 course of the growing season, particularly along habitat edges, because of aerial dispersal of
19 spores.

20 **Methods.** We monitored the richness and composition of foliar and airborne fungal communities
21 over the season, along transects perpendicular to edges between vineyard and forest patches,
22 using Illumina sequencing of the ITS2 region.

23 **Results.** In contrast to our expectation, foliar fungal communities in vineyards and forest patches
24 increasingly differentiate over the growing season, even along habitat edges. Moreover, the
25 richness of foliar fungal communities in grapevine drastically decreased over the growing
26 season, in contrast to that of forest trees. The composition of airborne communities did not differ
27 between habitats. The composition of oak foliar fungal communities change between forest edge
28 and centre.

29 **Discussion.** These results suggest that dispersal events between habitat patches are not major
30 drivers of foliar fungal communities at the landscape scale. Selective pressures exerted in each
31 habitat by the host plant, the microclimate and the agricultural practices play a greater role, and
32 might account for the differentiation of foliar fungal communities between habitats.

34 Introduction

35 Plant leaves provide one of the largest microbial habitats on Earth (Ruinen, 1956; Morris, 2001;
36 Vorholt, 2012). They harbour highly diverse microbial communities, including many genera of
37 bacteria and fungi (Lindow & Leveau, 2002; Vorholt, 2012; Turner et al., 2013). The eco-
38 evolutionary processes which shape these communities – dispersal, evolutionary diversification,
39 selection and drift – are increasingly well understood (Hanson et al., 2012; Nemergut et al.,
40 2013; Vacher et al., 2016). This new eco-evolutionary framework will undoubtedly have
41 important applications in agriculture. Indeed, crop performance depends on the balance and
42 interactions between pathogenic and beneficial microbial species (Newton et al., 2010a, 2010b).
43 Manipulating whole foliar microbial communities, by acting on the processes shaping them,
44 could thus greatly improve crop health (Newton et al., 2010a; Xu et al., 2011). However, to reach
45 this aim, a better understanding of the structure and dynamics of foliar microbial communities at
46 the landscape scale is required.

47 The landscape plays a key role in the dynamics of macro-organism populations interacting with
48 crops, such as arthropod pests or their natural enemies (Norris & Kogan, 2000; Chaplin-Kramer
49 et al., 2011). In ecology, the landscape is defined as an heterogeneous geographic area,
50 characterized by a dynamic mosaic of interacting habitat patches (Bastian, 2001). Species
51 movements between habitat patches - referred as dispersal (Vellend, 2010) - modulates the
52 richness, composition and function of macro-organism communities (Hurst et al., 2013; Ma
53 et al., 2013; Lacasella et al., 2014). In agricultural landscape, species dispersal between natural
54 and managed habitats can trigger detrimental or beneficial effects in crops (Chaplin-Kramer

et al., 2011; Blitzer et al., 2012), particularly along the edges (Thomson & Hoffmann, 2009; Lacasella et al., 2014).

The influence of dispersal events on the structure of foliar microbial communities at the landscape scale has hardly been studied. Many microbial species colonising plant leaves are horizontally transferred (i.e. from one adult plant to another) by airborne dispersal (Whipps et al., 2008; Bulgarelli et al., 2013), while others can come from the seeds, the rhizosphere or the twigs (Vorholt, 2012). The foliar microbial communities of a given plant can therefore be influenced by those of its neighbours. Plant pathogens, for instance, can be transmitted from a reservoir plant to neighbouring plants (Power & Mitchell, 2004; Beckstead et al., 2010; Wilson et al., 2014). These short-distance dispersal events could have a greater effect on the foliar microbial communities of annual or deciduous plants, because the leaves of those plants are colonised by micro-organisms every spring, after budbreak.

In this study, we analysed the structure and dynamic of foliar and airborne fungal communities in a heterogeneous landscape consisting of vineyard and forest patches in the south west of France. Vineyards are human-engineered agro-ecosystems, characterized by a low specific and genetic diversity, and where weeds, pests and pathogens are regularly controlled with different cultural practices and pesticides to preserve yield and to reduce infection of leaves and grapes. Conversely, deciduous forests in this area remain little managed and much less homogeneous. We expected the fungal communities of forest patches to be richer than those of vineyards, because the higher plant species richness and biomass in forests increase the diversity of micro-habitats available to foliar fungi. We also expected repeated dispersal events to homogenize foliar fungal communities between the two habitats over the course of the growing season, particularly along habitat edges. We thus tested the following hypotheses for both foliar and

airborne fungal communities: (1) community richness is higher in forests than in adjacent vineyards, (2) community similarity between the two habitats increase over the course of the growing season and (3) is higher along habitat edges.

Materials and methods

Sampling design

Three study sites, each consisting of a forest patch and an adjacent vineyard, were selected in the Bordeaux area (France). They were located in the domains of Châteaux Reignac (N44°54'03", O0°25'01"), Grand-Verdus (N44°47'21", O0°24'06") and Couhins (N44°45'04", O0°33'53") (Fig. 1a). At each site, the edge between the forest patch and the vineyard was at least 100 m long. The width of each habitat, perpendicular to the edge, was at least 200 m. The forest patches at all three sites contained mostly deciduous species, dominated by pedunculate oak (*Quercus robur* L.). The second most frequent tree species was European hornbeam (*Carpinus betulus* L.) in Reignac and Grand-Verdus, and sweet chestnut (*Castanea sativa* Mill.) in Couhins. In the vineyards, the grapevine (*Vitis vinifera* L.) cultivar was Cabernet Sauvignon in Reignac and Grand-Verdus, and Merlot in Couhins.

At each site, leaves were collected along three parallel transects perpendicular to the forest-vineyard edge and separated by a distance of about five meters (Fig. 1b). Leaves were sampled at four locations along each transect: in the centre of the forest (100 m away from the edge), at the edge of the forest, at the edge of the vineyard and in the centre of the vineyard (100 m away from the edge). In forest patches, leaves were sampled from the two most abundant tree species. For

each sampling location and each transect, a single tree of each species was selected. Three leaves oriented in different directions were collected from each tree, at a height of 7 m. In vineyards, three leaves were collected from three adjacent cloned grapevine stocks. Each of the sampled leaves was selected from the base of the cane (one-year-old shoot), to ensure the collection of leaves of the same age on each date. The leaves were removed with scissors that had been sterilised with 96 % ethanol, and all contact of the leaves with the hands was carefully avoided. The leaves were stored in clear plastic bags containing silica gel to ensure rapid drying. In addition, grapevine leaves were placed between two sheets of sterile paper filter to ensure good dessication despite their thickness. Leaves were sampled on three dates in 2013: in May (between the 15th and 23rd), July (between the 16th and 18th) and October (3rd). The sampling dates chosen were as far removed as possible from the last chemical treatment performed in the vineyard (Supporting Information Table S1).

Airborne particles were collected along the middle transect of each site, with two Coriolis air sampler devices positioned one meter above the ground. At each sampling location, three successive 10 minute sampling sessions were carried out, with a flow rate of 200 l/min.

DNA extraction and sequencing

Sample contamination was prevented by exposing all tools and materials required for sample processing and DNA extraction to UV light for 30 minutes in a laminar flow hood. Four discs (each 8.0 mm in diameter) were cut randomly from each leaf, in the flow hood, with a hole-punch sterilised by flaming with 95 % ethanol. The four discs were placed in a single well of an autoclaved DNA extraction plate. Three wells were left empty as negative controls. Two

autoclaved metallic beads were added to each well, and the plant material was ground into a homogeneous powder with a Geno/Grinder 2010 (SPEX Sample Prep, Metuchen, NJ).

The liquid used to collect airborne particles was transferred into sterile 15 ml centrifuge tubes. Each tube was then centrifuged for 30 minutes at 13000 RCF and the supernatant was removed with a sterile transfer pipette. The pellet was then transferred by resuspension to an autoclaved tube and freeze-dried. A tube of unused sampling liquid was treated in the same way and used as a negative control. Total DNA was extracted from each leaf and airborne sample with the DNeasy 96 Plant Kit (QIAGEN). Foliar DNA samples from the same tree were pooled, as were foliar DNA samples from the three adjacent grapevine stocks.

Fungal ITS2 (Internal Transcribed Spacer 2) was amplified with the fITS7 (forward) and ITS4 (reverse) primers (Ihrmark et al., 2012). Paired-end sequencing (300 bp) was then performed in a single run of an Illumina MiSeq sequencer, on the basis of V3 chemistry. PCR amplification, barcodes and MiSeq adapters addition, library sequencing and data preprocessing were carried out by the LGC Genomics sequencing service (Berlin, Germany). Sequences were deposited in the European Nucleotide Archive (ENA) database, under the PRJEB13880 project accession number.

Bioinformatic analysis

Sequences were first demultiplexed and filtered. All sequences with tag mismatches, missing tags, one-sided tags or conflicting tag pairs were discarded. Tags and Illumina TruSeq adapters were then clipped from all sequences, and sequences with a final length fewer than 100 bases

were discarded. All sequences with more than three mismatches with the ITS2 primers were discarded. Primers were then clipped and the sequence fragments were placed in a forward-reverse primer orientation. Forward and reverse reads were then combined, and read pair sequences that could not be combined were discarded.

The pipeline developed by Bálint et al. (2014) was used to process the sequences. The ITS2 sequence was first extracted from each sequence with the FungalITSextractor (Nilsson et al., 2010). All the sequences were then concatenated into a single fasta file, after adding the sample code in the label of each sequence. The sequences were dereplicated, sorted and singletons were discarded with VSEARCH (<https://github.com/torognes/vsearch>). The sequences were then clustered into molecular operational taxonomic units (OTUs) with the UPARSE algorithm implemented in USEARCH v8 (Edgar, 2013), with a minimum identity threshold of 97 %. Additional chimera detection was performed against the UNITE database (Kõljalg et al., 2013), with the UCHIME algorithm implemented in USEARCH v8 (Edgar et al., 2011). The OTU table, giving the number of sequences of each OTU for each sample, was created with USEARCH v8.

OTUs were taxonomically assigned using the online BLAST web interface (Madden, 2013) against the GenBank database, by excluding environmental and metagenome sequences. The assignment with the lowest e-value was retained. The full taxonomic lineage of each assignment was retrieved from the GI number information provided by NCBI. All the OTUs assigned to plants or other organisms, and all unassigned OTUs were removed, to ensure that only fungal OTUs were retained.

164 Statistical analyses

165 All statistical analyses were performed in the R environment. We computed 100 random rarefied
 166 OTU matrices, using the smallest number of sequences per sample as a threshold. The number of
 167 OTUs per sample (OTU richness) and the dissimilarity between samples (Bray-Curtis index
 168 based on abundances and Jaccard index based on occurrences) were calculated for each rarefied
 169 matrix and averaged (Cordier et al., 2012; Jakuschkin et al., 2016). However, because the
 170 relevance of rarefaction is debated in the scientific community (Hughes & Hellmann, 2005;
 171 McMurdie & Holmes, 2014), we also performed the analyses on the raw OTU matrix by
 172 including the square root of the total number of sequences per sample (abundance) as first
 173 explanatory variable in all the models.

174 Type III ANOVA, which tests for the presence of an effect, given the other effects and the
 175 interactions (Herr, 1986), was used to assess the effect of host plant species (grapevine, oak,
 176 hornbeam and chestnut), sampling date (May, July, October), edge (habitat centre or edge) and
 177 their interactions on foliar OTU richness. Sampling site was included in the model as a random
 178 factor. Marginal and conditional coefficients of determination were calculated to estimate the
 179 variance explained by fixed factors (R_m^2) and fixed *plus* random factors (R_c^2). Post-hoc pairwise
 180 comparisons were then performed for each level of each factor, with Tukey's adjustment
 181 method. A similar ANOVA was performed on airborne OTU richness, including habitat (forest
 182 and vineyard), sampling date, sampling site, and their interactions.

183 Dissimilarities in composition between samples were represented by non-metric
 184 multidimensional scaling analysis (NMDS) and were analysed by permutational multivariate
 185 analyses of variance (PERMANOVA), including the same fixed factors as the ANOVAs, with

sampling sites treated as strata. We dealt with complex interactions in PERMANOVA results by calculating post-hoc PERMANOVAs, including sampling date, sampling site and their interaction, separately for each host plant species (or habitat for airborne samples). We then corrected the P-values for multiple testing, as described by Benjamini & Yekutieli (2001).

Results

Taxonomic description of foliar and airborne fungal communities

In total, we obtained 7 946 646 high-quality sequences, which clustered into 4 360 OTUs. Overall, 867 OTUs, corresponding to 4 600 179 sequences (57.9% of the raw OTU table) were not taxonomically assigned to fungi by BLAST. Among them, 4 451 913 sequences were assigned to plant sequences (Tracheophyta division), principally *Vitis* (59%), and *Carpinus* (35%) genus, showing that fITS7-ITS4 primers are not specific of fungi. These OTUs were removed. The negative controls contained 29 857 fungal sequences clustering into 337 OTUs. There is no consensus on how to deal with OTUs found in negative controls (Nguyen et al., 2015; Galan et al., 2016). It is difficult to distinguish real contaminations - sequences originating from the people who performed the experiments, the laboratory environment and the DNA extraction kit – from cross-contaminations between samples, occurring during the DNA extraction, amplification and sequencing (Esling et al., 2015; Galan et al., 2016). It is highly probable that OTUs assigned to *Erysiphe alphitoides*, the agent responsible for the oak powdery mildew (1.5% of the negative control sequences; Jakuschkin et al., 2016) or *Botrytis cinerea*, responsible for the grey mold on grapes (1.2%; Jaspers et al., 2015) are likely cross-

contaminations because they are strongly related to a specific host. Moreover, the removal of very abundant OTUs strongly altered the taxonomic composition of the samples, and removed some species known to be abundant on leaves such as *Aureobasidium pullulans*, known as very abundant on grapevine (Pinto & Gomes, 2016). We thus decided to retain all these OTUs in the dataset. Two samples containing very few sequences (<300 sequences) were removed. These samples corresponded to grapevine leaves collected at the Couhins site, in May. The first was collected in the centre of the vineyard, and the other was collected at its edge. Finally, the OTU table used for the analyses contained 196 samples and 3 487 fungal OTUs, corresponding to 3 316 156 sequences. The number of sequences per sample ranged from 424 to 96 276, with a mean of 16 919. This OTU table was used for taxonomical description. Richness, Bray-Curtis and Jaccard averaged indices were calculated over 100 rarefactions of this OTU table, at a threshold of 420 sequences per sample.

The fungal communities of bioaerosols and leaves from forest trees and grapevines were dominated by ascomycetes (Fig. 2). The sequences assigned to Ascomycota division accounted for 85.7% of all the sequences, followed by Basidiomycota division (11.3%). Overall, 3.0% of the total sequences remained unassigned at the division level. Airborne and foliar samples shared 1440 OTUs (Fig. 3), but there was a significant difference in the composition of foliar and airborne fungal communities (PERMANOVA $F=20.15$, $p=0.001$). The ten most abundant fungal OTUs were shared by airborne, forest foliar and grapevine foliar communities, but their relative abundance differed between each compartment (Table 1).

Variations in the richness of foliar and airborne fungal communities at the landscape scale

ANOVA revealed a significant effect of the interaction between host plant species and sampling date on the richness of foliar fungal communities (Table 2). Differences in fungal community richness between plant species were not significant in May and July (Fig. 4 and Fig. S1). In October, grapevine stocks had significantly less rich foliar fungal communities than oak (post-hoc tests: $P < 0.0001$; Fig. 4) and hornbeam trees ($P < 0.0001$), but the richness of their fungal communities did not differ significantly from that of chestnut trees ($P = 0.147$; Fig. S1). Hornbeam leaves harboured the richest communities of all the plant species considered (post-hoc tests: $P < 0.0001$ between hornbeam and chestnut, $P = 0.0003$ between hornbeam and oak, $P < 0.0001$ between hornbeam and grapevine; Fig. S1).

ANOVA post-hoc tests also revealed a significant decrease in fungal species richness in grapevine over the course of the growing season ($P < 0.0001$ for each pairwise comparison; Fig. 4). Seasonal variations in fungal richness were less marked in oak ($P = 0.081$, $P = 0.999$ and $P = 0.004$, respectively between May and July, July and October, May and October), chestnut ($P = 0.011$, $P = 0.997$ and $P = 0.0002$, respectively) and hornbeam ($P = 1.00$, $P = 0.144$ and $P = 0.185$, respectively).

ANOVA also revealed a significant effect of the interaction between host plant species and edge on the richness of foliar fungal communities (Table 2). The richness of foliar fungal communities was significantly higher at the edge in oak ($P = 0.002$), but not in hornbeam ($P = 0.100$), chestnut ($P = 0.139$), or grapevine ($P = 0.790$) (Fig. S2).

Habitat had a significant effect on the richness of airborne fungal communities (Table 2), which was significantly higher in forests than in vineyards.

Conclusions were similar on models performed without rarefaction (Supporting information, Table SI4).

Variations in the composition of foliar and airborne fungal communities at the landscape scale

PERMANOVA revealed a significant effect of the interaction between host plant species and sampling date on the composition of foliar fungal communities (Table 3). Bray-Curtis dissimilarities between oak and grapevine foliar fungal communities increased over the course of the growing season (mean \pm SD; 0.47 ± 0.07 in May, 0.67 ± 0.09 in July and 0.91 ± 0.06 in October). These results are illustrated by non-metric multidimensional scaling (NMDS; Fig. 3a). Bray-Curtis dissimilarities also increased between each pair of host species (Supporting information, Table S2 and Fig. S3a). Similar results were obtained with the Jaccard dissimilarity index (Supporting information, Table S3 and Fig. S3b).

PERMANOVA also revealed significant edge effects on the composition of foliar fungal communities, in interaction with host plant species and sampling date. Post-hoc PERMANOVAs computed separately for each host species indicated differences in community composition between the edge and centre of the forest for oak and hornbeam, in interaction with sampling date ($F=1.68$, $P=0.031$ and $F=1.85$, $P=0.044$, respectively). The composition of the fungal community did not differ between the edge and the centre of the habitat for chestnut ($F=2.27$, $P=0.25$) or grapevine ($F=0.92$, $P=1$). Finally, PERMANOVA analysis of Bray-Curtis dissimilarities revealed a significant effect of sampling date on bioaerosol composition (Table 3 and Fig. 3b). Similar results were obtained for Jaccard dissimilarity (Supporting information,

Table S3). Overall, similar results were also obtained without rarefying (Supporting information, Table S5).

273

274 Discussion

To our knowledge, this is the first time that the spatial structure and the temporal dynamic of foliar and airborne fungal communities are assessed simultaneously at the landscape scale. We studied a landscape mosaic consisting of two main habitats, vineyard and forest patches. We expected that repeated dispersal events between habitat patches would homogenize the foliar communities over the course of the growing season. We expected the homogenization to be greater along habitat edges, where grapevine stocks and forest trees are closer to each other.

Accordingly, we found that 26% of the OTUs are shared between airborne and foliar fungal communities. The most abundant ones are principally generalist species, such as *Aureobasidium pullulans*, *Cladosporium sp.* or *Eppicoccum nigrum*, which were already found as abundant in the microbiome of many species (Jumpponen & Jones, 2009; Zambell & White, 2014; Pinto & Gomes, 2016). This result confirms that many fungal species disperse through the atmosphere (Lindemann et al., 1982; Brown & Hovmöller, 2002; Bulgarelli et al., 2013). Moreover, while the richness of airborne fungal communities was higher in forest patches than in adjacent vineyards, their composition did not differ significantly, whatever the season. This lack of spatial variation in airborne fungal communities could account for the high similarity between foliar fungal communities of grapevine and forest tree species at the beginning of the growing season. Flushing leaves in May receive similar pools of fungal species through airborne dispersal, whatever the habitat and the host plant species. Our results suggest that dispersal of foliar fungal

communities is not limited at the landscape scale. Similar patterns were already observed at far larger spatial scales. The atmosphere is indeed considered as a continental and inter-continental corridor for the dispersal of microorganisms (Finlay, 2002; Brown & Hovmøller, 2002; Womack et al., 2010; Barberán et al., 2014), resulting in global patterns across continents. However, our results contrast with the strong dispersal limitation observed at smaller scale (Bowers et al., 2013). Peay et al. (2010) evidenced that ectomycorrhizal richness is lower in small tree patches located 1km away from large tree patches than nearer ones. Dickie & Reich (2005) showed that the abundance and richness of ectomycorrhizal fungi decreased up to 20m away from the forest edge. Galante et al. (2011) also showed that 95% of ectomycorrhizal spores fell within 58cm from the source. While the dispersal of ectomycorrhizal fungi can differ from the foliar ones because of the difference in the height of spore emission (Schmale & Ross, 2015), our failure to detect such dispersal limitation at low spatial scale can be explained by the short time of sampling of airborne communities (30 minutes), which can be insufficient to characterize properly the airborne fungal composition of the whole season.

Against expectation, we found that the composition of the foliar fungal communities of forest tree species and grapevine increasingly diverged from May to October. Besides, a severe decline in the richness of foliar fungal communities was observed in grapevine over the course of the growing season, but not in forest tree species. Despite an identical pool of airborne fungi in vineyards and forests, the selective pressures exerted on foliar fungal communities therefore differ between both habitats. These selective pressures can be exerted by several factors, including the host species, the microclimate and the agricultural practices. Host-specificity has been demonstrated in foliar fungal communities (Kembel & Mueller, 2014; Lambais et al., 2014; Meiser et al., 2014). Our results paralleled these findings: in forest patches, foliar fungal

communities significantly differ among tree species at the end of the growing season. Seasonal variations in leaf physiology could also account for the observed temporal variations in foliar communities, especially the richness decline in grapevine fungal foliar communities. Older grapevine leaves indeed produce larger amounts of phytoalexins and tend to be more resistant to pathogens (Steimetz et al., 2012). Selection by the habitat can also be exerted by the microclimate (Vacher et al., 2016). Harsher microclimatic conditions in vineyards than in forests, especially in the summer, could account for the decline in fungal species richness in vineyards but not in forests. Particularly, greater exposure to UV and higher air temperatures in vineyards could decrease the survival of foliar fungi on grapevine leaves. By contrast, tree cover provides a milder microclimate which could be more suitable to foliar micro-organisms. Finally, selection by the habitat can be exerted by agricultural practices. A few studies showed that fungicide applications can reduce the diversity and alter the composition of the foliar microbial community (Gu et al., 2010; Moulas et al., 2013; Cordero-Bueso et al., 2014; Karlsson et al., 2014). However, several other studies showed that the foliar fungal communities of grapevine are highly resilient to some chemical or biological pesticides (Walter et al., 2007; Perazzolli et al., 2014; Ottesen et al., 2015). Further research is required to assess the influence of fungicide applications on the observed decline in the richness of foliar fungal communities.

Our study also showed, for the first time, significant edge effects on foliar fungal community assemblages. A higher level of foliar fungal community richness was found in oak trees growing at the edge of the forest than in oak trees growing 100 m away. Significant differences in community composition between the edge and the centre of the forest were also found for oak and hornbeam. Variations in microclimate and leaf physiology along the forest edge (Chen et al., 1993; Zheng et al., 2005; Kunert et al., 2015) are more likely to account for this result than

species dispersal from vineyards to forest patches, since the foliar fungal communities of the two habitats diverged over the course of the growing season. The absence of edge effect in grapevine foliar fungal communities suggests that dispersal of fungal species from forests to vineyards has little influence on community composition and richness. This result contrasts with the findings of many studies on macro-organisms, reporting that dispersal events between managed and non-managed habitats shape communities and influence ecosystem functioning and services (Thomson & Hoffmann, 2009; Rusch et al., 2010; Thomson et al., 2010; Chaplin-Kramer et al., 2011; Blitzer et al., 2012).

Conclusions

Our results suggest that dispersal events between habitat patches are not major drivers of foliar fungal communities at the landscape scale. Selective pressures exerted in each habitat by the plant host, the microclimate and the agricultural practices play a greater role, and might account for the differentiation of foliar fungal communities between habitats. However, our experimental design does not allow us to assess the relative influence of each factor in shaping foliar fungal communities. Our results suggest that the leaves of broad-leaf species are colonised by similar pools of airborne micro-organisms at the beginning of the growing season. The composition of foliar fungal communities then diverges between habitat patches and between plant species within the same habitat. In contrast, airborne communities remain similar between habitats. Overall, our results support those of Redford et al. (2010) and Morrison-Whittle & Goddard (2015) which indicated that selection predominates over dispersal in structuring plant microbial communities.

361

362 **Aknowledgements**

363 We thank the vineyard managers for allowing us to carrying out these experiments in their
364 domains: Nicolas Lesaint in Château Reignac, Mathieu Arroyo in Château Couhins, and Antoine
365 Le Grix de la Salle in Château Grand-Verdus. We thank Morgane Petitgenet for assistance with
366 sample collection, and Olivier Fabreguettes and Jessica Vallance for providing help and advice
367 for DNA extraction and bioinformatic analyses. We thank the entire LGC genomics team for
368 assistance and advice concerning sequencing.

369

370 **References**

- 371 Bálint M., Schmidt P-A., Sharma R., Thines M., Schmitt I. 2014. An Illumina metabarcoding
372 pipeline for fungi. *Ecology and Evolution* 4:2642–2653.
- 373 Barberán A., Henley J., Fierer N., Casamayor EO. 2014. Structure, inter-annual recurrence, and
374 global-scale connectivity of airborne microbial communities. *Science of The Total*
375 *Environment* 487:187–195.
- 376 Bastian O. 2001. Landscape Ecology – towards a unified discipline? *Landscape Ecology* 16:757–
377 766.
- 378 Beckstead J., Meyer SE., Connolly BM., Huck MB., Street LE. 2010. Cheatgrass facilitates
379 spillover of a seed bank pathogen onto native grass species. *Journal of Ecology* 98:168–177.

- 380 Benjamini Y., Yekutieli D. 2001. The control of the false discovery rate in multiple testing under
381 dependency. *Annals of Statistics* 29:1165–1188.
- 382 Blitzer EJ., Dormann CF., Holzschuh A., Klein A-M., Rand TA., Tschamntke T. 2012. Spillover
383 of functionally important organisms between managed and natural habitats. *Agriculture,
384 Ecosystems & Environment* 146:34–43.
- 385 Bowers RM., Clements N., Emerson JB., Wiedinmyer C., Hannigan MP., Fierer N. 2013.
386 Seasonal Variability in Bacterial and Fungal Diversity of the Near-Surface Atmosphere.
387 *Environmental Science & Technology* 47:12097–12106.
- 388 Brown JKM., Hovmøller MS. 2002. Aerial Dispersal of Pathogens on the Global and
389 Continental Scales and Its Impact on Plant Disease. *Science* 297:537–541.
- 390 Bulgarelli D., Schlaeppi K., Spaepen S., van Themaat EVL., Schulze-Lefert P. 2013. Structure
391 and functions of the bacterial microbiota of plants. In: Merchant SS ed. *Annual Review of
392 Plant Biology, Vol 64*. Palo Alto: Annual Reviews, 807–838.
- 393 Chaplin-Kramer R., O'Rourke ME., Blitzer EJ., Kremen C. 2011. A meta-analysis of crop pest
394 and natural enemy response to landscape complexity. *Ecology Letters* 14:922–932.
- 395 Chen J., Franklin JF., Spies TA. 1993. Contrasting microclimates among clearcut, edge, and
396 interior of old-growth Douglas-fir forest. *Agricultural and Forest Meteorology* 63:219–237.
- 397 Cordero-Bueso G., Arroyo T., Valero E. 2014. A long term field study of the effect of fungicides
398 penconazole and sulfur on yeasts in the vineyard. *International Journal of Food Microbiology*
399 189:189–194.

- 400 Cordier T., Robin C., Capdevielle X., Fabreguettes O., Desprez-Loustau M-L., Vacher C. 2012.
401 The composition of phyllosphere fungal assemblages of European beech (*Fagus sylvatica*)
402 varies significantly along an elevation gradient. *New Phytologist* 196:510–519.
- 403 Dickie IA., Reich PB. 2005. Ectomycorrhizal fungal communities at forest edges. *Journal of*
404 *Ecology* 93:244–255.
- 405 Edgar RC., Haas BJ., Clemente JC., Quince C., Knight R. 2011. UCHIME improves sensitivity
406 and speed of chimera detection. *Bioinformatics* 27:2194–2200.
- 407 Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads.
408 *Nature Methods* 10:996–998.
- 409 Esling P., Lejzerowicz F., Pawlowski J. 2015. Accurate multiplexing and filtering for high-
410 throughput amplicon-sequencing. *Nucleic Acids Research* 43:2513–2524.
- 411 Finlay BJ. 2002. Global Dispersal of Free-Living Microbial Eukaryote Species. *Science*
412 296:1061–1063.
- 413 Galan M., Razzauti M., Bard E., Bernard M., Brouat C., Charbonnel N., Dehne-Garcia A.,
414 Loiseau A., Tatard C., Tamisier L., Vayssier-Taussat M., Vignes H., Cosson JF. 2016. 16S
415 rRNA amplicon sequencing for epidemiological surveys of bacteria in wildlife: the
416 importance of cleaning post-sequencing data before estimating positivity, prevalence and co-
417 infection. *bioRxiv*:39826.
- 418 Galante TE., Horton TR., Swaney DP. 2011. 95% of basidiospores fall within 1 m of the cap: a
419 field-and modeling-based study. *Mycologia* 103:1175–1183.

- 420 Gu L., Bai Z., Jin B., Hu Q., Wang H., Zhuang G., Zhang H. 2010. Assessing the impact of
421 fungicide enostroburin application on bacterial community in wheat phyllosphere. *Journal of*
422 *Environmental Sciences* 22:134–141.
- 423 Hanson CA., Fuhrman JA., Horner-Devine MC., Martiny JB. 2012. Beyond biogeographic
424 patterns: processes shaping the microbial landscape. *Nature Reviews Microbiology* 10:497–
425 506.
- 426 Herr DG. 1986. On the History of ANOVA in Unbalanced, Factorial Designs: The First 30
427 Years. *The American Statistician* 40:265–270.
- 428 Hughes JB., Hellmann JJ. 2005. The Application of Rarefaction Techniques to Molecular
429 Inventories of Microbial Diversity. In: Enzymology B-M in ed. Environmental Microbiology.
430 Academic Press, 292–308.
- 431 Hurst ZM., McCleery RA., Collier BA., Fletcher RJ Jr., Silvy NJ., Taylor PJ., Monadjem A.
432 2013. Dynamic Edge Effects in Small Mammal Communities across a Conservation-
433 Agricultural Interface in Swaziland. *PLoS ONE* 8:e74520.
- 434 Ihrmark K., Bodeker ITM., Cruz-Martinez K., Friberg H., Kubartova A., Schenck J., Strid Y.,
435 Stenlid J., Brandström-Durling M., Clemmensen KE., Lindahl BD. 2012. New primers to
436 amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and natural
437 communities. *FEMS Microbiology Ecology* 82:666–677.
- 438 Jakuschkin B., Fievet V., Schwaller L., Fort T., Robin C., Vacher C. 2016. Deciphering the
439 Pathobiome: Intra- and Interkingdom Interactions Involving the Pathogen Erysiphe
440 alphitoides. *Microbial Ecology*:1–11.

- 441 Jaspers MV., Seyb AM., Trought MCT., Balasubramaniam R. 2015. Necrotic grapevine material
442 from the current season is a source of *Botrytis cinerea* inoculum. *European Journal of Plant*
443 *Pathology* 144:811–820.
- 444 Jumpponen A., Jones KL. 2009. Massively parallel 454 sequencing indicates hyperdiverse
445 fungal communities in temperate *Quercus macrocarpa* phyllosphere. *New Phytologist*
446 184:438–448. DOI: 10.1111/j.1469-8137.2009.02990.x.
- 447 Karlsson I., Friberg H., Steinberg C., Persson P. 2014. Fungicide effects on fungal community
448 composition in the wheat phyllosphere. *PLoS ONE* 9:e111786.
- 449 Kembel SW., O'Connor TK., Arnold HK., Hubbell SP., Wright SJ., Green JL. 2014.
450 Relationships between phyllosphere bacterial communities and plant functional traits in a
451 neotropical forest. *Proceedings of the National Academy of Sciences* 111:13715–13720.
- 452 Kõljalg U., Nilsson RH., Abarenkov K., Tedersoo L., Taylor AFS., Bahram M., Bates ST., Bruns
453 TD., Bengtsson-Palme J., Callaghan TM., Douglas B., Drenkhan T., Eberhardt U., Dueñas
454 M., Grebenc T., Griffith GW., Hartmann M., Kirk PM., Kohout P., Larsson E., Lindahl BD.,
455 Lücking R., Martín MP., Matheny PB., Nguyen NH., Niskanen T., Oja J., Peay KG., Peintner
456 U., Peterson M., Põldmaa K., Saag L., Saar I., Schüßler A., Scott JA., Senés C., Smith ME.,
457 Suija A., Taylor DL., Telleria MT., Weiss M., Larsson K-H. 2013. Towards a unified
458 paradigm for sequence-based identification of fungi. *Molecular Ecology* 22:5271–5277.
- 459 Kunert N., Aparecido LMT., Higuchi N., Santos J dos., Trumbore S. 2015. Higher tree
460 transpiration due to road-associated edge effects in a tropical moist lowland forest.
461 *Agricultural and Forest Meteorology* 213:183–192.

- 462 Lacasella F., Gratton C., Felici SD., Isaia M., Zapparoli M., Marta S., Sbordon V. 2014.
463 Asymmetrical responses of forest and “beyond edge” arthropod communities across a forest–
464 grassland ecotone. *Biodiversity and Conservation* 24:447–465.
- 465 Lambais MR., Lucheta AR., Crowley DE. 2014. Bacterial community assemblages associated
466 with the phyllosphere, dermosphere, and rhizosphere of tree species of the atlantic forest are
467 host taxon dependent. *Microbial Ecology*:1–8.
- 468 Lindemann J., Constantinidou HA., Barchet WR., Upper CD. 1982. Plants as Sources of
469 Airborne Bacteria, Including Ice Nucleation-Active Bacteria. *Applied and Environmental*
470 *Microbiology* 44:1059–1063.
- 471 Lindow SE., Leveau JH. 2002. Phyllosphere microbiology. *Current Opinion in Biotechnology*
472 13:238–243.
- 473 Ma M., Hietala R., Kuussaari M., Helenius J. 2013. Impacts of edge density of field patches on
474 plant species richness and community turnover among margin habitats in agricultural
475 landscapes. *Ecological Indicators* 31:25–34.
- 476 Madden T. 2013. The BLAST sequence analysis tool. In: *The NCBI Handbook*. National Center
477 for Biotechnology Information (US).
- 478 McMurdie PJ., Holmes S. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data Is
479 Inadmissible. *PLoS Computational Biology* 10.
- 480 Meiser A., Bálint M., Schmitt I. 2014. Meta-analysis of deep-sequenced fungal communities
481 indicates limited taxon sharing between studies and the presence of biogeographic patterns.

- 482 *New Phytologist* 201:623–635.
- 483 Morris CE. 2001. Phyllosphere. In: *Encyclopedia of Life Sciences*. London: Nature Publishing
484 Group,.
- 485 Morrison-Whittle P., Goddard MR. 2015. Quantifying the relative roles of selective and neutral
486 processes in defining eukaryotic microbial communities. *The ISME Journal* 9:2003–2011.
- 487 Moulas C., Petsoulas C., Rousidou K., Perruchon C., Karas P., Karpouzas DG. 2013. Effects of
488 systemic pesticides imidacloprid and metalaxyl on the phyllosphere of pepper plants. *BioMed*
489 *Research International* 2013:1–8.
- 490 Nemergut DR., Schmidt SK., Fukami T., O'Neill SP., Bilinski TM., Stanish LF., Knelman JE.,
491 Darcy JL., Lynch RC., Wickey P., Ferrenberg S. 2013. Patterns and processes of microbial
492 community assembly. *Microbiology and Molecular Biology Reviews* 77:342–356.
- 493 Newton AC., Fitt BDL., Atkins SD., Walters DR., Daniell TJ. 2010. Pathogenesis, parasitism
494 and mutualism in the trophic space of microbe–plant interactions. *Trends in Microbiology*
495 18:365–373.
- 496 Newton AC., Gravouil C., Fountaine JM. 2010. Managing the ecology of foliar pathogens:
497 ecological tolerance in crops. *Annals of Applied Biology* 157:343–359.
- 498 Nguyen NH., Smith D., Peay K., Kennedy P. 2015. Parsing ecological signal from noise in next
499 generation amplicon sequencing. *New Phytologist* 205:1389–1393.
- 500 Nilsson RH., Veldre V., Hartmann M., Unterseher M., Amend A., Bergsten J., Kristiansson E.,
501 Ryberg M., Jumpponen A., Abarenkov K. 2010. An open source software package for

- 502 automated extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-throughput
- 503 community assays and molecular ecology. *Fungal Ecology* 3:284–287.
- 504 Norris RF., Kogan M. 2000. Interactions between weeds, arthropod pests, and their natural
- 505 enemies in managed ecosystems. *Weed Science* 48:94–158.
- 506 Ottesen AR., Gorham S., Pettengill JB., Rideout S., Evans P., Brown E. 2015. The impact of
- 507 systemic and copper pesticide applications on the phyllosphere microflora of tomatoes.
- 508 *Journal of the Science of Food and Agriculture* 95:1116–1125.
- 509 Peay KG., Garbelotto M., Bruns TD. 2010. Evidence of dispersal limitation in soil
- 510 microorganisms: Isolation reduces species richness on mycorrhizal tree islands. *Ecology*
- 511 91:3631–3640.
- 512 Perazzolli M., Antonielli L., Storari M., Puopolo G., Pancher M., Giovannini O., Pindo M.,
- 513 Pertot I. 2014. Resilience of the natural phyllosphere microbiota of the grapevine to chemical
- 514 and biological pesticides. *Applied and Environmental Microbiology* 80:3585–3596.
- 515 Pinto C., Gomes AC. 2016. Vitis vinifera microbiome: from basic research to technological
- 516 development. *BioControl*.
- 517 Power AG., Mitchell CE. 2004. Pathogen spillover in disease epidemics. *The American*
- 518 *Naturalist* 164:S79–S89.
- 519 Redford AJ., Bowers RM., Knight R., Linhart Y., Fierer N. 2010. The ecology of the
- 520 phyllosphere: geographic and phylogenetic variability in the distribution of bacteria on tree
- 521 leaves: Biogeography of phyllosphere bacterial communities. *Environmental Microbiology*

- 522 12:2885–2893.
- 523 Ruinen J. 1956. Occurrence of Beijerinckia species in the “phyllosphere.” *Nature* 177:220–221.
- 524 Rusch A., Valantin-Morison M., Sarthou J-P., Roger-Estrade J. 2010. Biological control of insect
- 525 pests in agroecosystems: effects of crop management, farming systems, and seminatural
- 526 habitats at the landscape scale: a review. In: Sparks DL ed. *Advances in Agronomy, Vol 109*.
- 527 219–259.
- 528 Schmale DGI., Ross SD. 2015. Highways in the Sky: Scales of Atmospheric Transport of Plant
- 529 Pathogens. *Annual Review of Phytopathology* 53:591–611.
- 530 Steimetz E., Trouvelot S., Gindro K., Bordier A., Poinssot B., Adrian M., Daire X. 2012.
- 531 Influence of leaf age on induced resistance in grapevine against *Plasmopara viticola*.
- 532 *Physiological and Molecular Plant Pathology* 79:89–96.
- 533 Thomson LJ., McKenzie J., Sharley DJ., Nash MA., Tsitsilas A., Hoffmann AA. 2010. Effect of
- 534 woody vegetation at the landscape scale on the abundance of natural enemies in Australian
- 535 vineyards. *Biological Control* 54:248–254.
- 536 Thomson LJ., Hoffmann AA. 2009. Vegetation increases the abundance of natural enemies in
- 537 vineyards. *Biological Control* 49:259–269.
- 538 Turner TR., James EK., Poole PS. 2013. The plant microbiome. *Genome Biol* 14:209.
- 539 Vacher C., Hampe A., Porté AJ., Sauer U., Compant S., Morris CE. 2016. The phyllosphere:
- 540 microbial jungle at the plant-climate interface. *Annual Review of Ecology, Evolution, and*
- 541 *Systematics* 47.

- 542 Vellend M. 2010. Conceptual synthesis in community ecology. *The Quarterly review of biology*
543 85:183–206.
- 544 Vorholt JA. 2012. Microbial life in the phyllosphere. *Nature Reviews Microbiology* 10:828–840.
- 545 Walter M., Frampton CM., Boyd-Wilson KSH., Harris-Virgin P., Waipara NW. 2007.
546 Agrichemical impact on growth and survival of non-target apple phyllosphere
547 microorganisms. *Canadian Journal of Microbiology* 53:45–55.
- 548 Whipps J m., Hand P., Pink D., Bending G d. 2008. Phyllosphere microbiology with special
549 reference to diversity and plant genotype. *Journal of Applied Microbiology* 105:1744–1755.
- 550 Wilson HE., Carroll GC., Roy BA., Blaisdell GK. 2014. Tall fescue is a potential spillover
551 reservoir host for *Alternaria* species. *Mycologia* 106:22–31.
- 552 Womack AM., Bohannan BJM., Green JL. 2010. Biodiversity and biogeography of the
553 atmosphere. *Philosophical Transactions of the Royal Society of London B: Biological*
554 *Sciences* 365:3645–3653.
- 555 Xu X-M., Jeffries P., Pautasso M., Jeger MJ. 2011. Combined use of biocontrol agents to
556 manage plant diseases in theory and practice. *Phytopathology* 101:1024–1031.
- 557 Zambell CB., White JF. 2014. In the forest vine *Smilax rotundifolia*, fungal epiphytes show site-
558 wide spatial correlation, while endophytes show evidence of niche partitioning. *Fungal*
559 *Diversity* 75:279–297.
- 560 Zheng D., Chen J., LeMoine JM., Euskirchen ES. 2005. Influences of land-use change and edges
561 on soil respiration in a managed forest landscape, WI, USA. *Forest Ecology and Management*

562 215:169–182.

563

Tables

Table 1 Taxonomic assignment of the 10 most abundant OTUs by the online BLAST analysis against the GenBank database. The environmental and metagenome sequences were excluded. Identity is the percentage identity between the OTU representative sequence and the closest matching sequence in GenBank. Taxa shown as unassigned at the species level (*sp.*) indicate OTUs assigned to at least two species of the same genus with identical e-value. Relative abundance are percentage of abundance of each data subset and brackets contain the rank of the OTU in each data subset.

| GI number | Closest match | | Relative abundance in percent (rank) | | | |
|------------|---------------|--------------------------------------|--------------------------------------|----------|---------------|------------------|
| | Identity | Putative taxon | Total | Airborne | Forest leaves | Grapevine leaves |
| 1034220623 | 100 | <i>Aureobasidium pullulans</i> | 15.48 | 3.8 (4) | 12.6 (1) | 55.9 (1) |
| 1031917897 | 100 | <i>Cladosporium sp.</i> | 8.01 | 29.8 (1) | 2.7 (11) | 2.4 (5) |
| 1049480240 | 85.6 | <i>Collophora hispanica</i> | 5.64 | 1.7 (7) | 7.4 (2) | 1.1 (13) |
| 61619908 | 100 | <i>Ramularia endophylla</i> | 4.72 | 0.6 (20) | 6.4 (3) | 1.4 (12) |
| 1035371449 | 100 | <i>Cladosporium sp.</i> | 4.51 | 13.7 (2) | 2.3 (13) | 1.8 (7) |
| 530746702 | 100 | <i>Stromatoseptoria castaneicola</i> | 3.48 | 0.3 (31) | 4.8 (4) | 0.9 (15) |
| 626419142 | 99.5 | <i>Taphrina carpini</i> | 3.35 | 1.3 (9) | 4.3 (6) | 0.7 (19) |
| 1024249962 | 100 | <i>Erysiphe sp.</i> | 3.17 | 0.3 (33) | 4.4 (5) | 0.8 (16) |
| 61619940 | 100 | <i>Naevula minutissima</i> | 2.99 | 1.2 (10) | 3.8 (8) | 0.7 (20) |
| 961502090 | 91.0 | <i>Zeloasperisporium searsiae</i> | 2.93 | 0.2 (46) | 4.1 (7) | 0.6 (21) |

Table 2 Effect of sampling date (May, July or October), host species (oak, hornbeam, chestnut or grapevine) or habitat (vineyard or forest), edge (habitat centre or center) and their interaction on OTU richness in foliar and airborne fungal communities, assessed using a type III ANOVA. In both models, sampling site was included as a random variable. R_m^2 is the marginal coefficient of determination (for fixed effects) and R_c^2 the conditional coefficient of determination (for fixed and random effects). Bold values are the significant ones.

| | F | P-value | R_m^2 (R_c^2) |
|-----------------------|-------|------------------|---------------------|
| Foliar OTU richness | | | |
| Date | 44.49 | <0.001 | 0.64 (0.71) |
| Species | 14.97 | <0.001 | |
| Edge | 17.21 | <0.001 | |
| D x S | 23.42 | <0.001 | |
| D x E | 0.11 | 0.894 | |
| S x E | 6.72 | <0.001 | |
| D x S x E | 1.13 | 0.347 | |
| Airborne OTU richness | | | |
| Date | 1.07 | 0.362 | 0.34 (0.52) |
| Habitat | 10.19 | 0.004 | |
| Edge | 4.20 | 0.052 | |
| D x H | 0.86 | 0.436 | |
| D x E | 1.40 | 0.267 | |
| H x E | 0.01 | 0.912 | |
| D x H x E | 1.678 | 0.209 | |

579

Table 3 Effect of sampling date (May, July or October), host species (oak, hornbeam, chestnut or grapevine) or habitat (vineyard or forest), edge (habitat centre or center) and their interaction on the composition of foliar and airborne fungal communities, assessed using a PERMANOVA. In both models, sampling site was included as a stratification variable. Bold values are the significant ones.

| | F | R ² | P-value |
|---------------------------------------|-------|----------------|--------------|
| Foliar fungal community composition | | | |
| Date | 10.13 | 0.078 | 0.001 |
| Species | 13.70 | 0.158 | 0.001 |
| Edge | 3.94 | 0.015 | 0.001 |
| D x Sp | 6.92 | 0.160 | 0.001 |
| D x E | 2.05 | 0.016 | 0.007 |
| Sp x E | 2.22 | 0.026 | 0.001 |
| D x Sp x E | 1.08 | 0.025 | 0.239 |
| Airborne fungal community composition | | | |
| Date | 2.94 | 0.157 | 0.001 |
| Habitat | 1.54 | 0.041 | 0.062 |
| Edge | 0.68 | 0.018 | 0.827 |
| D x H | 0.95 | 0.051 | 0.418 |
| D x E | 0.66 | 0.035 | 0.938 |
| H x E | 0.77 | 0.020 | 0.684 |
| D x H x E | 0.71 | 0.038 | 0.878 |

Figures

Figure 1: Experimental design. (a) Geographical position of the three sampling sites, represented by red points. (b) Sampling design at each site. Leaf pictograms represent the sampling location of leaves in each site. Three leaves per plant species (i.e. grapevine in the vineyard and oak *plus* chestnut or hornbeam in the forest patch) were sampled at each location. Cloud pictograms represent the sampling location of airborne communities.






Figure 2 Taxonomic composition of the airborne and foliar fungal communities in forest and vineyard habitats. The inner disc shows the proportion of sequences assigned to each taxonomic division, and the outer disc the proportion of sequences assigned to each class of the Ascomycota and Basidiomycota divisions.



599 **Figure 3** Venn diagramm giving the number of OTUs shared between the airborne, forest foliar
600 and vineyard foliar communities.

601



602

603 **Figure 4** Richness of foliar fungal community in oak (green) and grapevine (red), depending on
604 the sampling date. Error bars represent the standard error.




Figure 5 NMDS representing dissimilarities in the composition of fungal communities. (a) Dissimilarities in the composition of foliar fungal communities between the host species (oak in green and grapevine in red), depending on the sampling date. The other two forest species are not shown here, to make the figure easier to read, and are presented in Fig. S2. The stress value associated with this representation was 0.170. (b) Airborne fungal communities between the habitat (forest in light-blue and vineyard in dark-blue), depending on the sampling date. The stress value associated with this representation was 0.188. Dissimilarities between samples were computed with the Bray-Curtis index, averaged over 100 random rarefactions of the OTU table. The confidence ellipsoid at the 0.68 level is shown, for all combinations of these two factors.

