

A peer-reviewed version of this preprint was published in PeerJ on 28 February 2017.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.3044) (peerj.com/articles/3044), which is the preferred citable publication unless you specifically need to cite this preprint.

O'Donnell JL, Kelly RP, Shelton AO, Samhouri JF, Lowell NC, Williams GD. 2017. Spatial distribution of environmental DNA in a nearshore marine habitat. PeerJ 5:e3044 <https://doi.org/10.7717/peerj.3044>

Spatial distribution of environmental DNA in a nearshore marine habitat

James L O'Donnell ^{Corresp., 1}, Ryan P Kelly ¹, Andrew Olaf Shelton ^{2,3}, Jameal F Samhoury ², Natalie C Lowell ^{1,4}, Gregory D Williams ⁵

¹ School of Marine and Environmental Affairs, University of Washington, Seattle, Washington, United States of America

² Northwest Fisheries Science Center, NOAA Fisheries, Seattle, Washington, United States of America

³ Earth resources technology

⁴ School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, United States of America

⁵ Pacific States Marine Fisheries Commission under contract to, Northwest Fisheries Science Center, Seattle, Washington, United States of America

Corresponding Author: James L O'Donnell

Email address: jimmyod@uw.edu

In the face of increasing threats to biodiversity, the advancement of methods for surveying biological communities is a major priority for ecologists. Recent advances in molecular biological technologies have made it possible to detect and sequence DNA from environmental samples (environmental DNA or eDNA); however, eDNA techniques have not yet seen widespread adoption as a routine method for biological surveillance primarily due to gaps in our understanding of the dynamics of eDNA in space and time. In order to identify the effective spatial scale of this approach in a dynamic marine environment, we collected marine surface water samples from transects ranging from the intertidal zone to 4 kilometers from shore. Using massively parallel sequencing of 16S amplicons, we identified a diverse community of metazoans and quantified their spatial patterns using a variety of statistical tools. We find evidence for multiple, discrete eDNA communities in this habitat, and show that these communities decrease in similarity as they become further apart. Offshore communities tend to be richer but less even than those inshore, though diversity was not spatially autocorrelated. Taxon-specific relative abundance coincided with our expectations of spatial distribution in taxa lacking a microscopic, pelagic life-history stage, though most of the taxa detected do not meet these criteria. Finally, we use carefully replicated laboratory procedures to show that laboratory treatments were remarkably similar in most cases, while allowing us to detect a faulty replicate, emphasizing the importance of replication to metabarcoding studies. While there is much work to be done before eDNA techniques can be confidently deployed as a standard method for ecological monitoring, this study serves as a first analysis of diversity at the fine spatial scales relevant to marine ecologists and confirms the promise of eDNA in dynamic environments.

Spatial distribution of environmental DNA in a nearshore marine habitat

James L. O'Donnell*¹, Ryan P. Kelly¹, Andrew O. Shelton², Jameal F. Samhouri², Natalie
C. Lowell^{1,3}, and Gregory D. Williams⁴

¹School of Marine and Environmental Affairs, University of Washington, 3707 Brooklyn
Ave NE, Seattle, Washington 98105, USA

²Northwest Fisheries Science Center, NOAA Fisheries, 2725 Montlake Blvd E, Seattle,
Washington 98112, USA

³School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat St,
Seattle, Washington 98105, USA

⁴Pacific States Marine Fisheries Commission, Under contract to the Northwest Fisheries
Science Center, NOAA Fisheries, 2725 Montlake Blvd E, Seattle, WA 98112

November 22, 2016

Keywords

metagenomics, metabarcoding, environmental monitoring, molecular ecology, marine, estuarine

*jodonnellbio@gmail.com

1 Abstract

2 In the face of increasing threats to biodiversity, the advancement of methods for surveying biological
3 communities is a major priority for ecologists. Recent advances in molecular biological technologies
4 have made it possible to detect and sequence DNA from environmental samples (environmental DNA
5 or eDNA); however, eDNA techniques have not yet seen widespread adoption as a routine method
6 for biological surveillance primarily due to gaps in our understanding of the dynamics of eDNA in
7 space and time. In order to identify the effective spatial scale of this approach in a dynamic marine
8 environment, we collected marine surface water samples from transects ranging from the intertidal
9 zone to 4 kilometers from shore. Using massively parallel sequencing of 16S amplicons, we identified
10 a diverse community of metazoans and quantified their spatial patterns using a variety of statistical
11 tools. We find evidence for multiple, discrete eDNA communities in this habitat, and show that
12 these communities decrease in similarity as they become further apart. Offshore communities tend
13 to be richer but less even than those inshore, though diversity was not spatially autocorrelated.
14 Taxon-specific relative abundance coincided with our expectations of spatial distribution in taxa
15 lacking a microscopic, pelagic life-history stage, though most of the taxa detected do not meet
16 these criteria. Finally, we use carefully replicated laboratory procedures to show that laboratory
17 treatments were remarkably similar in most cases, while allowing us to detect a faulty replicate,
18 emphasizing the importance of replication to metabarcoding studies. While there is much work to
19 be done before eDNA techniques can be confidently deployed as a standard method for ecological
20 monitoring, this study serves as a first analysis of diversity at the fine spatial scales relevant to
21 marine ecologists and confirms the promise of eDNA in dynamic environments.

22 Introduction

23 The patterns and causes of variability in ecological communities across space are both seminal and
24 contentious areas of study in ecology (Hubbell, 2001; Anderson et al., 2011). One consistently
25 observed pattern of community spatial heterogeneity is that communities close to one another tend
26 to be more similar than those that are farther apart (Nekola and White, 1999). This decrease
27 in community similarity with increasing spatial separation is called distance decay and has been
28 reported from communities of tropical trees (Condit, 2002; Chust et al., 2006), ectomycorrhizal fungi

29 (Bahram et al., 2013), salt marsh plants (Guo et al., 2015), and microorganisms (Martiny et al.,
30 2011; Chust et al., 2013; Wetzel et al., 2012; Bell, 2010). Typically, this relationship is assessed by
31 regressing a measure of community similarity against a measure of spatial separation for a set of
32 sites at which a set of species' abundances (or presences) is calculated. Yet no existing biodiversity
33 survey method completely censuses all of the organisms in a given area. The lack of a single 'silver
34 bullet' method of sampling contributes inconclusiveness to the study of spatial patterning in ecology
35 (Levin, 1992), and leaves open the possibility of new and more comprehensive methods.

36 From a boat or aircraft, scientists can count whales by sight, but not the krill on which they
37 feed. For example, towed fishing nets can efficiently sample organisms larger than the mesh and
38 slower than the boat, but overlook viruses and have undesirable effects on charismatic air-breathing
39 species. However, DNA-based surveys show great promise as an efficient technique for detecting a
40 previously unthinkable breadth of organisms from a single sample.

41 Microbiologists have used nucleic acid sequencing to quantify the composition and function of
42 microbial communities in a wide variety of habitats (Handelsman et al., 1998; Tyson et al., 2004;
43 Venter et al., 2004; Iverson et al., 2012). To do so, microorganisms are collected in a sample of
44 environmental medium (e.g. water), their DNA or RNA is isolated and sequenced, and the identity
45 and abundance of sequences is considered to reflect the community of organisms contained in the
46 sample, which indirectly estimates the quantity of organisms in an area.

47 Macroorganisms shed DNA-containing cells into the environment (environmental DNA or eDNA)
48 that can be sampled in the same way (Ficetola et al., 2008; Thomsen et al., 2012). Potentially, eDNA
49 methods allow a broad swath of macroorganisms to be surveyed from basic environmental samples.
50 However, the accuracy and reliability of indirect estimates of macroorganismal abundance has been
51 debated because the entire organisms are not contained within the sample (Coward et al., 2015).
52 Concern surrounding eDNA methods is rooted in uncertainty about the attributes of eDNA in the
53 environment relative to actual organisms (Shelton et al., 2016; Evans et al., 2016). Basic questions
54 such as how long DNA can persist in that environment and how far DNA can travel remain largely
55 unknown (but see Klymus et al. (2015); Turner et al. (2015); Strickler et al. (2015); Deiner and
56 Altermatt (2014)) and impede inference about local organismal presence from an environmental
57 sample. As a result, estimating the spatial and temporal resolution of eDNA studies in the field is
58 a key step in making these methods practical.

59 The relationship between local organismal abundance and eDNA is further complicated in habi-
60 tats where the environmental medium itself may transport eDNA away from its source. We know
61 that genetic material can move away from its source precisely because organisms can be detected
62 indirectly without being present in the sample (Kelly et al., 2016). One might reasonably expect
63 eDNA to travel farther in a highly dynamic fluid such as the open ocean or flowing river than it
64 would through the sediment at the bottom of a stagnant pond (Deiner and Altermatt, 2014; Shogren
65 et al., 2016). Yet even studies of extremely dynamic habitats such as coastlines with high wave en-
66 ergy have found remarkable evidence that eDNA transport is limited enough that DNA methods
67 can detect differences among communities separated by less than 100 meters (Port et al., 2016).

68 While rigorous laboratory studies have investigated the effects of some environmental factors on
69 eDNA persistence (Klymus et al., 2015; Barnes et al., 2014; Sassoubre et al., 2016) and the transport
70 of eDNA in specific contexts (Deiner and Altermatt, 2014), we suggest that field studies comparing
71 the spatial distribution of communities of eDNA with expectations based on prior knowledge of
72 organisms' distributions are also critical to developing a working understanding of eDNA in the real
73 world.

74 We apply methods derived from community ecology to understand spatial patterns and patchi-
75 ness of eDNA. The underlying mechanism thought to drive the slope of the distance decay relation-
76 ship in ecological communities is the rate of movement of individuals among sites, which may be
77 driven by underlying processes such as habitat suitability. Because eDNA is shed and transported
78 away from its source, the increased movement of eDNA particles should homogenize community
79 similarity, and thus erode the distance decay relationship of eDNA communities.

80 Puget Sound is a deep, narrow fjord in Washington, USA, where a narrow band of shallow
81 bottom hugs the shoreline and abruptly gives way to a central depth of up to 300 meters. This
82 form allows the juxtaposition of communities associated with distinctly different habitats: shallow,
83 intertidal benthos, and euphotic pelagic (Burns, 1985). At the upper reaches of the intertidal, the
84 shoreline substrate varies from soft, fine sediment to cobble and boulder rubble. Soft intertidal
85 sediments are inhabited by burrowing bivalves (Bivalvia), segmented worms (Annelida), and acorn
86 worms (Enteropneusta), and in some lower intertidal and high subtidal ranges by eelgrass (*Zostera*
87 *marina*) (Kozloff, 1973; Dethier, 2010). Eelgrass meadows harbor epifaunal and infaunal biota,
88 and attract transient species which use the meadows for shelter and to feed on resident organisms.

89 Hard intertidal surfaces support a well-documented biota including barnacles (Sessilia), mussels
90 (Bivalvia:Mytilidae), anemones (Actinaria), sea stars (Asteroidea), urchins (Echinoidea), Bryozoans
91 (Ectoprocta), crustaceans (Decapoda), and a variety of algae (Dethier, 2010). Hard bottoms of the
92 lower intertidal and high subtidal are home to macroalgae such as Laminariales and Desmarestiales
93 which provides habitat for a distinct community of fish and invertebrates. The upper pelagic is
94 home to a diverse assemblage of microscopic plankton including diatoms and larvae (Strickland,
95 1983), as well as transitory fish and marine mammals.

96 We took advantage of this setting to explore the spatial variation and distribution of marine
97 eDNA communities. Using PCR-based methods and massively parallel sequencing, we surveyed
98 mitochondrial 16S sequences from a suite of marine animals in water samples collected over a grid
99 of sites extending from the shoreline out to 4 kilometers offshore in Puget Sound, Washington, USA.
100 We leverage this sampling design to perform the first explicitly spatial analysis of eDNA-derived
101 community similarity. We investigate two primary objectives. First we examine the spatial pattern-
102 ing of eDNA and determine the degree to which eDNA community similarity can be predicted by
103 physical proximity. We expect that physical proximity will be a strong predictor of community sim-
104 ilarity, and that community differences can be detected over small distances. Second, we examine
105 the distribution of diversity from eDNA data, and compare it to our expectations based on distri-
106 butions of microbial communities. We expect that distinct eDNA communities exist in this setting,
107 and that their spatial distribution coincides with that of adult microbial organisms. Because of the
108 vastly different communities of benthic microbial metazoans as a function of distance from shore,
109 we expect that more than one eDNA community is present across our 4 kilometer sampling grid,
110 and that communities change as a function of distance from shore. For this reason, we examine two
111 diversity measures of eDNA communities that have been widely used to reveal broad scale patterns
112 based on macrobiota in many ecological systems. Finally, we identify the taxa represented in the
113 eDNA communities, which span a range of life-history characteristics, and we expect that the spatial
114 distribution of eDNA will most closely resemble the distribution of adults in taxa with low dispersal
115 potential.

116 **Methods**

117 There are seven discrete steps to our methodology: (1) Environmental sample collection, (2) isolation
118 of particulates from water via filtration, (3) isolation of DNA from filter membrane, (4) amplification
119 of target locus via PCR, (5) sequencing of amplicons, (6) bioinformatic translation of raw sequence
120 data into tables of sequence abundance among samples, and (7) community ecological analyses of
121 eDNA. We provide brief overviews of these steps here, and encourage the reader to review the fully
122 detailed methods presented in the supplementary material (Supplemental Material).

123 **Environmental Sampling**

124 Starting from lower-intertidal patches of *Zostera marina*, we collected water samples at 1 meter
125 depth from 8 points (0, 75, 125, 250, 500, 1000, 2000, and 4000 meters) along three parallel transects
126 separated by 1000 meters (24 sample locations total; Figure 1). Samples were collected by attaching
127 bottles to a PVC pole and lowering it over the side of a boat over the span of one hour on 27 June
128 2014. To destroy residual DNA on equipment used for field sampling and filtration, we washed
129 with a 1:10 solution of household bleach (8.25% sodium hypochlorite; 7.25% available chlorine) and
130 deionized water, followed by thorough rinsing with deionized water. Each environmental sample
131 was collected in a clean 1 liter high-density polyethylene bottle, the opening of which was covered
132 with 500 micrometer nylon mesh to prevent entry of larger particles. Immediately after collecting
133 the sample, the mesh was replaced with a clean lid and the sample was held on ice until filtering.

134 **Filtration**

135 One liter from each water sample was filtered in the lab on a clean polysulfone vacuum filter
136 holder fitted with a 47 millimeter diameter cellulose acetate membrane with 0.45 micrometer pores.
137 Filter membranes were moved into 900 microliters of Longmire buffer (Longmire et al., 1997) using
138 clean forceps and stored at room temperature (Renshaw et al., 2014). To test for the extent of
139 contamination attributable to laboratory procedures, we filtered three replicate 1 liter samples of
140 deionized water. These samples were treated identically to the environmental samples throughout
141 the remaining protocols.

142 DNA Purification

143 DNA was purified from the membrane following a phenol:chloroform:isoamyl alcohol protocol follow-
144 ing Renshaw (Renshaw et al., 2014). Preserved membranes were incubated at 65C for 30 minutes
145 before adding 900 microliters of phenol:chloroform:isoamyl alcohol and shaking vigorously for 60
146 seconds. We conducted two consecutive chloroform washes by centrifuging at 14,000 rpm for 5
147 minutes, transferring the aqueous layer to 700 microliters chloroform, and shaking vigorously for 60
148 seconds. After a third centrifugation, 500 microliters of the aqueous layer was transferred to tubes
149 containing 20 microliters 5 molar NaCl and 500 microliters 100% isopropanol, and frozen at -20C
150 for approximately 15 hours. Finally, all liquid was removed by centrifuging at 14000 rpm for 10
151 minutes, pouring off or pipetting out any remaining liquid, and drying in a vacuum centrifuge at
152 45C for 15 minutes. DNA was resuspended in 200 microliters of ultrapure water. Four replicates of
153 genomic DNA extracted from tissue of a species absent from the sampled environment (*Oreochromis*
154 *niloticus*) served as positive control for the remaining protocols.

155 PCR Amplification

156 From each DNA sample, we amplified an approximately 115 base pair (bp) region of the mito-
157 chondrial gene encoding 16S RNA using a two-step polymerase chain reaction (PCR) protocol
158 described by O'Donnell et al. (2016). In the first set of reactions, primers were identical in ev-
159 ery reaction (forward: AGTTACYYTAGGGATAACAGCG; reverse: CCGGTCTGAACTCAGAT-
160 CAYGT); primers in the second set of reactions included these same sequences but with 3 variable
161 nucleotides (NNN) and an index sequence on the 5' end (see Sequencing Metadata). We used the
162 program OligoTag (Coissac, 2012) to generate 30 unique 6-nucleotide index sequences differing by
163 a minimum Hamming distance of 3 (see Sequencing Metadata). Indexed primers were assigned to
164 samples randomly, with the identical index sequence on the forward and reverse primer to avoid
165 errors associated with dual-indexed multiplexing (Schnell et al., 2015). In a UV-sterilized hood,
166 we prepared 25 microliter reactions containing 18.375 microliters ultrapure water, 2.5 microliters
167 10x buffer, 0.625 microliters deoxynucleotide solution (8 millimolar), 1 microliter each forward and
168 reverse primer (10 micromolar, obtained lyophilized from Integrated DNA Technologies (Coralville,
169 IA, USA)), 0.25 microliters Qiagen HotStar Taq polymerase, and 1.25 microliter genomic eDNA

170 template at 1:100 dilution in ultrapure water. PCR thermal profiles began with an initialization
171 step (95C; 15 min) followed by cycles (40 and 20 for the first and second reaction, respectively) of
172 denaturation (95C; 15 sec), annealing (61C; 30 sec), and extension (72C; 30 sec). 20 identical PCRs
173 were conducted from each DNA extract using non-indexed primers; these were pooled into 4 groups
174 of 5 in order to ensure ample template for the subsequent PCR with indexed primers. In order to
175 isolate the fragment of interest from primer dimer and other spurious fragments generated in the
176 first PCR, we used the AxyPrep Mag FragmentSelect-I kit with solid-phase reversible immobiliza-
177 tion (SPRI) paramagnetic beads at 2.5x the volume of PCR product (Axygen BioSciences, Corning,
178 NY, USA). A 1:5 dilution in ultrapure water of the product was used as template for the second
179 reaction. PCR products of the second reaction were purified using the Qiagen MinElute PCR Pu-
180 rification Kit (Qiagen, Hilden, Germany). Ultrapure water was used in place of template DNA and
181 run along with each batch of PCRs to serve as a negative control for PCR; none of these produced
182 visible bands on an agarose gel. In total, four separate replicates from each of 31 DNA samples
183 were carried through the two-step PCR process for a total of 124 sequenced PCR products. These
184 were combined with additional samples from other projects, totaling 345 samples for sequencing.

185 DNA Sequencing

186 Up to 30 PCR products were combined according to their primer index in equal concentration into
187 one of 14 pools, and 150 nanograms from each were prepared for library sequencing using the KAPA
188 high-throughput library prep kit with real-time library amplification protocol (KAPA Biosystems,
189 Wilmington, MA, USA). Each of these ligated sequencing adapters included an additional 6 base
190 pair index sequence (NEXTflex DNA barcodes; BIOO Scientific, Austin, TX, USA). Thus, each
191 PCR product was identifiable via its unique combination of index sequences in the sequencing
192 adapters and primers. Fragment size distribution and concentration of each library was quantified
193 using an Agilent 2100 BioAnalyzer. Libraries were pooled in equal concentrations and sequenced
194 for 150 base pairs in both directions (PE150) using an Illumina NextSeq at the Stanford Functional
195 Genomics Facility (machine NS500615, run 115, flowcell H3LFLAFX), where 20% PhiX Control
196 v3 was added to act as a sequencing control and to enhance sequencing depth. Raw sequence data
197 in fastq format is publicly available (see Data Availability).

198 **Sequence Data Processing (Bioinformatics)**

199 Detailed bioinformatic methods are provided in the supplemental material, and analysis scripts
200 used from raw sequencer output onward can be found in the public project directory (see Analysis
201 Scripts). Briefly, we performed five steps to process the sequence data: (1) Merge paired-end
202 reads, (2) eliminate low-quality reads, (3) eliminate PCR artifacts (chimeras), (4) cluster reads by
203 similarity into operational taxonomic units (OTUs), and (5) match observed sequences to taxon
204 names. Additionally, we checked for consistency among PCR replicates, excluded extremely rare
205 sequences, and rescaled (rarefied) the data to account for differences in sequencing depth. The data
206 for input to further analyses are a contingency table of the mean count of unique sequences, OTUs,
207 or taxa present in each environmental sample.

208 **Ecological Analyses**

209 After gathering the data, we use the eDNA community observed at each location to make inferences
210 about the spatial patterning of eDNA communities. We use statistical tools from community ecology
211 to assess the spatial structure of eDNA communities. We report similarity (1- dissimilarity) rather
212 than dissimilarity in all cases for ease of interpretation.

213 **Objective 1: Community similarity as a function of distance**

214 **Distance Decay**

215 To address our first objective and determine whether or not nearby samples are more similar than
216 distant ones, we fit a nonlinear model to represent decreasing community similarity with distance.
217 We calculated the pairwise Bray-Curtis similarity (1 - Bray-Curtis dissimilarity) between eDNA
218 communities using the R package *vegan* (Oksanen et al., 2016) and the great circle distance between
219 sampling points using the Haversine method as implemented by the R package *geosphere* (Hijmans,
220 2016). This model is similar to the Michaelis-Menten function, but with an asymptote fixed at 0:

$$y_{ij} = \frac{AB}{B + x_{ij}} \quad (1)$$

221 Where the relationship between community similarity (y_{ij}) and spatial distance (x_{ij}) between

222 observations i and j is determined by the similarity of samples at distance 0 (A), and the distance at
223 which half the total change in similarity is achieved (B). This allows for a samples collected very close
224 together (near 0) to have similarity significantly less than one. We assessed model fit using the R
225 function `nls` (R Core Team, 2016), using the `nl2sol` algorithm from the `Port` library to solve separable
226 nonlinear least squares using analytically computed derivatives (<http://netlib.org/port/nsg.f>). We
227 set bounds of 0 and 1 for the intercept parameter and a lower bound of 0 for the distance at half
228 similarity; starting values of these parameters were 0.5 and $x_{max}/2$, respectively. We calculated
229 a 95% confidence interval for the parameters and the predicted values using a first-order Taylor
230 expansion approach implemented by the function `predictNLS` in the R package `propagate` (Spiess,
231 2014).

232 There are other conceptually reasonable forms to expect the space-by-similarity relationship
233 to take; we present these in the supplemental material along with alternative data subsets and
234 similarity indices (see Supplemental Material).

235 **Objective 2: Spatial distribution of diversity**

236 **Community Classification**

237 To determine the spatial distribution and variation of eDNA communities (objective 2), we used
238 multivariate classification algorithms. We simultaneously assessed the existence of distinct com-
239 munity types and the membership of samples to those community types using an unsupervised
240 classification algorithm known as partitioning around medoids (PAM; sometimes referred to as k -
241 medoids clustering) (Kaufman and Rousseeuw, 1990), as implemented in the R package `cluster`
242 (Maechler et al., 2016). The classification of samples to communities was made on the basis of
243 their pairwise Bray-Curtis similarity, calculated using the function `vegdist` in the R package `vegan`
244 (Oksanen et al., 2016). Other distance metrics were evaluated but had no appreciable effect on the
245 outcome of the analysis (Figure 8). In order to chose an optimal number of clusters (K), we evalu-
246 ated the distribution of silhouette widths, a measure of the similarity between each sample and its
247 cluster compared to its similarity to other clusters. We repeated the analysis using fuzzy clustering
248 (FANNY, (Kaufman and Rousseeuw, 1990); however, the results were qualitatively similar to the
249 results using PAM so we omit them here.

250 **Aggregate Measures of Diversity**

251 We calculated two measures of diversity, richness and evenness, to ask if aggregate metrics of the
252 eDNA community showed evidence of spatial patterning. Richness is a measure of the number of
253 distinct types of organisms present and so ranges from 1 (only one taxon observed) to S , the number
254 of taxa observed across all samples. To calculate the evenness of the distribution of abundance of
255 taxa in a sample, we used the complement of the Simpson (1949) index ($1 - \sum p_i^2$, where p_i is the
256 proportional abundance of taxon i). The values of this index ranges from 0 to 1, with the value
257 interpreted as the probability that two sequences randomly selected from the sample will belong to
258 different taxa; thus, larger values of the index indicate more evenly divided communities (Magurran,
259 2003). We calculated Moran's I for both diversity metrics to test for spatial autocorrelation. We
260 also tested for a linear effect of log-transformed distance from shore on each measure of diversity to
261 ask how diversity changes over this strong environmental gradient.

262 **Taxon and Life History Patterns**

263 After assigning taxon names to the abundance data, we plotted the distribution in space of a
264 selection of taxa to compare with our expectations on the basis of adult distributions (objective 2).
265 Our aim was to understand where each taxon occurred in the greatest proportional abundance, and
266 its distribution in space relative to that maximum. Thus, we rescaled each sample to proportional
267 abundance, extracted the data from a single taxon, and scaled those values between 0 and 1.
268 We collated life history characteristics for each of the major taxonomic groups recovered, including
269 dispersal range of the gametes, larvae, and adults, adult habitat type and selectivity, and adult body
270 size. Dispersal range was given as an order-of-magnitude approximation of the scale of dispersal:
271 for example, internally fertilized species were assigned a gamete range of 0 km, while broadcast
272 spawners were assigned a gamete range of 10 km. Similarly, adult range size was approximated as
273 0 km (sessile), 1 km (motile but not pelagic), or 10 km (highly mobile, pelagic). Variables were
274 specified as 'multiple' for groups known to span more than 1 magnitude of range size. For groups to
275 which sequences were annotated with high confidence, but for which life history strategy is diverse or
276 poorly known (e.g. families in the phylum Nemertea), we used conservative, coarse approximations
277 at a higher taxonomic rank (see Life History Data).

278 **Results**

279 **Sequence Data Processing (Bioinformatics)**

280 Preliminary sequence analysis strongly suggested that the observed variation among environmental
281 samples reflects true variation in the environment, rather than variability due to lab protocols, for
282 the following reasons (note that all value ranges are reported as mean plus and minus one standard
283 deviation). First, all libraries passed the FastQC per-base sequence quality filter, generating a total
284 of 371,576,190 reads passing filter generated in each direction. Second, samples in this study were
285 represented by an adequate number of reads ($333,537.9 \pm 112,200.5$), with no individual sample
286 receiving fewer than 130,402 reads. Third, there was a very low frequency of cross-contamination
287 from other libraries into those reported here ($5e-05 \pm 8e-05$; max proportion 0.00034). Fourth, after
288 scaling all samples to the same sequencing depth, OTUs with abundance greater than 178 reads
289 (0.14% of a sample's reads) experienced no turnover among PCR replicates within a sample. Fifth,
290 sequence abundances among PCR replicates within water samples were remarkably consistent. A
291 single sample had low similarity among PCR replicates (0.659) after removing this outlier, the
292 lowest mean similarity among replicates within a sample was 0.966. Overall similarities among
293 PCR replicates within a sample were extremely high (0.976 ± 0.013), and far higher than that of
294 than among samples (0.3 ± 0.16).

295 **Ecological Analyses**

296 **Distance Decay**

297 Physical proximity is a good predictor of eDNA community similarity: Similarity decreased from
298 0.40 (95%CI = 0.36, 0.45) to half that amount at 4500 meters (95%CI = 2900, 7500) (Figure 2).

299 **Community Classification**

300 Despite a clear trend in community similarity as a function of spatial separation, the results from
301 our classification analysis are difficult to interpret. The silhouette analysis indicated the presence
302 of 8 distinct communities; however, the gain in mean silhouette width from 2 was small (0.1), and
303 lacked a distinctive peak (Figure 4), indicating substantial uncertainty in the clustering algorithm.

304 Thus, we present the results of cluster assignment for both $K = 2$ and $K = 8$ to illustrate the
305 range of results (Figure 3). Excluding taxa which occur in only one site had no discernible effect
306 on the outcome of the PAM analysis (number of clusters, assignment to clusters). While there was
307 no distinct spatial divide indicating the presence of an inshore versus an offshore community, one
308 of the two communities (at $K = 2$) occurred in only 2 out of 18 samples inside 1000 meters from
309 shore, and never occurred within 125 meters of shore, suggesting the presence of an inshore and
310 offshore community.

311 Diversity in Space

312 Sites offshore tend to be less rich and more even than those inshore (Figure 6). Mean OTU richness
313 declined by 1.42 per 1000 meters from a mean of 17.6 taxa (95%CI = 2.15) inshore to 11.9 taxa
314 (95%CI = 4.31) at offshore locations ($p = 0.0415$; Figure 6). Evenness, the probability that two
315 reads chosen at random from a sample belong to different species, increased by .0666 per 1000
316 meters from 0.225 (95%CI = 0.0558) to 0.491 (95%CI = ± 0.112), indicating that sequence reads
317 were less evenly distributed among taxa in offshore samples ($p \ll 0.05$; Figure 6). There was no
318 evidence for spatial autocorrelation for any of the diversity metrics (Moran's I, $p > 0.05$; Figure 5).

319 Taxon and Life History Patterns

320 We were able to assign a taxon name with confidence to 136 of 146 OTU sequences. The vast ma-
321 jority of sequences (97.6%) and OTUs (96.9%) were matched to organisms that have high potential
322 for dispersal at either the gamete, larval, or adult stage, making it impossible to determine whether
323 the source of that DNA was adults with well-documented spatial patterns (e.g. sessile nearshore
324 specialists) or highly mobile early life history stages. Of the 6 OTUs for which dispersal is limited
325 during all life history stages, only 2 occurred in more than two samples, precluding a quantita-
326 tive comparison of spatial dispersion based on life history characteristics. These were assigned to
327 *Cymatogaster aggregata*, a viviparous nearshore fish with internal fertilization, and *Cupolaconcha*
328 *meroclista*, a sessile Vermetid gastropod with presumed internal fertilization and short larval dis-
329 persal (Strathmann and Strathmann, 2006; Phillips and Shima, 2010; Calvo and Templado, 2004).
330 *Cymatogaster aggregata* was distinctly more abundant close to shore, with no sequences occurring in
331 any sample beyond 250 meters (Figure 7). *Cupolaconcha meroclista* showed no such distinct spatial

332 trend, occurring in nearly equal abundance at three sites, 75, 500, and 2000 meters from shore. An
333 additional species that was highly abundant in the sequence data, the krill *Thysanoessa raschii*,
334 has pelagic adults, highly seasonal reproduction, and sinking eggs; their distribution was consistent
335 with our expectations based on a tendency of adults to aggregate offshore. Finally, the two most
336 abundant taxa in the dataset were the mussel genus *Mytilus* and the Barnacle order Sessilia; the
337 adults of both taxa are sessile and occur exclusively on hard intertidal substrata but have highly
338 motile larvae.

339 Discussion

340 Indirect surveys of organismal presence are a key development in ecosystem monitoring in the face
341 of increased anthropogenic pressure and dwindling resources for ecological research. Monitoring
342 of organisms using environmental DNA is an especially promising method, given the rapid pace
343 of advancement in technological innovation and cost efficiency in the field of DNA sequencing and
344 quantification. For the first time in a marine environment, we document four key patterns: (1) eDNA
345 communities far from one another tend to be less similar than those that are nearby, (2) distinct
346 eDNA communities exist and are distributed in a non-random fashion, (3) diversity declines with
347 distance from shore, and (4) spatial patterning of eDNA is associated with taxon-specific life history
348 characteristics.

349 (1) Communities far from one another tend to be less similar than those that are 350 nearby

351 We demonstrate that more distant locations have less similar eDNA communities than more proxi-
352 mate locations in Puget Sound, a dynamic marine environment. Our finding is in line with observa-
353 tions based on traditional surveys of terrestrial plants and fungi (Nekola and White, 1999; Bahram
354 et al., 2013; Condit, 2002; Chust et al., 2006) and of microorganisms in freshwater (Wetzel et al.,
355 2012), marine (Chust et al., 2013), and estuarine (Martiny et al., 2011) environments. To our knowl-
356 edge, it is the first to report such a pattern using massively parallel sequencing of environmental
357 DNA in the marine environment, and the first using any technique to describe this pattern from
358 microbial metazoans. We note that the theoretical expectation is that samples at very close distance

359 be nearly completely similar, while our samples separated by the 50 meters were only 40% similar.
360 We interpret this to reflect the highly dynamic nature of this environment, which could cause DNA
361 to be distributed quickly from its source, eroding the rise in similarity at small distances. At the
362 same time, community similarity decreased to very low levels at larger scales, indicating that DNA
363 distribution is not completely unpredictable. This finding implies that the effectively sampled area
364 of individual water samples for eDNA analysis is likely to be quite small (<100m) in this nearshore
365 environment. Our estimated distance-decay relationship does indicate that proximate samples are
366 more similar than distant samples, but we suggest this pattern is partially obscured by other factors,
367 including signal from mobile, microscopic life-stages.

368 **(2) Distinct eDNA communities exist and are distributed in a non-random fashion**

369 We demonstrate strong evidence for distinct community types and the non-random spatial pattern-
370 ing of those communities. While the spatial distributions of communities is surprising if one were
371 concerned only with the macroscopic life stages of metazoans, it indeed does align with the broader
372 view that even offshore pelagic communities are comprised of and influenced by nearshore organ-
373 isms. This result underscores the idea that areas immediately offshore act as ecotones, a mixing
374 zone of taxa characteristic of benthic and pelagic environments. While there was no distinct break
375 in community types between onshore and offshore sites, there was some clustering of community
376 types that may be explained by oceanographic features such as nearshore eddies generated by strong
377 tidal exchange in a steep bathymetric setting (Yang and Khangaonkar, 2010). It would be useful to
378 better understand such features during the period of sampling, by way of oceanographic monitoring
379 devices.

380 **(3) Richness declines and evenness increases with distance from shore**

381 We detected a general pattern of declining richness and increasing evenness with increasing distance
382 offshore. Such a pattern is consistent with many other ecosystems which show strong clines in
383 diversity metrics over environmental gradients. The coastal ocean is a highly productive and diverse
384 ecosystem (Ray, 1988). However, our study is novel in that it corroborates a cline well-known on
385 macroscales for macrobiota on a much smaller spatial scale for microscopic animals, suggesting that
386 there may be a self-similarity across scales in diversity patterning (Levin, 1992). Intriguingly, the

387 cline in diversity from inshore to offshore was not determined by shared changes in communities
388 as one moved offshore; the classification analysis suggested a fair amount of differences among
389 communities at a given offshore distance (Figure 3). Furthermore, the uncertainty in identification
390 of the number of distinct clusters to best characterize the community underlines the difficulty of
391 identifying community patterns with the number of taxonomic groups considered here. We suspect
392 that the signature of eDNA from microscopic life-stages may explain our inability to easily detect
393 spatial community level patterns that align with our initial expectations.

394 **(4) Spatial patterning of eDNA is associated with taxon-specific life history character-**
395 **istics.**

396 In contrast to our expectations, other taxa including species with sessile adult stages restricted
397 to benthic hard substrates (e.g. barnacles, mussels) are among the most abundant taxa at sites
398 furthest from shore. However, the larvae and gametes of these taxa are abundant, pelagic, and
399 can be transported long distances by water movement (Strathmann, 1987). This indicates that we
400 likely detected DNA of their pelagic phase gametes and larvae. It is always possible that DNA
401 of adults was advected over long distances and detected offshore but in light of our results with
402 krill and surfperch, we view this as unlikely. We interpret our results as evidence that the chaotic
403 spatial distribution of eDNA communities (Figure 3) results from our primers' affinity for many
404 species which at some point exist as microscopic pelagic gametes or larvae. Our results emphasize
405 that expected results based on easily visually observed individuals or detectable with traditional
406 sampling gear such as nets may be very different from results using eDNA. This does caution that
407 eDNA surveys may have different purposes and may not be directly comparable to existing surveys
408 (Shelton et al., 2016).

409 We acknowledge that sampling artifacts may have affected our results. For example if entire
410 multicellular individuals were captured in our samples, their DNA could be in much greater density
411 than eDNA, affecting the observed community. Our sampling bottles excluded particles larger than
412 500 micrometers, but gametes and very small larvae could have gained entry. It is possible that
413 even a single small individual, containing many thousand mitochondria, would overwhelm the signal
414 of another species from which hundreds of cells had been sloughed from many, larger individuals.
415 Data on larval size distribution at the time of sampling from each species in our data set would

416 allow us to estimate the frequency of such events. Nevertheless, it is precisely the sensitivity to
417 small particles that makes the eDNA approach powerful, so we are reluctant to recommend that
418 aquatic eDNA sampling use finer pre-filtering. Instead, we emphasize the importance of designing
419 and selecting primer sets that selectively amplify target organisms. In the case of the present study,
420 in order to recover patterns matching our expectations, this would be non-transient, benthic marine
421 organisms lacking any pelagic life stage.

422 Our results also highlight the need for curated life-history databases. As technological advances
423 increase the speed and throughput of DNA sequencing and sequence processing, making sense of
424 these data in a timely manner requires that natural history data be stored in standard formats in
425 centralized repositories. The rate at which we can make sense of high-throughput survey methods
426 will be limited by our ability to collate auxiliary data. Databases such as Global Biodiversity
427 Information Facility (GBIF), Encyclopedia of Life (EOL), and FishBase (Parr et al., 2014; Froese
428 and Pauly, 2016) contain records of taxonomy, occurrence, and other rudimentary data types, but
429 there is no centralized, standardized repository for even basic natural history data such as body
430 size. As NCBI's nucleotide and protein sequence database (GenBank) has facilitated transformative
431 studies in diverse fields, an ecological analog would be a boon for biodiversity science.

432 Surveys based on eDNA are intensely scrutinized because of the danger that the final data are
433 subject to complicated laboratory and bioinformatic procedures. Finding virtually no variability
434 among lab and bioinformatic treatments from the point of PCR onward, we were confident our
435 results represented actual field-based differences among samples. However, we note that one PCR
436 replicate had a clear signal of contamination in that the sequence community was extremely similar
437 to those from a different environmental sample. The source of this error is difficult to identify, but
438 seems most likely to be an error during PCR preparation, either in assignment or pipetting during
439 preparation of indexed primers. While the remainder of our results would be largely unchanged
440 had we sequenced a single replicate per environmental sample, we believe the sequencing of PCR
441 replicates is critical for ensuring data quality in eDNA sequencing studies.

442 While there is much work to be done before eDNA techniques can be confidently deployed as a
443 standard method for ecological monitoring, this study serves as a first analysis of diversity at the
444 fine spatial scales that are likely to be relevant to eDNA work in the field across a range of study
445 systems.

446 **Acknowledgements**

447 We wish to thank Robert Morris, E. Virginia Armbrust, and James Kralj.

448 **Funding**

449 This work was supported by a grant from the David and Lucile Packard Foundation to RPK (grant
450 2014-39827). The funders had no role in study design, data collection and analysis, decision to
451 publish, or preparation of the manuscript.

452 **Author Contributions**

453 Conceived and designed the experiments: JL O'Donnell, RP Kelly, AO Shelton; Collected the data:
454 JL O'Donnell, NC Lowell, GD Williams, RP Kelly, AO Shelton, JF Samhouri; Conducted the
455 analyses: JL O'Donnell; Wrote the first draft: JL O'Donnell; Edited the manuscript: JL O'Donnell,
456 AO Shelton, RP Kelly, JF Samhouri, GD Williams, NC Lowell

457 **Ethics Statement**

458 The authors declare no conflict of interest. No permits were required to do any of the research
459 described here.

460 **Data Availability**

461 **0.1 Sequence Data**

462 All sequence files and metadata are available from EMBL:

463 <http://www.ebi.ac.uk/ena/data/view/FIXME>

464 **0.2 Project Repository**

465 The following components are available from the project repository on GitHub:

466 https://github.com/jimmyodonnell/Carkeek_eDNA_grid

467 <http://dx.doi.org/FIXME>

468 0.2.1 Sequencing Metadata

469 Sequencing metadata is available in: `Data/metadata_spatial.csv`

470 0.2.2 Life History Data

471 Life history data is available in: `Data/life_history.csv`

472 0.2.3 Analysis Scripts

473 All analyses were performed using scripts available in the Analysis subdirectory.

474 References

475 Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A. L., Sanders,
476 N. J., Cornell, H. V., Comita, L. S., Davies, K. F., Harrison, S. P., Kraft, N. J. B., Stegen, J. C.,
477 and Swenson, N. G. (2011). Navigating the multiple meanings of beta diversity: A roadmap for
478 the practicing ecologist. *Ecology Letters*, 14(1):19–28.

479 Bahram, M., Kõljalg, U., Courty, P. E., Diédhiou, A. G., Kjølner, R., Põlme, S., Ryberg, M., Veldre,
480 V., and Tedersoo, L. (2013). The distance decay of similarity in communities of ectomycorrhizal
481 fungi in different ecosystems and scales. *Journal of Ecology*, 101(5):1335–1344.

482 Barnes, M. A., Turner, C. R., Jerde, C. L., Renshaw, M. A., Chadderton, W. L., and Lodge, D. M.
483 (2014). Environmental conditions influence eDNA persistence in aquatic systems. *Environmental*
484 *Science and Technology*, 48(3):1819–1827.

485 Bell, T. (2010). Experimental tests of the bacterial distance–decay relationship. *The ISME Journal*,
486 4(11):1357–1365.

487 Burns, R. E. (1985). *The shape and form of Puget Sound*. Washington Sea Grant, Seattle, 1 edition.

488 Calvo, M. and Templado, J. (2004). Reproduction and development in a vermetid gastropod,
489 *Vermetus triquetrus*. *Invertebrate Biology*, 123(4):289–303.

490 Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T. L.
491 (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, 10:421.

- 492 Chamberlain, S. a. and Szöcs, E. (2013). taxize: taxonomic search and retrieval in R. *F1000Research*,
493 2(0):191.
- 494 Chamberlain, S. A., Szöcs, E., Boettiger, C., Ram, K., Bartomeus, I., Foster, Z., and O'Donnell,
495 J. L. (2016). taxize: Taxonomic information from around the web. R package.
- 496 Chust, G., Chave, J., Condit, R., Aguilar, S., Lao, S., and Perez, R. (2006). Determinants and
497 spatial modeling of tree beta-diversity in a tropical forest landscape in Panama. *Journal of*
498 *Vegetation Science*, 17(1):83–92.
- 499 Chust, G., Irigoien, X., Chave, J., and Harris, R. P. (2013). Latitudinal phytoplankton distribution
500 and the neutral theory of biodiversity. *Global Ecology and Biogeography*, 22(5):531–543.
- 501 Coissac, E. (2012). OligoTag: A Program for Designing Sets of Tags for Next-Generation Sequencing
502 of Multiplexed Samples. In Pompanon, F. and Bonin, A., editors, *Data Production and Analysis in*
503 *Population Genomics SE - 2*, volume 888 of *Methods in Molecular Biology*, pages 13–31. Humana
504 Press.
- 505 Condit, R. (2002). Beta-Diversity in Tropical Forest Trees. *Science*, 295(5555):666–669.
- 506 Cowart, D. a., Pinheiro, M., Mouchel, O., Maguer, M., Grall, J., Miné, J., and Arnaud-Haond, S.
507 (2015). Metabarcoding Is Powerful yet Still Blind: A Comparative Analysis of Morphological and
508 Molecular Surveys of Seagrass Communities. *Plos One*, 10(2):e0117562.
- 509 Deiner, K. and Altermatt, F. (2014). Transport distance of invertebrate environmental DNA in a
510 natural river. *PLoS ONE*, 9(2).
- 511 Dethier, M. N. (2010). Overview of the ecology of Puget Sound beaches. In Shipman, H., Dethier,
512 M. N., Gelfenbaum, G., Fresh, K. L., and Dinicola, R. S., editors, *Puget Sound Shorelines and*
513 *the Impacts of Armoring—Proceedings of a State of the Science Workshop*, page 262.
- 514 Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*,
515 26(19):2460–2461.
- 516 Evans, N. T., Olds, B. P., Renshaw, M. A., Turner, C. R., Li, Y., Jerde, C. L., Mahon, A. R.,
517 Pfrender, M. E., Lamberti, G. A., and Lodge, D. M. (2016). Quantification of mesocosm fish and

- 518 amphibian species diversity via environmental DNA metabarcoding. *Molecular Ecology Resources*,
519 16(1):29–41.
- 520 Ficetola, G. F., Miaud, C., Pompanon, F., and Taberlet, P. (2008). Species detection using envi-
521 ronmental DNA from water samples. *Biology letters*, 4(4):423–425.
- 522 Froese, R. and Pauly, D. (2016). FishBase.
- 523 Guo, H., Chamberlain, S. A., Elhaik, E., Jalli, I., Lynes, A. R., Marczak, L., Sabath, N., Vargas,
524 A., Więski, K., Zelig, E. M., and Pennings, S. C. (2015). Geographic variation in plant commu-
525 nity structure of salt marshes: Species, functional and phylogenetic perspectives. *PLoS ONE*,
526 10(5):e0127781.
- 527 Handelsman, J., Rondon, M. R., Brady, S. F., Clardy, J., and Goodman, R. M. (1998). Molecular
528 biological access to the chemistry of unknown soil microbes: a new frontier for natural products.
529 *Chemistry & Biology*, 5(10):R245–R249.
- 530 Hijmans, R. J. (2016). *geosphere: Spherical Trigonometry*.
- 531 Hubbell, S. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography.*, volume 32.
- 532 Iverson, V., Morris, R. M., Frazar, C. D., Berthiaume, C. T., Morales, R. L., and Armbrust,
533 E. V. (2012). Untangling Genomes from Metagenomes: Revealing an Uncultured Class of Marine
534 Euryarchaeota. *Science*, 335(6068):587–590.
- 535 Kaufman, L. and Rousseeuw, P. J. (1990). *Finding Groups in Data: An Introduction to Cluster*
536 *Analysis*.
- 537 Kelly, R. P., O'Donnell, J. L., Lowell, N. C., Shelton, A. O., Samhour, J. F., Hennessey, S. M.,
538 Feist, B. E., and Williams, G. D. (2016). Genetic signatures of ecological diversity along an
539 urbanization gradient. *PeerJ*, 4:e2444.
- 540 Klymus, K. E., Richter, C. A., Chapman, D. C., and Paukert, C. (2015). Quantification of eDNA
541 shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypoph-*
542 *thalmichthys molitrix*. *Biological Conservation*, 183:77–84.

- 543 Kozloff, E. N. (1973). *Seashore life of Puget Sound, the Strait of Georgia, and the San Juan*
544 *Archipelago*. University of Washington Press, Seattle.
- 545 Levin, S. A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73(6):1943–1967.
- 546 Longmire, J. L., Maltbie, M., and Baker, R. J. (1997). Use of lysis buffer in DNA isolation and its
547 implication for museum collections. *Museum of Texas Tech University*, 163.
- 548 Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., and Hornik, K. (2016). *cluster: Cluster*
549 *Analysis Basics and Extensions*.
- 550 Magurran, A. E. (2003). *Measuring Biological Diversity*. Wiley.
- 551 Mahé, F., Rognes, T., Quince, C., de Vargas, C., and Dunthorn, M. (2014). Swarm: robust and
552 fast clustering method for amplicon-based studies. *PeerJ*, 2:e593.
- 553 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
554 *EMBnet.journal*, 17(1):10.
- 555 Martiny, J. B. H., Eisen, J. A., Penn, K., Allison, S. D., and Horner-Devine, M. C. (2011). Drivers of
556 bacterial beta diversity depend on spatial scale. *Proceedings of the National Academy of Sciences*,
557 108(19):7850–7854.
- 558 Nekola, J. C. and White, P. S. (1999). The distance decay of similarity in biogeography and ecology.
559 *Journal of Biogeography*, 26(4):867–878.
- 560 O'Donnell, J. L., Kelly, R. P., Lowell, N. C., and Port, J. A. (2016). Indexed PCR Primers Induce
561 Template-Specific Bias in Large-Scale DNA Sequencing Studies. *PLoS ONE*, 11(3):1–11.
- 562 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
563 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H.
564 (2016). *vegan: Community Ecology Package*.
- 565 Parr, C. S., Wilson, N., Leary, P., Schulz, K. S., Lans, K., Walley, L., Hammock, J. A., Goddard,
566 A., Rice, J., Studer, M., Holmes, J. T. G., and Corrigan, R. J. (2014). The Encyclopedia of
567 Life v2: Providing Global Access to Knowledge About Life on Earth. *Biodiversity Data Journal*,
568 2(2):e1079.

- 569 Phillips, N. E. and Shima, J. S. (2010). Reproduction of the vermetid gastropod *dendropoma*
570 maximum (Sowerby, 1825) in Moorea, French Polynesia. *Journal of Molluscan Studies*, 76(2):133–
571 137.
- 572 Port, J. A., O'Donnell, J. L., Romero-Maraccini, O. C., Leary, P. R., Litvin, S. Y., Nickols, K. J.,
573 Yamahara, K. M., and Kelly, R. P. (2016). Assessing vertebrate biodiversity in a kelp forest
574 ecosystem using environmental DNA. *Molecular Ecology*, 25(2):527–541.
- 575 R Core Team (2016). *R: A Language and Environment for Statistical Computing*. R Foundation
576 for Statistical Computing, Vienna, Austria.
- 577 Ray, G. C. (1988). Ecological diversity in coastal zones and oceans. In Wilson, E. O. and Peter,
578 F. M., editors, *Biodiversity*, chapter 4. National Academies Press (US), Washington, DC.
- 579 Renshaw, M. A., Olds, B. P., Jerde, C. L., Mcveigh, M. M., and Lodge, D. M. (2014). The room
580 temperature preservation of filtered environmental DNA samples and assimilation into a phenol-
581 chloroform-isoamyl alcohol DNA extraction. *Molecular Ecology Resources*.
- 582 Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. (2016). VSEARCH: a versatile open
583 source tool for metagenomics. *PeerJ*, 4:e2584.
- 584 Sassoubre, L. M., Yamahara, K. M., Gardner, L. D., Block, B. A., and Boehm, A. B. (2016).
585 Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three Marine
586 Fish. *Environmental Science & Technology*, 50(19):10456–10464.
- 587 Schnell, I. B., Bohmann, K., and Gilbert, M. T. P. (2015). Tag jumps illuminated - reducing
588 sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources*,
589 15(6):1289–1303.
- 590 Shelton, A. O., O'Donnell, J. L., Samhour, J. F., Lowell, N., Williams, G. D., and Kelly, R. P.
591 (2016). A framework for inferring biological communities from environmental DNA. *Ecological*
592 *Applications*, 26(6):1689–.
- 593 Shogren, A. J., Tank, J. L., Andruszkiewicz, E. A., Olds, B., Jerde, C., and Bolster, D. (2016).
594 Modelling the transport of environmental DNA through a porous substrate using continuous
595 flow-through column experiments. *Journal of The Royal Society Interface*, 13(119):423–425.

- 596 Simpson, E. H. (1949). Measurement of diversity. *Nature*, 163(688).
- 597 Spiess, A.-N. (2014). *propagate: Propagation of Uncertainty*.
- 598 Strathmann, M. F. (1987). *Reproduction and Development of Marine Invertebrates of the Northern*
599 *Pacific Coast: Data and Methods for the Study of Eggs, Embryos, and Larvae*. University of
600 Washington Press, Seattle.
- 601 Strathmann, M. F. and Strathmann, R. R. (2006). A Vermetid Gastropod with Complex Intracap-
602 sular Cannibalism of Nurse Eggs and Sibling Larvae and a High Potential for Invasion. *Pacific*
603 *Science*, 60(1):97–108.
- 604 Strickland, R. M. (1983). *The Fertile Fjord: Plankton in Puget Sound*. University of Washington
605 Press, Seattle.
- 606 Strickler, K. M., Fremier, A. K., and Goldberg, C. S. (2015). Quantifying effects of UV-B, temper-
607 ature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation*, 183:85–92.
- 608 Thomsen, P. F., Kielgast, J., Iversen, L. L., Møller, P. R., Rasmussen, M., and Willerslev, E. (2012).
609 Detection of a Diverse Marine Fish Fauna Using Environmental DNA from Seawater Samples.
610 *PLoS ONE*, 7(8):1–9.
- 611 Turner, C. R., Uy, K. L., and Everhart, R. C. (2015). Fish environmental DNA is more concentrated
612 in aquatic sediments than surface water. *Biological Conservation*, 183:93–102.
- 613 Tyson, G. W., Chapman, J., Hugenholtz, P., Allen, E. E., Ram, R. J., Richardson, P. M.,
614 Solovyev, V. V., Rubin, E. M., Rokhsar, D. S., and Banfield, J. F. (2004). Community structure
615 and metabolism through reconstruction of microbial genomes from the environment. *Nature*,
616 428(6978):37–43.
- 617 Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. a., Wu, D.,
618 Paulsen, I., Nelson, K. E., Nelson, W., Fouts, D. E., Levy, S., Knap, A. H., Lomas, M. W.,
619 Neelson, K., White, O., Peterson, J., Hoffman, J., Parsons, R., Baden-Tillson, H., Pfannkoch,
620 C., Rogers, Y.-H., and Smith, H. O. (2004). Environmental genome shotgun sequencing of the
621 Sargasso Sea. *Science*, 304(5667):66–74.

- 622 Wetzell, C. E., de Bicudo, D. C., Ector, L., Lobo, E. A., Soininen, J., Landeiro, V. L., and Bini,
623 L. M. (2012). Distance Decay of Similarity in Neotropical Diatom Communities. *PLoS ONE*,
624 7(9):e45071.
- 625 Yang, Z. and Khangaonkar, T. (2010). Multi-scale modeling of Puget Sound using an unstructured-
626 grid coastal ocean model: From tide flats to estuaries and coastal waters. *Ocean Dynamics*,
627 60(6):1621–1637.
- 628 Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A. (2014). PEAR: A fast and accurate Illumina
629 Paired-End reAd mergeR. *Bioinformatics*, 30(5):614–620.

630 Figures




Figure 1: Map of study area. Depth in meters below sea level is indicated by shading and 25 meter contours. Sampled locations are indicated by red points.




Figure 2: Distance decay relationship of environmental DNA communities. Each point represents the Bray-Curtis similarity of a site sampled along three parallel transects comprising a 3000 by 4000 meter grid. Blue dashed line represents fit of a nonlinear least squares regression (see Methods), and shading denotes the 95% confidence interval. Boxplot is comparisons within-sample across PCR replicates, separated by a vertical line at zero, where the central line is the median, the box encompasses the interquartile range, and the lines extend to 1.5 times the interquartile range. Boxplot outliers are omitted for clarity.




Figure 3: Cluster membership of sampled sites. Distance from onshore starting point is log scaled. Sites are colored and labeled by their assignment to a cluster by PAM analysis for number of clusters (K) chosen based on a priori expectations (2) and mean silhouette width (8).




Figure 4: Silhouette widths from PAM analysis. Points are the width of the PAM silhouette of each sample at each number of clusters (K). Red line is the mean, blue line is the median. Boxes encompass the interquartile range with a line at the median, and the whiskers extend to 1.5 times the interquartile range. Boxplot outliers are omitted for clarity.




Figure 5: Aggregate measures of diversity at each sample site.




Figure 6: Aggregate diversity metrics of each site plotted against distance from shore. Both Simpson's Index (left) and richness (right) are shown, and have been computed from the mean abundance of unique DNA sequences found across 4 PCR replicates at each of 24 sites. Lines and bands illustrate the fit and 95




Figure 7: Distribution of eDNA from select taxa. Circles are colored and scaled by the proportion of that taxon's maximum proportional abundance. That is, the largest circle is the same size in each of the panels, and occurs where that taxon contributed the greatest proportional abundance of reads to that sample.

631 Supplemental Material

632 Methods

633 Bioinformatics

634 Reads passing the preliminary Illumina quality filter were demultiplexed on the basis of the adapter
635 index sequence by the sequencing facility. We used fastqc to assess the fastq files output from the
636 sequencer for low-quality indications of a problematic run. Forward and reverse reads were merged
637 using PEAR v0.9.6 Zhang et al. (2014) and discarded if more than 0.01 of the bases were uncalled.
638 If a read contained two consecutive base calls with quality scores less than 15 (i.e. probability of
639 incorrect base call = 0.0316), these bases and all subsequent bases were removed from the read.
640 Paired reads for which the probability of matching by chance alone exceeded 0.01 were not assembled
641 and omitted from the analysis. Assembled reads were discarded if assembled sequences were not
642 between 50 and 168 bp long, or if reads did not overlap by at least 100 bp.

643 We used vsearch v2.1.1 (Rognes et al., 2016) to discard any merged reads for which the sum of the
644 per-base error probabilities was greater than 0.5 (“expected errors”) Edgar (2010). Sequences were
645 demultiplexed on the basis of the primer index sequence at base positions 4-9 at both ends using the
646 programming language AWK. Primer sequences were removed using cutadapt v1.7.1 Martin (2011),
647 allowing for 2 mismatches in the primer sequence. Identical duplicate sequences were identified,
648 counted, and removed in python to speed up subsequent steps by eliminating redundancy, and
649 sequences occurring only once were removed. We checked for and removed any sequence likely to be
650 a PCR artifact due to incomplete extension and subsequent mis-priming using a method described
651 by Edgar (2010) and implemented in vsearch v2.0.2. Sequences were clustered into operational
652 taxonomic units (OTUs) using the single-linkage clustering method implemented by swarm version
653 2.1.1 with a local clustering threshold (d) of 1 and fastidious processing (Mahé et al., 2014).

654 Cross-contamination of environmental, DNA, or PCR samples can result in erroneous inference
655 about the presence of a given DNA sequence in a sample. However, other processes can contribute
656 to the same signature of contamination. For example, errors during oligonucleotide synthesis or
657 sequencing of the indexes could cause reads to be erroneously assigned to samples. The frequency
658 of such errors can be estimated by counting the occurrence of sequences known to be absent from

659 a given sample, and of reads that do not contain primer index sequences in the expected position
660 or combinations. These occurrences indicate an error in the preparation or sequencing procedures.
661 We estimated a rate of incorrect sample assignment by calculating the maximum rate of occur-
662 rence of index sequences combinations we did not actually use, as well as the rates of cross-library
663 contamination by counting occurrences of primer sequences from 12S amplicons prepared in a lab
664 more than 1000 kilometers away, but pooled and sequenced alongside our samples. This represents
665 a general minimum rate at which we can expect that sequences from one environmental sample
666 could be erroneously assigned to another, and so we considered for further analysis only those reads
667 occurring with greater frequency than this across the entire dataset.

668 We checked for experimental error by evaluating the Bray-Curtis similarity (1 - Bray-Curtis
669 dissimilarity) among replicate PCRs from the same DNA sample. We calculated the mean and
670 standard deviation across the dataset, and excluded any PCR replicates for which the similarity
671 between itself and the other replicates was less than 1.5 standard deviations from the mean.

672 To account for variation in the number of sequencing reads (sequencing depth) recovered per
673 sample, we rarefied the within-sample abundance of each OTU by the minimum sequencing depth
674 (Oksanen et al., 2016).

675 Because each step in this workflow is sensitive to contamination, it is possible that some se-
676 quences are not truly derived from the environmental sample, and instead represent contamination
677 during field sampling, filtration, DNA extraction, PCR, fragment size selection, quantitation, se-
678 quencing adapter ligation, or the sequencing process itself. We take the view that contaminants
679 are unlikely to manifest as sequences in the final dataset in consistent abundance across replicates;
680 indeed, our data show that the process from PCR onward is remarkably consistent. Thus, after
681 scaling to correct for sequencing depth variation, we calculated from our data the maximum number
682 of sequence counts for which there is turnover in presence-absence among PCR replicates within an
683 environmental sample. We use this number to determine a conservative minimum threshold above
684 which we can be confident that counts are consistent among replicates and not of spurious origin,
685 and exclude from further analysis observations where the mean abundance across PCR replicates
686 within samples does not reach this threshold. For further analyses we use the mean abundance
687 across PCR replicates for each of the 24 environmental samples.

688 In order to determine the most likely taxon from which each sequence originated, the representa-

689 tive sequence from each OTU was then queried against the NCBI nucleotide collection (GenBank;
690 version October 7, 2015; 32,827,936 sequences) using the blastn command line utility (Camacho
691 et al., 2009). In order to maximize the accuracy of this computationally intensive step, we imple-
692 mented a nested approach whereby each sequence was first queried using strict parameters (e-value
693 = 5e-52), and if no match was found, the query was repeated with decreasingly strict e-values (5e-48
694 5e-44 5e-40 5e-36 5e-33 5e-29 5e-25 5e-21 5e-17 5e-13). Other parameters were unchanged among
695 repetitions (word size: 7; maximum matches: 1000; culling limit: 100; minimum percent identity:
696 0). Each query sequence can be an equally good match to multiple taxa either because of invariabil-
697 ity among taxa or errors in the database (e.g. human sequences are commonly attributed to other
698 organisms when they in fact represent lab contamination). In order to guard against these spurious
699 results, we used an algorithm to find the lowest common taxon for at least 80% of the matched
700 taxa, implemented in the R package taxize 0.7.8 (Chamberlain and Szöcs, 2013; Chamberlain et al.,
701 2016). Similarly, we repeated analyses using the dataset consolidated at the same taxonomic rank
702 across all queries, for the rank of both family and order.

703 **Alternative distance decay model formulations**

704 **Linear:** We fit a straight line through the points after log-transforming the spatial distances
705 to estimate the intercept and slope. This model ignores the bounds of our response variable of
706 community similarity.

707 **Michaelis-Menten:** We fit a Michaelis-Menten-like curve to our data. Our formulation can be
708 thought of as a modification of the Michaelis-Menten equation, but with the addition of a parameter
709 in the numerator which modifies the intercept.

$$y = \frac{AB + Cx}{B + x} \quad (2)$$

710 Where C is the asymptote of minimum similarity. This formulation allows us to estimate the
711 maximum similarity in the system, and the rate at which it is achieved. If the value of the parameter
712 (AB) is 0 (i.e. if the intercept is 0), the form is identical to the Michaelis-Menten equation:

$$y = \frac{Cx}{B+x} \quad (3)$$

713 This is conceptually satisfying in that a fit through [0,1] reflects the theoretical expectation that
714 samples at zero distance from one another are necessarily identical. Given an efficient sampling
715 technique, replicate samples taken at the same position in space should be identical, and thus the
716 intercept of the regression of similarity against distance should be 1, and deviation from 1 is an
717 indicator of the efficiency of the sampling method.

718 Finally, we considered a model which estimates an asymptote as the total change in similarity
719 (D):

$$y = \frac{A + Dx}{B + x} \quad (4)$$

720 However, this model failed to converge and produced uninformative estimates of all parameters.

721 Supplemental Figures




Figure 8: Distance decay relationship of environmental DNA communities using a variety of models, metrics, and data subsets. Each point represents the similarity of a site sampled along three parallel transects comprising a 3000 by 4000 meter grid. Each row of plots represents a different data subset indicated in the right margin, including the final filtered data reported in the main text (a-d), the unfiltered data including all rare OTUs (e-h), log-transformed ($\log(x+1)$) data (i-l), OTU abundance scaled relative to within-taxon maximum (m-p), and exclusion of OTUs found at only one site (q-t). Columns indicate the similarity index used (Bray Curtis or Morisita-Horn) and whether the input was full abundance data or binary (0,1) transformed data. Lines and bands illustrate the fit and 95% confidence interval of both the main nonlinear model (red, dashed line) and a simple linear model (blue, solid line). Results using the Jaccard distance are omitted because of its similarity to Bray-Curtis.




Figure 9: Aggregate diversity metrics of each site plotted against distance from shore. Both Simpson's Index (top) and richness (bottom) are shown for a variety of data subsets and transformations (left to right: mean, unfiltered mean, $\log(x + 1)$, transformed, scaled, spatially variable, and taxon clustered). Lines and bands illustrate the fit and 95% confidence interval of a linear model. See methods text for detailed data descriptions.