

A peer-reviewed version of this preprint was published in PeerJ on 15 April 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.348) (peerj.com/articles/348), which is the preferred citable publication unless you specifically need to cite this preprint.

Valdivia A, Bruno JF, Cox CE, Hackerott S, Green SJ. 2014. Re-examining the relationship between invasive lionfish and native grouper in the Caribbean. PeerJ 2:e348 <https://doi.org/10.7717/peerj.348>

Re-examining the relationship between invasive lionfish and native grouper in the Caribbean

Biotic resistance is the idea that native species negatively affect the invasion success of introduced species, but whether this can occur at large spatial scales is poorly understood. Here we re-evaluated the hypothesis that native large-bodied grouper and other predators are controlling the abundance of exotic lionfish (*Pterois volitans*/miles) on Caribbean coral reefs. We assessed the relationship between the biomass of lionfish and native predators at 71 reefs in three biogeographic regions while taking into consideration several cofactors that may affect fish abundance, including among others, proxies for fishing pressure and habitat structural complexity. Our results indicate that the abundance of lionfish, large-bodied grouper and other predators were not negatively related. Lionfish abundance was instead controlled by several physical site characteristics, and possibly by culling. Taken together, our results suggest that managers cannot rely on current native grouper populations to control the lionfish invasion.

1 Abel Valdivia^{1*}, John F. Bruno¹, Courtney E. Cox¹, S. Hackerott¹, Stephanie J. Green²

2 ¹Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, NC

3 27599, USA

4 ²Department of Zoology, Oregon State University, Oregon, USA

5 *corresponding author: abel.valdivia@unc.edu

6 **Introduction**

7 Biotic resistance describes the capacity of native or resident species in a community to
8 constrain the success of invasive species (Elton, 1958). While there are several examples of
9 native species controlling invasive populations, especially invasive plants (Reusch & Williams,
10 1999; Mazia et al., 2001; Magoulick & Lewis, 2002; Levine et al., 2004; Mitchell et al., 2006),
11 less clear are the ecological mechanisms that allow heterogeneous communities to resist invasion
12 (Lockwood et al., 2005; Melbourne et al., 2007), and whether these processes are strong enough
13 to compromise invasion success on a large scale (Byers & Noonburg, 2003; Davies et al., 2005).
14 Especially elusive is whether native predators or competitors can constrain the expansion of
15 exotic predator species at large spatial scales (but see, deRivera et al., 2005). Although biotic
16 resistance substantially reduces the establishment of invaders, there is little evidence that species
17 interactions such as predation completely prevent invasion (Levine et al., 2004; Bruno et al.,
18 2005)

19 The invasion of Pacific lionfishes (*Pterois volitans* and *Pterois miles*) into the Caribbean
20 basin (Schofield, 2009) over the past ten years provides an example of biotic interactions within a
21 system that have been unable to reduce exotic invasion at a regional scale (Hackerott et al.,
22 2013). Lionfish have spread to every shallow and deep habitat of the Western North Atlantic and
23 the Caribbean (Whitfield et al., 2007; Betancur-R et al., 2011) including fore reef and patch reef
24 environments (Green & Côté, 2009; Albins & Hixon, 2011), seagrass meadows (Claydon et al.,

2012), mangrove root forests (Barbour et al., 2010), estuarine habitats (Jud et al., 2011), and even depths of 300 feet (Green, pers. obs.). Lionfish dissemination in the region has added additional stress (Albins & Hixon, 2011; Lesser & Slattery, 2011; Côté et al., 2013) to an already disturbed coral reef ecosystem (Paddock et al., 2009; Schutte et al., 2010). Their voracious appetite threatens small reef fish and juveniles of depleted fish populations including commercially important species such as groupers and snappers, and keystone grazers such as parrotfishes (Albins & Hixon, 2008; Green et al., 2012). The failure of the system to constrain invasion success may be associated in part to the lack of native predatory capacity due to overfishing (Carlsson et al., 2009; Mumby et al., 2011), or weak biotic resistance by the native predators and competitors (Levine et al., 2004).

The first study to investigate the potential for biotic control of lionfish by native predators found an inverse relationship between the biomass of native groupers and lionfish on reefs at the Exuma Cays Land and Sea Park (ECLSP) in the Bahamas (Mumby et al., 2011). Specifically, Mumby et al. (2011) found that grouper biomass could explain ~56% of the variability in lionfish biomass, and concluded that large-bodied groupers can constrain lionfish abundance if a series of cofactors at the site level are kept constant (i.e., reef complexity, larval supply, habitat characteristics). To examine whether this relationship holds true at a scale that reflects the heterogeneity of Caribbean reefs, Hackerott et al. (2013) gathered data on lionfish and grouper abundance from 71 sites across multiple regions in the Caribbean. When accounting for several site-specific covariates, Hackerott et al. (2013) did not find a relationship between the abundance of lionfish and native predators/competitors at a broad spatial scale in the Caribbean.

Aside from the suite of variables considered by Hackerott et al. (2013), several other covariates that are known to affect fish community structure, but vary across the region, could mask the effect native predators have on lionfish abundance. Accounting for spatial scale and potential cofactors is essential when evaluating the importance of any single variable in a spatial

comparative study (MacNeil et al., 2009). In particular, fishing mortality, larval dispersal, habitat quality, connectivity, reef structural complexity, depth, ecological interactions, and a myriad of other factors control the population dynamics of reef fish species (Sale, 2002). Here we re-evaluated the relationship between large-bodied grouper and other predators and lionfish abundance, accounting for a broader set of covariates than those included by Hackerott et al. (2013) that may mediate the interaction between predators and the invader (Mumby et al., 2013). We also evaluated the grouper bio-control hypothesis proposed by Mumby et al. (2011) and provide new insights into how such biotic resistance is unlikely at the scale of the Caribbean reef system. The issue still remains how to best manage and/or reduce numbers of lionfish where they are currently found, and the only effective solution to date is direct removal by fisherman and divers (Barbour et al., 2011; Frazer et al., 2012; Green et al., 2013 in press).

Materials and Methods

Sites and fish surveys

Survey methods are explained in detail in Hackerott et al., (2013). In summary, we surveyed 71 coral reefs (3-15 m deep) across three distinct reef habitats (spur-and-grove, slope, and patch reef) in three regions of the Caribbean: The Bahamas, Cuba, and the Mesoamerican Barrier Reef (Belize and Mexico) from 2009 to 2012 (Fig. S1, Table S1). All these habitats were once dominated by the coral complex *Montastraea/Orbicella* (Edmunds & Elahi, 2007). Reef sites were selected to cover a wide range of reef fish abundance. To survey fish abundance, we conducted underwater visual censuses at each site using belt transects (for spur-and-grove and slopes) or roving survey dives (for patch reef) (see details in Hackerott et al., 2013). Fish biomass was calculated through the allometric length-weight conversion formula (Froese & Pauly, 2013) and scaling parameters for lionfish were obtained elsewhere (Green et al., 2011). Grouper was defined as the combined biomass of relatively large-bodied species such as Nassau (*Epinephelus*

striatus), tiger (*Mycteroperca tigris*), black (*Mycteroperca bonaci*), and yellowfin grouper (*Epinephelus interstitialis*) as defined also by Mumby et al., (2011). These species could potentially prey on lionfish (Maljković et al., 2008; Mumby et al., 2011) and are relatively more abundant than other potential predators in the region (Hackerott et al., 2013). Other predators considered in this study included any species that could potentially prey on lionfish (see Table S2 in Hackerott et al., 2013). To directly compare our study with the generality of the results by Mumby et al. (2011), we overlaid their values of fish biomass on our main biomass plot and added boxplots that described the distribution of both data sets.

Covariates

The site-specific parameters included as covariates in our statistical model were wind exposure, habitat type, protection status, depth, and time since invasion which are described in detail in Hackerott et al. (2013). We added two new variables to the models that are hypothesized to strongly modulate lionfish abundance (Mumby et al., 2013): human population density/reef area (humans/reef) which is a proxy for fishing effects (Newton et al., 2007; Mora, 2008), and is predicted to be negatively correlated with lionfish density; and reef complexity, which is a proxy for habitat heterogeneity within sites, predicted to have a positive effect on lionfish density (Green et al., 2012). Human population density was calculated as the number of humans within 50 km (maximum number of people living within 50 km radius of each site). We chose 50 km because it is a reasonable range of human influence on Caribbean reefs (Mora, 2008). Estimates of human population counts for the year 2010 were obtained from the Gridded Population of the World V.3 at 0.25 degree resolution (SEDAC, 2010). Reef area was calculated within 10 km radius of each site, well below the average home range for certain predators species (Farmer & Ault, 2011). Reef area was calculated from the Global Distribution of Coral Reefs (2010) database as available at the Ocean Data Viewer (<http://data.unep-wcmc.org/datasets/13>). This database represents the global distribution of warm-water coral reefs compiled mostly from the

99 Millennium Coral Reef Mapping Project (UNEP-WCMC et al., 2010). All spatial calculations
100 were done in ArcGIS v10.0. Humans/Reef Area (humans/km² of reef) was defined as:

101 Number of humans within 50 km / Reef area within 10 km / ($\pi 10^2$) (km²)

102 To estimate reef complexity we used a rugosity index (0-5) estimated at the transect level,
103 where “0” was a flat substrate with no vertical relief and “5” was an exceptionally complex
104 substrate with numerous caves and overhangs (Polunin & Roberts, 1993). Relief complexity for
105 Eleuthera and New Providence sub-regions was estimated by averaging measurements of reef
106 height (i.e., the vertical distance between the lowest and highest point of the reef structure in cm),
107 taken at five haphazard points within the survey area (either transect or rover diver area) (Wilson
108 et al., 2007). To make reef complexity estimates homogenous for all sites, we transformed the
109 relief complexity estimates taken in Eleuthera and New Providence to the rugosity index,
110 described by Polunin & Roberts (1993), by assigning a gradient of 0 cm to “0” and over 300 cm
111 to “5”. This resulted in a continuous rugosity index for these two sub-regions that was
112 comparable with the rest of the sites.

113 **Data analysis**

114 Before applying the statistical model, we explored the data and determined that a negative
115 binomial or Poisson were the most plausible distributions for lionfish counts (Appendix).
116 Additionally, we checked for collinearity among covariates. We ran a logistic regression model
117 with all the covariates and examined the variance inflation factor (VIF) for each variable. We
118 used a VIF > 2 as a threshold to determine collinearity (Graham, 2003). Depth was correlated
119 with reef habitat type as shallower sites tended to be dominated by patch reefs. Thus we modeled
120 these two factors separately. However, we found that keeping depth in the full model, together
121 with habitat type, did not compromise fitting or the magnitude of the effects (Appendix).

122 We ran a generalized linear mixed-effect model using the Automatic Differentiation
123 Model Builder (glmmADMB) package (Skaug et al., 2013) in R 3.0.2 (R Core Team, 2013). As

the lionfish data was over-dispersed and with excess of zeroes (Hackerott et al., 2013), a glmmADMB which accommodates zero inflation was the most adequate model structure (Bolker et al., 2012). We modeled lionfish counts with a negative binomial type 1 distribution and log link because this model performed better than a Poisson distribution based on the Akaike Information Criterion (AIC) (Appendix). Since a negative binomial is a discrete distribution we included an offset in the model to account for survey area (sampling unit level), thus we could effectively analyze the relationship between the density of lionfish and grouper biomass, i.e.:

$$\text{Log (LF Density)} = \text{Log (LF Counts)} - \text{Log (Survey Area)}$$

Because lionfish density and biomass were highly correlated (Pearson's product moment correlation ~ 0.96 , $p < 0.0001$, Appendix), the results of the model should be applicable to biomass as well. The rest of the covariates were considered fixed. We standardized and centered the numerical covariates to aid in comparison of the coefficient estimates. To account for spatial autocorrelation we nested sites within sub-regions and used them as random effects (see Table S1 for sub-regions). To validate the model we corroborated that no patterns were found on the plot of the model residuals versus fitted values.

Moran's I similarity spline correlograms constructed from the residuals of the glmmADMB model (Zuur et al., 2009) graphically indicated that our mixed-effect modeling framework successfully accommodated the spatial autocorrelation observed in the raw data (Fig S2). Additionally, we used Mantel tests (Mantel, 1967) to confirm the lack of spatial autocorrelation between the Pearson residuals of the model and the lag distance (in km) between sites (i.e., whether sites that are closer together were more similar), and found that the overall correlation coefficient for the model was low ($r = 0.073$, $p = 0.0001$). We performed the autocorrelation analyses using the spatial nonparametric covariance function (ncf) package version 1.1-5 (Bjørnstad, 2013). All analyses were performed in R version 3.0.2 (R Core Team, 2013). Additionally, we provide the entire workflow R code (Appendix) and the master data

summary by site level (FigShare, <http://dx.doi.org/10.6084/m9.figshare.899210>).

Results and Discussion

Even when including proxies for fishing and habitat structure in our statistical model, we found no support for an effect of large-bodied grouper or other predator biomass on lionfish abundance (Fig. 1, Table S3). As in Hackerott et al. (2013), the effects of other covariates in our analysis (namely wind exposure, habitat type, and protection status) (Fig. 1) remained the principal factors that appear to influence lionfish abundance. Our analyses suggest that variation in lionfish density across the region is driven by environmental processes and human activity and not by biotic resistance from native predators.

The absence of a relationship between lionfish and native grouper biomass across a large scale suggests that the results of Mumby et al. (2011), which found a negative association across 12 sites – 5 inside and 7 adjacent to a no-take reserve (ECLSP) – represented a subset of a much broader and complicated relationship driven by other factors (Fig. 1 and 2). The average biomass of large-bodied grouper in our study of the Caribbean region ($7.6 \pm 0.8 \text{ gm}^{-2}$, mean $\pm$ standard error) was slightly lower (Wilcoxon test, $W = 1197$, $p = 0.002$) than that found by Mumby et al. (2011) at Exuma ($10.0 \pm 2.6 \text{ gm}^{-2}$) (Fig. 2). In contrast, the average biomass of lionfish in our study ($7.8 \pm 0.5 \text{ gm}^{-2}$) was ~ 20 times higher (or ~ 2 times higher excluding patch reefs, i.e., $0.7 \pm 0.1 \text{ gm}^{-2}$) than those found at Exuma ($0.4 \pm 0.1 \text{ gm}^{-2}$) by Mumby et al. (2011) (Fig. 2). In that study, relatively low lionfish biomass ($\sim 0.3 \text{ gm}^{-2}$) was associated with relatively high grouper biomass ($\sim 25 \text{ gm}^{-2}$). However, across 71 sites in our study, lionfish biomass ranged widely (0-50 gm^{-2}) at sites with equivalent grouper abundance (Fig. 2). Thus, while predators may negatively impact lionfish under a particular set of local conditions (Mumby et al., 2011), the underlying relationship between lionfish and predator biomass was undetectable on a wide range of heterogeneous sites across the Caribbean region.

173 In this study, we assume that high predator biomass is indicative of high predatory
174 capacity resulting from a high frequency of large individuals (Fig. 3a). Grouper at protected sites
175 were, on average, larger (48.6 ± 1.5 cm TL, mean $\pm$ standard error total length) than those at
176 unprotected sites (34.7 ± 1.1 cm) ($t = -7.68$, $p < 0.001$, Fig. 3a). It is unlikely that sites with
177 relatively high grouper biomass have low predatory capacity as a result of more abundant, but
178 smaller, individual fishes. Indeed, the exact opposite pattern is well documented in a wide range
179 of habitat types for several fish species (Gust et al., 2001; Friedlander & DeMartini, 2002;
180 McClanahan et al., 2007). This seems to also be the case for groupers in our study (Fig. 3b). At
181 sites with grouper biomass of at least 10 gm^{-2} , which was the minimum biomass per site in the
182 ECLSP (Mumby et al., 2011), there were relatively high frequencies of medium/large individuals
183 (Fig. 3b). Medium/large groupers (>30 cm TL) have been classified as having potentially high
184 predatory capacity (Mumby et al., 2011). We found relatively lower frequencies ($<50\%$) of small
185 individuals (<30 cm TL) across all protected sites. Therefore, it is unlikely that a lack of
186 predatory capacity at sites with the highest grouper biomass (Fig. 2 and Fig. 3b) explains the
187 absence of a relationship between lionfish and grouper in our results.

188 While we did not find evidence for an effect of native predators on invasion status,
189 lionfish biomass varied significantly between the reef types we examined. All of our fore-reef
190 sites (slope and spur-and-groove) constituted high-profile habitats and we also included a set of
191 patch reefs, a reef habitat common in the region. In particular, slope and spur-and-groove habitat
192 had a negative effect on lionfish abundance (Fig. 1, Table S3) with higher average lionfish
193 abundance in patch reef habitats ($27.5 \pm 2.1 \text{ gm}^{-2}$ vs. $0.7 \pm 0.1 \text{ gm}^{-2}$). However, both lionfish and
194 large-bodied grouper and predators were frequently observed in each of these habitats (Fig. 3c).
195 The class size distribution for groupers among reef habitats were similar (Fig. 3c). Almost 90% of
196 the patch reef sites had groupers in the 21-40 cm class size range, while $\sim 60\%$ of slope and spur-
197 and-groove sites had groupers within 31-50 cm total length (Fig. 3c). Although, the size

198 distribution of our study sites indicates that grouper >30cm TL (deemed 'large-bodied' by Mumby
199 et al. 2011) were frequently (over 50%) observed in patch reef habitats (Figure 3c), we caution
200 that other patch reefs across the Caribbean must be surveyed in order to make meaningful
201 extrapolations of the observed patterns in this habitat.

202 Other variables may also partly explain the variability of lionfish abundance in the region.
203 Wind exposure, specifically whether sites were located on the windward side, had a weak
204 negative effect on lionfish abundance (Fig. 1). However, the mechanism behind this association is
205 not well understood and a premature explanation may be misleading. Larval supply, which we
206 did not measure, may contribute to the lack of biotic resistance. As with other reef fish species
207 (James et al., 2002; Cowen & Sponaugle, 2009), differential larval supply could influence site-
208 specific lionfish recruitment (Ahrenholz & Morris, 2010). However, such data is not available for
209 our sites. While measuring larval supply would have been interesting, it was outside the scope of
210 our study due to the large number of sites included and the regional scale of the analysis.
211 Additionally, though larval supply can be predicted by biophysical models that describe
212 oceanographic features such as wind direction, surface temperature, or tidal amplitude, these
213 relationships are often taxon-dependent (Wilson & Meekan, 2001; Vallès et al., 2009).

214 The question from a management point of view is whether native predators can actually
215 constrain lionfish abundance across the Caribbean, given the heterogeneity of the systems and the
216 factors that seemingly affect lionfish abundance. While we found no evidence that large-bodied
217 grouper or any other large-bodied predators influence lionfish invasion success across the region,
218 this finding is expected based on other systems and examples of invasive predators. For example,
219 there is weak support in the literature for the biotic resistance hypothesis of native species
220 constraining exotic predators in natural ecosystems, and rarely can resident predators constrain
221 the distribution expansion of the invader (Harding, 2003; deRivera et al., 2005). In fact, the exact
222 opposite is typical in systems where native predators are abundant. For example, the successful

invasion of the Burmese python (*Python molurus bivittatus*) in the Everglades of South Florida has not been constrained by potential and abundant predators such as alligators (*Alligator mississippiensis*) (Willson et al., 2011). Moreover, it is common that invasive predators feed on the juveniles of the resident predators and competitors (Snyder & Evans, 2006; MacDonald et al., 2007; Doody et al., 2009; Kestrup et al., 2011; Willson et al., 2011; Côté et al., 2013), further weakening the potential resistance capacity of the system. Ecological interactions, such as predation and competition, seldom enable communities to resist invasion, but instead constrain the abundance of invasive species once they have successfully established (Levine et al., 2004). However, the abundance of lionfish across the region does not appear to be constrained by ecological interactions (Hackerott et al., 2013). In the one published record of grouper eating lionfish (Maljkovic et al., 2008), it could not be determined whether the lionfish were dead or alive when consumed. It is common for divers and tour operators to feed speared lionfish to native predators, including sharks (Busiello, 2011). However, there is no evidence that this practice has changed the natural predatory instincts of resident predators towards the invader and feeding speared lionfish to native predators is now being discouraged due to safety concerns for divers (Whittaker, 2013).

Our results indicate that protection status (i.e., whether sites were located within a marine reserve or not) also had a negative effect on lionfish abundance (Fig. 1). This is most likely due to targeted culling in protected areas. Morris and Whitfield (2009) suggested that lionfish removals should be focused on ecologically important areas, including marine protected areas and reserves. Lionfish removals have since occurred in many marine reserves through organized citizen programs (Biggs & Olden, 2011; López-Gómez et al., 2013) and by reef managers (author pers. comm. with Belize Audubon Society). This effort is paying off and has the potential to greatly reduce lionfish abundance, at least temporarily (Barbour et al., 2011; Frazer et al., 2012; Côté et al., 2013). In our dataset, of the six sites with grouper biomass over 20 gm⁻², five were in

248 protected areas where culling is very likely occurring (Fig. 2). This pattern supports the results of
249 our statistical analysis that lionfish abundance is reduced in marine protected areas due to some
250 factor other than predator abundance. The negative effect of protection status on lionfish
251 abundance and lack of effect of grouper or other predator biomass on lionfish abundance indicate
252 that culling within protected areas most likely explains the observed pattern.

253 This analysis expands our original statistical model of the relationship between invasive
254 lionfish and native grouper species (Hackerott et al., 2013) to include two additional covariates
255 hypothesized to moderate the relationship between these species (Mumby et al. (2013). After
256 accounting for these additional processes, we find that: (a) the biomasses of lionfish and large-
257 bodied grouper (or other predators) are not negatively related, and (b) lionfish biomass is
258 controlled by a number of physical site characteristics, as well as by culling within marine
259 reserves. Our study was motivated by the desire to explore whether the findings and solutions
260 from local case studies will be effective elsewhere, which is key to informed management
261 decisions about the invasion. We conclude that removals are most likely the only feasible
262 mechanism for controlling lionfish at a Caribbean-wide scale.

263 **Grant Disclosures**

264 National Science Foundation (OCE-0940019 to JFB), National Geographic Society Committee
265 for Research and Exploration (grant 8514-08 to JFB)

266 **Funding**

267 This work was funded in part by the National Science Foundation, the National Geographic
268 Society Committee for Research and Exploration, the Royster Society Carol and Edward
269 Smithwick Dissertation Fellowship (to AV), the Rufford Small Grants Foundation (to CEC), the

270 David H. Smith Conservation Research Fellowship (to SJG), and the University of North
271 Carolina at Chapel Hill. The funders had no role in study design, data collection and analysis,
272 decision to publish, or preparation of the manuscript.

273 **Acknowledgements**

274 We thank F. Pina for logistic support in Cuba. C. Layman, D. Knowles, The Bahamas National
275 Trust and Friends of the Environment provided logistic support in the Bahamas. We thank J.
276 Pawlik, M. Marti, and L. Deagan for their help and facilitation of a research expedition to
277 Mexico. We thank the Belize Fisheries Department, the Southern Environmental Association,
278 Healthy Reefs Initiative and the Toledo Institute for Development and Environment for support in
279 Belize.

280 **Permits**

281 Bahamas: Department of Marine Resources, Ministry of Agriculture and Marine Resources.
282 Permit MAF/FIS/17. Cuba: Centro de Control y Inspección Ambiental, Cuba via Fabian Pina.
283 Mexico: Dirección General de Ordenamiento Pesquero y Acuicola de la Comisión Nacional de
284 Acuicultura y Pesca (CONAPESCA) de la Secretaría de Agricultura, Ganaderia, Desarrollo
285 Rural, Pesca y Alimentación (SAGARPA). Permiso DAPA/2/06504/110612/1608. Belize: Belize
286 Fisheries Department. Permit # 000028-11.

287 **References**

288 Ahrenholz DW, Morris JA. 2010. Larval duration of the lionfish, *Pterois volitans* along the
289 Bahamian Archipelago. *Environmental Biology of Fishes*. 88(4):305–309.
290 Albins M, Hixon M. 2011. Worst case scenario: potential long-term effects of invasive predatory
291 lionfish (*Pterois volitans*) on Atlantic and Caribbean coral-reef communities. *Environmental*
292 *Biology of Fishes*. 1–7.

- 293 Albins MA, Hixon MA. 2008. Invasive Indo-Pacific lionfish *Pterois volitans* reduce recruitment
294 of Atlantic coral-reef fishes. *Marine Ecology Progress Series*. 367:233–238.
- 295 Barbour AB, Allen MS, Frazer TK, Sherman KD. 2011. Evaluating the Potential Efficacy of
296 Invasive Lionfish (*Pterois volitans*) Removals. *PLoS ONE*. 6(5):e19666.
- 297 Barbour AB, Montgomery ML, Adamson AA, Díaz-Ferguson E, Silliman BR. 2010. Mangrove
298 use by the invasive lionfish *Pterois volitans*. *Mar Ecol Prog Ser*. 401:291–294.
- 299 Betancur-R R, Hines A, Acero P. A, Ortí G, Wilbur AE, Freshwater DW. 2011. Reconstructing the
300 lionfish invasion: insights into Greater Caribbean biogeography. *Journal of Biogeography*.
301 38(7):1281–1293.
- 302 Biggs CR, Olden JD. 2011. Multi-scale habitat occupancy of invasive lionfish (*Pterois volitans*)
303 in coral reef environments of Roatan, Honduras. *Aquatic Invasions*. 6(3):447–453.
- 304 Bjørnstad ON. 2013. ncf: Spatial nonparametric covariance functions. *R package version 1.1-5*.
- 305 Bolker B, Skaug H, Magnusson A, Nielsen A. 2012. Getting started with the glmmADMB
306 package.
- 307 Bruno J, Fridley J, Bromberg K, Bertness M. 2005. Insights into biotic interactions from studies
308 of species invasions. *Species invasions: insights into ecology, evolution and biogeography*. 13–
309 40.
- 310 Busiello A. 2011. Pictures: Sharks Taught to Hunt Alien Lionfish. *National Geographic*.
- 311 Byers JE, Noonburg EG. 2003. Scale dependent effects of biotic resistance to biological invasion.
312 *Ecology*. 84(6):1428–1433.
- 313 Carlsson NO, Sarnelle O, Strayer DL. 2009. Native predators and exotic prey –an acquired taste?
314 *Frontiers in Ecology and the Environment*. 7(10):525–532.
- 315 Claydon JAB, Calosso MC, Traiger SB. 2012. Progression of invasive lionfish in seagrass,
316 mangrove and reef habitats. *Marine Ecology Progress Series*. 448:119–129.
- 317 Côté IM, Green SJ, Hixon MA. 2013. Predatory fish invaders: Insights from Indo-Pacific lionfish
318 in the western Atlantic and Caribbean. *Biological Conservation*. 164:50–61.
- 319 Cowen RK, Sponaugle S. 2009. Larval Dispersal and Marine Population Connectivity. *Annual*
320 *Review of Marine Science*. 1(1):443–466.
- 321 Davies KF, Chesson P, Harrison S, Inouye BD, Melbourne BA, Rice KJ. 2005. Spatial
322 heterogeneity explains the scale dependence of the native–exotic diversity relationship. *Ecology*.
323 86(6):1602–1610.
- 324 deRivera CE, Ruiz GM, Hines AH, Jivoff P. 2005. Biotic resistance to invasion: native predator
325 limits abundance and distribution of an introduced crab. *Ecology*. 86(12):3364–3376.
- 326 Doody JS, Green B, Rhind D, Castellano CM, Sims R, Robinson T. 2009. Population-level
327 declines in Australian predators caused by an invasive species. *Animal Conservation*. 12(1):46–

328 53.

329 Edmunds PJ, Elahi R. 2007. The demographics of a 15-year decline in cover of the Caribbean
330 reef coral *Montastraea annularis*. *Ecological Monographs*. 77(1):3–18.

331 Elton CS. 1958. The ecology of invasions by plants and animals. *Methuen, London*. 18.

332 Farmer NA, Ault JS. 2011. Grouper and snapper movements and habitat use in Dry Tortugas,
333 Florida. *Marine Ecology Progress Series*. 433:169–184.

334 Frazer TK, Jacoby CA, Edwards MA, Barry SC, Manfrino CM. 2012. Coping with the Lionfish
335 Invasion: Can Targeted Removals Yield Beneficial Effects? *Reviews in Fisheries Science*.
336 20(4):185–191.

337 Friedlander AM, DeMartini EE. 2002. Contrasts in density, size, and biomass of reef fishes
338 between the northwestern and the main Hawaiian islands: the effects of fishing down apex
339 predators. *Marine Ecology Progress Series*. 230:253–264.

340 Froese R, Pauly D. 2013. FishBase. *World Wide Web electronic publication (version 10/2013)*.

341 Graham MH. 2003. Confronting multicollinearity in ecological multiple regression. *Ecology*.
342 84(11):2809–2815.

343 Green S, Côté I. 2009. Record densities of Indo-Pacific lionfish on Bahamian coral reefs. *Coral*
344 *Reefs*. 28(1):107–107.

345 Green SJ, Akins JL, Ct IM. 2011. Foraging behaviour and prey consumption in the Indo-Pacific
346 lionfish on Bahamian coral reefs. *Marine Ecology Progress Series*. 433:159–167.

347 Green SJ, Akins JL, Maljković A, Côté IM. 2012. Invasive Lionfish Drive Atlantic Coral Reef
348 Fish Declines. *PLoS ONE*. 7(3):e32596.

349 Green SJ, Dulvy NK, Brooks ALM, Akins JL, Cooper AB, Miller S, Côté IM. 2013. Linking
350 removal targets to the ecological effects of invaders: a predictive model and field test. *Ecological*
351 *Applications*.

352 Green SJ, Tamburello N, Miller SE, Akins JL, Côté IM. 2012. Habitat complexity and fish size
353 affect the detection of Indo-Pacific lionfish on invaded coral reefs. *Coral Reefs*.

354 Gust N, Choat J, McCormick M. 2001. Spatial variability in reef fish distribution, abundance,
355 size and biomass: a multi-scale analysis. *Marine Ecology Progress Series*. 214:237–251.

356 Hackerott S, Valdivia A, Green SJ, Côté IM, Cox CE, Akins L, Layman CA, Precht WF, Bruno
357 JF. 2013. Native Predators Do Not Influence Invasion Success of Pacific Lionfish on Caribbean
358 Reefs. (F. Guichard, Ed.) *PLoS ONE*. 8(7):e68259.

359 Harding JM. 2003. Predation by blue crabs, *Callinectes sapidus*, on rapa whelks, *Rapana venosa*:
360 possible natural controls for an invasive species? *Journal of Experimental Marine Biology and*
361 *Ecology*. 297(2):161–177.

362 James MK, Armsworth PR, Mason LB, Bode L. 2002. The structure of reef fish metapopulations:

- 363 modelling larval dispersal and retention patterns. *Proceedings of the Royal Society of London.*
 364 *Series B: Biological Sciences.* 269(1505):2079–2086.
- 365 Jud ZR, Layman CA, Lee JA, Arrington DA. 2011. NOTE Recent invasion of a Florida (USA)
 366 estuarine system by lionfish *Pterois volitans* / *P. miles*. *Aquatic Biology.* 13(1):21–26.
- 367 Kestrup ÅM, Dick JTA, Ricciardi A. 2011. Interactions between invasive and native crustaceans:
 368 differential functional responses of intraguild predators towards juvenile hetero-specifics.
 369 *Biological Invasions.* 13(3):731–737.
- 370 Lesser MP, Slaterry M. 2011. Phase shift to algal dominated communities at mesophotic depths
 371 associated with lionfish (*Pterois volitans*) invasion on a Bahamian coral reef. *Biological*
 372 *Invasions.* 13(8):1855–1868.
- 373 Levine JM, Adler PB, Yelenik SG. 2004. A meta-analysis of biotic resistance to exotic plant
 374 invasions. *Ecology Letters.* 7(10):975–989.
- 375 Lockwood JL, Cassey P, Blackburn T. 2005. The role of propagule pressure in explaining species
 376 invasions. *Special issue: Invasions, guest edited by Michael E. Hochberg and Nicholas J. Gotelli.*
 377 20(5):223–228.
- 378 López-Gómez MJ, Aguilar-Perera A, Perera-Chan L. 2013. Mayan diver-fishers as citizen
 379 scientists: detection and monitoring of the invasive red lionfish in the Parque Nacional Arrecife
 380 Alacranes, southern Gulf of Mexico. *Biological Invasions.* 1–7.
- 381 MacDonald JA, Roudez R, Glover T, Weis JS. 2007. The invasive green crab and Japanese shore
 382 crab: behavioral interactions with a native crab species, the blue crab. *Biological Invasions.*
 383 9(7):837–848.
- 384 MacNeil MA, Graham NAJ, Polunin NVC, Kulbicki M, Galzin R, Harmelin-Vivien M, Rushton
 385 SP. 2009. Hierarchical drivers of reef-fish metacommunity structure. *Ecology.* 90(1):252–264.
- 386 Magoulick DD, Lewis LC. 2002. Predation on exotic zebra mussels by native fishes: effects on
 387 predator and prey. *Freshwater Biology.* 47(10):1908–1918.
- 388 Maljković A, Leeuwen TEV, Cove SN. 2008. Predation on the invasive red lionfish, *Pterois*
 389 *volitans* (Pisces: Scorpaenidae), by native groupers in the Bahamas. *Coral Reefs.* 27(3):501–501.
- 390 Maljkovic A, van Leeuwen TE, Cove SN. 2008. Predation on the invasive red lionfish, *Pterois*
 391 *volitans* (Pisces : Scorpaenidae), by native groupers in the Bahamas. *Coral Reefs.* 27:501–501.
- 392 Mantel N. 1967. The Detection of Disease Clustering and a Generalized Regression Approach.
 393 *Cancer Research.* 27(2 Part 1):209 –220.
- 394 Mazia NC, Chaneton EJ, Ghera CM, León RJ. 2001. Limits to tree species invasion in pampean
 395 grassland and forest plant communities. *Oecologia.* 128(4):594–602.
- 396 McClanahan TR, Graham NAJ, Calnan JM, MacNeil MA. 2007. Toward pristine biomass: reef
 397 fish recovery in coral reef marine protected areas in kenya. *Ecological Applications.* 17(4):1055–
 398 1067.

- 399 Melbourne BA, Cornell HV, Davies KF, Dugaw CJ, Elmendorf S, Freestone AL, Hall RJ,
400 Harrison S, Hastings A, Holland M, Holyoak M, Lambrinos J, Moore K, Yokomizo H. 2007.
401 Invasion in a heterogeneous world: resistance, coexistence or hostile takeover? *Ecology Letters*.
402 10(1):77–94.
- 403 Mitchell CE, Agrawal AA, Bever JD, Gilbert GS, Hufbauer RA, Klironomos JN, Maron JL,
404 Morris WF, Parker IM, Power AG, Seabloom EW, Torchin ME, Vázquez DP. 2006. Biotic
405 interactions and plant invasions. *Ecology Letters*. 9(6):726–740.
- 406 Mora C. 2008. A clear human footprint in the coral reefs of the Caribbean. *Proceedings of the*
407 *Royal Society B: Biological Sciences*. 275 (1636):767–773.
- 408 Morris JA, Akins JL. 2009. Feeding ecology of invasive lionfish (*Pterois volitans*) in the
409 Bahamian archipelago. *Environmental Biology of Fishes*. 86(3):389–398.
- 410 Mumby PJ, Brumbaugh DR, Harborne AR, Roff G. 2013. On the relationship between native
411 grouper and invasive lionfish in the Caribbean. *PeerJ PrePrints*.
- 412 Mumby PJ, Harborne AR, Brumbaugh DR. 2011. Grouper as a natural biocontrol of invasive
413 lionfish. *PLoS ONE*. 6(6):e21510.
- 414 Newton K, Côté IM, Pilling GM, Jennings S, Dulvy NK. 2007. Current and Future Sustainability
415 of Island Coral Reef Fisheries. *Current Biology*. 17(7):655–658.
- 416 Paddack MJ, Reynolds JD, Aguilar C, Appeldoorn RS, Beets J, Burkett EW, Chittaro PM, Clarke
417 K, Esteves R, Fonseca AC, Forrester GE, Friedlander AM, García-Sais J, González-Sansón G,
418 Jordan LKB, McClellan DB, Miller MW, Molloy PP, Mumby PJ, Nagelkerken I, Nemeth M,
419 Navas-Camacho R, Pitt J, Polunin NVC, Reyes-Nivia MC, Robertson DR, Rodríguez-Ramírez A,
420 Salas E, Smith SR, Spieler RE, Steele MA, Williams ID, Wormald CL, Watkinson AR, Côté IM.
421 2009. Recent region-wide declines in Caribbean reef fish abundance. *Current biology : CB*.
422 19(7):590–595.
- 423 Polunin NVC, Roberts CM. 1993. Greater biomass and value of target coral-reef fishes in two
424 small Caribbean marine reserves. *Marine Ecology-Progress Series*. 100:167.
- 425 R Core Team. 2013. R: A Language and Environment for Statistical Computing, Vienna, Austria.
- 426 Reusch TB, Williams SL. 1999. Macrophyte canopy structure and the success of an invasive
427 marine bivalve. *Oikos*. 398–416.
- 428 Sale PF. 2002. Coral Reef Fishes: Dynamics and Diversity in a Complex Ecosystem. Academic
429 Press.
- 430 Schofield P. 2009. Geographic extent and chronology of the invasion of non-native lionfish
431 (*Pterois volitans* [Linnaeus 1758] and *P. miles* [Bennett 1828]) in the Western North Atlantic and
432 Caribbean Sea. *Aquatic Invasions*. 4(3):473–479.
- 433 Schutte VGW, Selig ER, Bruno JF. 2010. Regional spatio-temporal trends in Caribbean coral reef
434 benthic communities. *Mar Ecol Prog Ser*. 402:115–122.
- 435 SEDAC (Socioeconomic Data and Applications Center). 2010. Global rural-urban mapping

- 436 project – settlement points. Columbia University, New York.
- 437 Skaug H, Fournier D, Nielsen A, Magnusson A, Bolker B. 2013. glmmADMB: Generalized
- 438 linear mixed models using AD model builder. *R package version 0.7.7*. 4.
- 439 Snyder WE, Evans EW. 2006. Ecological Effects of Invasive Arthropod Generalist Predators.
- 440 *Annual Review of Ecology, Evolution, and Systematics*. 37:95–122.
- 441 UNEP-WCMC, WorldFish Centre, WRI, TNC. 2010. Global distribution of warm-water coral
- 442 reefs, compiled from multiple sources, including the Millennium Coral Reef Mapping Project.
- 443 UNEP World Conservation Monitoring Centre. Cambridge (UK).
- 444 Vallès H, Hunte W, Kramer DL. 2009. Variable temporal relationships between environment and
- 445 recruitment in coral reef fishes. *Mar Ecol Prog Ser*. 379:225–240.
- 446 Whitfield P, Hare J, David A, Harter S, Muñoz R, Addison C. 2007. Abundance estimates of the
- 447 Indo-Pacific lionfish *Pterois volitans*-miles complex in the Western North Atlantic. *Biological*
- 448 *Invasions*. 9(1):53–64.
- 449 Whittaker J. 2013. Hunters warned not to feed lionfish to predators. *CayCompass*.
- 450 Willson JD, Dorcas ME, Snow RW. 2011. Identifying plausible scenarios for the establishment of
- 451 invasive Burmese pythons (*Python molurus*) in Southern Florida. *Biological Invasions*.
- 452 13(7):1493–1504.
- 453 Wilson DT, Meekan MG. 2001. Environmental influences on patterns of larval replenishment in
- 454 coral reef fishes. *Marine Ecology Progress Series*. 222:197–207.
- 455 Wilson S, Graham N, Polunin N. 2007. Appraisal of visual assessments of habitat complexity and
- 456 benthic composition on coral reefs. *Marine Biology*. 151(3):1069–1076.
- 457 Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Violation of Independence – Part
- 458 II. In: *Mixed effects models and extensions in ecology with R*. Springer New York, 161–191.

459 Figure and tables legends

460 Figure 1 **Coefficient estimates ($\pm$ 95% confident intervals) showing the effect of different**
461 **variables on lionfish abundance.** Lionfish counts were modeled with a generalized linear
462 mixed effect model using the automatic differentiation model builder (glmmADMB) based on a
463 negative binomial distribution type 1 and log link. Abundance values were obtained by adding the
464 log of survey area as offset in the model. Numerical variables (top axis, circles) and categorical
465 variables (bottom axis, squares) are on different scale for easy visual representation as the
466 magnitude effects of the former are relatively smaller. For full summary of the model see Table
467 S3.

468 Figure 2 **Relationship between mean grouper and lionfish biomass.** In this study, 71 fore reefs
469 (black dots protected sites, grey dots non-protected sites) were surveyed and analyzed across the
470 Caribbean. For comparison, we included 12 sites (red squares) surveyed at Exuma Cays Land and
471 Sea Park by Mumby et al., (2011). Red fitted line is for the linear regression model by Mumby et
472 al., (2011) that explain 56 % of the variability of lionfish biomass due to grouper abundance.
473 Note that red squares represent ~16 % of all sites. Boxplots are median (vertical or horizontal
474 line), 50 and 90 percentiles for lionfish biomass (right) and grouper biomass (top). Boxplots with
475 black dots (general mean) correspond to our study and boxplots with red squares (general mean)
476 to Mumby et al., (2011). Empty circle are outliers. Axes are in log scale.

477 Figure 3 **Histograms of grouper class size (total length in cm) by categories.** A) Class size
478 distribution for protected and non-protected sites, B) for sites with over and under 10 gm⁻² of
479 grouper biomass, and C) for reef habitat types. Note that over 90% of protected sites and sites
480 with >10 gm⁻² of grouper biomass have individuals >30 cm in total length. Only every other class
481 size has a label for clarity.

482 Figure S1 **Location of survey sites.** For site abbreviations, surveys dates and coordinates refer to
483 Table S1

484 Figure S2 **Moran's I similarity spline correlograms for lionfish and grouper raw data across**
485 **all sites (top two panels) and for the glmmADMB model residuals (bottom panel).** Note the
486 strong spatial autocorrelation of the raw data (i.e., swirling lines around zero) and how the
487 hierarchical structure of the random effects (sites nested in regions) of the full glmmADMB
488 model eliminated this correlation in the model residuals. A Mantel test of the model Pearson
489 residuals ($r = 0.073$) corroborates the lack of spatial correlation of the residuals. Lines are the
490 mean $\pm$ 95% confidence interval.

491 Table S1 **Reef site detailed information.** Location names, coordinates, and site characteristics
492 used in the study. S&G, spur-and-groove.

493 Table S2 **Summary of the glmmADMB results.** Lionfish abundance (ind. 100 m⁻²) on grouper
494 biomass (g 100 m⁻²), predators, and other co-factors.

Figure 1

Coefficient estimates ($\pm$ 95% confident intervals) showing the effect of different variables on lionfish abundance.

Lionfish counts were modeled with a generalized linear mixed effect model using the automatic differentiation model builder (glmmADMB) based on a negative binomial distribution type 1 and log link. Abundance values were obtained by adding the log of survey area as offset in the model. Numerical variables (top axis, circles) and categorical variables (bottom axis, squares) are on different scale for easy visual representation as the magnitude effects of the former are relatively smaller. For full summary of the model see Table S3.




Figure 2

Relationship between mean grouper and lionfish biomass.

In this study, 71 fore reefs (black dots protected sites, grey dots non-protected sites) were surveyed and analyzed across the Caribbean. For comparison, we included 12 sites (red squares) surveyed at Exuma Cays Land and Sea Park by Mumby et al., (2011). Red fitted line is for the linear regression model by Mumby et al., (2011) that explain 56 % of the variability of lionfish biomass due to grouper abundance. Note that red squares represent ~16 % of all sites. Boxplots are median (vertical or horizontal line), 50 and 90 percentiles for lionfish biomass (right) and grouper biomass (top). Boxplots with black dots (general mean) correspond to our study and boxplots with red squares (general mean) to Mumby et al., (2011). Empty circle are outliers. Axes are in log scale.




Figure 3

Histograms of grouper class size (total length in cm) by categories.

A) Class size distribution for protected and non-protected sites, B) for sites with over and under 10 gm⁻² of grouper biomass, and C) for reef habitat types. Note that over 90% of protected sites and sites with >10 gm⁻² of grouper biomass have individuals >30 cm in total length. Only every other class size has a label for clarity.

