

A peer-reviewed version of this preprint was published in PeerJ on 5 February 2021.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.10873) (peerj.com/articles/10873), which is the preferred citable publication unless you specifically need to cite this preprint.

Eng F, Marin JE, Zienkiewicz K, Gutiérrez-Rojas M, Favela-Torres E, Feussner I. 2021. Jasmonic acid biosynthesis by fungi: derivatives, first evidence on biochemical pathways and culture conditions for production. PeerJ 9:e10873 <https://doi.org/10.7717/peerj.10873>

Jasmonic acid biosynthesis by microorganisms: Derivatives, first evidences on biochemical pathways and culture conditions for production

Felipe Eng^{1,2}, Krzysztof Zienkiewicz¹, Mariano Gutiérrez-Rojas³, Ernesto Favela-Torres³, Ivo Feussner^{Corresp. 1}

¹ Plant Biochemistry, University of Goettingen, Goettingen, Germany

² Biotechnology Division, Cuban Research Institute on Sugar Cane Byproducts, Havana, Cuba

³ Campus Iztapalapa, Biotechnology Department, Universidad Autónoma Metropolitana, Mexico City, Mexico

Corresponding Author: Ivo Feussner

Email address: ifeussn@uni-goettingen.de

Jasmonic acid (JA) and its derivatives (called jasmonates) are lipid-derived signalling molecules that are produced by certain bacteria, fungi and plants. Beside this function, jasmonates have a great variety of applications in the flavour and fragrances production. In addition, they may have a high potential in agriculture. JAs protect plant against infections and may suppress the growth of cancer cells in humans and animals. Although a lot of information on the biosynthesis and function of JA exists from plants, knowledge on these aspects is still scarce for microorganisms. Taking into account the practical importance of JA, the objective of this review is to summarize knowledge on the occurrence of jasmonates from microbial culture media, their biosynthetic pathways and the culture conditions for optimal JA production as an alternative source for the production of these valuable metabolites

1 Review Article

2 **Jasmonic acid biosynthesis by microorganisms: Derivatives, first evidences on biochemical**
3 **pathways and culture conditions for production**

4

5 **Felipe Eng^{1,2}, Krzysztof Zienkiewicz², Mariano Gutiérrez-Rojas³, Ernesto Favela-Torres³,**
6 **Ivo Feussner^{*,2,4,5}**

7

8 ¹ Biotechnology Division, Cuban Research Institute on Sugar Cane Byproducts, Havana, Cuba

9 ² Department of Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences, University
10 of Göttingen, Göttingen, Germany

11 ³ Biotechnology Department, Universidad Autónoma Metropolitana, Campus Iztapalapa, Mexico
12 city, Mexico

13 ⁴ Department of Plant Biochemistry, Göttingen Center for Molecular Biosciences (GZMB),
14 University of Göttingen, Göttingen, Germany

15 ⁵ Department of Plant Biochemistry, International Center for advanced Studies of Energy
16 Conversion (ICASEC), University of Göttingen, Göttingen, Germany

17

18

19

20 ***Correspondence:** Ivo Feussner, Department of Plant Biochemistry, Albrecht-von-Haller-
21 Institute for Plant Sciences, University of Göttingen, Justus-von-Liebig-Weg 11 37077
22 Göttingen, Germany; **Fax:**+49-551-39-5749; **Email:** ifeussn@uni-goettingen.de

23

Running Title: Microbial biosynthesis of jasmonates

Keywords: culture medium, fungi, jasmonic acid, metabolic pathway, oxylipin, submerged fermentation

27

Abbreviations: **10,11-OHJA**, 10,11-hydroxy jasmonic acid, **12-HSO₄-JA**, 12-hydroxy jasmonic acid sulfate, **16:3**, hexadecatrienoic acid, **18:3**, α -linolenic acid, **ABA**, abscisic acid, **AOC**, allene oxide cyclase, **AOS**, allene oxide synthase, **CA**, curcubic acid, **CAMe**, methyl curcubate, **CJ**, *cis*-jasmone, **ddh-JA**, 4,5-didehydro jasmonic acid, **dn-OPDA**, dinor-12-oxo-phytodienoic acid, **GA₃**, gibberellic acid, **JA**, jasmonic acid, **JA-Leu**, jasmonoyl leucine, **JMT**, JA carboxyl methyltransferase, **JAMe**, methyl jasmonate, **LOX**, lipoxygenase, **OPC-4**, 3-oxo-2(2'-pentenyl)-cyclopentane-1-butanoic acid, **OPC-6**, 3-oxo-2(2'-pentenyl)-cyclopentane-1-hexanoic acid, **OPC-8**, 10,11-dihydro-12-oxo-phytodienoic acid, **OPDA**, 12-oxo-phytodienoic acid, **SSF**, solid state fermentation, **TA**, tuberonic acid.

37

Abstract

Jasmonic acid (JA) and its derivatives (called jasmonates) are lipid-derived signalling molecules that are produced by certain bacteria, fungi and plants. Beside this function, jasmonates have a great variety of applications in the flavour and fragrances production. In addition, they may have a high potential in agriculture. JAs protect plant against infections and may suppress the growth of cancer cells in humans and animals. Although a lot of information on the biosynthesis and function of JA exists from plants, knowledge on these aspects is still scarce for microorganisms. Taking into account the practical importance of JA, the objective of this review is to summarize knowledge on the occurrence of jasmonates from microbial culture media, their biosynthetic

pathways and the culture conditions for optimal JA production as an alternative source for the production of these valuable metabolites.

1 Introduction

Jasmonic acid (JA) and its derivatives belong to a group of plant growth regulators called jasmonates (Wasternack & Feussner 2018). They belong to the large group oxidized lipid signalling molecules, so-called oxylipins (Gerwick et al. 1991). In plants, jasmonates derive either from α -linolenic acid (18:3n-3) or raughanic acid (16:3n-3) and their major representatives are the isomers (+)-7-iso-JA and (-)-JA. These compounds are widely distributed in algae (Andreou et al. 2009; Ueda et al. 1991; Vick & Zimmerman 1989), angiosperms (Wasternack & Hause 2013) and microorganisms (Abdala et al. 1999; Forchetti et al. 2007; Hause et al. 2007; Miersch et al. 1993b). They belong to the group of phytohormones playing important roles as growth inhibitors and by regulating plants defence responses (Pieterse et al. 2009; Wasternack et al. 2006).

Methyl jasmonate (JAMe) was firstly isolated as an odoriferous constituent of the essential oil of *Jasminun grandiflorum* and other plant species (Crabalona 1967; Demole et al. 1962). It is recognized as an important ingredient in high-grade perfumes, cosmetics and in the preparation of detergents, soaps and food aromas with floral notes (Asamitsu et al. 2006; Dhandhukia & Thakkar 2007c). JA was first isolated as plant growth inhibitor from cultures of the fungus *Lasiodiplodia theobromae* (synonym *Botryodiplodia theobromae*) (Aldridge et al. 1971).

With the development of efficient methods for detection and quantification of metabolites about thirty years ago, JA and JAMe attracted the attention of plant physiologists. The presence of these compounds in different parts of plants was initially correlated with their strong promotion

of senescence and inhibition growth in angiosperms when applied exogenously (Wasternack & Hause 2002). Although, these compounds act as growth inhibitors or senescence promoters at high concentration, they induce the expression of defensive genes at much lower levels. For instance, they promote the synthesis of proteinase inhibitors, enzymes of phytoalexin synthesis, thionins, defensins and the vegetative storage protein genes in plants against pathogen attack or wounding (Howe & Jander 2008).

Jasmonates however play an important role in agriculture nowadays by regulating the defensive systems of plants against pests and pathogens (Gális et al. 2009; Gavin et al. 2012; Hawkins et al. 2007; Heil et al. 2001; Rohwer & Erwin 2008; Sanches et al. 2017; Stout et al. 2002; Wasternack 2014). Their application seems to be in line with the principles of sustainable agriculture since they may be less aggressive to the environment than pesticides and mineral fertilizers (Secatto 2013).

Jasmonates may be also used in medicine in the future because they are able to suppress so far *in vitro* the growth of some cancerous cells lines from humans and rats (Fingrut et al. 2005; Fleisher 2005; Fleisher 2007; Goldin et al. 2007; Kniazhanski et al. 2008; Li et al. 2017; Zhang et al. 2015). When applied *in vitro* it was observed that they induce cancer cell death in a wide range of organs, mainly colon, lymphatic tissue, breast, skin, prostate and lung. It has been also observed in preliminary studies *in vitro*, that the application of JAMe to parasites such as *Plasmodium falciparum*, *Shistosoma mansoni* (Gold et al. 2003) and *Trichomonas vaginalis* (Ofer et al. 2008) had a cytotoxic effect leading to cell death. Jasmonates have also been evaluated as antidepressants as well as anti-aggressive and anti-inflammatory agents (Ghasemi Pirbalouti et al. 2014).

Furthermore, it has been observed that the addition of exogenous JAMe stimulates the production of many secondary metabolites, such as taxane derivatives in plant cell cultures of *Taxus* sp. These metabolites are also very promising anticancer drugs in humans. Studies have been conducted to optimize the production of these substances; focusing on their metabolic pathways, selecting more productive cell lines, optimizing cell culture processes, product purification, and up scaling of the whole process (Bai et al. 2004; Miller et al. 2008; Onrubia et al. 2013; Syklowska-Baranek et al. 2009; Tabata 2006; Wilson & Roberts 2012).

Currently most of the aroma compounds including jasmonates are extracted from natural plant sources. However, recent advances in metabolic engineering have generated a great interest to produce these substances from alternative sources (Gupta et al. 2015). An alternative and attractive route for producing jasmonates will be based on microbial biosynthesis and biotransformation. Microorganisms such as bacteria and yeast can be used at variable scales as a safe producers of flavours and fragrances (Gill & Valivety 1997). Most importantly, these microorganisms can be metabolically and genetically modified to enhance the production of the desired metabolites. Moreover, the production of aroma compounds from microbial cultures or their enzyme preparations offers several advantages over traditional methods. The microbial metabolites can be produced in large quantities by the use of a fermentation process and can give high yields in very good qualities with better product characteristics along with low economical costs (Gupta et al. 2015).

Currently there are numerous projects for sequencing the genomes of ascomycete fungi ongoing (<http://genome.jgi-psf.org/pages/fungi-1000-projects.jsf>) and one of them is the jasmonate producing fungus *L. theobromae*. From this project, valuable information will be available in the near future that will help to continue the analysis of fungal JA biosynthesis and other related

metabolites using a reverse genetic approach. In fact, the lasiodiplodin biosynthetic gene cluster from the genome of *L. theobromae* strain NBRC 31059 was expressed in *Saccharomyces cerevisiae* strain BJ5464 to obtain a phytotoxic polyketide that inhibited human blood coagulation factor XIIIa, mineral corticoid receptors and prostaglandin biosynthesis (Xu et al. 2014).

2 Survey Methodology

Scientific reports and patents dealing to the production and properties of JA are still steadily increasing (Ghasemi Pirbalouti et al. 2014; Raviv et al. 2013; Wasternack 2015). However, there are few reports related to the production of JA by microorganisms. Therefore, the objective of this review is to discuss the existing reports related to the microbial production of jasmonates with a focus on the type of microorganisms, biosynthetic pathways, and culture conditions. By screening the publicly available databases Free Patents Online (<http://www.freepatentsonline.com/>), Google Patents (<https://patents.google.com/>), Espacent (<https://worldwide.espacenet.com/>), Google Scholar (<https://scholar.google.de/>), PubMed (<https://www.ncbi.nlm.nih.gov/>) and Web of Science (<https://apps.webofknowledge.com/>), we aimed to cover the current status of the field and apologize to scientists whose work we overlooked.

3 Jasmonates from fungi

L. theobromae is a common phytopathogenic fungus capable of producing jasmonates at high level, as a result of its primary and secondary metabolism (Alves et al. 2008; Eng et al. 2016). Although, JA is produced as the main product, other jasmonates such as 9,10-didehydro JA, 11-

hydroxy JA and 12-hydroxy JA sulphate (12-HSO₄-JA) were formed to a lesser extent (Figure 1, Table 1) (Eng 2012; Miersch et al. 1987). Cucurbitic acid (CA) which may also be recognized as a phytohormone and synthesized by a so far unknown pathway has been also detected in trace amounts (Eng 2012; Miersch et al. 1987).

Overall eight hydroxy JAs (11-hydroxy JA, 12-hydroxy JA or tuberonic acid (TA), 8-hydroxy JA, 3-oxo-2(1-hydroxy-2'-pentenyl)-cyclopentane-1-butanoic acid and 3-oxo-2(4-hydroxy-2'-pentenyl)-cyclopentane-1-butanoic acid) were detected in the culture medium and biomass of *L. theobromae* strain D7/2 growing in a medium containing sucrose, soy flour, corn steep liquor and a mineral salt solution (Miersch et al. 1991). Twenty-two jasmonates were identified after 8 weeks of culture of *Fusarium oxysporum* f sp *matthiole* strain 247.61 grown in liquid potato-dextrose medium under static conditions (Miersch et al. 1999a). Among the metabolites produced, 9,10-dihydro-7-iso-jasmonoyl-isoleucine, jasmonoyl-isoleucine, 9,10-dihydro jasmonoyl-isoleucine, 3-oxo-2-(2-pentenyl)cyclopentane-1-butyric acid, 3-oxo-2-(2-pentenyl)cyclopentane-1-hexanoic acid and 3-oxo-2-pentylcyclopentane-1-octanoic acid were identified. These isoleucine conjugates were also produced during the culture of *Gibberella fujikuroi* (Miersch et al. 1992). Interestingly, *F. oxysporum* f sp *mattiole* was unable to accumulate any hydroxylated-JAs as shown for *L. theobromae* (Miersch et al. 1993b).

The occurrence of the JA-serine and JA-threonine conjugates was confirmed in the fermentation broth from *L. theobromae* strain 2334 using HPLC-ESI tandem mass spectrometry in negative ionization mode, while JA-glycine and JA-isoleucine conjugates were identified with the same technique but with positive ionization (Castillo et al. 2014). In higher plants, JA amino conjugates are regular constituents accumulating upon sorbitol treatment or wounding (Miersch et al. 1999a).

While the conjugating enzyme was first isolated from the flowering plant *Arabidopsis thaliana* (Staswick et al. 2002), the corresponding peptidase activity was isolated from *L. theobromae* strain D 7/2 (Hertel et al. 1997). This enzyme was capable of hydrolysing JA-conjugates with α -amino acids. The enzyme was purified by gel filtration, ion exchange and hydrophobic interaction chromatography. It was characterized as glycoprotein with a molecular mass of about 107 kDa and its amidohydrolase activity was very specific with regard to (-)-JA and α -amino acids with (*S*)-configuration. Therefore, the authors suggested that this fungus may need this enzyme during infection of the host plant for starting or modifying plant processes, e.g. senescence or the release of nutrients, probably being beneficial for the fungal growth.

JA, JAMe and three JA esters, named lasiojasmonates (botryosphaerilactone A, (3*S*,4*R*,5*R*)-4-hydroxymethyl-3,5-dimethyldihydro-2-furanone and (3*R*,4*S*)-botryodiplodin) were detected from culture filtrates of *Lasiodiplodia* sp strain BL101 isolated from declining grapevine plants that showed wedge-shaped cankers (Andolfi et al. 2014). However, phytotoxic assays recording necrotic lesions on grapevine and cork oak leaves demonstrated that only JA was found to be active.

The diversity of octadecanoid and jasmonoyl compounds found in the culture filtrate of these fungi rise the question whether the compounds are formed only or at least primarily during the interaction with plants and, if so, what the function of these compounds might be. Evidences suggest that fungal pathogens exploit host oxylipins to facilitate their development via inducing plant lipid metabolism to utilize plant oxylipins in order to promote G-protein-mediated regulation of sporulation and mycotoxin production in the fungus and use of host-ligand mimicry to manipulate plant defence responses from which the fungus benefits (Christensen & Kolomiets 2011). However, in others cases *F. oxysporum* colonization remains symptomless or even has

beneficial effects on plant growth and/or stress tolerance. Also in pathogenic interactions a lengthy asymptomatic phase usually precedes disease development. All this suggests for a sophisticated and fine-tuned interaction between *F. oxysporum* and its host and the molecular mechanisms underlying this balance are poorly understood (Di et al. 2016).

Recently, (i) phytotoxic metabolites were identified in the culture media of six species of *Lasiodiplodia* isolated in Brazil causing *Botryosphaeria* dieback of grapevine (Cimmino et al. 2017). As ascertained by LC-MS, only four of these strains (*L. brasiliense*, *L. crassispota*, *L. jatrophiicola* and *L. pseudotheobromae*) produced JA. *L. brasiliense* synthesized also (3*R*,4*S*)-4-hydroxymellein. This was the first report on JA production from these species. (ii) Recently, fungal-derived *cis*-jasmonone (CJ) was detected in *L. theobromae* strain MAFF 306027 (Matsui et al. 2017). These authors carried out studies of the metabolism of deuterium-labelled of 18:3-d₅, OPC:4-d₆, OPC:6-d₆, OPC:8-d₆ and *cis*-OPDA-d₅ to JAME-d₅ and/or CJ-d₅ in feeding experiments with this strain, revealing that the fungus produced CJ through a single biosynthetic pathway via iso-12-oxophytodienoic acid (*iso*-OPDA). Interestingly, it was suggested that the previously predicted decarboxylation step of 3,7-didehydroJA to afford CJ might be not involved in CJ biosynthesis in *L. theobromae* (Matsui et al. 2017). However, in plants CJ is synthesized from 18:3 via two biosynthetic pathways using JA and *iso*-OPDA as key intermediates.

4 Jasmonic acid biosynthetic pathway

3.1 Plants

Many reviews have summarized the developments on the biosynthetic pathway of JA in plants and our knowledge will be briefly summarized in the following section (Agrawal et al. 2004; Creelman & Mullet 1997; Goepfert & Poirier 2007; Hamberg & Gardner 1992; Schaller et al.

2005; Vick & Zimmerman 1984; Wasternack & Feussner 2018; Wasternack & Hause 2002; Wasternack & Hause 2013).

JA biosynthesis in plants starts with the liberation of α -linolenic acid (18:3(n-3)) or roughanic acid (16:3(n-3)) from the plastid envelope membranes by lipases (shown in Figure 1A for α -linolenic acid). This reaction as well as the next three steps of the pathway are localized in plastid ending with the formation of either *cis*-(+)-12-oxophytodienoic acid (OPDA) or dinor-oxophytodienoic acid (dn-OPDA), respectively. This is the result of the sequential action of the enzyme lipoxygenase (LOX), allene oxide synthase (AOS) and allene oxide cyclase (AOC) on 18:3(n-3) or 16:3(n-3). The next steps take place in peroxisomes where OPDA and dn-OPDA are activated and reduced to 10,11-dihydro-12-oxophytodienoic acid (OPC-8) and 3-oxo-2(2'-pentenyl)-cyclopentane-1-hexanoic acid (OPC-6) by 12-oxophytodienoate reductase isoenzyme 3 (OPR3), respectively. These reactions are followed by two or three rounds of β -oxidation, yielding OPC-6; 3-oxo-2(2'-pentenyl)-cyclopentane-1-butanoic acid (OPC-4) and finally (+)-7-*iso*-JA that rearranges into the (-)-JA isomer (with an molar ratio of 9:1 for (-)-JA/(+)-7-*iso*JA) (Wasternack & Hause 2013). JA can be further metabolized into its methyl ester (JAMe) JA carboxyl methyltransferase (JMT) (Cheong & Choi 2003; Seo et al. 2001), or by conjugation with amino acids (such as leucine and isoleucine) or sugars, respectively (Sembdner & Parthier 1993; Wasternack 2016)}.

3.2 Microorganisms

Till today knowledge about metabolic pathways leading to the production of JAs by fungi and other microorganisms is scarce. Therefore, more physiological and biochemical studies are required and the existing knowledge will be summarized throughout the next paragraphs.

230 Starting with the products formed, the same ratio of isomers (-)-JA:(+)-7-isoJA that was found in
 231 plants was measured in the culture filtrate of *F. oxysporum* strain 247.61 (Miersch et al. 1999a).
 232 By contrast, only the (+)-7-isoJA isomer was found in a culture of *L. theobromae* strain D7/2
 233 (Miersch et al. 1987), but later both isomers, with a ratio of ~15:1 and 1:1 in two different
 234 experiments in the culture medium filtrate from *L. theobromae* strain 2334 were described
 235 (Jernerén et al. 2012).
 236 For *L. theobromae* it was shown in addition that JA production derived from 18:3(n-3) by using
 237 a culture medium that was supplemented either with ¹³C-sodium acetate or [²H₆]-18:3 (Tsukada
 238 et al. 2010). Appreciable amounts of [¹³C]-JA and [²H₅]-JA were detected in culture
 239 supernatants, and the methyl ester of OPDA was detected in mycelium extracts. Recently, by
 240 incubating mycelia from the JA-producing fungus *F. oxysporum* f. sp. *tulipae* with labelled 18:3,
 241 the plants-like intermediates allene oxide and 12-OPDA of the JA pathway were detected (Oliw
 242 & Hamberg 2017). The allene oxide was likely formed by a CYP enzyme or a catalase-related
 243 hydroperoxidase. These results suggest that JA is synthesised by this strain of *L. theobromae*
 244 starting from 18:3 via OPDA and that the enzymes being involved may be similar to those
 245 governing JA biosynthesis in higher plants. However, there are probably also some differences in
 246 the genes and enzymes of the JA pathway between plants and fungi. For example, although
 247 higher plants and the fungus *G. fujikuroi* produce structurally identical gibberellins (GAs) using
 248 similar steps, there are important differences in pathways and enzymes involved (Hedden et al.
 249 2002). These profound differences suggest that higher plants and fungi have evolved their
 250 complex biosynthetic pathways to GAs independently and not by horizontal gene transfer.
 251 In fact, the fatty acid composition in *L. theobromae* strain 2334 showed that the mycelium
 252 contained polyunsaturated C18 fatty acids, including 18:3(n-3) as probable substrate for JA

253 biosynthesis (Eng 2012; Eng et al. 2016; Jernerén et al. 2012). However, polyunsaturated C16
 254 fatty acids were not detected (Jernerén et al. 2012). OPDA and OPC:4 were also detected in
 255 culture filtrates from this fungus as probable intermediates on fungal JA pathway (Eng 2012;
 256 Eng et al. 2016). In addition, the JA precursors 3-oxo-2-pentylcyclopentane-1-butyric acid, 3-
 257 oxo-2-(2-pentenyl)cyclopentane-1-hexanoic and 3-oxo-2-(2-pentenyl)cyclopentane-1-octanoic
 258 acid were detected in a culture filtrate from *F. oxysporum* f sp *matthiole* strain 247.61 (Miersch
 259 et al. 1993a; Miersch et al. 1989).
 260 Interestingly, three bacterial strains producing JA, OPDA and ABA in a control culture medium
 261 were isolated from soil of sunflower cultures (Forchetti et al. 2007). Beside this observation
 262 nothing is known till now on how and under which conditions these mentioned strains form these
 263 phytohormones.
 264 Studies aiming at identifying single steps in fungal JA biosynthesis have been reported using
 265 different exogenously applied substrates (Jernerén et al. 2012), a reverse genetic approach
 266 (Brodhun et al. 2013) and enzyme purification (Patel et al. 2014). In the first case, a fatty acid
 267 dioxygenase activity from three strains of *Lasiodiplodia* was described (Jernerén et al. 2012).
 268 Two of the strains revealed low secretion of JA ($\sim 0.2 \text{ mg L}^{-1}$). These strains oxygenated 18:3(n-
 269 3) to 5,8-dihydroxy linolenic acid as well as to 9*R*-hydroperoxy linolenic acid, which was further
 270 metabolized by an AOS activity into 9-hydroxy-10-oxo-12*Z*,15*Z*-octadecadienoic acid.
 271 Analogous conversions were observed with linoleic acid (18:2(n-6)) as a substrate. Studies using
 272 [^{11}S - ^2H]18:2 revealed that the putative 9*R*-dioxygenase catalysed the stereospecific removal of
 273 the 11*R* hydrogen followed by a suprafacial attack of dioxygen at C-9. Mycelia from these
 274 strains contained 18:2 as the major polyunsaturated fatty acid but lacked 18:3(n-3). The third
 275 strain however secreted high amounts of JA ($\sim 200 \text{ mg L}^{-1}$). It contained 18:3(n-3) as major fatty

acid and produced 5,8-dihydroxy linolenic acid from exogenously added 18:3(n-3). Together, from these three strains no enzyme activity pointing to a JA pathway and being similar to that of higher plants could be identified.

Since no sequence information on the *L. theobromae* genome is yet available, a reverse genetic strategy focused on a 13-LOX from *F. oxysporum* that may initiate JA production was used as second approach. It was based on using sequences similar to those found from enzymes being part of the JA biosynthetic pathway of plants (Brodhun et al. 2013). One of the sequences called FoxLOX was cloned and expressed in *E. coli*. FoxLOX was found to be the first non-heme Fe-LOX, which oxidizes polyunsaturated C18 fatty acids to 13S-hydroperoxy derivatives by an antarafacial reaction mechanism where the *bis*-allylic hydrogen abstraction is the rate-limiting step. With 18:3 as substrate, FoxLOX was found to exhibit a multifunctional activity, because the hydroperoxy derivatives formed were further converted to dihydroxy-, keto-, and epoxy alcohol derivatives. The identification of FoxLOX as a specific linoleate 13S-LOX might hint towards a JA biosynthetic pathway in *F. oxysporum*, which is analogous to that in plants.

A LOX enzyme was purified from the mycelium of *L. theobromae* strain MTCC 3068 by chromatography (Patel et al. 2015). It was found that this fungus contains two LOXs isoenzymes, one of 93 kDa (LOX1) and another of 45 kDa (LOX2) with the later being most likely a degradation product of LOX1. Both LOX isozymes oxidized linoleic acid to produce a mixture of 9- and 13-hydroperoxy linoleic acid. Therefore, this LOX may be another candidate enzyme being involved in fungal JA production.

In summary, these are first data suggesting, that JA may be synthesised from 18:3(n-3) via OPDA in fungi. Since fungi do not have plastids, the reactions leading to the formation of OPDA most likely takes place in the cytosol or associated to a membrane leaflet facing the cytosol

(Figure 1B). Whether this pathway may be initiated by LOX enzymes or other dioxygenases is still unclear, just like the identity of the following enzymatic activities. The reactions downstream from OPDA however may follow a recently discovered pathway in *Arabidopsis thaliana* via direct β -oxidation of OPDA leading to formation of 4,5-didehydro jasmonic acid (ddh-JA) which is then reduced by a fungal OPR2 homologue to JA (Chini et al. 2018).

5 Chemical synthesis of jasmonates

Chemical synthesis and isolation of jasmonates from microorganisms and plants started in the 70s of last the century (Aldridge et al. 1971). JA is traditionally isolated from plants; mainly from jasmine and tea flowers; where JA is found in trace concentrations. A large number of flowers produce small amounts of essential oils. For instance, it takes about 500 Kg of petals to obtain approximately 1 Kg of rose oil. Therefore, this is a very expensive and time-consuming process that accounts for the high price of these oils (Dhandhukia & Thakkar 2007c).

Therefore, numerous chemical synthesis strategies for obtaining JA, JAME and other derivatives have been developed. In that way, the synthesis of JAME and methyl curcubate (CAME; Figure 2B, free fatty acid is shown as compound 1) have been reported by using 2-allylcyclohexan-1,3-dione as starting compound and hydroboration-oxidation followed of seven or eight steps for the first and second product, respectively (Kitahara et al. 1987). Moreover, the same authors improved the total yield for JAME to up to 20% in twelve reaction steps by improving the stereoselectivity of the hydroboration-oxidation by using 3-hydroxy methylcyclopentanone as starting compound (Kitahara et al. 1991).

Shortly after these reports, racemic 7-substituted derivatives of JAME have been synthesized (Taapken et al. 1994). 7-Methyl JAME was also synthesized in enantiomerically pure form in 7

steps from the Hajos-Wichert ketone. In addition, the biological activity of the prepared compounds has been investigated for the induction of tendril coiling in *Bryonia dioica* and the elicitation of the phytoalexin production in *Eschscholtzia californica*. However, beside 7-methyl JAMe all synthesized compounds showed poor activity in the bioassays (Taapken et al. 1994).

10 years later, Suzuki et al. developed a new method of JAMe and CAMe synthesis using a chiral tricyclic lactone as starting compound via a new type of tandem retro-Diels-Alder-ene reaction activated by a trimethylsilyl substituent as the key step, followed at seven reaction steps (Suzuki et al. 2004).

Other authors have dedicated their efforts to the synthesis of β -oxidation intermediates of JA, such as 10,11-dihydro-12-OPDA (OPC:8) by chemical or enzymatic means with good yields (Nonaka et al. 2010; Takayuki et al. 2003; Zerbe et al. 2007).

JA and TA (Figure 2B, free fatty acid is shown as compound 6) were synthesized from the key aldehyde, all *cis*-2-(2-hydroxy-5-vinylcyclopentyl)acetaldehyde, which was in turn prepared stereoselectively from the (1*R*)-acetate of 4-cyclopentene-1,3-diol through a S_N2 -type allylic substitution with $CH_2-CHMgBr$ followed by Mitsunobu inversion, Eschenmoser–Claisen rearrangement, and regioselective Swern oxidation of the corresponding bis-TES ether. Wittig reaction of the aldehyde with $[PH_3P(CH_2)Me]^+Br^-$ followed by oxidation afforded JA stereoselectivity over the *trans* isomer (Nonaka et al. 2010). Similarly, TA was synthesized.

Secatto proposed a racemic synthesis of JA involving additional steps to obtain higher yields (Secatto 2013). This would envisage an application at industrial scale. This synthetic route consisted of 7 steps with an overall yield of 30%. The improvement of this route is due to the use of an available starting compounds without hygroscopic characteristics and no requirement for

any pretreatment and easy handling. Moreover, the starting materials (adipic acid and cyclohexane and ethanol as solvents) are not expensive, leading overall to low production costs. Two macrolactones (JA-Ile-lactones) derived from 12-OH-JA-Ile were synthesized in 7 steps with an overall yield of 33% from commercially available JAMe (Jimenez-Aleman et al. 2015b). The biological activity of macrolactones was tested for their ability to elicit nicotine production, a well-known jasmonate dependent secondary metabolite. Both macrolactones showed strong biological activity, inducing nicotine accumulation to a similar extent as JAMe does in *Nicotiana attenuata* leaves. Surprisingly, the highest nicotine contents were found in plants treated with the JA-Ile-lactone, which has (3*S*,7*S*) configuration at the cyclopentanone ring and is not known from natural jasmonates.

A new synthetic route to JA-Ile-lactones was developed recently using the *Z*-selective cross-metathesis of (±)-MeJA and 3-butenyl acetate (both compounds commercially available and inexpensive) resulting in the (±)-1-acetate derivative in excellent yield (>80%) and *Z*-selectivity (> 90%) (Jimenez-Aleman et al. 2015a). Saponification of the (±)-1-acetate derivative (> 85 % yield) and conjugation to L-Ile resulted in the 1-hydroxy-12-L-Ile derivative. Finally, this derivative was exposed to macrolactonization resulting in enantiomerically pure macrolactones in only three steps. In agreement with the previous studies (Jimenez-Aleman et al. 2015b), these macrolactones also induced the accumulation of nicotine suggesting that these compounds open the possibility of uncoupling defence and growth in plants by using small molecules.

6 Microorganisms as producers of jasmonates

The first report about JA production by microbes was published already 50 years ago (Broadbent et al. 1968). These authors obtained JA from a culture of *L. theobromae* in a culture medium

367 containing glucose, glycerol or a mixture of both as carbon source, sodium nitrate, potassium
 368 nitrate or ammonium nitrate as nitrogen source. JA reached a concentration of 475 mg L⁻¹ and a
 369 productivity of 36.6 mg L⁻¹ d⁻¹. In order to purify the produced JA, biomass was removed by
 370 filtration and the filtrate was acidified and further extracted with ethyl acetate. Three years later
 371 JA biosynthesis was reported in a concentration of 500 mg L⁻¹ and a productivity of 38.4 mg L⁻¹
 372 d⁻¹ from *L. theobromae*, using a surface culture in 1L ceramic vessels with Czapek medium
 373 (Aldridge et al. 1971). These authors also observed that the culture supernatant inhibited the
 374 growth of higher plants and that the active component was JA. Similar results were obtained by
 375 *L. theobromae* strain D7/2 isolated from orange and cacao residues (Miersch et al. 1987). This
 376 strain was grown in a liquid medium based on sucrose, soybean meal, corn steep liquor and salt
 377 solution with a JA concentration and productivity of 500 mg L⁻¹ and 71 mg L⁻¹ d⁻¹, respectively.
 378 The same authors performed a screening for JA production using 46 species of *Ascomycetes* and
 379 *Basidiomycetes* belonging to 23 different genera (*Agrocybe*, *Aspergillus*, *Collybia*, *Coprinus*,
 380 *Cunninghamella*, *Daedalea*, *Fomes*, *Fusarium*, *Gleoporus*, *Homoconis*, *Marasmius*, *Mucor*,
 381 *Mycena*, *Paecilomyces*, *Phellinus*, *Penicillium*, *Pleurotus*, *Polyporus*, *Rhizoctonia*, *Stropharia*,
 382 *Talaromyces*, *Trametes* and *Trichoderma*) that were grown under the same conditions as *L.*
 383 *theobromae*. *Collybia*, *Coprinus* and *Mycena* were the best producers of JA. However, JA
 384 concentrations were four to eight times lower than produced by *L. theobromae* (Miersch et al.
 385 1993b). In addition, some mutants of *G. fujikuroi* were also able to produce free JA in culture
 386 supernatants but in trace amounts. Similarly, mycorrhizal fungi such as *Laccaria laccata* and
 387 *Pisolithus tinctorius* were identified as JA producers but again only in trace amounts (Miersch et
 388 al. 1999b).

A mutant approach was applied to obtain better JA producers of *L. theobromae* (Patel & Thakkar 2015). The mutants were generated using ethylmethanesulfonate and two mutants were isolated having the capacity to produce JA with 70 mg L⁻¹ and 78 mg L⁻¹ compared to wild type 32 mg L⁻¹.

The highest rates for JA production were described however for *Diplodia gossypina* strain ATCC 10936 (Farbood et al. 2001). Under optimal culture conditions JA concentration and productivity were 1200 mg L⁻¹ and 171 mg L⁻¹d⁻¹, respectively. This study even included the up scaling of JA production up to a volume of 150 L.

In case of bacteria a strain of *E. coli*, some rhizospheric bacteria such as *Azospirillum brasilense*, *Bradyrhizobium* sp. and *Rhizobium meliloti* (isolated from soils, which have been widely studied for their direct relationship with higher plants) and yeasts were capable of JA synthesis in concentrations as low as ng.L⁻¹ (Abdala et al. 1999; Forchetti et al. 2007). An endophytic diazotrophic bacterium was isolated from roots of the halophyte shrub *Prosopiss trombulifera* and probably identified as *Arthrobacter* sp (Piccoli et al. 2011). This strain produced abscisic acid, indole-3-acetic acid, gibberellins and JA in a chemically defined culture medium.

6 Culture conditions for JA production

Although the annual demand for JA increases primarily for applications in perfume production and flavourings (Dhandhukia & Thakkar 2007c), there are still only few reports published related to the practical aspects of the commercial production of JA (Farbood et al. 2001; Ghasemi Pirbalouti et al. 2014; Miersch et al. 1987).

The ability of fungi to produce JA varies between strains from 1 mg L⁻¹ to 1000 mg L⁻¹ of JA (Dhandhukia & Thakkar 2007c; Eng et al. 1998; Farbood et al. 2001). Therefore, at first strains

412 of *L. theobromae* or *D. gossypina* were screened for JA production in order to obtain better
413 strains (Altuna et al. 1996; Eng 2012; Farbood et al. 2001), because these two fungi seem to be
414 the species with the highest potential for JA production.

415 Next different culturing conditions were tested. Batch fermentation in static conditions using a
416 stationary Fernbach flask culture, an aseptic stationary tray culture or Erlenmeyers flasks were
417 tested between 5 to 10 days at temperature between 27 and 30 °C and slightly acidic initial pH
418 values between 5 and 6 of the culture medium (Altuna et al. 1996; Farbood et al. 2001; Miersch
419 et al. 1987). As a carbon source for producing JA soybean meal, citrus pulp, corn steep liquor
420 and milk serum were used and supplemented with oilseed meal, which can supply sources of
421 protein, minerals and water soluble vitamins (Miersch et al. 1987). However, the use of more
422 complex media had the drawback of needing more complicated processes for purifying JA for
423 some applications such as in perfumery, a removal of malodorous compounds and allergens is
424 required. Another drawback is that the composition thereof is not constant and therefore this
425 result is difficult to reproduce. Therefore, primarily synthetic media were used that are based on
426 sucrose or glucose as carbon source and mineral salts as potassium nitrate as nitrogen source,
427 with the addition of monobasic potassium phosphate, ammonium molybdate, and sulphate of
428 magnesium, iron, zinc and copper (Almeida et al. 1999; Miersch et al. 1987). However, also only
429 one type of carbon source (glucose or sucrose) can be used for JA production (Eng et al. 1998).
430 Already an early study showed that the addition of an inductor is not required to produce JA in
431 synthetic culture medium (Miersch et al. 1987). However, the addition of yeast extract and/or
432 soy peptone as a source of vitamins and cofactors to the culture medium stimulated the rate of JA
433 biosynthesis (Dhandhukia & Thakkar 2007b; Eng 2012; Eng et al. 2008; Farbood et al. 2001).

434 Under these standard culture conditions JA production took place at the late exponential growth
435 phase or stationary phase showing a behaviour similar to the accumulation of secondary
436 metabolites (Eng et al. 2016) and may only partially be associated with the growth phase of the
437 culture (Dhandhukia & Thakkar 2007c). Using these optimized culture conditions JA production
438 levels reached 500-1300 mg L⁻¹ and productivities of 28-170 mg L⁻¹ d⁻¹ (Dhandhukia & Thakkar
439 2007a; Dhandhukia & Thakkar 2007b; dos Santos et al. 2014a; dos Santos et al. 2014b; Eng
440 2012; Eng et al. 2016; Farbood et al. 2001; Inho et al. 2006).

441 Under static conditions, some *Lasiodiplodia* strains formed a mat on the surface of the culture
442 medium (Eng 1996). Therefore, the effect of the available surface area by increasing the vessel
443 size may be another critical aspect for JA production. This was confirmed by a study on JA
444 production by *L. theobromae* strain MTCC 3068 using the same amount of culture medium with
445 Erlenmeyer flasks of 250, 500 and 1000 mL in which the authors could show, that increasing the surface
446 area of the culture up to 1000 mL flasks lead to an increase JA yield (Dhandhukia & Thakkar 2007c).

447 In another study, the surface of the culture (100-500 mL) was simultaneously increased with the volume of
448 the culture medium (25-100 mL). Here, JA production was highest at the largest surface area in
449 combination with the highest volume of culture medium (Eng et al. 2016). However, an increase of
450 the flask volume to 5 or even to 50 L and for the culture medium volume up to 10 L did not lead to further
451 increases in JA yield (Eng 2012).

452 However, scaling up JA production in a fermenter or in a shaking incubator at 190 rpm and 30 °C
453 and dissolved oxygen saturation in the culture medium of up to 150 L (Farbood et al. 2001) as
454 well as using a fixed inoculation ratio of 0.5 g L⁻¹ of dry biomass of culture medium was shown
455 to improve JA yields (Miersch et al. 1987). In addition, it was of advantage to use homogenized
456 mycelium and not spores (Almeida et al. 2001; dos Santos et al. 2014a; dos Santos et al. 2014b).

Agitation turned out to be another critical aspect for JA production, because shaking speeds above 200 rpm lead to increased synthesis of extracellular polysaccharides (Selbmann et al. 2004), which had negative effect on JA production (Eng et al. 1998; Miersch et al. 1987). JA was also obtained by solid state fermentation (SSF) from *L. theobromae* strain 2334 using columns with sugar cane bagasse impregnated as support, at 30 °C and with a similar culture medium that was used in liquid fermentation. JA productivity was 2 times higher in solid state fermentation probably due to growth conditions that were more similar to the natural environment of this fungus (Eng 1996). Using similar conditions, JA productivity of a strain of *B. theobromae* isolated from cacao tissue was reported to be 4.8 times higher by SSF as with submerged fermentation (Laredo-Alcalá et al. 2016).

Finally, it should be noted that JA production was possible with *D. gossypina* strain ATCC 10936 in stirred fermenters of 150 L at an agitation velocity of 450 rpm, but productivity decreased at about two times with respect to the production in 500 mL Erlenmeyer flasks agitated at a speed of 200 rpm (Farbood et al. 2001).

7 Patents

There is a growing number of patents describing the production and application of jasmonates since the 60's and their quantity have increased during the last decades showing a growing interest in these substance class (Ghasemi Pirbalouti et al. 2014) (Figure 3). At first, the patents dealing with the isolation, detection and culture conditions for production of jasmonates in microorganisms such as *L. theobromae* will be discussed (Aldridge et al. 1971; Broadbent et al. 1968; Farbood et al. 2001; Günther et al. 1989; Miersch et al. 1984).

479 Other topics deal with agricultural applications of jasmonates in order to improve plant yield by
 480 inducing plants defence against herbivores and pathogens (Dathe et al. 1990; Ryan & Farmer
 481 1991). Recently these effects were combined with new formulations for jasmonates in water in
 482 combination with herbicides, pesticides, bioactive or biological seed treatment components and
 483 semiochemicals (Ghasemi Pirbalouti et al. 2014; Marks 2012). The application of JAMe to
 484 grapes in order to improve the quality of machine-harvested raisin grapes allowed the harvest
 485 without damages to the fruit or plants associated with traditional mechanical harvesting and
 486 thereby eliminating the need for expensive hand picking (Ghasemi Pirbalouti et al. 2014).
 487 Meanwhile a method was also reported for improving the turf grass quality (Mcelroy 2011).
 488 There is an only one patent claiming the use of a JA extract to inhibit the growth of the
 489 bacterium *Leuconostoc sp.* and dextran production during the juice processing of the sugar cane
 490 industry (Michelena et al. 2010).
 491 Nowadays, patents about jasmonates have expanded to medicinal, cosmetic, and flavouring
 492 applications (Ghasemi Pirbalouti et al. 2014). During the last 20 years, the vast majority of
 493 studies and inventions claim that JA, JAMe and dihydroJAMe have anticancer activity against
 494 various forms of cancer (Fleischer & Fingrut 2007; Fleischer et al. 2012; Fleischer et al. 2010;
 495 Ghasemi Pirbalouti et al. 2014; Herzberg et al. 2006; Martinez et al. 2010). Additional patents
 496 focus on improving the convenience and safety of their administration and on expanding the
 497 applications for the treatment, for example, the use of nanocarriers in order to increase the
 498 solubility of jasmonates, because these compounds are poorly water-soluble, not allowing an
 499 application by an intravenous route without an efficient nanostructured carrier system. In
 500 addition, they are not easily delivered to the cancerous cells, usually being degraded before they
 501 reach the tumour cells (da Silva et al. 2014; Katona et al. 2015; Lopes 2014).

Jasmonates are also used as skin care and hair care products, e.g., for treating hair, the scalp, dry and greasy skin (Bababunmi 2005; Broady 2012; Dalko 2006; Ghasemi Pirbalouti et al. 2014; Malik 2006) and also in Bladder dysfunction (Ghasemi Pirbalouti et al. 2014).

Jasmone, JAMe, CJ and γ -Jasmolactone are considered as the main odorous principles in the essential oil of jasmine flowers (jasmine oil) used in perfumes (Steinegger & Hansel 1988). Other author described the use of dihydromethyl-JA as enhancer or imparter fragrances in or to a perfume composition, perfumed articles and colognes (Boden et al. 1993). In addition, jasmonates are also used to flavour fruit beverages, confectionery like sweets and candy, food products like cocoa and tooth cleansing products like toothpaste (Hurst et al. 2015; Hurst et al. 2011; Johnson et al. 1977; Mookherjee et al. 1981).

8 Conclusions

Beside plants, bacteria and fungi are additional producers of jasmonates, and those providing the highest yields for JA production are *Ascomycetes* from the genus *Lasiodiplodia* and *Diplodia*. There is a great of diversity of JAs that are produced by microorganisms, but JA, JAMe and dihydroJAMe are the one being most intensively studied because of their value in numerous applications. In the fungus *L. theobromae*, plant-type jasmonate derivatives such as hydroxy and amino acids conjugates, as methyl and sulphate ester occur. In addition, derivatives being specific for fungi such as hydroxy-lactones, didehydro or dihomom-JAs are found. However, till today the function of jasmonates being produced by these microorganisms is not known. However, it is tempting to assume that they are involved regulating the interaction between plants and microorganisms.

Strategies to produce jasmonates via chemical synthesis suffer still from low yields. In case of microbial production strategies, a number of promising strains from the genus *Lasiodiplodia* and *Diplodia* have been selected, but they suffer from producing jasmonate mixtures and elaborated product purification strategies are required to develop an industrial processes for jasmonate production.

The knowledge gained so far provides a promising basis for additional research on the interaction of these microorganisms with plants, the chemical nature of JA biosynthesis in fungi, mechanisms that regulate this pathway in fungi and other microorganisms and design simpler and viable technological strategies to produce jasmonates in these fungi in order to satisfy the high demand for these products.

Nowadays, it is envisioned that JA and its derivatives continue to be used in the biomedicine, cosmetic, food industries and in agriculture, with new biomedical applications and patents emerging with a better understanding of their mechanisms of action and their molecular interactions with biological targets.

Acknowledgements: This work was supported by Deutscher Akademischer Austauschdienst (DAAD) through a short stay scholarship to F.E. Studies in the lab of I.F. were financially supported by the Deutsche Forschungsgemeinschaft (DFG) in frame of the GRK-1422.

The authors have declared no conflict of interest.

References

Abdala G, Miersch O, Correa N, and Rosas S. 1999. Detection of jasmonic acid in cultures of *Escherichia coli* and *Saccharomyces cerevisiae*. *Natural Product Letters* 14:55-63. doi: 10.1080/10575639908045435.

- 548 Agrawal GK, Tamogami S, Han O, Iwahashi H, and Rakwal R. 2004. Rice octadecanoid
549 pathway. *Biochemical and Biophysical Research Communications* 317:1-15.
- 550 Aldridge DC, Galt S, Giles D, and Turner WB. 1971. Metabolites of *Lasiodiplodia theobromae*.
551 *Journal of the Chemical Society C*:1623-1627. doi: 10.1039/j39710001623.
- 552 Almeida G, Altuna B, and Michelena G. 2001. Espoluración en *Botryodiplodia theobromae*.
553 Influencia de la luz en la producción de picnidios. *Laboract Acta* 13:89-92.
- 554 Almeida G, Klibansky M, Altuna B, Eng F, Legrá S, and Armenteros S. 1999. Some
555 considerations of the carbon sources used in jasmonic acid production. *Revista*
556 *Iberoamericana de Micología* 16:146-148.
- 557 Altuna B, Klibansky M, Almeida G, Eng F, Legra S, and Armenteros S. 1996. Jasmonic acid
558 detection as a new plant growth regulator in strains of *Botryodiplodia theobromae*. *Revista*
559 *sobre los derivados de la cana de azucar* 30:16-21.
- 560 Alves A, Crous PW, Correia A, and Phillips AJL. 2008. Morphological and molecular data
561 reveal cryptic speciation in *Lasiodiplodia theobromae*. *Fungal Diversity* 28:1-13.
- 562 Andolfi A, Maddau L, Cimmino A, Linaldeddu BT, Basso S, Deidda A, Serra S, and Evidente A.
563 2014. Lasiojasmonates A-C, three jasmonic acid esters produced by *Lasiodiplodia* sp., a
564 grapevine pathogen. *Phytochemistry* 103:145-153. doi: 10.1016/j.phytochem.2014.03.016.
- 565 Andreou A, Brodhun F, and Feussner I. 2009. Biosynthesis of oxylipins in non-mammals.
566 *Progress in Lipid Research* 48:148-170. doi: 10.1016/j.plipres.2009.02.002.
- 567 Asamitsu Y, Nakamura Y, Ueda M, Kuwahara S, and Kiyota H. 2006. Synthesis and odor
568 description of both enantiomers of methyl 4,5-didehydrojasmonate, a component of jasmin
569 absolute. *Chemistry & Biodiversity* 3:654-659.
- 570 Bababunmi E. 2005. Formulation and method for treating skeletal muscle degeneration caused
571 by malnutrition and disease.
- 572 Bai J, Kitanabe M, Toyozumi K, Fu L, Zhang S, Dai J, Sakai J, Hiroe K, Yamori T, Turuo T,
573 and Ando M. 2004. Production of biologically active taxoids by a callus culture of *Taxus*
574 *cuspidata*. *Journal of Natural Products* 67:58-63.
- 575 Boden R, Fujioka F, and Hanna M. 1993. Fragrance use of hydromethyl jasmonic acid.
- 576 Broadbent D, Hemming HG, and Turner WB. 1968. Preparation of jasmonic acid.
- 577 Broady B. 2012. Use of jasmone for modulating melatonin production and calcification of the
578 pineal gland.
- 579 Brodhun F, Cristobal-Sarramian A, Zabel S, Newie J, Hamberg M, and Feussner I. 2013. An iron
580 13S-lipoxygenase with an α -linolenic acid specific hydroperoxidase activity from *Fusarium*
581 *oxysporum*. *PLoS ONE* 8:e64919. doi: 10.1371/journal.pone.0064919.
- 582 Castillo G, Torrecillas A, Nogueiras C, Michelena G, Sanchez-Bravo J, and Acosta M. 2014.
583 Simultaneous quantification of phytohormones in fermentation extracts of *Botryodiplodia*
584 *theobromae* by liquid chromatography-electrospray tandem mass spectrometry. *World*
585 *Journal of Microbiology & Biotechnology* 30:1937-1946. doi: 10.1007/s11274-014-1612-5.
- 586 Cheong JJ, and Choi YD. 2003. Methyl jasmonate as a vital substance in plants. *Trends in*
587 *Genetics* 19:409-413.
- 588 Chini A, Monte I, Zamarreño AM, Hamberg M, Lassueur S, Reymond P, Weiss S, Stintzi A,
589 Schaller A, Porzel A, García-Mina JM, and Solano R. 2018. An OPR3-independent pathway
590 uses 4,5-didehydrojasmonate for jasmonate synthesis. *Nature Chemical Biology* 14:171-178.
591 doi: 10.1038/nchembio.2540.
- 592 Christensen SA, and Kolomiets MV. 2011. The lipid language of plant-fungal interactions.
593 *Fungal Genetics and Biology* 48:4-14. doi: 10.1016/j.fgb.2010.05.005.

- Cimmino A, Cinelli T, Masi M, Reveglia P, da Silva MA, Mugnai L, Michereff SJ, Surico G, and Evidente A. 2017. Phytotoxic lipophilic metabolites produced by grapevine strains of *Lasioidiplodia* species in Brazil. *Journal of Agricultural and Food Chemistry* 65:1102-1107. doi: 10.1021/acs.jafc.6b04906.
- Cole SJ, Yoon AJ, Faull KF, and Diener AC. 2014. Host perception of jasmonates promotes infection by *Fusarium oxysporum* formae speciales that produce isoleucine- and leucine-conjugated jasmonates. *Molecular Plant Pathology* 15:589-600. doi: 10.1111/mpp.12117.
- Crabalona L. 1967. Presence of levorotatory methyl jasmonate, methyl cis-2(2-penten1-yl)-3-oxocyclopentenyl acetate in the essential oil of Tunisian rosemary. *Comptes rendus de l'Académie des sciences (Paris)* 264:2074-2076.
- Creelman RA, and Mullet JE. 1997. Biosynthesis and action of jasmonates in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 48:355-381.
- Cross BE, and Webster GRB. 1970. New metabolites of *Gibberella fujikuroi*. Part XV. *N*-jasmonoyl-isoleucine and *N*-dihydrojasmonoyl-isoleucine. *Journal of the Chemical Society C*:1839-1942. doi: 10.1039/j39700001839.
- da Silva IBRF, Scarpa MV, Rossanezi G, do Egito EST, and de Oliveira AG. 2014. Development and characterization of biocompatible isotropic and anisotropic oil-in-water colloidal dispersions as a new delivery system for methyl dihydrojasmonate antitumor drug. *International Journal of Nanomedicine* 9:867-876.
- Dalko M. 2006. Use of a (dihydro)jasmonic acid derivative for dry skin treatment.
- Dathe W, Miersch O, Sembdner G, Kochmann W, Kunchnel F, and Reinhardt P. 1990. Mittel zur Ertragssteigerung bei Erdbeeren.
- Demole E, Lederer E, and Mercier D. 1962. Isolement et détermination de la structure du jasmonate de méthyle, constituant odorant caractéristique de l'essence de jasmin. *Helvetica Chimica Acta* 45:675-685. doi: 10.1002/hlca.19620450233.
- Dhandhukia PC, and Thakkar VR. 2007a. Response surface methodology to optimize the nutritional parameters for enhanced production of jasmonic acid by *Lasioidiplodia theobromae*. *J Appl Microbiol* 105:636-643.
- Dhandhukia PC, and Thakkar VR. 2007b. Significant medium components for enhanced jasmonic acid production by *Lasioidiplodia theobromae* using plackett-burman design. *Current Trends in Biotechnology & Pharmacy* 1:79-86.
- Dhandhukia PC, and Thakkar VR. 2007c. Standardization of growth and fermentation criteria of *Lasioidiplodia theobromae* for production of jasmonic acid. *African Journal of Biotechnology* 6:707-712.
- Di X, Takken FLW, and Tintor N. 2016. How phytohormones shape interactions between plants and the soil-borne fungus *Fusarium oxysporum*. *Frontiers in Plant Science* 7:170.
- dos Santos AZ, Araujo Leite VL, de Mello Innocentini MD, and Lourenco MV. 2014a. Evaluation of process parameters in the production of jasmonic acid. *BMC Proceedings* 8:244. doi: 10.1186/1753-6561-8-S4-P244.
- dos Santos AZ, Innocentini MDM, and Lourenco MV. 2014b. Optimization of jasmonates bioproduction. *BMC Proceedings* 8:208. doi: 10.1186/1753-6561-8-S4-P208.
- Eng F. 1996. Jasmonic acid production from *Botryodiplodia theobromae* by liquid and solid state fermentation. M.Sc. Thesis. University of Autonomous Metropolitan, Campus Iztapalapa, Mexico.

- Eng F. 2012. Characterization of a new strain of fungus *Lasiodiplodia* sp 2334 for obtaining jasmonic acid and its application in agriculture. PhD Dissertation. University of Havana, Cuba.
- Eng F, Gutierrez-Rojas M, and Favela-Torres E. 2008. Studies of the effects of carbon:nitrogen ratio inoculum type and yeast extract addition on jasmonic acid production by *Botryodiplodia theobromae* Pat. strain RC1. *Revista Iberoamericana de Micología* 25:188-192.
- Eng F, Gutiérrez-Rojas M, and Favela-Torres E. 1998. Culture conditions for jasmonic acid and biomass production by *Botryodiplodia theobromae* in submerged fermentation. *Process Biochemistry* 33:715-720.
- Eng F, Haroth S, Feussner K, Meldau D, Rekhter D, Ischebeck T, Brodhun F, and Feussner I. 2016. Optimized jasmonic acid production by *Lasiodiplodia theobromae* reveals formation of valuable plant secondary metabolites. *PLoS ONE* 11:e0167627. doi: 10.1371/journal.pone.0167627.
- Farbood MI, Blocker RW, McLean LB, Sprecker MA, McLean MP, Kossiakoff N, Kim AY, and Hagedorn M. 2001. Bioprocess for the high-yield production of food flavor-acceptable jasmonic acid and methyl jasmonate, United States Patent 6333180 (International Flavors & Fragrances Inc, IC:C12P 017/02, New York USA).
- Fingrut O, Reischer D, Rotem R, Goldin N, Altboum I, Zan-Bar I, and Fleisher E. 2005. Jasmonates induce nonapoptotic death in high-resistance mutant p53-expressing b-lymphoma cells. *British Journal Pharmacology* 146:800-808.
- Fleischer E, and Fingrut O. 2007. A jasmonate pharmaceutical composition for treatment of cancer.
- Fleischer E, Herzberg M, and Kashman Y. 2012. Chemical derivatives of jasmonate, pharmaceutical compositions and methods of use thereof.
- Fleischer F, Kashman Y, Reischer D, and Shimony S. 2010. Jasmonate derivative compounds pharmaceutical compounds and methods of use thereof.
- Fleisher E. 2005. Jasmonates, a new family of anti-cancer agents. *Anticancer Drugs* 16:911-916.
- Fleisher E. 2007. Jasmonates in cancer therapy. *Cancer Lett* 245:1-10.
- Forchetti G, Masciarelli O, Alemano S, Alvarez D, and Abdala G. 2007. Endophytic bacteria in sunflower (*Helianthus annuus* L.): isolation, characterization, and production of jasmonates and abscisic acid in culture medium. *Applied Microbiology Biotechnology* 76:1145-1152.
- Gális I, Gaquerel E, Pandey SP, and Baldwin IT. 2009. Molecular mechanisms underlying plant memory in JA-mediated defence responses. *Plant, Cell & Environment* 32:617-627.
- Gavin AS, Faggion SA, Hernandez C, Lourenco MV, Franca SC, and Beleboni RO. 2012. Nematocidal effects of natural phyto regulators jasmonic acid and methyl-jasmonate against *Pratylenchus zae* and *Helicotylenchus* spp. *Natural Product Research* 10:1-8.
- Gerwick WH, Moghaddam M, and Hamberg M. 1991. Oxylin metabolism in the red alga *Gracilariopsis lemaneiformis*: Mechanism of formation of vicinal dihydroxy fatty acids. *Archives of Biochemistry and Biophysics* 290:436-444.
- Ghasemi Pirbalouti A, Sajjadi SE, and Parang K. 2014. A review (research and patents) on jasmonic acid and its derivatives. *Archiv der Pharmazie* 347:229-239. doi: 10.1002/ardp.201300287.
- Gill I, and Valivety R. 1997. Polyunsaturated fatty acids, part 2: Biotransformations and biotechnological applications. *Trends in Biotechnology* 15:470-478.
- Goepfert S, and Poirier Y. 2007. β -Oxidation in fatty acid degradation and beyond. *Current Opinion in Plant Biology* 10:245-251.

- Gold D, Pankova-Kholmyansky I, Fingrut O, and Flescher E. 2003. The antiparasitic actions of plant jasmonates. *Journal of Parasitology* 89:1242–1244.
- Goldin N, Heyfets A, Reischer D, and Flescher E. 2007. Mitochondria-mediated ATP depletion by anti-cancer agents of the jasmonate family. *J Bioenerg Biomembr* 39:51-57.
- Günther T, Miersch O, Fritsche W, and Sembdner G. 1989. Synthetic medium for manufacture of 7-iso-jasmonic acid with *Botryodiplodia theobromae*.
- Gupta C, Prakash D, and Gupta S. 2015. A biotechnological approach to microbial based perfumes and flavours. *Journal of Microbiology & Experimentation* 2:00034.
- Hamberg M, and Gardner HW. 1992. Oxylin pathway to jasmonates: biochemistry and biological significance. *Biochimica et Biophysica Acta* 1165:1-18.
- Hause B, Mrosk C, Isayenkov S, and Strack D. 2007. Jasmonates in abuscular mycorrhizal interactions. *Phytochemistry* 68:101-110.
- Hawkins LK, Luthe DS, Windham GL, and Williams WP. 2007. Effect of exogenous jasmonic acid application on *Aspergillus flavus* kernel infection and aflatoxin production in two maize hybrids (*Zea mays* L.). *Proceedings of the 2006 Multicrop Aflatoxin/Fumonisin Elimination & Fungal Genomics Workshop, Ft Worth (Texas):*45-48.
- Hedden P, Phillips AJL, Rojas CM, Carrera E, and Tudzynski B. 2002. Gibberellin biosynthesis in plants and fungi: a case of convergent evolution? *Journal of Plant Growth Regulators* 20:319-331.
- Heil M, Koch T, Hilpert A, Fiala B, Boland W, and Linsenmair KE. 2001. Extrafloral nectar production of the ant-associated plant, *Macaranga tanarius*, is an induced, indirect, defense elicited by jasmonic acid. *Proceedings of the National Academy of Sciences USA* 98:1083-1088.
- Hertel SC, Knöfel H-D, Kramell R, and Miersch O. 1997. Partial purification and characterization of a jasmonic acid conjugate cleaving amidohydrolase from the fungus *Botryodiplodia theobromae*. *FEBS Letters* 407:105-110.
- Herzberg M, Harel A, and Mang C. 2006. Jasmonate derivatives, pharmaceutical compositions and methods of use thereof.
- Howe GA, and Jander G. 2008. Plant immunity to insect herbivores. *Annual Review of Plant Biology* 59:41-66. doi: 10.1146/annurev.arplant.59.032607.092825.
- Hurst W, Stuart D, Calderon A, and Van Breemen R. 2015. Jasmonic acid compounds in cocoa products.
- Hurst W, Stuart D, Justavino A, and Van Breemen R. 2011. Jasmonic acid compounds in cocoa products.
- Inho G, Kyoungju K, and Yonghwi K. 2006. Optimal conditions for the production of (+)-jasmonic acid by *Diplodia gossypina* ATCC 10936. *The Korean Journal of Microbiology* 42:210-215.
- Jernerén F, Eng F, Hamberg M, and Oliw E. 2012. Linolenate 9R-dioxygenase and allene oxide synthase activities of *Lasiodiplodia theobromae*. *Lipids* 47:65-73. doi: 10.1007/s11745-011-3622-5.
- Jimenez-Aleman GH, Machado RAR, Gorls H, Baldwin IT, and Boland W. 2015a. Synthesis, structural characterization and biological activity of two diastereomeric JA-Ile macrolactones. *Organic & Biomolecular Chemistry* 13:5885-5893. doi: 10.1039/C5OB00362H.
- Jimenez-Aleman GH, Scholz SS, Heyer M, Reichelt M, Mithöfer A, and Boland W. 2015b. Synthesis, metabolism and systemic transport of a fluorinated mimic of the endogenous

- jasmonate precursor OPC-8:0. *Biochimica et Biophysica Acta* 1851:1545-1553. doi: 10.1016/j.bbalip.2015.09.002.
- Johnson F, Paul K, and Favara D. 1977. Process for preparing methyl jasmonate and related compounds.
- Katona DL, Vempati S, and Savir OS. 2015. Brazilian scientist is part of elite group of researchers fighting cancer, obtains an unprecedented patent in the United States. *Braz J Biol* 75:256-258.
- Kitahara T, Miura K, Warita Y, Takagi Y, and Mori K. 1987. Synthesis of (±)-methyl epijasmonate and (±)-methyl cucurbate. *Agricultural and Biological Chemistry* 51:1129-1133. doi: 10.1271/bbb1961.51.1129.
- Kitahara T, Warita Y, Abe M, Seya M, Takagi Y, and Mori K. 1991. Stereoselective synthesis of methyl epijasmonate. *Agricultural and Biological Chemistry* 55:1013-1017. doi: 10.1271/bbb1961.55.1013.
- Kniazhanski T, Jackman A, Heyfets A, Gonen P, Fleisher E, and Sherman L. 2008. Methyl jasmonate induces cell death with mixed characteristics of apoptosis and necrosis in cervical cancer cells. *Cancer Letters* 271:34-46.
- Koch T, Bandemer K, and Boland W. 1997. Biosynthesis of *cis*-jasmonate: A pathway for the inactivation and the disposal of the plant stress hormone jasmonic acid to the gas phase? *Helvetica Chimica Acta* 80:838-850.
- Laredo-Alcalá E, Michelena-Álvarez G, Iliná A, Borjas-Ramos J, Hernández-Castillo F, and Martínez-Hernández J. 2016. Uso de *Botryodiplodia theobromae* como productor de ácido jasmónico en dos sistemas de fermentación. *Revista Mexicana de Micología* 44:1-9.
- Li J, Chen K, Wang F, Dai W, Li S, Feng J, Wu L, Liu T, Xu S, Xia Y, Lu J, Zhou Y, Xu L, and Guo C. 2017. Methyl jasmonate leads to necrosis and apoptosis in hepatocellular carcinoma cells via inhibition of glycolysis and represses tumor growth in mice. *Oncotarget* 8:45965-45980. doi: 10.18632/oncotarget.17469.
- Lopes J. 2014. Compositions of jasmonate compounds.
- Malik S. 2006. Wound and skin care compositions.
- Marks M. 2012. Agricultural compositions containing a jasmonic acid or dihydrojasmonic acid salt.
- Martinez R, Pedreño G, Navarro S, and Romero L. 2010. Combined use of methyl jasmonate and cyclodextrins for the production of resveratrol.
- Matsui R, Amano N, Takahashi K, Taguchi Y, Saburi W, Mori H, Kondo N, Matsuda K, and Matsuura H. 2017. Elucidation of the biosynthetic pathway of *cis*-jasmonate in *Lasiodiplodia theobromae*. *Scientific Reports* 7:6688. doi: 10.1038/s41598-017-05851-7.
- Mcelroy S. 2011. Improved turfgrass quality.
- Michelena G, Bell A, Martínez A, Carrera E, Cerrutti G, López-Munguía A, and Olvera C. 2010. Biocida compuesto por una sal de jasmonato y otros aditivos de uso en la industria azucarera.
- Miersch O, Bohlmann H, and Wasternack C. 1999a. Jasmonates and related compounds from *Fusarium oxysporum*. *Phytochemistry* 50:517-523.
- Miersch O, Brückner B, Schmidt J, and Semblner G. 1992. Cyclopentane fatty acids from *Gibberella fujikuroi*. *Phytochemistry* 31:3835-3837. doi: 10.1016/S0031-9422(00)97537-X.
- Miersch O, Günther T, Frishe W, and Semblner G. 1993a. Jasmonates from different fungal species. *Natural Products Letters* 2:293-299.
- Miersch O, Günther T, Fritsche W, and Semblner G. 1993b. Jasmonates from different fungal species. *Natural Product Letters* 2:293-299. doi: 10.1080/10575639308043825.

- Miersch O, Preiss A, Sembdner G, and Schreiber K. 1987. (+)-7-iso-jasmonic acid and related compounds from *Botryodiplodia theobromae*. *Phytochemistry* 26:1037-1039.
- Miersch O, Regvar M, and Wasternack C. 1999b. Metabolism of jasmonic acid in *Pisolithus tinctorius* cultures. *Phyton - Annales Rei Botanicae* 39:243-247.
- Miersch O, Schmidt J, Sembdner G, and Schreiber K. 1989. Jasmonic acid-like substances from the culture filtrate of *Botryodiplodia theobromae*. *Phytochemistry* 28:1303-1305. doi: 10.1016/S0031-9422(00)97735-5.
- Miersch O, Schneider G, and Sembdner G. 1991. Hydroxylated jasmonic acid and related compounds from *Botryodiplodia theobromae*. *Phytochemistry* 30:4049-4051. doi: 10.1016/0031-9422(91)83464-V.
- Miersch O, Sembdner G, Schreiber K, Richter K, Kochmann W, and Fritche W. 1984. Verfahren zur Herstellung von Jasmonsäureanaloge.
- Miller K, Neilan BA, and Sze DM. 2008. Development of taxol and other endophyte produced anticancer agents. *Recent Patents in Anticancer Drug Discovery* 3:14-19.
- Mookherjee B, Wilson R, Schmitt F, and Vock M. 1981. Flavoring with mixture of nor-methyl jasmonate and dihydro-nor-methyl jasmonate.
- Nonaka H, Ogawa N, Maeda N, Yong-Gang W, and Kobayashi Y. 2010. Stereoselective synthesis of epi-jasmonic acid, tuberonic acid, and 12-oxo-PDA. *Organic & Biomolecular Chemistry* 8:5212-5223. doi: 10.1039/C0OB00218F.
- Ofer K, Gold D, and E. F. 2008. Methyl jasmonate induces cell cycle block and cell death in the amitochondriate parasite *Trichomonas vaginalis*. *International Journal for Parasitology* 38:959-968.
- Oliw EH, and Hamberg M. 2017. An allene oxide and 12-oxophytodienoic acid are key intermediates in jasmonic acid biosynthesis by *Fusarium oxysporum*. *Journal of Lipid Research* 58:1670-1680. doi: 10.1194/jlr.M077305.
- Onrubia M, Cusidó RM, Ramirez K, Hernandez-Vazquez L, Moyano E, Bonfill M, and Palazon J. 2013. Bioprocessing of plant in vitro systems for the mass production of pharmaceutically important metabolites: paclitaxel and its derivatives. *Current Medicinal Chemistry* 20:880-891.
- Patel D, Patel R, and Thakkar V. 2014. Purification, Characterization and Application of Lipxygenase Isoenzymes from *Lasioidiplodia theobromae*. *Applied Biochemistry and Biotechnology*:1-13. doi: 10.1007/s12010-014-1278-3.
- Patel DD, Patel RR, and Thakkar VR. 2015. Purification, characterization and application of lipxygenase isoenzymes from *Lasioidiplodia theobromae*. *Applied Biochemistry and Biotechnology* 175:513-525. doi: 10.1007/s12010-014-1278-3.
- Patel DD, and Thakkar VR. 2015. Screening and characterization of hyper jasmonic acid producing mutant of *Lasioidiplodia theobromae*. *The FASEB Journal* 29:Supplement 884.822.
- Piccoli P, Travaglia C, Cohen A, Sosa L, Cornejo P, Masuelli R, and Bottini R. 2011. An endophytic bacterium isolated from roots of the halophyte *Prosopis strombulifera* produces ABA, IAA, gibberellins A₁ and A₃ and jasmonic acid in chemically-defined culture medium. *Plant Growth Regulation* 64:207-210. doi: 10.1007/s10725-010-9536-z.
- Pieterse CMJ, Leon-Reyes A, Van der Ent S, and Van Wees SCM. 2009. Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology* 5:308-316.
- Raviv Z, Cohen S, and Reischer-Pelech D. 2013. The anti-cancer activities of jasmonates. *Cancer Chemother Pharmacol* 71:275-285. doi: 10.1007/s00280-012-2039-z.

- 820 Rohwer CL, and Erwin JE. 2008. Horticultural applications of jasmonates: a review. *Journal of*
821 *Horticultural Science & Biotechnology* 83:283-304.
- 822 Ryan CA, and Farmer EE. 1991. Method of inducing plant defense mechanisms.
- 823 Sanches PA, Santos F, Peñaflor MFGV, and Bento JMS. 2017. Direct and indirect resistance of
824 sugarcane to *Diatraea saccharalis* induced by jasmonic acid. *Bulletin of Entomological*
825 *Research* 107:828-838. doi: 10.1017/S0007485317000372.
- 826 Schaller F, Schaller A, and Stintzi A. 2005. Biosynthesis and metabolism of jasmonates. *Journal*
827 *Plant Growth Regulation* 23:179-199.
- 828 Secatto M. 2013. Desenvolvimento da síntese total do (±)-ácido jasmônico visando aplicação en
829 escala industrial. *MSc Dissertation, Universidade Federal São Carlos, Centro de Ciências*
830 *Exatas e de Tecnologia, Departamento de Química*.
- 831 Selbmann L, Crognale C, and Petruccioli M. 2004. Beta-glucan production by *Botryosphaeria*
832 *rhodina* in different bench-top bioreactors. *Journal of Applied Microbiology* 95:1074-1081.
- 833 Sembdner G, Atzorn R, and Schneider G. 1994. Plant hormone conjugation. *Plant Molecular*
834 *Biology* 26:1459-1481.
- 835 Sembdner G, and Parthier B. 1993. The biochemistry and the physiological and molecular
836 actions of jasmonates. *Annual Review of Plant Physiology and Plant Molecular Biology*
837 44:569-589.
- 838 Seo HS, Song JT, Cheong JJ, Lee YH, Lee YW, Hwang I, Lee JS, and Choi YD. 2001. Jasmonic
839 acid carboxyl methyltransferase: a key enzyme for jasmonate- regulated plant responses.
840 *Proceedings of the National Academy of Sciences USA* 98:4788-4793.
- 841 Staswick PE, Tiryaki I, and Rowe ML. 2002. Jasmonate response locus *JAR1* and several related
842 *Arabidopsis* genes encode enzymes of the firefly luciferase superfamily that show activity on
843 jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation. *The Plant Cell*
844 14:1405-1415.
- 845 Steinegger E, and Hansel R. 1988. *Lehrbuch der Pharmakognosie und Phytopharmazie*. Berlin:
846 Springer-Verlag Berlin.
- 847 Stout MJ, Zehnder GW, and Baur ME. 2002. Potential for the use of elicitors of plant resistance
848 in arthropod management programs. *Archives of Insect Biochemistry and Physiology* 51:222-
849 235.
- 850 Suzuki K, Inomata K, and Endo Y. 2004. Enantiocontrolled synthesis of jasmonates via tandem
851 retro-Diels–Alder–ene reaction activated by a silyl substituent. *Organic Letters* 6:409-411.
852 doi: 10.1021/ol036253p.
- 853 Syklovska-Baranek K, Pietrosiuk A, Kokoszka A, and Furmanowa M. 2009. Enhancement of
854 taxane production in hairy root culture of *Taxus media* var. *Hicksii*. *Journal of Plant*
855 *Physiology* 166:1950-1954.
- 856 Taapken T, Bleichert S, Weiler EW, and Zenk MH. 1994. Stereoselective synthesis of 7-
857 substituted jasmonic acid derivatives and investigation of their biological activity. *Journal of*
858 *the Chemical Society, Perkin Transactions 1* 11:1439-1442. doi: 10.1039/p19940001439.
- 859 Tabata H. 2006. Production of paclitaxel and the related taxanes by cell suspension cultural of
860 *Taxus* species. *Current Drug Target* 7:453-461.
- 861 Takayuki A, Michitaka M, and Yuichi K. 2003. Efficient total synthesis of 12-oxo-PDA and
862 OPC-8:0. *Journal of Organic Chemistry* 68:7825-7832.
- 863 Tsukada K, Takahashi K, and Nabeta K. 2010. Biosynthesis of jasmonic acid in a plant
864 pathogenic fungus, *Lasiodiplodia theobromae*. *Phytochemistry* 71:2019-2023.

- 865 Ueda J, Miyamoto K, Aoki M, Hirata T, Sato T, and Momotani Y. 1991. Identification of
866 jasmonic acid in *Chlorella* and *Spirulina*. *Bulletin of the University of Osaka Prefecture Ser B*
867 43:103-108.
- 868 Vick BA, and Zimmerman DC. 1984. Biosynthesis of jasmonic acid by several plant species.
869 *Plant Physiology* 75:458-461.
- 870 Vick BA, and Zimmerman DC. 1989. Metabolism of fatty acid hydroperoxides by *Chlorella*
871 *pyrenoidosa*. *Plant Physiology* 90:125-132.
- 872 Wasternack C. 2006. Oxylipins: biosynthesis, signal traduction and action. In: Hedden P, and
873 Thomas S, eds. *Annual Plant Reviews*, 185-222.
- 874 Wasternack C. 2014. Action of jasmonates in plant stress responses and development - Applied
875 aspects. *Biotechnology Advances* 32:31-39. doi: 10.1016/j.biotechadv.2013.09.009.
- 876 Wasternack C. 2015. How jasmonates earned their laurels: Past and present. *Journal of Plant*
877 *Growth Regulation*:1-34. doi: 10.1007/s00344-015-9526-5.
- 878 Wasternack C. 2016. Jasmonates: Synthesis, metabolism, signal transduction and action. *eLS*:1-
879 9. doi: 10.1002/9780470015902.a0020138.pub2.
- 880 Wasternack C, and Feussner I. 2018. The oxylipin pathways: Biochemistry and function. *Annual*
881 *Review of Plant Biology* 69:in press. doi: 10.1146/annurev-arplant-042817-040440.
- 882 Wasternack C, and Hause B. 2002. Jasmonates and octadecanoids: signals in plant stress
883 responses and development. *Progress in Nucleic Acid Research and Molecular Biology*
884 72:165-221.
- 885 Wasternack C, and Hause B. 2013. Jasmonates: biosynthesis, perception, signal transduction and
886 action in plant stress response, growth and development. An update to the 2007 review in
887 *Annals of Botany*. *Annals of Botany* 111:1021-1058. doi: 10.1093/aob/mct067.
- 888 Wasternack C, Stenzel I, Hause B, Hause G, Kutter C, Maucher H, Neumerkel J, Feussner I, and
889 Miersch O. 2006. The wound response in tomato - Role of jasmonic acid. *Journal of Plant*
890 *Physiology* 163:297-306.
- 891 Wilson SA, and Roberts SC. 2012. Recent advances towards development and
892 commercialization of plant cell culture processes for synthesis of biolmolecules. *Plant*
893 *Biotechnol J* 10:249-268.
- 894 Xu Y, Zhou T, Espinosa-Artiles P, Tang Y, Zhan J, and Molnar I. 2014. Insights into the
895 biosynthesis of 12-membered resorcylic acid lactones from heterologous production in
896 *Saccharomyces cerevisiae*. *ACS Chemical Biology* 9:1119-1127. doi: 10.1021/cb500043g.
- 897 Zerbe P, Weiler EW, and Schaller F. 2007. Preparative enzymatic solid phase synthesis of cis(+)-
898 12-oxo-phytodienoic acid - physical interaction of AOS and AOC is not necessary.
899 *Phytochemistry* 68:229-236. doi: 10.1016/j.phytochem.2006.10.010.
- 900 Zhang M, Zhang MW, Zhang L, and Zhang L. 2015. Methyl jasmonate and its potential in
901 cancer therapy. *Plant Signaling & Behavior* 10. doi: 10.1080/15592324.2015.1062199.

903 Legends to the figures

904 **Figure 1.** Synthesis of JA and its amino acid-conjugate JA-Ile in plants (A) and fungi (B). Some
 905 known enzymes for Arabidopsis are indicated in yellow circles. Abbreviations: AOC, allene
 906 oxide cyclase; AOS, allene oxide synthase; ddh-JA, 4,5-didehydro jasmonic acid; JA, jasmonic
 907 acid; JA-Ile, jasmonic acid isoleucine conjugate; JAR1, jasmonoyl amino acid conjugate
 908 synthase; LOX, lipoxygenase; OPC-8, 3-oxo-2-(2-pentenyl)-cyclopentane-1-octanoic acid;
 909 OPDA, *cis*-(+)-12-oxo-phytodienoic acid; OPR, 12-oxo-phytodienoic acid reductase.

910 **Figure 2.** Chemical structure of the most important jasmonates found in fungi: **A:** **1**, jasmonic
 911 acid; **2**, jasmonoyl isoleucine, glycine, serine and threonine conjugates; **3**, 3-oxo-2-
 912 pentylcyclopentane-1-butyric acid; **4**, 3-oxo-2-(2-pentenyl)cyclopentane-1-hexanoic acid; **5**, 3-
 913 oxo-2-(2-pentenyl)cyclopentane-1-octanoic acid; **6**, 9,10-didehydro-JA; **7**, 9,10-dihydro-7-iso-
 914 jasmonoyl-isoleucine; **8**, 3-oxo-2-(2-pentenyl)cyclopentane-1-butyric acid; **9**, 3-oxo-2-(2-
 915 pentenyl)cyclopentane-1-hexanoic acid; **10**, 3-oxo-2-(2-pentenyl)cyclopentane-1-octanoic acid
 916 (all of them was found with *trans*- or *cis*-attached side chains). **B:** **1**, curcubic acid; **2**, 8-hydroxy
 917 jasmonic acid; **3**, 3-oxo-2-(1-hydroxy-2'-pentenyl)-1-butanoic-cyclopentenyl acid; **4**, 11-hydroxy
 918 jasmonic acid; **5**, 3-oxo-2-(4-hydroxy-2'-pentenyl)-cyclopentenyl-1-butanoic acid; **6**: tuberonic
 919 acid; **7**: 12-hydroxy jasmonic acid sulphate. **C:** Possible biosynthetic pathways for jasmonates
 920 detected in the culture filtrate of fungi (where A.1-10 and B.1-7 belong to the structure given
 921 under A and B, respectively)

922 **Figure 3.** Number of patents about applications for jasmonates reported in the agricultural
 923 literature (**A**), to obtain fragrances and flavours (**FF**), in medicine (**M**), in sugar cane industry (**I**)
 924 and for the isolation, detection and production of jasmonates (**P**).

Figure 1(on next page)


Synthesis of JA and its amino acid-conjugate JA-Ile in plants and fungi.

Synthesis of JA and its amino acid-conjugate JA-Ile in plants (A) and fungi (B). Some known enzymes for Arabidopsis are indicated in yellow circles. Abbreviations: AOC, allene oxide cyclase; AOS, allene oxide synthase; ddh-JA, 4,5-didehydro jasmonic acid; JA, jasmonic acid; JA-Ile, jasmonic acid isoleucine conjugate; JAR1, jasmonoyl amino acid conjugate synthase; LOX, lipoxygenase; OPC-8, 3-oxo-2-(2-pentenyl)-cyclopentane-1-octanoic acid; OPDA, *cis*-(+)-12-oxo-phytodienoic acid; OPR, 12-oxo-phytodienoic acid reductase


A

Chloroplast


PeerJ Preprints



Peroxisome




Cytosol



OPR2


JAR1



B

Cytosol

NOT REVEN

3 cycles of β -oxidation3 cycles of β -oxidation

ddh-JA





Figure 2(on next page)

Chemical structure of the most important jasmonates found in fungi.

Chemical structure of the most important jasmonates found in fungi: **A:** **1**, jasmonic acid; **2**, jasmonoyl isoleucine, glycine, serine and threonine conjugates; **3**, 3-oxo-2-pentylcyclopentane-1-butyric acid; **4**, 3-oxo-2-(2-pentenyl)cyclopentane-1-hexanoic acid; **5**, 3-oxo-2-(2-pentenyl)cyclopentane-1-octanoic acid; **6**, 9,10-didehydro-JA; **7**, 9,10-dihydro-7-iso-jasmonoyl-isoleucine; **8**, 3-oxo-2-(2-pentenyl)cyclopentane-1-butyric acid; **9**, 3-oxo-2-(2-pentenyl)cyclopentane-1-hexanoic acid; **10**, 3-oxo-2-(2-pentenyl)cyclopentane-1-octanoic acid (all of them was found with *trans*- or *cis*-attached side chains). **B:** **1**, curcubic acid; **2**, 8-hydroxy jasmonic acid; **3**, 3-oxo-2-(1-hydroxy-2'-pentenyl)-1-butanoic-cyclopentenyl acid; **4**, 11-hydroxy jasmonic acid; **5**, 3-oxo-2-(4-hydroxy-2'-pentenyl)-cyclopentenyl-1-butanoic acid; **6**: tuberonic acid; **7**: 12-hydroxy jasmonic acid sulphate. **C:** Possible biosynthetic pathways for jasmonates detected in the culture filtrate of fungi (where A.1-10 and B.1-7 belong to the structure given under A and B, respectively)

Figure 2

A




1 1 OH

2 1 (S)-Ile, -Gly, -Ser, -
Thr

3 3 OH

4 5 OH

5 7 OH



6 1 OH








7 1 (S)-Ile

8 3 OH

9 5 OH

10 7 OH

B



C




Figure 3(on next page)

Number of patents about applications for jasmonates reported in the agricultural literature.

Number of patents about applications for jasmonates reported in the agricultural literature (**A**), to obtain fragrances and flavours (**FF**), in medicine (**M**), in sugar cane industry (**I**) and for the isolation, detection and production of jasmonates (**P**)

Figure 3




Table 1(on next page)

Occurrence of jasmonic acid and other jasmonates from plants and microorganisms.

1 **Table 1.** Occurrence of jasmonic acid and other jasmonates from plants and microorganisms.

2

Jasmonates	Plant	Microorganism
Jasmonoyl isoleucine, glycine, serine, threonine, phenylalanine, tyrosine, tryptophan, leucine, isoleucine conjugates	(Hamberg & Gardner 1992)	(Castillo et al. 2014; Cole et al. 2014; Cross & Webster 1970; Miersch et al. 1999; Miersch et al. 1992)
9,10-didehydro-JA	(Hamberg & Gardner 1992)	(Eng 2012)
9,10-dihydro-7-iso-jasmonoyl-isoleucine	(Sembdner et al. 1994)	(Cross & Webster 1970; Miersch et al. 1999; Miersch et al. 1992)
3-oxo-2-(2-pentenyl)cyclopentane-1-butyric acid, 3-oxo-2-(2-pentenyl)cyclopentane-1-hexanoic acid, 3-oxo-2-(2-pentenyl)cyclopentane-1-octanoic acid	-	(Miersch et al. 1999)
Curcubic acid	(Sembdner & Parthier 1993)	(Eng 2012; Miersch et al. 1987)
8-hydroxy jasmonic acid	(Hamberg & Gardner 1992)	(Miersch et al. 1991)
11-hydroxy jasmonic acid	(Wasternack 2006)	(Miersch et al. 1991)
12-hydroxy jasmonic acid or tuberonic acid	(Hamberg & Gardner 1992; Wasternack 2006)	(Miersch et al. 1991)
12-hydroxy jasmonic acid lactone, tuberonic acid-O- β -glucopyranoside, curcubic acid-O- β -glucopyranoside	(Hamberg & Gardner 1992)	-

3-oxo-2-(1-hydroxy-2'-pentenyl)-1-butanoic-cyclopentenyl acid, 3-oxo-2-(4-hydroxy-2'-pentenyl)-cyclopentenyl-1-butanoic acid	-	(Miersch et al. 1991)
12-hydroxy jasmonic acid sulphate	Gidida et al., 2003	(Eng 2012)
4,5 didehydro-7-isojasmonic acid, 3,7-didehydrojasmonic acid, 6-epi-curcubic acid lactone, Homo-7-isojasmonic acid, Dihomo-7-isojasmonic acid, 11-hydroxy-dihomojasmonic acid, 8-hydroxy-dihomojasmonic acid	(Asamitsu et al. 2006; Hamberg & Gardner 1992)	-
<i>cis</i> -Jasmone	(Koch et al. 1997)	-
Methyl jasmonate	(Cheong & Choi 2003; Seo et al. 2001)	(Andolfi et al. 2014)