

A peer-reviewed version of this preprint was published in PeerJ on 31 January 2019.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.6338) (peerj.com/articles/6338), which is the preferred citable publication unless you specifically need to cite this preprint.

Colson V, Cousture M, Damasceno D, Valotaire C, Nguyen T, Le Cam A, Bobe J. 2019. Maternal temperature exposure impairs emotional and cognitive responses and triggers dysregulation of neurodevelopment genes in fish. PeerJ 7:e6338 <https://doi.org/10.7717/peerj.6338>

Maternal temperature exposure triggers emotional and cognitive disorders and dysregulation of neurodevelopment genes in fish

Violaine Colson^{Corresp., 1}, Morgane Cousture¹, Danielle Zanerato-Damasceno¹, Claudiane Valotaire¹, Thaovi Nguyen¹, Aurélie Le Cam¹, Julien Bobe^{Corresp., 1}

¹ Fish Physiology and Genomics, INRA LPGP UR1037, Rennes, France

Corresponding Authors: Violaine Colson, Julien Bobe
Email address: violaine.colson@inra.fr, julien.bobe@inra.fr

Fish are sensitive to temperature, but the intergenerational consequences of maternal exposure to high temperature on offspring adaptive behaviour and underlying mechanisms are unknown. Here we show that a thermal maternal stress induces emotional and cognitive disorders in offspring. Thermal stress in mothers triggered the inhibition of fear responses and decreased spatial learning abilities in progeny. Impaired behavioural phenotypes were associated with the dysregulation of several genes known to play major roles in neurodevelopment, including *auts2*, a key gene for neurodevelopment, more specifically neuronal migration and neurite extension, and critical for the acquisition of neurocognitive function. In addition, our analysis revealed the dysregulation of another neurodevelopment gene (*dpysl5*) as well as genes associated with human cognitive disorders (*arv1*, *plp2*). We observed major differences in maternal mRNA abundance in the eggs following maternal exposure to high temperature indicating that some of the observed intergenerational effects are mediated by maternally-inherited mRNAs accumulated in the egg. Together, our observations shed new light on the intergenerational determinism of fish behaviour and associated underlying mechanisms. They also stress the importance of maternal history on fish adaptive capacities in a context of global climate changes.

Maternal temperature exposure triggers emotional and cognitive disorders and dysregulation of neurodevelopment genes in fish

Violaine Colson¹, Morgane Cousture, Danielle Zanerato-Damasceno, Claudiane Valotaire, Thaovi Nguyen, Aurélie Le Cam, Julien Bobe

INRA, UR1037 Fish Physiology and Genomics (LPGP), F-35000 Rennes, France

Corresponding Author:

Violaine Colson¹

Email address: violaine.colson@inra.fr

13 Abstract

14 Fish are sensitive to temperature, but the intergenerational consequences of maternal
15 exposure to high temperature on offspring adaptive behaviour and underlying mechanisms
16 are unknown. Here we show that a thermal maternal stress induces emotional and
17 cognitive disorders in offspring. Thermal stress in mothers triggered the inhibition of fear
18 responses and decreased spatial learning abilities in progeny. Impaired behavioural
19 phenotypes were associated with the dysregulation of several genes known to play major
20 roles in neurodevelopment, including *auts2*, a key gene for neurodevelopment, more
21 specifically neuronal migration and neurite extension, and critical for the acquisition of
22 neurocognitive function. In addition, our analysis revealed the dysregulation of another
23 neurodevelopment gene (*dpysl5*) as well as genes associated with human cognitive
24 disorders (*arv1*, *plp2*). We observed major differences in maternal mRNA abundance in the
25 eggs following maternal exposure to high temperature indicating that some of the observed
26 intergenerational effects are mediated by maternally-inherited mRNAs accumulated in the
27 egg. Together, our observations shed new light on the intergenerational determinism of
28 fish behaviour and associated underlying mechanisms. They also stress the importance of
29 maternal history on fish adaptive capacities in a context of global climate changes.

30

31 Introduction

32 In the current context of global climate warming, wild and aquaculture fish are exposed to
33 varying environmental factors including suboptimal temperatures at specific periods of
34 their lifecycle. Fish are highly sensitive to extreme or abnormal (i.e. outside of the normal
35 physiological range) temperatures throughout their lifecycle, even for short periods of
36 time. This is especially true for key periods such as the reproductive period, during which
37 the female gamete undergoes final oocyte maturation. The direct impact on gamete quality
38 has been thoroughly investigated in many temperate species (see (Bobe & Labbe 2010;
39 Kjorsvik et al. 1990; Migaud et al. 2013) for review). Exposure of mature female fish to high
40 temperature during reproductive season (i.e. prior or around the time of ovulation) has a
41 dramatic impact on egg size (Jonsson & Jonsson 2016), egg viability and subsequent
42 embryonic success (Aegerter & Jalabert 2004) including reduced survival throughout
43 development. Despite this well documented negative impact on egg quality and subsequent
44 embryonic development, the long-term effects of maternal exposure to suboptimal
45 temperature on progeny behaviour and adaptive capacities remain unknown. More
46 specifically, the intergenerational consequences of mother exposure to abnormal
47 temperature on offspring emotional responses and cognitive performances – two key
48 components of fish adaptation and welfare – have never been investigated.

49 Several studies have however shown that maternal history can impact offspring behaviour
50 and adaptive capacities (i.e. ability of an organism to change its morphology, physiology, or
51 behaviour according to stressful environmental conditions (Bijlsma & Loeschcke 2005)).
52 This intergenerational effect on offspring behaviour was observed in salmonid fish in
53 which stress during reproductive season, or at least artificial exposure to stress hormones,

has a significant intergenerational impact on offspring adaptive capacities, including modifications of cognitive abilities (Sloman 2010) and emotional reactivity (Colson et al. 2015b; Eriksen et al. 2011; Espmark et al. 2008). In contrast, the underlying mechanisms mediating these effects remain poorly documented. In mammals, profound long lasting behavioural deficits have been observed in mice originating from stressed mothers, possibly due to epigenetic modifications occurring in the mother and transmitted to offspring (Weiss et al. 2011). In fish, a recent study has demonstrated the existence of the programming of stress axis function in zebrafish (*Danio rerio*) offspring by maternal social status (Jeffrey & Gilmour 2016). Another study showed that three-spined stickleback (*Gasterosteus aculeatus*) embryos respond to maternal exposure to predation risk via changes in gene expression (Mommer & Bell 2014).

The aim of this study was to thoroughly characterize the impact of high temperature exposure of female rainbow trout (*Oncorhynchus mykiss*) during reproductive season on offspring emotional and cognitive phenotypes, using specific behavioural tests previously validated in the laboratory (Colson et al. 2015a; Poisson et al. 2017; Sadoul et al. 2016). We also aimed at deciphering the molecular mechanisms mediating such intergenerational effects by analysing genome-wide gene expression in eggs and developing embryos following maternal exposure to high temperature.

73 **Material and methods**

74 **Ethics statement**

75 Fish were reared in INRA LPGP facilities, which hold full approval for animal
76 experimentation (C35-238-6). All fish were reared and handled in strict accordance with
77 French and European policies and guidelines of the INRA LPGP Institutional Animal Care
78 and Use Committee, which specifically approved this study (n° T-2016-55-VC-CV).

79 **Maternal treatment and fertilization**

80 Two-year old females rainbow trout were exposed to either 12°C (12°C group, standard
81 reproduction conditions) or 17°C (17°C group, high suboptimal temperature) for six weeks
82 before ovulation. The temperature of 17°C was selected because it is known to induce a
83 dramatic decrease in embryonic survival (Aegerter & Jalabert 2004). For each group, 30
84 marked (external tag placed on the dorsal fin) females were kept in 2.5 m³ tanks (2 x 2 x
85 0.62 m, length × width × water height). In the 17°C group, females initially reared at 12°C
86 were acclimated for five days to an increase of 1°C/day until 17°C. For three weeks
87 preceding ovulation, females were checked every two-three days to detect ovulation. In
88 both experimental group, eggs originating from four simultaneously ovulating females of
89 each group were collected and fertilized using a pool of sperm collected from four males
90 held at 12°C. Fertilization was performed immediately in both groups in order to avoid any
91 bias on subsequent behavioural phenotypes that would have been induced by differences
92 during embryo development. For each of the eight females, fertilizations of 800 eggs were
93 performed at 10°C using the medium ActiFish (IMV, L'Aigle, France; 100 ml ActiFish + 400
94 ml water) and fertilized eggs were distributed within a tray (20 x 50 cm) in two incubators

(10 x 10 cm) (approximately 400 eggs/incubator and two incubators/tray) supplied with 10°C flow-through recycled water. Each tray was covered with a lid to avoid exposure to light.

Monitoring of developmental success

Developmental success was monitored at eyeing stage, i.e. 19 days post-fertilization (dpf), hatching (32-33 dpf), and completion of yolk-sac resorption (YSR, 55 dpf) by counting dead embryos that were removed from incubators. The occurrence of malformations was obtained by taking a picture of euthanized malformed fry in each incubator at YSR. The types of malformed fry observed in this study were: torsion (T), yolk sac resorption defects (YSD) and other malformations (O) as described in (Bonnet et al. 2007). For each female, the occurrence of each type of malformations was calculated in comparison to the total number of malformed fry. Percentages of mortalities and malformations per incubator were obtained by counting the final number of live fry at swim-up stage, before transfer into rearing tanks.

Sample collection during embryo development

In both experimental groups, and in all egg clutches, biological samples were collected at four different stages: unfertilized eggs, around zygotic genome activation (i.e. 5 days post fertilization), hatching after removing yolk-sac, and YSR, which also corresponded to the stage of behavioural phenotyping. Entire (i.e. whole body) embryos were sampled. All samples were frozen in liquid nitrogen and held at -80°C until further processing.

115 Fry rearing

116 After vitellus resorption, at 55 dpf, swim-up fry from the two incubators per female were
 117 combined and transferred to seven distinct tanks (50 x 60 x 28 cm) (approximately 200
 118 fry/84 L), corresponding to seven different females (four from the 12°C group and three
 119 from the 17°C group). The mortality rate of one of the 17°C female was 98.4% and we did
 120 not obtain enough offspring to perform behavioural phenotyping. For this female, we only
 121 sampled the remaining fry to perform transcriptome analyses. Water temperature was
 122 maintained at 12°C. Fry received manually four meals per day with a commercial diet
 123 (Biomar, 48% protein and 22% lipid, 0.5 mm diameter pellets). Tanks were automatically
 124 illuminated from 8:00 to 20:00. Before each behavioural test, fish were starved for 24h. At
 125 the end of each test, fish were netted and transferred into individual bowls containing 250
 126 ml of the circuit water to which a lethal dose of anaesthetic (tricaine: 4.5 ml + bicarbonate
 127 of sodium: 5 ml) had been added.

128 Phenotyping of offspring behaviour

129 For each female (three 17°C females and four 12°C females), different hatchlings were
 130 subjected to the following behavioural tests thoroughly described in (Poisson et al. 2017):

131 Assessment of offspring emotional reactivity:

132 Fish propensity to express adapted fear-related behaviour (e.g. emotional reactivity) was
 133 evaluated individually in a novel-tank test at 75-76 dpf (social isolation in a novel tank).
 134 The novel tank (30 x 19 x 16 cm) was supplied with 12°C flow-through recycled water.
 135 Fifteen fish per female were observed. The treatment order was randomly chosen.

Behavioural responses were video-recorded for 30 minutes, divided into six 5-min intervals and analysed with EthovisionXT software (Noldus, Netherland). The following behavioural parameters were calculated for each individual: total distance travelled (cm), maximum swimming velocity (cm/sec), angular velocity ($^{\circ}$ /sec) (i.e. erratic swim), and time spent (%) in the border zone (i.e. thigmotaxis). At the end of the test, each fish was measured and weighed.

Assessment of offspring spatial learning abilities and memory:

Offspring propensity to locate a food-rewarded arm was assessed in a T-maze supplied with 12°C flow-through recycled water (see (Poisson et al. 2017) for a complete setup description). Five fish per female were tested. After 24h food deprivation and 30min of acclimation in the start-box of the T-maze, a remote-controlled guillotine door was pulled-up and fish occupation of the maze was video-recorded for a maximum of 1800 seconds per trial. A visual cue (black cross) was located on the wall of the T-maze at the entrance of the reward arm. When the fish crossed an invisible line separating the rewarded arm from the rest of the maze, a mechanic ridge, remotely-controlled by an experimenter observing live videos in an adjacent control room, released pellets. Then the fish was left to eat the pellets for at least 5min before being gently netted and introduced in its individual holding tank until the next trial. Eleven successive trials were run for four consecutive days (two trials on the first day and three on the other days). The treatment order was randomly chosen on the first day. We measured the latency to leave the start-box (Latency SB) and the latency to reach the reward arm after the fish had left the start-box (e.g. right choice, Latency RC). Retention of the acquired information (i.e. memory) was evaluated three days after the last

trial by another trial (resulting in a 2-day break). We measured Latency SB and latencies to make either the right (Latency RC) or the wrong choice (Latency WC).

Egg cortisol contain

Six eggs per female were homogenized in 600mL of deionized water using Precellys Evolution (Bertin Technologies, France). The program used was: 2mL CK14 tubes (work 4x20s 6800rpm + 30s break). Extraction of cortisol was performed using 200μL of homogenate (after short centrifugation at 3000g) and adding 2 mL of ethylacetate/cyclohexane (50/50, vol/vol) at room temperature. The supernatant was recuperated after strong mix and freezing at -20°C for at least 1 hour followed by another round of ethylacetate/cyclohexane extraction. After solvent evaporation, extracted cortisol was re-dissolved in 100μL ethanol. After evaporation, the dry residue was dissolved in 500μL buffer from cortisol ELISA kit purchased from Cayman Chemical (USA). Cortisol levels were determined following the ELISA kit manufacturer's instructions. The absorbance of each well was measured at 412 nm using a Synergy-2 microplate reader from BioTek (USA) instruments. Cortisol levels of eggs were calculated based on the calibration curve of absorbance. The assay has a range from 6.6-4000 pg/mL and a sensitivity (80% B/B0) of approximately 35 pg/mL.

Gene expression profiling

Transcriptome analysis was conducted using four egg batches originating from female held at 12°C and four egg batches originating from females held at 17°C, with the exception of YSR/12°C for which only three RNA samples of sufficient quality could be obtained. RNA was extracted from 20 eggs sampled at fertilization, 20 eggs at 5 dpf, six embryos at

hatching and six alevins sampled at YSR. Frozen tissues were lysed with Precellys Evolution Homogenizer (Ozyme, bertin technologies) in TRI Reagent (TR118, Euromedex) and total RNA was extracted according to the reagent's method followed by Nucleospin RNA isolation kit (740955, Macherey Nagel). Gene expression profiling was conducted using an Agilent 8x60K microarray (GPL24910) as previously described (Żarski et al. 2017). Samples were randomly distributed on the microarray for hybridization. The data were processed using the GeneSpring software (Agilent v.14.5) using gMedianSignal values. After data processing, one sample from the hatching/17°C group, which behaved differently from the other samples, even after normalization, was removed from subsequent analysis. Corresponding data were deposited in Gene Expression Omnibus (GEO) database (<https://www.ncbi.nlm.nih.gov/geo/>) under the reference GSE113377.

Statistics

Due to low number of synchronized females per treatment, percentage mortalities and malformations were compared between treatments using nonparametric Mann-Whitney tests (R, Mann-Whitney-Wilcoxon non-paired tests). Fish weights were analysed after taking into account the temperature as a fixed factor (two levels: 12°C and 17°C) and the females as a random factor. A generalized linear mixed model (GLMM) was fitted using the nlme package in R 3.3.1 (<http://cran.r-project.org/>), and by assuming a normal distribution. Significance of the random effect was checked using the 95% confidence interval of the variance, 0 being excluded of the interval in case of significance. The analyses of the novel tank test consisted in testing the effect of the temperature, the effect of the interval (two levels: 0-5 min and 25-30 min) and their interaction on each

203 dependant variable. The females and individuals (repeated measures) tested within
 204 treatments and intervals were defined as random factors in our statistical modelling.
 205 Distance travelled and maximum velocity were square root transformed, while angular
 206 velocity was log-transformed in order to reach normality and to fix GLMMs models using
 207 the nlme package. Significance of the random effect was checked using the 95% confidence
 208 interval of the variance, 0 being excluded of the interval in case of significance. When
 209 models were significant, post-hoc analyses were performed using HSD-Tukey tests. For
 210 thigmotaxis, data were too far from a normal distribution so we fixed a GLMM using the
 211 lme4 package, assuming a gamma distribution with inverse function. With this R package, a
 212 low variance associated with the random factor female indicated non-significant random
 213 effects.

214 The analyses of the spatial learning test consisted in testing the effects of the temperature,
 215 the trial (considered as a fixed factor with 11 levels, which are dependant data throughout
 216 the time), and their interaction on each dependant variable. The female and individuals
 217 (repeated measures) were defined as random factors. We fitted a GLMM model (using the
 218 lme4 package), assuming a normal distribution for Latency SB (after log transformation).
 219 Latency RC data did not reach normal distributions so we fitted a GLMM assuming a
 220 gamma distribution with inverse function. We fitted the same models for memory data
 221 considering the fixed factor trial with two levels (trial 11 vs. three days later). For latencies
 222 to make either the right or the wrong choice three days after the 11th trial, we also fixed a
 223 GLMM model (gamma distribution with inverse function) to test the effects of the
 224 temperature, the entry choice (Latencies RC and WC) made by the fish (fixed factor with

two levels, which are dependent data), and their interaction. When models were significant,
 the summary was considered for pairwise comparisons.
 For all models, if there were non-significant effects on factors or interactions, stepwise
 backward eliminations were performed to sequentially simplify the full model. The models
 were validated using analysis of residuals: normality assessment.
 Egg cortisol contains were compared between treatments using nonparametric Mann-
 Whitney tests.
 Differences were found to be significant when $P < 0.05$ and tendencies were considered for
 $0.05 < P < 0.1$. We indicated results of post-hoc analyses by different letters or by daggers
 symbols in the figures but they are not described in the Results section.
 For microarray analysis, gene expression data was scale normalized and $\log(2)$
 transformed before statistical analysis. The differences between the groups were analyzed
 using a two-way ANOVA with two factors (temperature, stage and their interaction), with a
 corrected P -value < 0.05 after Benjamini-Hochberg correction. For the four individual
 genes presented in Fig. 4C, non-parametric Mann-Whitney tests were performed between
 12°C and 17°C groups within egg and 5 dpf stages to reveal any significant differential
 expression.

Results

Influence of maternal exposure to high temperature on developmental success and growth

Maternal exposure to high temperature had a major impact on offspring survival. A dramatic increase in mortality was observed throughout early development when eggs originated from females held at 17°C even though this difference was not significant until hatching due to a high variability (Fig. 1). The overall median (quartiles: 25 and 75%) mortality rate was below 10% in the 12°C group, while it was over 40% in the 17°C group, with 6.62(5.30-7.59)% and 40.77(13.49-73.86)%, respectively ($W = 0$, $P < 0.05$). In contrast, no difference in the median (quartiles: 25 and 75%) occurrence of malformed fry was observed at yolk-sac resorption between the 12°C and the 17°C groups, with 7.57(6.44-8.76)% and 5.45(4.48-7.14)%, respectively ($W = 11$, $P = 0.48$). Similarly, the occurrence of the different types of malformation did not significantly vary among the experimental groups (Fig. S1). At 75 days post-fertilization, fish mean (\pm SEM) weight tended to be lower in 17°C than in 12°C (0.39 ± 0.02 g vs. 0.46 ± 0.05 g), although not significantly ($F_{1,5} = 4.88$, $P = 0.08$).

Offspring behaviour in the novel-tank test

Offspring from thermally stressed mothers displayed weaker emotional responses than controls when individually introduced into a novel-tank (Fig. 2). When considering distance travelled, the temperature X interval interaction tended to be significant ($F_{1,201} = 3.80$, $P = 0.05$, Fig. 2A), and a significant global increase was found between the first 5 min and the last 5 min (interval effect: $F_{1,201} = 4.90$, $P = 0.03$). Distance travelled did not differ

264 between 12°C and 17°C alevins (non significant temperature effect: $F_{1,5} = 0.002$, $P = 0.96$).
 265 Post-hoc tests are detailed in the Fig. 2 legend. The variance associated with the random
 266 factor female (4.63) was included in a confidence interval excluding 0, indicating that the
 267 random factor was significant. Maximum velocity did not differ between 12°C and 17°C
 268 ($F_{1,5} = 0.08$, $P = 0.79$, Fig. 2B). A significant global decrease was observed between the two
 269 intervals ($F_{1,201} = 22.89$, $P < 0.001$). No significant interaction was found ($P = 0.56$). The
 270 random factor female was significant. When considering angular velocity, the temperature
 271 X interval interaction was significant ($F_{1,201} = 5.58$, $P = 0.02$, Fig. 2C). No temperature or
 272 interval effects were found. Post-hoc tests are detailed in Fig. 2 legend. No temperature
 273 effect, interval effect or significant interaction was found for time spent in thigmotaxis (Fig.
 274 2D). The low variance (0.41) associated with the random factor female indicated a non-
 275 significant random effect.

276 Offspring spatial learning and memory

277 Offspring from thermally stressed mothers were slower to locate the rewarded arm than
 278 controls, when tested in a T-maze (Fig. 3). During the acquisition phase, we did not find any
 279 temperature ($\chi^2 = 1.65$, $df = 1$, $P = 0.20$) or trial effect ($\chi^2 = 12.01$, $df = 10$, $P = 0.28$) when
 280 considering the latency to leave the start-box. The temperature X trial interaction was not
 281 significant ($P = 0.15$). The low variances (< 0.001) associated with random factors female
 282 and individual indicated non-significant random effects. When considering the latency to
 283 make the right choice, we found a significant temperature X trial interaction ($\chi^2 = 19.49$, df
 284 $= 10$, $P = 0.03$, Fig. 3A). The summary of the model indicated a decrease in the latency from
 285 the 5th trial compared to trial 1 ($P < 0.05$), in the 12°C group only. These post-hoc effects
 286 are detailed in Fig. 3 legend. The fixed factors temperature and trial were not significant (P

287 = 0.20 and $P = 0.22$, respectively). The random factors female and individual were not
288 significant.

289 Memory data were considered between trial 11 and three days after. No temperature
290 effect, trial effect or significant interaction was found for the latency to leave the start-box
291 ($P > 0.1$). In 12°C group, the mean (\pm SEM) elapsed time between opening of the door and
292 fish exit was 164.95 ± 41.48 seconds three days after trial 11 (vs. 300.65 ± 92 sec, in trial
293 11), whereas in 17°C Latency SB reached 407.60 ± 145.15 sec (vs. 591.13 ± 152.87 sec in
294 trial 11). The variances associated with random factors female (0.10) and individual (1.58)
295 indicated significant random effects.

296 The latency to make the right choice tended to decrease between trial 11 and three days
297 later (trial effect: $\chi^2 = 3.25$, $df = 1$, $P = 0.07$, Fig. 3B). No temperature effect or significant
298 interaction was found. In 12°C group, the mean (\pm SEM) elapsed time between fish exit
299 from the start-box and entry into the rewarded arm was 296.8 ± 74.57 sec (vs. $625 \pm$
300 151.14 sec, in trial 11), whereas in 17°C, Latency RC was 607.27 ± 152.16 sec (vs. $1009.4 \pm$
301 193.62 sec in trial 11). The low variances associated with random factors female (< 0.001)
302 and individual (< 0.01) indicated non-significant random effects.

303 Three days after the 11th trial, we found a significant temperature X entry choice
304 interaction ($\chi^2 = 5.60$, $df = 1$, $P = 0.02$, Fig. 3C). A significant choice effect was found ($\chi^2 =$
305 16.02 , $df = 1$, $P < 0.001$), the elapsed time before entering into the right arm being lower
306 than into the wrong arm. The factor temperature was not significant.

307 Egg cortisol content

308 We did not observe any significant difference in the cortisol content of eggs originating
309 from different maternal groups. Median (quartiles: 25 and 75%) cortisol levels did not

310 differ between 17°C and 12°C unfertilized eggs, with 2.37 (1.84-3.88) ng/g vs. 4.11 (1.90-
311 7.13) ng/g, respectively ($W = 9$, $P = 0.89$).

312 **Gene expression profiling in embryos with different maternal history**

313 Gene expression profiling was performed in eggs and throughout development after
314 maternal exposure to either 12°C or 17°C. The ANOVA resulted in the identification of
315 47,711 differentially expressed genes throughout development. In contrast, a much lower
316 number of genes were differentially expressed in response to maternal exposure to high
317 temperature (Fig. 4A). Twelve genes exhibited a differential expression in response to
318 temperature while only 5 genes were differentially expressed in response to temperature
319 and among the developmental stages analysed (temperature X stage significant interaction:
320 $P < 0.05$). A total of sixteen genes were thus significantly dysregulated during development
321 in response to maternal exposure to high temperature, one gene (*srsf2a*) being present in
322 both groups. Among these genes, several were of particular interest due to their role in
323 neurodevelopment (*auts2*, *dpysl5*), neural disorder (*arv1*), and X-linked cognitive disability
324 (*plp2*), as discussed below. Interestingly the expression profiling analysis (Fig. 4B) revealed
325 that the differential expression between groups was especially marked in eggs, and to a
326 lower extend at 5 dpf, while differences were more limited during further development (i.e.
327 hatching and yolk-sac resorption stages). For *auts2* and *dpysl5* maternal mRNA abundance
328 was dramatically lower when females were exposed to high temperature ($W = 16$, $P < 0.05$;
329 Fig 4C), while *arv1* exhibited an opposite pattern ($W = 0$, $P < 0.05$). Similarly, *plp2*
330 abundance appeared higher in the 17°C group in eggs and 5 dpf embryos ($W = 0$, $P < 0.05$;
331 Fig.4C).

Discussion

Our aim was to investigate the effect of a thermal stress, applied to female rainbow trout during the peri-ovulatory period, on offspring behavioural phenotypes. As expected, the thermal stress triggered an increase in embryonic mortality, but not in the occurrence of malformed fry. In addition, fear responses to a novel environment were inhibited in 17°C offspring, which indicates emotional blunting. The thermal maternal stress also impaired spatial learning abilities in progeny. In consistency with these impaired behaviours, we observed a dysregulated expression of embryonic genes involved in neural and cognitive development revealed by a large-scale transcriptomic analysis.

Maternal effects on embryonic survival and development

Our results are in full agreement with previous reports on the deleterious effect of high temperature exposure in peri-ovulatory period on offspring survival in salmonids (rainbow trout: (Aegerter & Jalabert 2004), Atlantic salmon, *Salmo salar*: (King & Pankhurst 2004; King et al. 2003; Taranger & Hansen 1993), Arctic charr, *Salvelinus alpinus*: (Atse et al. 2002)). Despite small sample size (e.g. four females per treatment), differences between treatments were significant at hatching and yolk-sac resorption, but not at eyeing, which is also consistent with the results obtained by Aegerter et al. (2004). In addition, body weight measured at 75 dpf tended to be lower in offspring originating from high temperature-exposed females. This is consistent with previous studies performed on fish, which showed lower offspring survival rates and impaired growth after maternal cortisol administration (Eriksen et al. 2007; Eriksen et al. 2015) or maternal stress exposure (Campbell et al. 1994; McCormick 2009).

354 Maternal effects on emotional responses

355 The novel-tank test consisted in observing immediate fish behavioural responses when
356 individually transferred into a novel environment, which is a context known to elicit acute
357 stress responses in various vertebrates including salmonid fish species (Colson et al. 2018;
358 Colson et al. 2015a; Kittilsen et al. 2009; Overli et al. 2005; Rouger et al. 1998; Winberg et
359 al. 2007). Our results show that fishes originating from thermally stressed females were
360 less reactive to the challenging situation than controls. Angular velocity, which represents
361 erratic swimming and is commonly considered as an expression of fish anxiety (Blaser et al.
362 2010; Egan et al. 2009), tended to be lower in 17°C fish during the first 5 minutes of the
363 test. In a previous study performed on wild largemouth bass (*Micropterus salmoides*),
364 mature females were cortisol-injected (Redfern et al. 2017). In line with our results related
365 to lower angular velocity, offspring of treated females exhibited less anxiety, as indicated
366 by decreased thigmotaxis behaviour (e.g. close to the tank walls). The maximum velocity,
367 which is the first escape response commonly observed in isolated fish subjected to the
368 novel-tank test (Champagne et al. 2010; Colson et al. 2015a), was dramatically increased in
369 both groups immediately after the introduction into the novel tank (first 5 minutes). This
370 observed ceiling effect is certainly due to the strength of induced fear, avoiding any
371 possible discrepancy between the two groups for this parameter.

372 While mean distance travelled increased at the end of the test for control fish suggesting a
373 return to normal swimming pattern in this group, 17°C fish exhibited a constant low
374 swimming activity from the start to the end of the test. Existing studies reporting activity
375 levels in prenatally stressed individuals when subsequently subjected to challenging
376 situations are often contradictory. Increased or decreased activity is reported but this

377 discrepancy can be due to different contexts, intensity or duration of the challenge, species,
 378 age, and sex of the individuals tested. Our findings are however consistent with the
 379 majority of studies performed in mammals, which showed reduced activity in the offspring
 380 of females subjected to different stressors during pregnancy (Fride et al. 1986; Fujioka et
 381 al. 2001; Masterpasqua et al. 1976; Patin et al. 2004; Suchecki & Palermo Neto 1991), even
 382 though the stress was applied before fertilization in the present work. Interestingly, similar
 383 results were also found in fish (Eriksen et al. 2011; Espmark et al. 2008; Sopinka et al.
 384 2014; Tierney et al. 2009). For instance, Sockeye salmon (*Oncorhynchus nerka*) reared from
 385 mothers exposed to a chase stressor swam for shorter periods of time (Sopinka et al.
 386 2014). In Atlantic salmon, maternal cortisol exposure increased time spent non-swimming
 387 in juveniles (Espmark et al. 2008), and 1.5-year offspring from cortisol-implanted females
 388 also exhibited a reduction in the time spent moving compared to the controls during an
 389 acute confinement stress (Eriksen et al. 2011). These last studies focussed on the maternal
 390 endocrine status at spawning affecting several aspects of progeny behaviour and the
 391 results are consistent with the behavioural phenotypes observed here. In fish, thermal
 392 stress is known to trigger an increased plasma cortisol level (Quigley & Hinch 2006; Ryan
 393 1995; Zubair et al. 2012). We however did not find any increase in cortisol concentration
 394 into the 17°C eggs sampled before fertilization, which is in agreement with observations
 395 made by (Sopinka et al. 2014) and (Redfern et al. 2017). This finding rules out the
 396 participation of egg cortisol and indicates that the maternal observed effects are triggered
 397 by other mechanisms than the direct deposition of cortisol into the egg. In oviparous
 398 species, the external embryonic development implies that maternal stress transmission is
 399 only possible before fertilization through either egg content in molecules of various nature

400 (Lubzens et al. 2017), or epigenetic mechanisms. The intergenerational effects reported
 401 here are thus more likely due to genomic effects mediated by epigenetics mechanisms
 402 and/or specific features of the female gamete including maternally inherited nucleic acids
 403 and proteins.
 404 In mammals, there is growing evidence that stress during pregnancy causes attention
 405 deficits and depressive disorders (Ronald et al. 2010; Talge et al. 2007), as well as impaired
 406 emotional behaviours of adult offspring (Fride et al. 1986; Shiota & Kayamura 1989; Vallée
 407 et al. 1999; Zagron & Weinstock 2006). The lack of behavioural reaction to the challenge
 408 observed in 17°C fish suggests a global emotional blunting and an attention deficit,
 409 resembling the depressive-like symptoms described in prenatally stressed rodents
 410 (Morley-Fletcher et al. 2003; Poltyrev et al. 2005). In these studies, animals do not further
 411 respond to stressful stimuli, decrease explorative behaviour and their activity implying a
 412 form of resignation to an adverse uncontrollable situation. In a previous experiment, we
 413 noticed the absence of fear from a novel object (e.g. neophobia) in offspring from thermally
 414 stressed females (V. Colson, unpublished data). The absence of neophobia was likewise
 415 observed in suffering rainbow trout after being exposed to a nociceptive stimulus
 416 (Sneddon et al. 2003) and can be interpreted as a lack of attention for the environment. In
 417 the present study, the weaker emotional responses, as indicated by a decrease in angular
 418 velocity upon initial exposure to the novel tank and an absence of resumed ambulation at
 419 the end of the test might be explained by attention alterations due to maternal stress. Fish
 420 originating from thermally stressed mothers may be predicted to display a reduced ability
 421 to cope with their environment, since emotion and attention deficits might be major
 422 disadvantages in adverse or changing environments (Bijlsma & Loeschcke 2005). In

rainbow trout, first feeding is a key-stage during which fear-related behaviour, such as fast-start swimming, 'freezing', hiding and exploring are essential traits for fry survival. Therefore, hypo-active behaviour, as shown in the present study, could have direct impacts on fish survival chances under natural conditions.

Maternal effects on cognition

In the present experiment, we observed cognitive disorders in 17°C fish. Fry from mothers exposed to suboptimal temperature during late oogenesis were slower to locate the rewarded area in the spatial learning task. This finding is consistent with studies performed in other oviparous species (birds: (Guibert et al. 2013; Lindqvist et al. 2007) and fish: (Eaton et al. 2015; Roche et al. 2012)), showing cognitive disorders in offspring of mothers stressed before fertilization compared to offspring of non-stressed animals. In three-spined sticklebacks, offspring of predator-exposed mothers located the food reward more slowly than offspring of unexposed mothers (Roche et al. 2012). Female guppies (*Poecilia reticulata*) exposed to routine husbandry procedures that induced only a minimal elevation of cortisol, produced offspring that failed to associate a colour cue and food reward (Eaton et al. 2015). Conversely, in brook trout *Salvelinus fontinalis*, maternal cortisol consumption and handling did not impact spatial learning or memory in 6 month-old offspring (Cortez Ghio et al. 2016). This inconsistency might indicate that maternal effects on fish cognition are context-dependent or different depending on the type of stress used. Except for the above examples, very few studies have investigated intergenerational effects on fish cognition, and to our knowledge our findings are the first to show that a thermal maternal stress is linked to emotional and cognitive disorders.

445 When the learnt association was recalled 3 days after the last trial, 12°C fish were quicker
 446 to reach the right arm than 17°C fish, while the wrong arm was reached in both groups
 447 after a longer latency. Interestingly, the duration needed to reach the correct arm tended to
 448 be shortened during the memory test comparing to the latency measured at the last
 449 training trial within the two groups. This indicates that impaired cognition due to the
 450 thermal maternal stress concerned only the acquisition phase but not the retention
 451 pathway. Indeed, although 12°C fish were quicker to obtain the reward, fish from both
 452 groups remembered the correct location. During the 2-day break between the last trial and
 453 the memory test, fish were not fed as well as between each trial, the food being obtained
 454 only as a reward during the entire experimental procedure. Thus a high feeding motivation
 455 was observed at the moment of the recall, as also indicated by the short latency to leave the
 456 start-box, although the effects were not significant. Therefore, when highly motivated,
 457 rainbow trout demonstrated a memory capacity of at least 3 days in this spatial learning
 458 paradigm, which is consistent with a previous experiment showing that this fish species
 459 likewise remembered for 3 days an association by appetitive classical conditioning
 460 (Nordgreen et al. 2010).

461 In summary, fish originating from thermally stressed mothers were slower than controls in
 462 the spatial learning task. Although they were able to remember the food location after a 2-
 463 day break, they seemed to be less motivated, as indicated by the longer latency to reach the
 464 goal, which is consistent with the global blunting emotional responses observed in the
 465 novel-tank test in 17°C group. Cognitive abilities are critical for aquaculture fish since they
 466 need to anticipate specific events (e.g. food delivery) in order to reduce stress triggered by
 467 an unpredictable environment (Jones et al. 2012). Moreover, the ability to habituate to

repeated and fearful, but harmless, stimuli (e.g. repeated fishing linked to aquaculture practices) (Lieberman 2000) can be extremely useful for fish in order to avoid chronic stress under aquaculture conditions. It is thus highly beneficial for cultured fish to enhance or at least to preserve learning abilities (e.g. conditioning and habituation). Adaptive capacities remain a key component of fish welfare under breeding condition and this study reveals detrimental effects of heat maternal exposure on these capacities in the offspring.

Maternal effects on embryonic gene expression

Our results on the impact of thermal stress during the peri-ovulatory period (i.e. before fertilization) on offspring behaviour are similar to results obtained in mammals during pregnancy (Szuran et al. 1994; Talge et al. 2007; Vallee et al. 1999; Weinstock 2005; Zagron & Weinstock 2006). In humans, studies have shown that if a mother is stressed while pregnant, her child is at increased risk of having a range of problems, including emotional problems, attention deficits, and impaired cognitive development. These behavioural patterns are very similar to those observed in the present experiment. There is growing evidence for non-genetic effects of maternal experience on offspring in rodents (Weiss et al. 2011), and more recently in fish (Mommer & Bell 2014). Here, we used a robust methodology (i.e. microarray) and a conservative statistical approach to reveal the most relevant molecular players despite the low number of females that we simultaneously ovulated in both experimental groups. Among the sixteen differentially expressed genes, four genes are known to participate in neurodevelopment (*auts2*, *dpysl5*) or associated with neural/cerebral disorders (*arv1*) and X-linked cognitive disability (*plp2*). In humans, *AUTS2* is officially named *activator of transcription and developmental regulator* according to the official gene nomenclature (HGNC:14262 <https://www.genenames.org/>). It was

491 previously named *autism susceptibility candidate 2* (Sultana et al. 2002). The human *AUTS2*
 492 locus is associated with a wide diversity of neurological disorders, indicating that *AUTS2* is
 493 involved in neurodevelopment (see (Hori & Hoshino 2017a) for review). Several forms
 494 (splice variants) of the genes are expressed in the mouse during development, including
 495 during *in utero* development (Gao et al. 2014; Hori et al. 2014). In zebrafish, *auts2* is also
 496 embryonically expressed and found in the forebrain, midbrain and hindbrain at 24 hours
 497 post-fertilization (Oksenberg et al. 2013). This early embryonic pattern in zebrafish and
 498 mouse is consistent with the embryonic expression profile reported here throughout
 499 rainbow trout development. Interestingly, *Auts2* expression in the mouse brain is especially
 500 high in regions associated with higher cognitive functions, including in the prenatal brain
 501 (Bedogni et al. 2010). Functional analyses conducted in zebrafish (*Danio rerio*) confirmed
 502 the major role played by *auts2* in fish neurodevelopment (Oksenberg et al. 2013). Knock
 503 down of *auts2* in zebrafish resulted in considerably less developing neurons in the optic
 504 tectum, retina, and cerebellum. Interestingly, observed phenotypes were less severe when
 505 the morpholino (MO) used was directed against a splice junction rather than the
 506 translation initiation site, indicating that maternally-inherited *auts2* mRNA played an
 507 important role in *Auts2*-mediated neurodevelopment. Together, these observations are
 508 fully consistent with our data, especially the profiles of *auts2* maternal RNA shown in Fig.
 509 4B and suggest that the intergenerational effects of maternal exposure to high temperature
 510 could be mediated, at least in part, by differences in egg content in *auts2* messenger RNA.
 511 Data in mouse and zebrafish indicate that *Auts2* acts as a transcriptional regulator for
 512 neural development through interactions with several genes related to brain development
 513 and neurological disorders. More specifically, *Auts2* appears to be participating in neuronal

514 migration and neurite extension and is critical for the acquisition of neurocognitive
515 function (see (Hori & Hoshino 2017b) for review). Behavioural phenotypes observed in
516 *Auts2* heterozygous mutant mice are characterized by lower anxiety-like behaviour and
517 impaired memory (Gao et al. 2014; Hori et al. 2015). These phenotypes are strikingly
518 similar to the phenotypes observed here after maternal exposure to high temperature and
519 characterized by emotional numbing (i.e. lower angular velocity and absence of locomotor
520 activity modifications under stressful situation) and impaired learning abilities (i.e. slower
521 to locate a food-reward than controls in a T-maze). In addition to *auts2*, we also observed
522 the dysregulation of *dpysl5*, (*dihydropyrimidinase-like 5*) a member of CRMP (collapsing
523 response mediator protein) family thought to be involved in neural development (Veyrac et
524 al. 2011). Together these observations strongly suggest that the dysregulation of
525 neurodevelopment genes expression, especially *auts2*, but also *dpysl5*, in eggs and embryos
526 participate in mediating the intergenerational effects on offspring behaviour observed here
527 after exposing rainbow trout females to high temperature.
528 The transcriptomic analysis also revealed the differential expression, in response to
529 maternal exposure to high temperature, of genes associated with neural/cerebral disorders
530 (*arv1*) and X-linked cognitive disability (*plp2*). In an attempt to better understand genes
531 affecting human brain function, a recent whole-exome sequencing study in 143 families
532 resulted in the identification of 68 recessive genes associated with neurological disorders
533 (Alazami et al. 2015). Among those genes was *ARV1*, which was also associated with
534 autosomal recessive epileptic encelopathy in another study (Palmer et al. 2016). We also
535 observed a dysregulation of *plp2* in response to maternal exposure to high temperature. In
536 humans a polymorphism in *PLP2* (*Proteolipid protein 2*) promoter was associated with X-

537 linked Mental Retardation (XLMR) (Zhang et al. 2007). While the roles of *arv1* and *plp2* in
538 fish are currently unknown, the identity and suspected roles of these genes in humans are
539 consistent with the differential abundance of the gene and the emotional blunting and
540 impaired learning abilities observed in the present study.

541 The genome-wide transcriptome analysis also revealed the dysregulation of several other
542 genes, including a so far uncharacterized gene (*cxxcl1l*) that exhibits a strong differential
543 expression in eggs from different maternal origin. These genes are likely to mediate, or at
544 least to participate, in the intergenerational effect of maternal exposure to high
545 temperature observed here. Further analyses are needed to decipher the specific
546 contribution of these genes to the phenotypes reported here.

547 Together, our results revealed the dysregulation of several genes that are important for the
548 development of cognitive abilities in response to maternal exposure to high temperature.
549 The identity of these genes is consistent with the behavioural phenotypes observed in fry
550 originating from thermally stressed mothers. Additional studies aiming at characterizing
551 possible epigenetic modifications, gene expression and neurotransmitters activity in target
552 brain structures are still needed to further understand the mechanisms mediating the
553 observed behavioural modifications subsequent to thermal maternal stress in rainbow
554 trout.

555 Conclusions

556 Here we show that fish originating from thermally stressed mothers exhibit emotional and
557 cognitive disorders, which would be a major disadvantage under suboptimal or fluctuating
558 environments. These impaired behavioural phenotypes are associated with the

dysregulation of several genes known to play a major role in neurodevelopment. This is especially true for *auts2*, a key gene for neurodevelopment, more specifically neuronal migration and neurite extension, and critical for the acquisition of neurocognitive function in fish and mammals. In addition to *auts2*, our analysis revealed the dysregulation of another neurodevelopment gene (*dpysl5*) as well as genes associated with cognitive disorders in humans (*arv1*, *plp2*). Our study also revealed that some of the observed intergenerational effects are associated with a major dysregulation of several maternally-inherited mRNAs accumulated into the egg. Together, our observations shed new light on the intergenerational determinism of fish behaviour and associated underlying mechanisms. Our results address an important question for wild or cultured fish adaptive capacities in the context of climate warming.

Acknowledgments

The authors would like to thank F. Borel and C. Duret for their precious technical assistance and care of the animals. This work was supported in part by the French National Research Agency (ANR) under grant agreements ANR-13-BSV7-0002 PReSTO'Cog to VC and ANR-13-BSV7-0015 Maternal Legacy to JB.

References

- Aegerter S, and Jalabert B. 2004. Effects of post-ovulatory oocyte ageing and temperature on egg quality and on the occurrence of triploid fry in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 231:59-71. 10.1016/j.aquaculture.2003.08.019
- Alazami AM, Patel N, Shamseldin HE, Anazi S, Al-Dosari MS, Alzahrani F, Hijazi H, Alshammari M, Aldahmesh MA, Salih MA, Fageih E, Alhashem A, Bashiri FA, Al-Owain M, Kentab AY, Sogaty S, Al Tala S, Temsah MH, Tulbah M, Aljelaify RF, Alshahwan SA, Seidahmed MZ, Alhadid AA, Aldhalaan H, AlQallaf F, Kurdi W,

- Alfadhel M, Babay Z, Alsogheer M, Kaya N, Al-Hassnan ZN, Abdel-Salam GM, Al-Sannaa N, Al Mutairi F, El Khashab HY, Bohlega S, Jia X, Nguyen HC, Hammami R, Adly N, Mohamed JY, Abdulwahab F, Ibrahim N, Naim EA, Al-Younes B, Meyer BF, Hashem M, Shaheen R, Xiong Y, Abouelhoda M, Aldeeri AA, Monies DM, and Alkuraya FS. 2015. Accelerating novel candidate gene discovery in neurogenetic disorders via whole-exome sequencing of prescreened multiplex consanguineous families. *Cell Rep* 10:148-161. 10.1016/j.celrep.2014.12.015
- Atse CB, Audet C, and de la Noüe J. 2002. Effects of temperature and salinity on the reproductive success of Arctic charr, *Salvelinus alpinus* L.: egg composition, milt characteristics and fry survival. *Aquaculture Research* 33:299-309.
- Bedogni F, Hodge RD, Nelson BR, Frederick EA, Shiba N, Daza RA, and Hevner RF. 2010. Autism susceptibility candidate 2 (Auts2) encodes a nuclear protein expressed in developing brain regions implicated in autism neuropathology. *Gene Expr Patterns* 10:9. 10.1016/j.gep.2009.11.005
- Bijlsma R, and Loeschcke V. 2005. Environmental stress, adaptation and evolution: an overview. *Journal of Evolutionary Biology* 18:744-749. 10.1111/j.1420-9101.2005.00962.x
- Blaser RE, Chadwick L, and McGinnis GC. 2010. Behavioral measures of anxiety in zebrafish (*Danio rerio*). *Behav Brain Res* 208:56-62. 10.1016/j.bbr.2009.11.009
- Bobe J, and Labbe C. 2010. Egg and sperm quality in fish. *General and Comparative Endocrinology* 165:535-548. 10.1016/j.ygcen.2009.02.011
- Bonnet E, Fostier A, and Bobe J. 2007. Characterization of rainbow trout egg quality: a case study using four different breeding protocols, with emphasis on the incidence of embryonic malformations. *Theriogenology* 67:786-794. 10.1016/j.theriogenology.2006.10.008
- Campbell PM, Pottinger TG, and Sumpter JP. 1994. PRELIMINARY EVIDENCE THAT CHRONIC CONFINEMENT STRESS REDUCES THE QUALITY OF GAMETES PRODUCED BY BROWN AND RAINBOW-TROUT. *Aquaculture* 120:151-169. 10.1016/0044-8486(94)90230-5
- Champagne DL, Hoefnagels CCM, de Kloet RE, and Richardson MK. 2010. Translating rodent behavioral repertoire to zebrafish (*Danio rerio*): Relevance for stress research. *Behav Brain Res* 214:332-342. 10.1016/j.bbr.2010.06.001
- Colson V, Mure A, Valotaire C, Le Calvez J-M, Goardon L, Labbe L, Leguen I, and Prunet P. 2018. A novel emotional and cognitive approach to welfare phenotyping in rainbow trout exposed to poor water quality. *bioRxiv*. 10.1101/297747
- Colson V, Sadoul B, Valotaire C, Prunet P, Gaumé M, and Labbé L. 2015a. Welfare assessment of rainbow trout reared in a Recirculating Aquaculture System: Comparison with a Flow-Through System. *Aquaculture* 436:151-159. <http://dx.doi.org/10.1016/j.aquaculture.2014.10.047>
- Colson V, Valotaire C, Geffroy B, and Kiilerich P. 2015b. Egg Cortisol Exposure Enhances Fearfulness in Larvae and Juvenile Rainbow Trout. *Ethology* 121:1191-1201. 10.1111/eth.12437
- Cortez Ghio S, Boudreau Leblanc A, Audet C, and Aubin-Horth N. 2016. Effects of maternal stress and cortisol exposure at the egg stage on learning, boldness and neophobia in brook trout. *Behaviour* 153:1639-1663. 10.1163/1568539x-00003377

- 629 Eaton L, Edmonds E, Henry TB, Snellgrove DL, and Sloman KA. 2015. Mild maternal stress
630 disrupts associative learning and increases aggression in offspring. *Horm Behav*
631 71:10-15. 10.1016/j.yhbeh.2015.03.005
- 632 Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavella PR, Elegante MF, Elkhayat SI, Bartels BK,
633 Tien AK, Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z, and Kalueff
634 AV. 2009. Understanding behavioral and physiological phenotypes of stress and
635 anxiety in zebrafish. *Behav Brain Res* 205:38-44. 10.1016/j.bbr.2009.06.022
- 636 Eriksen MS, Espmark A, Braastad BO, Salte R, and Bakken M. 2007. Long-term effects of
637 maternal cortisol exposure and mild hyperthermia during embryogeny on survival,
638 growth and morphological anomalies in farmed Atlantic salmon *Salmo salar*
639 offspring. *Journal of Fish Biology* 70:462-473. 10.1111/j.1095-8649.2007.01317.x
- 640 Eriksen MS, Faerevik G, Kittilsen S, McCormick MI, Damsgard B, Braithwaite VA, Braastad
641 BO, and Bakken M. 2011. Stressed mothers - troubled offspring: a study of
642 behavioural maternal effects in farmed *Salmo salar*. *Journal of Fish Biology* 79:575-
643 586. 10.1111/j.1095-8649.2011.03036.x
- 644 Eriksen MS, Poppe TT, McCormick M, Damsgård B, Salte R, Braastad BO, and Bakken M.
645 2015. Simulated maternal pre-spawning stress affects offspring's attributes in
646 farmed Atlantic salmon *Salmo salar* (Linnaeus, 1758). *Aquaculture Research*
647 46:1480-1489. 10.1111/are.12301
- 648 Espmark AM, Eriksen MS, Salte R, Braastad BO, and Bakken M. 2008. A note on pre-
649 spawning maternal cortisol exposure in farmed Atlantic salmon and its impact on
650 the behaviour of offspring in response to a novel environment. *Applied Animal*
651 *Behaviour Science* 110:404-409. 10.1016/j.applanim.2007.04.003
- 652 Fride E, Dan Y, Feldon J, Halevy G, and Weinstock M. 1986. Effects of prenatal stress on
653 vulnerability to stress in prepubertal and adult rats. *Physiology & Behavior* 37:681-
654 687. [https://doi.org/10.1016/0031-9384\(86\)90172-1](https://doi.org/10.1016/0031-9384(86)90172-1)
- 655 Fujioka T, Fujioka A, Tan N, Chowdhury GMI, Mouri H, Sakata Y, and Nakamura S. 2001.
656 Mild prenatal stress enhances learning performance in the non-adopted rat
657 offspring. *Neuroscience* 103:301-307. [https://doi.org/10.1016/S0306-](https://doi.org/10.1016/S0306-4522(00)00582-0)
658 [4522\(00\)00582-0](https://doi.org/10.1016/S0306-4522(00)00582-0)
- 659 Gao Z, Lee P, Stafford JM, von Schimmelmänn M, Schaefer A, and Reinberg D. 2014. An
660 AUTS2-Polycomb complex activates gene expression in the CNS. *Nature* 516:349-
661 354. 10.1038/nature13921
- 662 Guibert F, Lumineau S, Kotrschal K, Mostl E, Richard-Yris MA, and Houdelier C. 2013.
663 Trans-generational effects of prenatal stress in quail. *Proceedings of the Royal*
664 *Society B-Biological Sciences* 280. 10.1098/rspb.2012.2368
- 665 Hori K, and Hoshino M. 2017a. Neuronal Migration and AUTS2 Syndrome. *Brain Sciences* 7.
666 10.3390/brainsci7050054
- 667 Hori K, and Hoshino M. 2017b. Neuronal Migration and AUTS2 Syndrome. *Brain Sciences*
668 7:54. 10.3390/brainsci7050054
- 669 Hori K, Nagai T, Shan W, Sakamoto A, Abe M, Yamazaki M, Sakimura K, Yamada K, and
670 Hoshino M. 2015. Heterozygous Disruption of Autism susceptibility candidate 2
671 Causes Impaired Emotional Control and Cognitive Memory. *PLoS One* 10:e0145979.
672 10.1371/journal.pone.0145979
- 673 Hori K, Nagai T, Shan W, Sakamoto A, Taya S, Hashimoto R, Hayashi T, Abe M, Yamazaki M,
674 Nakao K, Nishioka T, Sakimura K, Yamada K, Kaibuchi K, and Hoshino M. 2014.

- 675 Cytoskeletal regulation by AUTS2 in neuronal migration and neuritogenesis. *Cell Rep*
676 9:2166-2179. 10.1016/j.celrep.2014.11.045
- 677 Jeffrey JD, and Gilmour KM. 2016. Programming of the hypothalamic-pituitary-interrenal
678 axis by maternal social status in zebrafish (*Danio rerio*). *J Exp Biol* 219:1734-1743.
679 10.1242/jeb.138826
- 680 Jones HAC, Noble C, Damsgard B, and Pearce GP. 2012. Investigating the influence of
681 predictable and unpredictable feed delivery schedules upon the behaviour and
682 welfare of Atlantic salmon parr (*Salmo salar*) using social network analysis and fin
683 damage. *Applied Animal Behaviour Science* 138:132-140.
684 10.1016/j.applanim.2012.01.019
- 685 Jonsson B, and Jonsson N. 2016. Trans-generational maternal effect: temperature
686 influences egg size of the offspring in Atlantic salmon *Salmo salar*. *J Fish Biol.*
687 10.1111/jfb.13040
- 688 King HR, and Pankhurst NW. 2004. Effect of maintenance at elevated temperatures on
689 ovulation and luteinizing hormone releasing hormone analogue responsiveness of
690 female Atlantic salmon (*Salmo salar*) in Tasmania. *Aquaculture* 233:583-597.
691 <https://doi.org/10.1016/j.aquaculture.2003.09.049>
- 692 King HR, Pankhurst NW, Watts M, and Pankhurst PM. 2003. Effect of elevated summer
693 temperatures on gonadal steroid production, vitellogenesis and egg quality in
694 female Atlantic salmon. *Journal of Fish Biology* 63:153-167. 10.1046/j.1095-
695 8649.2003.00137.x
- 696 Kittilsen S, Ellis T, Schjolden J, Braastad BO, and Overli O. 2009. Determining stress-
697 responsiveness in family groups of Atlantic salmon (*Salmo salar*) using non-invasive
698 measures. *Aquaculture* 298:146-152. 10.1016/j.aquaculture.2009.10.009
- 699 Kjorsvik E, Mangorjensen A, and Holmefjord I. 1990. EGG QUALITY IN FISHES. *Advances in*
700 *Marine Biology* 26:71-113. 10.1016/s0065-2881(08)60199-6
- 701 Lieberman DA. 2000. *Learning: Behavior and cognition*. Belmont, CA, US:
702 Wadsworth/Thomson Learning.
- 703 Lindqvist C, Janczak AM, Natt D, Baranowska I, Lindqvist N, Wichman A, Lundeberg J,
704 Lindberg J, Torjesen PA, and Jensen P. 2007. Transmission of Stress-Induced
705 Learning Impairment and Associated Brain Gene Expression from Parents to
706 Offspring in Chickens. *PLoS One* 2. 10.1371/journal.pone.0000364
- 707 Lubzens E, Bobe J, Young G, and Sullivan CV. 2017. Maternal investment in fish oocytes and
708 eggs: The molecular cargo and its contributions to fertility and early development.
709 *Aquaculture* 472:107-143. 10.1016/j.aquaculture.2016.10.029
- 710 Masterpasqua F, Chapman RH, and Lore RK. 1976. The effects of prenatal psychological
711 stress on the sexual behavior and reactivity of male rats. *Developmental*
712 *Psychobiology* 9:403-411. 10.1002/dev.420090503
- 713 McCormick MI. 2009. Indirect effects of heterospecific interactions on progeny size through
714 maternal stress. *Oikos* 118:744-752. 10.1111/j.1600-0706.2008.17410.x
- 715 Migaud H, Bell G, Cabrita E, McAndrew B, Davie A, Bobe J, Herraes MP, and Carrillo M. 2013.
716 Gamete quality and broodstock management in temperate fish. *Reviews in*
717 *Aquaculture* 5:S194-S223. 10.1111/raq.12025
- 718 Mommer BC, and Bell AM. 2014. Maternal Experience with Predation Risk Influences
719 Genome-Wide Embryonic Gene Expression in Threespined Sticklebacks
720 (*Gasterosteus aculeatus*). *PLoS One* 9. 10.1371/journal.pone.0098564

- 721 Morley-Fletcher S, Darnaudery M, Koehl M, Casolini P, Van Reeth O, and Maccari S. 2003.
722 Prenatal stress in rats predicts immobility behavior in the forced swim test. Effects
723 of a chronic treatment with tianeptine. *Brain Res* 989:246-251.
- 724 Nordgreen J, Janczak AM, Hovland AL, Ranheim B, and Horsberg TE. 2010. Trace classical
725 conditioning in rainbow trout (*Oncorhynchus mykiss*): what do they learn? *Animal*
726 *Cognition* 13:303-309. 10.1007/s10071-009-0267-3
- 727 Oksenberg N, Stevison L, Wall JD, and Ahituv N. 2013. Function and Regulation of AUTS2, a
728 Gene Implicated in Autism and Human Evolution. *PLOS Genetics* 9:e1003221.
729 10.1371/journal.pgen.1003221
- 730 Overli O, Winberg S, and Pottinger TG. 2005. Behavioral and neuroendocrine correlates of
731 selection for stress responsiveness in rainbow trout - a review. *Integr Comp Biol*
732 45:463-474. 10.1093/icb/45.3.463
- 733 Palmer EE, Jarrett KE, Sachdev RK, Al Zahrani F, Hashem MO, Ibrahim N, Sampaio H,
734 Kandula T, Macintosh R, Gupta R, Conlon DM, Billheimer JT, Rader DJ, Funato K,
735 Walkey CJ, Lee CS, Loo C, Brammah S, Elakis G, Zhu Y, Buckley M, Kirk EP, Bye A,
736 Alkuraya FS, Roscioli T, and Lagor WR. 2016. Neuronal deficiency of ARV1 causes an
737 autosomal recessive epileptic encephalopathy. *Hum Mol Genet* 25:3042-3054.
738 10.1093/hmg/ddw157
- 739 Patin V, Vincent A, Lordi B, and Caston J. 2004. Does prenatal stress affect the motoric
740 development of rat pups? *Developmental Brain Research* 149:85-92.
741 <https://doi.org/10.1016/j.devbrainres.2003.12.008>
- 742 Poisson A, Valotaire C, Borel F, Bertin A, Darmaillacq A-S, Dickel L, and Colson V. 2017.
743 Embryonic exposure to a conspecific alarm cue triggers behavioural plasticity in
744 juvenile rainbow trout. *Animal Behaviour* 133:35-45.
745 <https://doi.org/10.1016/j.anbehav.2017.09.013>
- 746 Poltyrev T, Gorodetsky E, Bejar C, Schorer-Apelbaum D, and Weinstock M. 2005. Effect of
747 chronic treatment with ladostigil (TV-3326) on anxiogenic and depressive-like
748 behaviour and on activity of the hypothalamic-pituitary-adrenal axis in male and
749 female prenatally stressed rats. *Psychopharmacology (Berl)* 181:118-125.
750 10.1007/s00213-005-2229-z
- 751 Quigley JT, and Hinch SG. 2006. Effects of rapid experimental temperature increases on
752 acute physiological stress and behaviour of stream dwelling juvenile chinook
753 salmon. *Journal of Thermal Biology* 31:429-441. 10.1016/j.jtherbio.2006.02.003
- 754 Redfern JC, Cooke SJ, Lennox RJ, Nannini MA, Wahl DH, and Gilmour KM. 2017. Effects of
755 maternal cortisol treatment on offspring size, responses to stress, and anxiety-
756 related behavior in wild largemouth bass (*Micropterus salmoides*). *Physiol Behav.*
757 10.1016/j.physbeh.2017.08.001
- 758 Roche DP, McGhee KE, and Bell AM. 2012. Maternal predator-exposure has lifelong
759 consequences for offspring learning in threespined sticklebacks. *Biol Lett* 8:932-935.
760 10.1098/rsbl.2012.0685
- 761 Ronald A, Pennell CE, and Whitehouse AJO. 2010. Prenatal Maternal Stress Associated with
762 ADHD and Autistic Traits in early Childhood. *Front Psychol* 1:223.
763 10.3389/fpsyg.2010.00223
- 764 Rouger Y, Aubin J, Breton B, Fauconneau B, Fostier A, Le Bail PY, Loir M, Prunet P, and
765 Maisse G. 1998. Réponse au stress induit par le transport chez la truite arc-en-ciel

- 766 (Oncorhynchus mykiss). *Bulletin Français de la Pêche et de la Pisciculture*:511-519.
767 10.1051/kmae:1998020
- 768 Ryan SN. 1995. THE EFFECT OF CHRONIC HEAT-STRESS ON CORTISOL-LEVELS IN THE
769 ANTARCTIC FISH PAGOTHENIA-BORCHGREVINKI. *Experientia* 51:768-774.
770 10.1007/bf01922428
- 771 Sadoul B, Foucard A, Valotaire C, Labbe L, Goardon L, LeCalvez JM, Medale F, Quillet E,
772 Dupont-Nivet M, Geurden I, Prunet P, and Colson V. 2016. Adaptive capacities from
773 survival to stress responses of two isogenic lines of rainbow trout fed a plant-based
774 diet. *Sci Rep* 6:35957. 10.1038/srep35957
775 <https://http://www.nature.com/articles/srep35957-supplementary-information>
- 776 Shiota K, and Kayamura T. 1989. Effects of Prenatal Heat Stress on Postnatal Growth,
777 Behavior and Learning Capacity in Mice. *Neonatology* 56:6-14.
- 778 Sloman KA. 2010. Exposure of ova to cortisol pre-fertilisation affects subsequent behaviour
779 and physiology of brown trout. *Horm Behav* 58:433-439.
780 10.1016/j.yhbeh.2010.05.010
- 781 Sneddon LU, Braithwaite VA, and Gentle MJ. 2003. Novel object test: Examining nociception
782 and fear in the rainbow trout. *Journal of Pain* 4:431-440. 10.1067/s1526-
783 5900(03)00717-x
- 784 Sopinka NM, Hinch SG, Middleton CT, Hills JA, and Patterson DA. 2014. Mother knows best,
785 even when stressed? Effects of maternal exposure to a stressor on offspring
786 performance at different life stages in a wild semelparous fish. *Oecologia* 175:493-
787 500. 10.1007/s00442-014-2915-9
- 788 Suchecki D, and Palermo Neto J. 1991. Prenatal stress and emotional response of adult
789 offspring. *Physiol Behav* 49:423-426.
- 790 Sultana R, Yu CE, Yu J, Munson J, Chen D, Hua W, Estes A, Cortes F, de la Barra F, Yu D,
791 Haider ST, Trask BJ, Green ED, Raskind WH, Distech CM, Wijsman E, Dawson G,
792 Storm DR, Schellenberg GD, and Villacres EC. 2002. Identification of a novel gene on
793 chromosome 7q11.2 interrupted by a translocation breakpoint in a pair of autistic
794 twins. *Genomics* 80:129-134.
- 795 Szuran T, Zimmermann E, and Welzl H. 1994. WATER MAZE PERFORMANCE AND
796 HIPPOCAMPAL WEIGHT OF PRENATALLY STRESSED RATS. *Behav Brain Res*
797 65:153-155. 10.1016/0166-4328(94)90100-7
- 798 Talge NM, Neal C, and Glover V. 2007. Antenatal maternal stress and long-term effects on
799 child neurodevelopment: how and why? *J Child Psychol Psychiatry* 48:245-261.
800 10.1111/j.1469-7610.2006.01714.x
- 801 Taranger GL, and Hansen B. 1993. Ovulation and egg survival following exposure of
802 Atlantic salmon, *Salmo salar* L., broodstock to different water temperatures.
803 *Aquaculture Research* 24:151-156.
- 804 Tierney KB, Patterson DA, and Kennedy CJ. 2009. The influence of maternal condition on
805 offspring performance in sockeye salmon *Oncorhynchus nerka*. *Journal of Fish*
806 *Biology* 75:1244-1257. 10.1111/j.1095-8649.2009.02360.x
- 807 Vallee M, Maccari S, Dellu F, Simon H, Le Moal M, and Mayo W. 1999. Long-term effects of
808 prenatal stress and postnatal handling on age-related glucocorticoid secretion and
809 cognitive performance: a longitudinal study in the rat. *European Journal of*
810 *Neuroscience* 11:2906-2916. 10.1046/j.1460-9568.1999.00705.x

- 811 Vallée M, Maccari S, Dellu F, Simon H, Le Moal M, and Mayo W. 1999. Long-term effects of
812 prenatal stress and postnatal handling on age-related glucocorticoid secretion and
813 cognitive performance: a longitudinal study in the rat. *European Journal of*
814 *Neuroscience* 11:2906-2916. 10.1046/j.1460-9568.1999.00705.x
- 815 Veyrac A, Reibel S, Sacquet J, Mutin M, Camdessanche JP, Kolattukudy P, Honnorat J, and
816 Jourdan F. 2011. CRMP5 regulates generation and survival of newborn neurons in
817 olfactory and hippocampal neurogenic areas of the adult mouse brain. *PLoS One*
818 6:e23721. 10.1371/journal.pone.0023721
- 819 Weinstock M. 2005. The potential influence of maternal stress hormones on development
820 and mental health of the offspring. *Brain, Behavior, and Immunity* 19:296-308.
821 <http://dx.doi.org/10.1016/j.bbi.2004.09.006>
- 822 Weiss IC, Franklin TB, Vizi S, and Mansuy IM. 2011. Inheritable effect of unpredictable
823 maternal separation on behavioral responses in mice. *Front Behav Neurosci* 5:3.
824 10.3389/fnbeh.2011.00003
- 825 Winberg S, Schjolden J, Overli O, and Pottinger T. 2007. Stress and stress coping in fish,
826 behavioural correlates and neuroendocrine mechanisms. *Comparative Biochemistry*
827 *and Physiology a-Molecular & Integrative Physiology* 146:S77-S77.
828 10.1016/j.cbpa.2007.01.719
- 829 Zagron G, and Weinstock M. 2006. Maternal adrenal hormone secretion mediates
830 behavioural alterations induced by prenatal stress in male and female rats. *Behav*
831 *Brain Res* 175:323-328. <http://dx.doi.org/10.1016/j.bbr.2006.09.003>
- 832 Źarski D, Nguyen T, Le Cam A, Montfort J, Dutto G, Vidal MO, Fauvel C, and Bobe J. 2017.
833 Transcriptomic Profiling of Egg Quality in Sea Bass (*Dicentrarchus labrax*) Sheds
834 Light on Genes Involved in Ubiquitination and Translation. *Mar Biotechnol (NY)*
835 19:102-115. 10.1007/s10126-017-9732-1
- 836 Zhang L, Jie C, Obie C, Abidi F, Schwartz CE, Stevenson RE, Valle D, and Wang T. 2007. X
837 chromosome cDNA microarray screening identifies a functional PLP2 promoter
838 polymorphism enriched in patients with X-linked mental retardation. *Genome Res*
839 17:641-648. 10.1101/gr.5336307
- 840 Zubair SN, Peake SJ, Hare JF, and Anderson WG. 2012. The effect of temperature and
841 substrate on the development of the cortisol stress response in the lake sturgeon,
842 *Acipenser fulvescens*, Rafinesque (1817). *Environmental Biology of Fishes* 93:577-
843 587. 10.1007/s10641-011-9951-7
- 844

845

Figure 1(on next page)

Embryonic mortality

Effects of rearing temperature before ovulation (12°C and 17°C) on the occurrence of embryonic mortality (%) at different developmental stages (eyeing, hatching and yolk-sac resorption). Values are medians (quartiles: 25 and 75%). * $P < 0.05$: significant difference between treatments (n= 4).




Figure 2 (on next page)

Swimming behaviour in novel-tank test

Swimming behaviour of 75-76 dpf progeny from mothers exposed to 12°C and 17°C before ovulation, video-filmed for 30 min in social isolation in a novel environment. Behaviours were recorded during the first 5-min interval of the test (5 min) and the last 5-min interval of the test (30 min). (A) Total distance travelled (cm). (B) Maximum velocity (cm/sec). (C) Angular velocity (°/sec). (D) Time spent in the border over the 5 minutes (% of time). Values are means and their associated mean standard error (SEM) (n=15). Significant main effects and interactions are indicated (NS: non significant, * $P < 0.05$, *** $P < 0.001$). Random female effect is indicated (* $P < 0.05$). Different letters indicate significant differences shown by post-hoc tests ($P < 0.05$) or a tendency ($0.05 < P < 0.1$).




Figure 3(on next page)

Spatial learning and memory

Latency (seconds) to make the right choice by reaching the rewarded arm of a T-maze in progeny from mothers exposed to 12°C and 17°C before ovulation. Latencies were measured (A) during the acquisition phase within 11 successive trials, lasting 1800 seconds each, (B) three days after trial 11, to measure fish memory, and (C) before fish entry into either the right or the wrong arm, three days after the 11th trial. Values are means and their associated mean standard error (SEM) (n= 5). Main effects and significant interactions are indicated (NS: non significant, * $P < 0.05$). Non-significant random female effect is indicated. † $P < 0.05$, ‡ $P < 0.01$, # $P < 0.001$: significant differences from Trial 1, within 12°C. Different letters indicate significant differences between 12°C and 17°C shown by post-hoc tests ($P < 0.05$).







Figure 4(on next page)


Microarray analysis of gene expression in eggs and progeny originating from mothers exposed to either 12°C or 17°C during the peri-ovulatory period.

A.Venn diagram representing the number of differentially expressed genes. Two-way-ANOVA performed using maternal temperature and developmental stage as fixed factors. Benjamini-Hochberg corrected p values < 0.05. Gene symbols are shown when a significant effect was obtained for Temperature and Temperature X Stage interaction. All corresponding data are presented in supplementary data file 1.**B.**Supervised clustering analysis of the expression profiles of the 12 genes significantly dysregulated due to the temperature effect (panel A). Data were median-centered and an average linkage clustering was performed. Neurodevelopment genes and genes related to human cognitive disorders are shown in purple. **C.**Boxplot representation of gene expression profiles of neurodevelopment genes (*auts2*and *dpyls5a*) and genes related to human cognitive disorders (*arv1*and *plp2*) corresponding to the data delineated in purple on panel B. AU: arbitrary units.

A



B



C

