

75 Years ago *Arabidopsis* was first suggested as a Model Plant – But how did *Arabidopsis* Col-0 become the standard Natural Accession?

A short history of *Arabidopsis thaliana* (L.) Heynh. Columbia-0

Marc Somssich

Persson Lab, School of BioSciences, the University of Melbourne, Parkville 3010 VIC, Australia

Email: marc.somssich@unimelb.edu.au ; Twitter: [@somssichm](https://twitter.com/somssichm)

The Origin of *Arabidopsis thaliana* Research (1905 – 1943)

Modern work with *Arabidopsis thaliana* goes back to the German botanist Friedrich Laibach who, while working as a Ph.D. student in the laboratory of Eduard Strasburger in Bonn, analyzed the number of chromosomes in different plants that he had collected around Bonn and his hometown Limburg^{1,2}. The first *Arabidopsis* plants to be experimented on were collected by Laibach in 1905, and belonged to the natural accession Limburg (Laibach introduced a system of naming the natural accessions after the places he collected them from)². Laibach found that they carried 5 pairs of chromosomes, one of the smallest numbers known at the time (he published his results in 1907, even though *Arabidopsis* was only included in the written thesis, but not specifically mentioned in the paper)¹⁻³. Unfortunately, the natural habitat of the Limburg population was destroyed shortly after to make way for the new “Autobahn” (highway), connecting the cities of Frankfurt and Köln². At the time, *Arabidopsis* was ‘only known to florists and taxonomists, who had nothing better to do than constantly change its name and systematic positioning’, as Laibach put it in 1965². However, he became interested in the little weed, and between 1930 and 1950 collected seeds from over 150 different natural accessions (or races, as he called them) of *Arabidopsis* from anywhere he or his colleagues travelled to^{2,4}. Laibach kept all of these individual seed lines meticulously organized and maintained in his Department at Frankfurt University, and his collection eventually formed the foundation of the *Arabidopsis* Information Service (AIS) seed bank in the 1960s, which itself served as the basis for the modern Columbus (ABRC), Nottingham (NASC) and Tsukuba (RIKEN) stock centres decades later^{2,5,6}.

Arabidopsis thaliana First Proposed as a Plant Model (1943 – 1957)

Laibach's interest and preliminary studies of *Arabidopsis* eventually resulted in a now famous publication titled ‘*Arabidopsis Thaliana* (L.) Heynh. als Objekt für genetische und

entwicklungsphysiologische Untersuchungen' (*Arabidopsis Thaliana* (L.) Heynh. as an object for genetic, developmental and physiological analyses'), in which Laibach points out the benefits of working with *Arabidopsis* (easy to grow, small genome, short lifecycle, high seed yield, can be crossed and mutated...)³. Based on these observations he proposed to adopt *Arabidopsis* as a model organism for plant science, pointing out how comparable it is in its suitability to the 'prime example' of other models, *Drosophila melanogaster*³. This proposal however, was largely ignored by the scientific community at the time, who needed almost another 40 years to finally see the light and adopt *Arabidopsis* as a plant model system⁷. One academic who shared Laibach's enthusiasm for *Arabidopsis* was György P. Rédei from Hungary, who in 1955 had just finished his Ph.D. thesis, working on tomato and wheat⁸. After reading Laibach's article, Rédei recognized the potential of *Arabidopsis* for genetic studies, and with the help of his supervisor, Prof. Györffy, he asked Laibach for some *Arabidopsis* seeds to start his own work on this new model⁸. The seeds he obtained were the four natural accessions Graz, Limburg, Estland and Landsberg⁹. Rédei took these four lines with him, when he left Europe to start his own laboratory at the University of Missouri in Columbia, Mo⁹. For the next 20 years Rédei remained the only researcher working on *Arabidopsis* in the United States; or, as his former colleague Prof. Doug Randall put it, "George was 20 to 30 years ahead of his time"¹⁰. This situation, however, made it incredibly hard for Rédei to receive funding⁹. In fact, one of his funding applications to the National Science Foundation was now famously rejected on the basis that '*the genetics panel does not believe that it is worthwhile to develop Arabidopsis as a new model organism for genetic studies because only prokaryotes can contribute significantly to new knowledge*'⁹. But Rédei refused to give up on *Arabidopsis* and from the four seed lines he had received from Laibach, chose Landsberg as his model for future work. This choice was due to that Estland phenotypically did not match its description and Graz was late flowering, while Landsberg matched the description and seemed vigorous and healthy (it is not clear on which grounds Limburg was dropped)⁹.

The Columbia and Landsberg *erecta* lines Emerge (1957 – 1965)

In 1957 Rédei used his Landsberg seeds in a mutagenesis experiment, where he irradiated the seeds with X-rays and then screened for mutants with interesting phenotypes (meanwhile, in Australia, John Langridge was doing the same for Estland seeds he had received from Laibach)^{9,11-13}. Gene mutagenesis by X-ray irradiation had been described in the 1920s for

Drosophila and *Antirrhinum*, and one of Laibach's students, Erna Reinholz, went on to establish this technique for *Arabidopsis* seeds^{4,14,15}. One of the first mutants Rédei recovered was the *erecta* mutant, which, with its stunted growth, appeared to be quite sturdy, and he thought it might come in handy for further experimentation^{9,16}. He published the Landsberg *erecta* mutant in a paper dealing with heterosis, despite not being sure if the importance of his observation warranted a full publication¹⁶. His paper therefore opens with the paragraph '*The author feels somewhat hesitant to add to the large volume of the literature on the subject but its practical importance and theoretical interest prompt the decision in favor of this brief account*'¹⁶. However, in his mutagenesis screen Rédei also realized that the original Landsberg population was actually not a homogenous line, but appeared to be a mix of different lines^{9,11}. Therefore, he chose a single plant from the batch that he had not irradiated, to establish a new, clean line for all further studies^{9,11}. Following Laibach's example of naming the different natural accessions after the location where he found them, he named his new line Columbia^{9,11}. So interestingly, Columbia is an American plant by name, but a central European plant by genetic heritage – something that can be demonstrated experimentally, when analysing its genetic polymorphisms¹⁷. In 1959, another plant biologist, Willem Feenstra from the University of Groningen in the Netherlands, visited Rédei in Columbia and took the Landsberg *erecta* line with him for his own research, establishing this line as a standard in Europe, while Rédei concentrated his work on his own Columbia line^{9,11,18}.

***Arabidopsis thaliana* gets its Breakthrough (1965 – 1996)**

In the following two decades, interest in *Arabidopsis* research slowly increased. By the mid-1960s, the AIS (<https://www.arabidopsis.org/ais/newaisvols.jsp>) was established as a yearly newsletter to connect the small *Arabidopsis* research community, and in 1965 the first International Arabidopsis Symposium in Göttingen, Germany, already attracted a full 25 participants^{19,20}. The AIS would eventually evolve into the now invaluable The Arabidopsis Information Resource (TAIR) database²¹. As a result of this increased interest, György Rédei decided to take up Laibach's suggestion from 1943, and published the second article calling for the acceptance of *Arabidopsis* as a plant model in 1975, simply titled '*Arabidopsis* as a genetic tool' (where he pointed out the same benefits Laibach had already pointed out 30 years earlier)²². Following this publication and a couple of highly influential papers from people like Maarten Koornneef (who worked with Will Feenstra), or Chris R. Somerville and Elliott M. Meyerowitz

(converts from the model organisms *Escherichia coli* and *Drosophila melanogaster*, respectively), *Arabidopsis* finally got its break in the early 1980s^{7,23–25}. With *Arabidopsis* now finally established, the third article discussing its role as a model (published in 1985 and pointing out the same benefits that Rédei and Laibach had pointed out 10 and 40 years earlier) was now published in the prestigious *Science* journal⁷.

Col-0 takes over as the Standard Accession (1996 – today)

During the next decade, *Arabidopsis* research was mostly done using the Landsberg *erecta* accession, even though Columbia also regularly appeared, especially in US laboratories or from groups that had obtained seeds directly from Rédei. However, this was about to change when, in 1996, Columbia was chosen as the natural accession for the sequencing and annotation of the complete *Arabidopsis* genome²⁶. Despite Landsberg *erecta* being more commonly used at the time, this choice was the obvious one in this case, because the Landsberg *erecta* line had previously been subjected to X-ray irradiation, and therefore carried several unnatural mutations, while Columbia had been maintained as a clean homozygous line^{11,26}. Shortly after the genome was eventually published in the year 2000, Columbia was also chosen as the natural accession for a genome-wide mutagenesis project at the SALK institute in San Diego, resulting in the SALK collection of T-DNA insertion lines – still the biggest resource of ready-to-order *Arabidopsis* mutants²⁷. Following these two massive projects, it was clear that Columbia was firmly established as the number one natural accession for *Arabidopsis* research, while the use of Landsberg *erecta* has been declining ever since. And this all just because the Landsberg batch that György Rédei received from Friedrich Laibach in 1955 was not a homogenous line.

Addendum> What about the ‘(L.)’ and the ‘Heynh.’ behind *Arabidopsis thaliana*, and the ‘-0’ behind Col?

The ‘(L.)’ and ‘Heynh.’, which are often found after *Arabidopsis thaliana*, are so-called ‘authorities’ - the official author abbreviation of the person who gave the plant its name²⁸. Though *Arabidopsis thaliana* was first described by Johannes Thal, who gave it the name *Pilosella siliquosa minor*, it was Carl Linnaeus who named it *Arabis thaliana* (*thaliana* in honour of Johannes Thal)^{29,30}. Therefore, the ‘(L.)’ behind genus and species is the author abbreviation for Carl Linnaeus^{29,30}. Botanist Gustav Heynhold then merged similar plants into one new genus, *Arabidopsis*, signifying *Arabis-like*, and added his own author abbreviation, ‘Heynh.’, behind the

one from Linnaeus (Heynholds book '*Flora von Sachsen*' is generally cited here, though I could only find *Arabidopsis* in his book '*Nomenclator botanicus hortensis*')^{29,31,32}. The '0' behind the Col name, on the other hand, signifies the source of an individual seed line³³. Over the years, different laboratories that received Col seeds from György Rédei have propagated and maintained their own inbred lines of the original batch. When all these lines were later donated to the seed centres, a numbering system was developed to be able to distinguish these individual lines³³. In this system, George Rédei's Columbia line in the ABRC stock centre would be named Col-1/CS3176, or Col-1 in short³³. The name is made up of [wild type]-[originator]/[maintainer stock-#], with the wild type being 'Col', the originator George Rédei, who was designated the number 1, and the maintainer, the ABRC stock centre, carrying it under the stock number 3176³³. The line donated by Shauna Somerville to the ABRC, a direct descendent of Rédei's Col-1, is Col-2/CS907, or in short, Col-2³³. Confusingly, the Col-0 line (Col-0/CS1092) is actually a descendent of Rédei's Col-1 line³³. It received the lower originator number 0 because it was already maintained and propagated in the original AIS-seed bank by Albert Kranz, and is therefore an 'older' stock⁵.

More 'History of *Arabidopsis*' Resources:

- Friedrich Laibach - 60 Jahre Arabidopsis-Forschung, 1905-1965²
- György P. Rédei - *Arabidopsis thaliana* (L.) Heynh. A review of the genetics and biology²⁹
- Elliot M. Meyerowitz – *Arabidopsis thaliana*³⁴
- György P. Rédei - A heuristic glance at the past of Arabidopsis genetics⁹
- Elizabeth Pennisi - Arabidopsis Comes of Age³⁵
- Elliot M. Meyerowitz – Prehistory and history of Arabidopsis research³⁶
- Chris R. Somerville, Maarten Koornneef - A fortunate choice¹⁹
- Maarten Koornneef, David Meinke - The development of Arabidopsis as a model plant³⁷
- Ute Krämer - Planting molecular functions in an ecological context with Arabidopsis thaliana³⁸
- Nicholas J. Provart et al. - 50 years of Arabidopsis research³⁹

Acknowledgments

Thanks to Imre E. Somssich, Benjamin Schwessinger, Magnus Nordborg, Detlef Weigel, Rüdiger Simon, Kelsey L. Picard and Staffan Persson for helpful comments and support, and the Deutsche Forschungsgemeinschaft (DFG) for funding (project 344523413).

References

- 157 1. **Laibach F.** Zur Frage nach der Individualität der Chromosomen im Pflanzenreich. **Beih**
158 **Bot Zentralbl.** **1907**;22: 191–210. Available:
159 <https://www.biodiversitylibrary.org/item/27073#page/233/mode/1up>
- 160 2. **Laibach F.** 60 Jahre Arabidopsis-Forschung, 1905-1965. **Arab Inf Serv.** **1965**;1: 16.
161 Available: <http://www.arabidopsis.org/ais/1965/laiba-1965-aagle.html>
- 162 3. **Laibach F.** Arabidopsis Thaliana (L.) Heynh. als Objekt für genetische und
163 entwicklungsphysiologische Untersuchungen. **Bot Arch.** **1943**;44: 439–455. Available:
164 <http://131.130.57.230/clarotest190/claroline/backends/download.php?url=L0xhaWJhY2gtMTk0My5wZGY=&cidReset=true&cidReq=300415WS14>
165
- 166 4. **Reinholz E.** Röntgenmutationen bei Arabidopsis thaliana (L) Heynh.
167 **Naturwissenschaften.** **1947**;1: 26–28. Available:
168 <https://link.springer.com/article/10.1007/BF00633319>
- 169 5. **Kranz AR.** Demonstration of new and additional population samples and mutant lines of
170 the AIS-seed bank. **Arab Inf Serv.** **1978**;15: 2–4. Available:
171 <https://www.arabidopsis.org/ais/1978/kranz-1978-aabgw.html>
- 172 6. **Röbbelen G.** The LAIBACH Standard Collection of Natural Races. **Arab Inf Serv.**
173 **1965**;2. Available: <http://www.arabidopsis.org/ais/1965/roebb-1965-xxxxx.html>
- 174 7. **Meyerowitz EM, Pruitt RE.** Arabidopsis thaliana and Plant Molecular Genetics. **Science.**
175 **1985**;229: 1214–8. Available at doi:10.1126/science.229.4719.1214
- 176 8. **Koncz C.** Dedication: George P. Rédei Arabidopsis Geneticist and Polymath. Plant
177 Breeding Reviews. Oxford, UK: **John Wiley & Sons, Inc.**; **2010.** pp. 1–33. Available at
178 doi:10.1002/9780470650325.ch1
- 179 9. **Rédei GP.** A heuristic glance at the past of Arabidopsis genetics. **Methods in**
180 **Arabidopsis Research.** **1992.** pp. 1–15. Available at doi:10.1142/9789814439701_0001
- 181 10. **Potter E.** From Apathy to Apogee - Hardly anyone believed George Rédei’s research
182 mattered — until it changed everything. **Mizzou.** **2014**; Available:
183 <https://mizzoumag.missouri.edu/2014/08/from-apathy-to-apogee/>

- 184 11. **Rédei GP.** Supervital Mutants of Arabidopsis. **Genetics.** **1962**;47: 443–60. Available:
185 <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1210343>
- 186 12. **Langridge J.** Biochemical Mutations in the Crucifer Arabidopsis thaliana (L.) Heynh.
187 **Nature.** **1955**;176: 260–261. Available at doi:10.1038/176260b0
- 188 13. **Langridge J.** Arabidopsis thaliana, a plant Drosophila. **BioEssays.** **1994**;16: 775–778.
189 Available at doi:10.1002/bies.950161014
- 190 14. **Muller HJ.** Artificial transmutation of the gene. **Science.** **1927**;66: 84–87. Available at
191 doi:10.1126/science.66.1699.84
- 192 15. **Reinholz E.** Auslösung von Röntgen-Mutationen bei Arabidopsis thaliana (L.) Heynh. und
193 ihre Bedeutung für die Pflanzenzüchtung und Evolutionstheorie. **FIAT Report No. 1006.**
194 **1945.** Available: [https://www.tib.eu/de/suchen/id/TIBKAT:778643786/X-ray-mutations-](https://www.tib.eu/de/suchen/id/TIBKAT:778643786/X-ray-mutations-in-Arabidopsis-Thaliana-L-Heynh/)
195 [in-Arabidopsis-Thaliana-L-Heynh/](https://www.tib.eu/de/suchen/id/TIBKAT:778643786/X-ray-mutations-in-Arabidopsis-Thaliana-L-Heynh/)
- 196 16. **Rédei GP.** Single locus heterosis. **Z Vererbungsl.** **1962**;93: 164–170. Available:
197 <https://link.springer.com/article/10.1007/BF00897025>
- 198 17. **Nordborg M, Hu TT, Ishino Y, Jhaveri J, Toomajian C, Zheng H, et al.** The pattern of
199 polymorphism in Arabidopsis thaliana. **PLOS Biol.** **2005**;3: 1289–1299. Available at
200 doi:10.1371/journal.pbio.0030196
- 201 18. **Feenstra WJ.** Isolation of nutritional mutants in Arabidopsis thaliana. **Genetica.** **1964**;35:
202 259–269. Available at doi:10.1007/BF01804894
- 203 19. **Somerville CR, Koornneef M.** A fortunate choice: the history of Arabidopsis as a model
204 plant. **Nat Rev Genet.** **2002**;3: 883–9. Available at doi:10.1038/nrg927
- 205 20. **Röbbelen G.** Preface. **Arab Inf Serv.** **1964**;1: 1. Available:
206 <https://www.arabidopsis.org/ais/1964/preface.html>
- 207 21. **Huala E, Dickerman AW, Garcia-Hernandez M, Weems D, Reiser L, LaFond F, et al.**
208 The Arabidopsis Information Resource (TAIR): a comprehensive database and web-based
209 information retrieval, analysis, and visualization system for a model plant. **Nucleic Acids**
210 **Res.** **2001**;29: 102–5. Available at doi:10.1093/nar/29.1.102

- 211 22. **Rédei GP.** Arabidopsis as a Genetic Tool. **Annu Rev Genet.** **1975**;9: 111–127. Available
212 at doi:10.1146/annurev.ge.09.120175.000551
- 213 23. **Koornneef M, van Eden J, Hanhart CJ, Stam P, Braaksma FJ, Feenstra WJ.** Linkage
214 map of Arabidopsis thaliana. **J Hered.** **1983**;74: 265–272. Available at
215 doi:10.1093/oxfordjournals.jhered.a109781
- 216 24. **Leutwiler LS, Hough-Evans BR, Meyerowitz EM.** The DNA of Arabidopsis thaliana.
217 **Mol Gen Genet.** **1984**;194: 15–23. Available at doi:10.1007/BF00383491
- 218 25. **Somerville CR, Ogren WL.** Inhibition of photosynthesis in Arabidopsis mutants lacking
219 leaf glutamate synthase activity. **Nature.** **1980**;286: 257–259. Available at
220 doi:10.1038/286257a0
- 221 26. **Arabidopsis Genome Initiative.** Analysis of the genome sequence of the flowering plant
222 Arabidopsis thaliana. **Nature.** **2000**;408: 796–815. Available at doi:10.1038/35048692
- 223 27. **Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, et al.** Genome-wide
224 insertional mutagenesis of Arabidopsis thaliana. **Science.** **2003**;301: 653–7. Available at
225 doi:10.1126/science.1086391
- 226 28. **McNeill J, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, et al.**
227 International Code of Nomenclature for algae, fungi, and plants (Melbourne Code). **Koeltz**
228 **Sci Books.** **2012**;: 1–140. Available at doi:10.1111/j.1365-2699.2010.02341.x
- 229 29. **Rédei GP.** Arabidopsis thaliana (L.) Heynh. A review of the genetics and biology.
230 **Bibliogr Genet.** **1969**;20: 1–151.
- 231 30. **Linnaeus C.** Species Plantarum. **Impensis G. C. Nauk. Holmiae; 1753.** Available:
232 <https://www.biodiversitylibrary.org/item/13830#page/1/mode/1up>
- 233 31. **Heynhold G.** Nomenclator botanicus hortensis. **Arnoldische Buchhandlung.** Dresden
234 und Leipzig; **1840.** Available: <https://archive.org/details/nomenclatorbota00heyngoog>
- 235 32. **Holl F, Heynhold G.** Flora von Sachsen. **Verlag von Justus Naumann.** Dresden; **1842.**
236 Available: [https://books.google.com.au/books/about/Flora_von_Sachsen.html?id=pEI-](https://books.google.com.au/books/about/Flora_von_Sachsen.html?id=pEI-AAAACAAJ&redir_esc=y)
237 [AAAACAAJ&redir_esc=y](https://books.google.com.au/books/about/Flora_von_Sachsen.html?id=pEI-AAAACAAJ&redir_esc=y)

33. **ABRC**. Arabidopsis Natural Accessions (Ecotypes). **TAIR**. **2018**; Available:
https://www.arabidopsis.org/abrc/catalog/natural_accession_5.html
34. **Meyerowitz EM**. Arabidopsis Thaliana. **Annu Rev Genet**. **1987**;21: 93–111. Available at
[doi:10.1146/annurev.ge.21.120187.000521](https://doi.org/10.1146/annurev.ge.21.120187.000521)
35. **Pennisi E**. Arabidopsis Comes of Age. **Science**. **2000**;290: 32–35. Available at
[doi:10.1126/science.290.5489.32](https://doi.org/10.1126/science.290.5489.32)
36. **Meyerowitz EM**. Prehistory and history of Arabidopsis research. **Plant Physiol**.
2001;125: 15–9. Available at [doi:10.1104/pp.125.1.15](https://doi.org/10.1104/pp.125.1.15)
37. **Koornneef M, Meinke D**. The development of Arabidopsis as a model plant. **Plant J**.
2010;61: 909–21. Available at [doi:10.1111/j.1365-313X.2009.04086.x](https://doi.org/10.1111/j.1365-313X.2009.04086.x)
38. **Krämer U**. Planting molecular functions in an ecological context with Arabidopsis
thaliana. **Elife**. **2015**;4: 1–13. Available at [doi:10.7554/eLife.06100](https://doi.org/10.7554/eLife.06100)
39. **Provart NJ, Alonso J, Assmann SM, Bergmann DC, Brady SM, Brkljacic J, et al**. 50
years of Arabidopsis research: highlights and future directions. **New Phytol**. **2016**;209:
921–944. Available at [doi:10.1111/nph.13687](https://doi.org/10.1111/nph.13687)