

A peer-reviewed version of this preprint was published in PeerJ on 19 April 2019.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.6670) (peerj.com/articles/6670), which is the preferred citable publication unless you specifically need to cite this preprint.

Melo-Santos G, Figueiredo Rodrigues AL, Tardin RH, de Sá Maciel I, Marmontel M, Da Silva ML, May-Collado LJ. 2019. The newly described Araguaian river dolphins, *Inia araguaiaensis* (Cetartiodactyla, Iniidae), produce a diverse repertoire of acoustic signals. PeerJ 7:e6670 <https://doi.org/10.7717/peerj.6670>

The newly described Araguaian river dolphins, *Inia araguaiaensis*
(Cetartiodactyla, Iniidae), produce a diverse repertoire of acoustic signals

Gabriel Melo-Santos^{1,2, 3, 4}; Angélica Lúcia Figueiredo Rorigues¹; Rodrigo Hipólito Tardin^{5,6};
Israel de Sá Maciel^{6,7}; Miriam Marmontel³; Maria Luisa da Silva⁴; Laura Johanna May-Collado⁸

¹BioMA - Biology and Conservation of Amazonian Aquatic Mammals, Belém, Brazil

²Graduate program in Theory and Research of Behavior, Universidade Federal do Pará, Belém, Pará, Brazil

³Mamirauá Institute for Sustainable Development, Tefé, Amazonas, Brazil

⁴Laboratory of Ornithology and Bioacoustics, Institute of Biological Sciences, Universidade Federal do Pará, Belém, Pará, Brazil

⁵Department of Ecology, Universidade do Estado do Rio de Janeiro, Brazil

⁶Laboratory of Bioacoustics and Cetacean Ecology, Universidade Federal Rural do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

⁷Graduate Program in Animal Biology, Universidade Federal Rural do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

⁸Department of Biology, University of Vermont, Vermont, USA

Corresponding author:

Gabriel Melo-Santos

E-mail address: botogabriel@gmail.com

The newly described Araguaian river dolphins, *Inia araguaiaensis* (Cetartiodactyla, Iniidae), produce a diverse repertoire of acoustic signals

Abstract: The recent discovery of the Araguaian river dolphin (*Inia araguaiaensis*) highlights how little we know about the diversity and biology of river dolphins. In this study, we described the acoustic repertoire of this newly discovered species in concert with their behaviour in free-ranging, human-habituated individuals. We analysed 393 signals that we classified into 13 types of tonal sounds (n=15) and 66 types of pulsed calls (n=378). The most common sounds were short two-component calls. Thirty-five percent (n=140) of these calls were emitted by calves as they reunited with their mothers suggesting a key role in mother-calf communication. Our findings show that the acoustic repertoire of river dolphins is far from simple. Furthermore, the calls described here are similar in acoustic structure to those produced by social delphinids, such as orcas and pilot whales. Uncovering the context in which these signals are produced may help understand the social structure of this species and contribute to our understanding of the evolution of acoustic communication in whales.

Introduction

River dolphins of the genus *Inia* - commonly known as botos - are evolutionary relics found exclusively in the Amazon, Orinoco, and Tocantins River Basins of South America (Best & Da Silva, 1989, 1993; Hrbek et al., 2014; May-Collado & Agnarsson, 2011; Pilleri & Gühr, 1977; Santos et al., 2012, 2014). Like the franciscana dolphin (*Pontoporia blainvillei*), the baiji (*Lipotes vexillifer*), and the Ganges and Indus river dolphins (*Platanista* spp.), botos have flexible necks

and backbones, a low and large-based dorsal fin, and a slender rostrum (Best & Da Silva, 1989, 1993; Brownwell & Herald, 1972; Cassens, et al. 2000; Crespo et al., 1998; Da Silva et al., 2009; Hamilton et al., 2001; Leatherwoods & Reeves, 1994; Reeves & Martin, 2009;; Zhou 2009). Botos have a preference for habitats with slow currents and high prey concentration such as bays, confluences, small streams, and channels and island margins (Gomez-Salazar et al., 2012a, 2012b; Martin & Da Silva, 2004; Martin, Da Silva & Salmon, 2004; Pavanato et al., 2016). However, residency patterns vary within locations from long-term residency to occasional visitors (Martin & Da Silva, 2004). Although, botos are traditionally considered solitary, with long-term social interactions restricted to mothers and their calves, large aggregations have been documented during foraging and mating events (Best & Da Silva 1989, 1993; Martin, Da Silva & Rothery, 2008).

The acoustic repertoire of botos has traditionally been thought to be limited to a few sounds (Podos et al., 2002), however, studies of free-ranging and captive botos suggest otherwise. Among some of the sounds reported for botos are burst-pulsed sounds, jaw-snaps, low-frequency sounds, pulsed sounds, echolocation clicks, and whistles (Amorin et al., 2016; Cadwell et al., 1966; Diazgranados & Trujillo, 2002; May-Collado & Wartzok, 2007; Ding et al., 1995; Ding, Würsig & Leatherwoods, 2001; Kamminga et al., 1993; Podos, Da Silva & Rossi-Santos., 2002; Penner & Murchison, 1970). Ding, Würsig & Leatherwoods (2001) also described the emission of low-frequency whistles (up to 5 kHz) for Peruvian botos. However, this discovery was disputed (Podos, Da Silva & Rossi-Santos, 2002) due to the presence of sympatric tucuxi dolphins (*Sotalia fluviatilis*) known to emit whistles. Later, May-Collado & Wartzok (2007) confirmed that botos do emit whistles, but at much higher frequencies (up to 48 kHz) than previously thought. These high frequency whistles were recorded from botos at the Yasuni and Napo rivers in Ecuador. Today,

there is a consensus that, while botos do emit whistles, these sounds are emitted rarely (Diazgranados & Trujillo, 2002; May-Collado & Wartzok, 2007; Ding, Würsig & Leatherwoods, 2001) and likely play a different social role as the one described for delphinids (May-Collado & Wartzok, 2007). Podos, Da Silva & Rossi-Santos (2002) found that the acoustic repertoire of Amazonian botos consisted primarily of pulsed calls with a low emission rate. However, these results were likely limited by the sampling rate of the recorders used by the authors. Amorin *et al.* (2016) studied the same population using a broadband frequency recording system and described a high emission of a variety of pulsed calls.


A major constraint in studying river dolphins is that they usually do not perform conspicuous surface displays combined with the difficulty of identifying individuals in the field (Best & Da Silva, 1989, 1993; Brownwell & Herald, 1972; Cassens, et al. 2000; Crespo, Harris & González, 1998; Da Silva, 2009; Hamilton et al., 2001; Leatherwoods & Reeves, 1994; Reeves & Martin, 2009; Zhou 2009). Here we studied the acoustic repertoire of free-ranging Arauguaian botos that regularly visit a fish market in Mocajuba in Northern Brazil (Santos et al., 2014). This semi-controlled setting gave us the unique opportunity to combine acoustic technology with underwater behavioural observations.

Material & Methods

Study area

This study took place along the Tocantins River in the town of Mocajuba in Pará State, Brazil (Fig. 1). The Tocantins River is classified as a clearwater river with a small floodplain as the river runs through a narrow valley. There are large sandbanks in the river's main channel where herbaceous vegetation may occur, there are floating vegetation, and submerged aquatic macrophytes where there is light penetration (Junk et al., 2011). At its lower reaches, water cycles in this region are very dynamic with a greater rainfall from November to April, with the highest

95 waters on March, rainfall declines from May to October, with lowest waters on September
 96 (Ribeiro, Pretere & Juras, 1995). There is also a daily cycle of tide pulses (Goulding et al., 2003;
 97 Ribeiro, Petrere & Juras, 1995). Mocajuba has a fish market that serves as the main place to acquire
 98 fish products for the city and the riverside communities. The wastes of the market and the provision
 99 of fish by locals attracts botos to the pier. This set up together with low turbidity waters during the
 100 dry season allows great proximity to botos enabling us to identify individuals and observe their
 101 behaviour in detail (Fig. 2).



102

103 **Figure 1.** Location of the Mocajuba fish market at the margins of the Tocantins River

104

105

106



107 **Figure 2.** Mocajuba's fish market, clear and shallow waters allow for detailed observation of
 108 Araguaian botos' behaviour. (A): Taken from drone footage by our team led by Gabriel Melo-
 109 Santos. (B) Photo by Rodrigo Tardin.

110

111 *Data collection*

112 Acoustic and behavioural data was collected in visits that ranged from three to 15 days
 113 during October to December 2013, March 2014, June 2015, July, September and December 2016.
 114 The presence of botos at the market depends on the market opening hours, which is the time when
 115 the animals are fed (Santos et al., 2014). Therefore our observations took place in the morning.
 116 Behavioural observations were collected in a continuous all-events sampling (Mann, 1999). For
 117 each session, we collected the following data: number of individuals present, age class (adult,
 118 juvenile, calf), and sex (based on the presence of mammary slits). In addition, animals were
 119 identified based on natural marks on the dorsal and ventral parts of the body, given that the botos
 120 in the market frequently swim upside down (Santos et al., 2014). Photographs of their bodies were
 121 taken with a Nikon 3200 SLR Camera (Nikon Corp., Tokyo, Japan) and a 70x300mm zoom lens
 122 (Nikon Corp., Tokyo, Japan). Underwater video was collected with a GoPro Hero 4 (GoPro Inc.,

San Mateo, USA) held on hand (Fig. 5). Notes and drawings of the marks and their locations were also taken if we were unable to take pictures. We held permits to perform this study issued by SISBio (number 52892) from the Brazilian Ministry of Environment.

Sound recordings were taken continuously in synchrony with behavioural observations. We used three recording systems along the study: (1) an Aquarian hydrophone (Anacortes, Washington, USA) connected to a Tascam DR1 digital recorder (22kHz sampling rate), (2) a CR1 hydrophone (Cetacean Research Technology, Seattle, USA) connected to a pre-amplifier and a Tascam DR-44WL (96kHz sampling rate) and (3) a Soundtrap (Ocean Instruments, New Zealand, 576kHz, sampling rate).

Data analysis

All recorded signals were inspected using a spectrogram analysis in Raven Pro 1.5 (Cornell Laboratory of Ornithology, New York, USA). Only whistles and pulsed calls with good signal to noise ratio were selected for further analysis. Signals were assigned visually *a posteriori* into categories based on their shape of their contours (Fig.6). Other defining characteristics of the classification of boto sounds were duration - short (<200ms) versus long (>200ms) signals - and the presence of nonlinear phenomena: (a) subharmonics - signals with additional spectral components in the harmonic stack, generally in multiples of 1/2 or 1/3 of the fundamental frequency- and (b) biphonation - signals with the presence of two independent fundamental frequencies (Tokuda et al., 2002; Wielden et al., 1998). Pulsed calls produced in a succession where considered a single signal if they were separated by less than 200 ms. Using a rarefaction curve (Magurran, 2004), we evaluated how much of the acoustic repertoire was registered during

our sampling period. Using a Whittaker diagram (Magurran, 2004), we assessed the occurrence of the signals recorded as part of these animal's repertoire. Analyses were conducted in R.

Results

Botos were observed on 32 days (sightings happened everyday) and we collected 20.2 hours of acoustic recordings. Group sizes ranged between 3 to 12 individuals. These animals repeatedly visited the market allowing us to photo-identify the animals and often following specific individuals. We were able to identify nine individuals by their natural marks, five adult females, one adult male, one juvenile female, one female calf and one male calf. Only two behaviours were observed: socialization and feeding (Fig.3). Social interactions consisted of animals having physical contact with one another and swimming alongside each other. Although, occasionally animals would bite the neck of another when waiting to be fed. While we did not specifically test for associations between individuals, the most stable associations appeared to be between mothers and their calves. Feeding behaviour consisted of animals soliciting food with the head out of the water and open-mouthed or poking humans with their snout. However, with the help of underwater cameras we were able to match some of the observations to the vocalizing animals (see below).





Figure 3. The two behaviours observed during this study were socialization and feeding. (A) and (B): botos engaged in social activity, slow-swimming and physical contact, (C): boto feeding on a fish and (D): waiting to be fed. Photos (A), (B) and (C) by Gabriel Melo-Santos and (D) by Luiza Pereira.

Based on signal quality we selected at total 393 acoustic signals and classify them into six major sound types and 78 sub-categories. The six sound types were: long-two-component calls, long calls with subharmonics, short calls with biphonation (short-two-component calls), short calls without non-linear phenomena, short-calls with subharmonics, and tonal sounds (Fig. 4, Table 1). While we found a diversity of calls the rarefaction curve indicates that 20 hours of acoustic sampling was not sufficient to capture most of the acoustic repertoire of these animals (Fig.5).

174 However, botos do seem to produce some signals more abundantly than others (Fig. 6). The long-
175 calls with subharmonics (n=11) and whistles (n=15) were rarest and the short-two-component calls
176 were the most commonly produced (n=246). Interestingly, 35% (n=140) of these were short two-
177 component calls. Some of these calls were produced by calves as observed by the bubble emission
178 from the blowhole, and their emission was followed by physical contact with their mothers (Fig.7).

179



181 **Figure 4.** Examples of acoustic signals recorded during our study:(A)- Long call with
182 subharmonics followed by two short two-component calls; (B) -Long two-component call; (C) -
183 Narrow-banded frequency modulated whistle and the longest sound registered in this study; (D) –
184 Short call with no non-linear phenomena. E – A short call with subharmonics.

185


186

187

188


Table 1. Major classification categories of signals recorded from the Araguaian river dolphin at the Mocajuba fish market.

Vocalization type	Number of sub-categories	Number of sounds produced
Long two-component calls - pulsed calls longer than 0.200s, with biphonation as the most marking characteristic, may exhibit subharmonics	9	18
Long calls with subharmonics- calls longer than 0.200s, may exhibit biphonation	5	11
Short two-component calls - pulsed calls shorter than 0.200s, with biphonation as the most marking characteristic, may exhibit subharmonics	29	246
Short calls - signals shorter than 0.200s with no non-linear phenomena	7	29
Short calls with subharmonics - calls shorter than 0.200s, may exhibit biphonation	15	74
Whistles - tonal narrow-banded signals longer than 0.100s, may exhibit harmonics	13	15
Total	78	393



193

194 **Figure 5.** Rarefaction curve showing the cumulative number of call types detected with
 195 increasing sampling time (hours of recordings analysed). The curve suggests that 20 hours of
 196 acoustic sampling is not enough to capture most of the acoustic repertoire of the Araguaian river
 197 dolphin.



198

199 **Figure 6.** Whittaker diagram displaying the occurrence of calls emitted by Araguaian botos with
 200 most calls produced rarely.




Figure 7. (A): Vocalizing calf as indicated by the bubbles blowing from its blowhole and associated short-two-component call, taken from video footage by our team. (B) Spectrogram of calls produced by a calf of Araguaian boto, followed by a bubble stream.

Discussion

Our results show that the Araguaian river dolphin has a more diverse acoustic repertoire than previously documented for the genus *Inia* (Amorin et al., 2016; Cadwell et al., 1966; Diazgranados & Trujillo, 2002; May-Collado & Wartzok, 2007; Ding et al., 1995, 2001; Kamminga et al., 1993; Podos, Da Silva & Rossi-Santos, 2002). Early studies described members of the genus *Inia* as silent animals (Podos et al., 2002) or with a limited acoustic repertoire (Amorin et al., 2016; Cadwell et al., 1966; Diazgranados & Trujillo, 2002; May-Collado & Wartzok, 2007; Ding et al., 1995; Ding, Würsig & Leatherwoods, 2001; Kamminga et al., 1993; Podos, Da Silva & Rossi-Santos, 2002). However, here we show that the acoustic repertoire of *Inia* dolphins is diverse and likely as complex as the acoustic repertoire of delphinids.

While pulsed calls were the most commonly produced sounds, tonal sounds were also produced. These findings agree with previous descriptions of whistle emission from Peruvian

(Ding, Würsig & Leatherwoods, 2001) and Ecuadorian botos (May-Collado & Wartzok, 2007) and other river dolphins like the Franciscana and the baiji (Cremer et al., 2017; Wang, Wang & Akamatsu, 1999; Wang et al., 2006). In these studies, the function of emitted whistles was unclear. May-Collado & Wartzok (2007) found that botos emitted whistles in a different social context than in delphinids, keeping distance between each other, rather than promoting social interactions as in marine dolphins.

Among pulsed calls, the short-two-component call was the most commonly produced sound. These calls were emitted in what appear to be mother-calf interactions. Our video footage and some underwater follows show bubbles emanating from calves' blowholes while they emitted these calls as they approached their mothers after a short separation (see Supplementary Material). Bubble streams are often used as a cue to identify vocalizing animals (Bebus & Herzing, 2015; Fripp, 2005; Jones, 2014) and in this case the bubble stream revealed that the calves were producing the calls and did so in a repetitive fashion. These vocal patterns are similar to what has been described for calves of bottlenose dolphins, which use signature whistle as contact calls, where calves increase whistle emission as they approach their mothers (Smolker, Mann & Smuts, 1993). Given the strength of mother-calf associations in botos (Best & Da Silva, 1989, 1993) and the characteristics of their habitat, a shared signal that enhances mother-calf recognition may be key as they move through murky waters and complex underwater vegetation. The complex structure of botos' might also have led to evolution towards signals with short duration, longer signals might suffer interference of echoes caused by obstacles (sandbanks, underwater vegetation, riverbed, even the water surface). Notwithstanding, social signals produced by *Inia* sister taxa *Pontoporia* who also evolved in riverine environments are also short (Cremer et al., 2017).

Several species of toothed whales emit calls of similar acoustic nature as the ones described here for botos (Filatova et al., 2012; Fitch et al., 2002; Ford, 1989; Deecke et al., 1999, 2010, 2011; Garland et al., 2015; Marcoux et al., 2012; Miller & Bain, 2000; Papale et al., 2015; Pérez et al., 2017; Sjare & Smith, 1986; Vergara & Barrett-Lennard, 2008; Vergara, Michaud & Barrett-Lennard, 2010; Yurk et al., 2002; Zwamborn & Whitehead, 2017). For example, the calls of killer (*Orcinus orca*), pilot (*Globicephala* spp.) whales have been shown to contain non-linear features suggesting they may carry information on group identity and maintaining social cohesion (Deecke et al., 2010; Pérez et al., 2017; Yurk et al., 2002; Zwamborn & Whitehead, 2017). Similarly, Marcoux, Auger- Méthé & Humphries (2012) show evidence that narwhal (*Monodon Monoceros*) calls might be related to specific groups or individuals. Non-linear calls have also been reported to convey individuals' identity and/or emotional state (Fitch, Neubauer & Hetzel, 2002; Papale et al., 2015). Given these similarities we propose these two-component signals evolved early in the evolutionary history of toothed whales as social contact signals, likely for mother-calf interactions and later in the lineage leading to delphinids it evolved into a group recognition signal.

Conclusions

We show that the acoustic repertoire of botos is far from being simple. The Araguaian river dolphins studied at Mocajuba fish market produce a diverse acoustic repertoire, as we found 78 sound types, mostly pulsed calls, and our analysis indicates that there is more to discover. Notwithstanding, these sounds are mostly complex in structure presenting nonlinear phenomena. The animals we studied are habituated to humans, which provided a unique opportunity to shed light on the acoustic and social behaviour of this understudied species. Under relatively controlled

conditions we identified more than half of the studied animals and recorded their acoustic and underwater behaviour. When possible, we matched recordings with video footage of calves as they reunited with their mothers. During this reunions calves appeared to use the two-component calls as contact calls, nevertheless further investigation is needed to understand the importance of these calls for mother-calf interactions. Furthermore, given that Araguaian river dolphin calls are similar in acoustic structure to those of delphinids, we proposed that these signals evolved early in the evolutionary history of toothed whales as social calls, likely as mother-calf contact calls, and that later in the lineage leading to dolphins its function evolved to group/family call recognition.

Acknowledgments

We are thankful to the Rufford Foundation and Cetacean Society International for financing our field work. The first author was granted a scholarship from the CAPES foundation (Brazilian Ministry of Education). We are also thankful to the community of Mocajuba for the support to our work in the market. We are also thankful to Volker Deecke by the contributions to this manuscript.

References

- Amorim, T.O.S., Andriolo, A., Reis, S.S. and dos Santos, M.E., 2016. Vocalizations of Amazon river dolphins (*Inia geoffrensis*): Characterization, effect of physical environment and differences between populations. *The Journal of the Acoustical Society of America*, 139(3), pp.1285-1293.
- Bebus, S.E., Herzing, D.L., 2015. Mother-offspring signature whistle similarity and patterns of association in Atlantic spotted dolphins (*Stenella frontalis*). *Animal Behavior and Cognition*, 2(1), pp.71-87.
- Best, R.C., da Silva, V.M., 1989. Amazon river dolphin, boto *Inia geoffrensis* (de Blainville, 1817). *Handbook of marine mammals*, 4, pp.1-23.
- Best, R.C. and da Silva, V.M., 1993. *Inia geoffrensis*. *Mammalian Species*, (426), pp.1-8.
- Brownell, R.L., Herald, E.S., 1972. *Lipotes vexillifer*. *Mammalian Species*, (10), pp.1-4.

- 289 Caldwell, M.C., Caldwell, D.K., Evans, W.E., 1966. *Sounds and behavior of captive Amazon*
290 *freshwater dolphins, Inia geoffrensis*. Los Angeles County Museum of Natural History.
- 291 Cassens, I., Vicario, S., Waddell, V.G., Balchowsky, H., Van Belle, D., Ding, W., Fan, C., Mohan,
292 R.L., Simões-Lopes, P.C., Bastida, R., Meyer, A., 2000. Independent adaptation to riverine
293 habitats allowed survival of ancient cetacean lineages. *Proceedings of the National Academy of*
294 *Sciences*, 97(21), pp.11343-11347.
- 295 Cremer, M.J., Holz, A.C., Bordino, P., Wells, R.S. and Simões-Lopes, P.C., 2017. Social sounds
296 produced by franciscana dolphins, *Pontoporia blainvillei* (Cetartiodactyla, Pontoporiidae). *The*
297 *Journal of the Acoustical Society of America*, 141(3), pp.2047-2054.
- 298 Crespo, E.A., Harris, G., González, R., 1998. Group size and distributional range of the
299 franciscana, *Pontoporia blainvillei*. *Marine Mammal Science*, 14(4), pp.845-849.
- 300 Da Silva, V.M., 2009. Amazon river dolphin: *Inia geoffrensis*. In Perrin, W. F., Wursig, B.,
301 Thewissen, J. G. M., eds. *Encyclopedia of Marine Mammals (Second Edition)* (pp. 26-28).
- 302 Deecke, V.B., Ford, J.K., Spong, P., 1999. Quantifying complex patterns of bioacoustic variation:
303 Use of a neural network to compare killer whale (*Orcinus orca*) dialects. *The Journal of the*
304 *Acoustical Society of America*, 105(4), pp.2499-2507.
- 305 Deecke, V.B., Barrett-Lennard, L.G., Spong, P. and Ford, J.K., 2010. The structure of stereotyped
306 calls reflects kinship and social affiliation in resident killer whales (*Orcinus*
307 *orca*). *Naturwissenschaften*, 97(5), pp.513-518.
- 308 Deecke, V.B., Nykänen, M., Foote, A.D., Janik, V.M., 2011. Vocal behaviour and feeding ecology
309 of killer whales *Orcinus orca* around Shetland, UK. *Aquatic Biology*, 13(1), pp.78-88.
- 310 Diazgranados, M.C., Trujillo, F., 2002. Vocal repertoire of the freshwater dolphins *Inia geoffrensis*
311 and *Sotalia fluviatilis* in Colombia, South America. *The Journal of the Acoustical Society of*
312 *America*, 112(5), pp.2400-2400.
- 313 Ding, W., Würsig, B., Evans, W., 1995. Comparisons of whistles among seven odontocete
314 species. *Sensory systems of aquatic mammals*, 25.
- 315
- 316 Ding, W., Würsig, B., Leatherwoods, S., 2001. Whistles of boto, *Inia geoffrensis*, and tucuxi,
317 *Sotalia fluviatilis*. *The Journal of the Acoustical Society of America*, 109(1), pp.407-411.
- 318 Filatova, O.A., Deecke, V.B., Ford, J.K., Matkin, C.O., Barrett-Lennard, L.G., Guzeev, M.A.,
319 Burdin, A.M., Hoyt, E., 2012. Call diversity in the North Pacific killer whale populations:
320 implications for dialect evolution and population history. *Animal Behaviour*, 83(3), pp.595-603.

- 321 Fitch, W.T., Neubauer, J, Herzel, H., 2002. Calls out of chaos: the adaptive significance of
322 nonlinear phenomena in mammalian vocal production. *Animal behaviour*, 63(3), pp.407-418.
- 323 Ford, J.K., 1989. Acoustic behaviour of resident killer whales (*Orcinus orca*) off Vancouver
324 Island, British Columbia. *Canadian Journal of Zoology*, 67(3), pp.727-745.
- 325 Fripp, D., 2005. Bubblestream whistles are not representative of a bottlenose dolphin's vocal
326 repertoire. *Marine Mammal Science*, 21(1), pp.29-44.
- 327 Garland, E.C., Castellote, M., Berchok, C.L., 2015. Beluga whale (*Delphinapterus leucas*)
328 vocalizations and call classification from the eastern Beaufort Sea population. *The Journal of the*
329 *Acoustical Society of America*, 137(6), pp.3054-3067.
- 330 Gomez-Salazar, C., Trujillo, F., Portocarrero-Aya, M., Whitehead, H., 2012a. Population, density
331 estimates, and conservation of river dolphins (*Inia* and *Sotalia*) in the Amazon and Orinoco river
332 basins. *Marine Mammal Science*, 28(1), pp.124-153.
- 333 Gomez-Salazar, C., Trujillo, F., Whitehead, H., 2012b. Ecological factors influencing group sizes
334 of river dolphins (*Inia geoffrensis* and *Sotalia fluviatilis*). *Marine Mammal Science*, 28(2).
- 335 Goulding, M., Barthem, R., Ferreira, E.J.G., Duenas, R., 2003. *The Smithsonian atlas of the*
336 *Amazon* (p. 253). Washington: Smithsonian Books.
- 337 Hamilton, H., Caballero, S., Collins, A.G., Brownell, R.L., 2001. Evolution of river
338 dolphins. *Proceedings of the Royal Society of London B: Biological Sciences*, 268(1466), pp.549-
339 556.
- 340 Hrbek, T., da Silva, V.M.F., Dutra, N., Gravena, W., Martin, A.R., Farias, I.P., 2014. A new
341 species of river dolphin from Brazil or: how little do we know our biodiversity. *PLoS one*, 9(1),
342 p.e83623.
- 343 Jones, B.L., 2014. The ontogeny of whistle production in infant Atlantic Bottlenose Dolphins
344 (*Tursiops truncatus*) during the first thirty days of life. Masters Thesis, The University of Southern
345 Mississippi.
- 346 Junk, W.J., Piedade, M.T.F., Schöngart, J., Cohn-Haft, M., Adeney, J.M., Wittmann, F., 2011. A
347 classification of major naturally-occurring Amazonian lowland wetlands. *Wetlands*, 31(4),
348 pp.623-640.
- 349 Kamminga, C., Van Hove, M.T., Englesma, F.J. and Terry, R.P., 1993. Investigations on cetacean
350 sonar X: a comparative analysis of underwater echolocation clicks of *Inia* spp. and *Sotalia*
351 spp. *Aquatic Mammals*, 19, pp.31-31.
- 352 Leatherwoods, S., Reeves, R.R., 1994. River dolphins: a review of activities and plans of the
353 Cetacean Specialist Group. *Aquatic Mammals*, 20(3), pp.137-154.

- 354 Magurran, A.E., 2004. *Measuring biological diversity*. John Wiley & Sons.
- 355 Mann, J., 1999. Behavioral sampling methods for cetaceans: a review and critique. *Marine*
356 *mammal science*, 15(1), pp.102-122.
- 357 Marcoux, M., Auger-Méthé, M., Humphries, M.M., 2012. Variability and context specificity of
358 narwhal (*Monodon monoceros*) whistles and pulsed calls. *Marine Mammal Science*, 28(4),
359 pp.649-665.
- 360 Martin, A.R. and da Silva, V.M., 2004. Number, seasonal movements, and residency
361 characteristics of river dolphins in an Amazonian floodplain lake system. *Canadian Journal of*
362 *Zoology*, 82(8), pp.1307-1315.
- 363 Martin, A.R., Silva, V., Salmon, D.L., 2004. Riverine habitat preferences of botos (*Inia*
364 *geoffrensis*) and tucuxis (*Sotalia fluviatilis*) in the central Amazon. *Marine Mammal*
365 *Science*, 20(2), pp.189-200.
- 366 Martin, A.R., Da Silva, V.M.F., Rothery, P., 2008. Object carrying as socio-sexual display in an
367 aquatic mammal. *Biology letters*, 4(3), pp.243-245.
- 368 May-Collado, L.J., Wartzok, D., 2007. The freshwater dolphin *Inia geoffrensis* geoffrensis
369 produces high frequency whistles. *The Journal of the Acoustical Society of America*, 121(2),
370 pp.1203-1212.
- 371 May-Collado, L.J., Agnarsson, I., 2011. Phylogenetic analysis of conservation priorities for
372 aquatic mammals and their terrestrial relatives, with a comparison of methods. *PLoS One*, 6(7),
373 p.e22562.
- 374 Miller, P.J. and Bain, D.E., 2000. Within-pod variation in the sound production of a pod of killer
375 whales, *Orcinus orca*. *Animal Behaviour*, 60(5), pp.617-628.
- 376 Papale, E., Buffa, G., Filiciotto, F., Maccarrone, V., Mazzola, S., Ceraulo, M., Giacomini, C. and
377 Buscaino, G., 2015. Biphonic calls as signature whistles in a free-ranging bottlenose
378 dolphin. *Bioacoustics*, 24(3), pp.223-231.
- 379 Pavanato, H.J., Melo-Santos, G., Lima, D.S., Portocarrero-Aya, M., Paschoalini, M., Mosquera,
380 F., Trujillo, F., Meneses, R., Marmontel, M., Maretti, C., 2016. Risks of dam construction for
381 South American river dolphins: a case study of the Tapajós River. *Endangered Species*
382 *Research*, 31, pp.47-60.
- 383 Pérez, J.M., Jensen, F.H., Rojano-Doñate, L., Aguilar de Soto, N., 2017. Different modes of
384 acoustic communication in deep-diving short-finned pilot whales (*Globicephala*
385 *macrorhynchus*). *Marine Mammal Science*, 33(1), pp.59-78.

- 386 Pilleri, G., Gahr, M., 1977. Observations on the Bolivian, *Inia boliviensis*, (D'Orbigny, 1834) and
387 the Amazonian bufeo, *Inia geoffrensis* (Blainville, 1817), with a description of a new subspecies
388 (*Inia geoffrensis humboldtiana*). *Investigations on CETACEA*, 8(1), pp.1-76.
- 389 Podos, J., Da Silva, V.M., Rossi-Santos, M.R., 2002. Vocalizations of Amazon river dolphins, *Inia*
390 *geoffrensis*: insights into the Evolutionary origins of delphinid whistles. *Ethology*, 108(7), pp.601-
391 612.
- 392 Reeves, R.R., Martin, A.R., 2009. River dolphins. In Perrin, W. F., Wursig, B., Thewissen, J. G.
393 M., eds. *Encyclopedia of Marine Mammals (Second Edition)* (pp. 976-979).
- 394 Ribeiro, M.C.L.D.B., Petrere, M., Juras, A.A., 1995. Ecological integrity and fisheries ecology of
395 the Araguaia—Tocantins River Basin, Brazil. *River Research and Applications*, 11(3-4), pp.325-
396 350.
- 397 Santos, G.M.A., Quaresma, A.C., Barata, R.R., Martins, B.M., Siciliano, S., e Silva, J.D.S., Emin-
398 Lima, R., 2012. Etho-ecological study of the Amazon River dolphin, *Inia geoffrensis* (Cetacea:
399 Iniidae), and the dolphins of the genus *Sotalia* (Cetacea: Delphinidae) in Guamá River,
400 Amazonia. *Marine Biodiversity Records*, 5.
- 401 Santos, G.M.A., Rodrigues, A.L.F., Arcoverde, D.L., Ramos, I., Sena, L., Silva, M.L., 2014.
402 Unusual records of the behavior of boto *Inia* sp.(Cetartiodactyla, Iniidae) in the lower reaches of
403 the Tocantins and Guamá River, Amazônia. *Dolphins: ecology, behavior and conservation*
404 *strategies*. Nova Science Publishers, New York, NY, pp.1-17.
- 405 Sjare, B.L., Smith, T.G., 1986. The vocal repertoire of white whales, *Delphinapterus leucas*,
406 summering in Cunningham Inlet, Northwest Territories. *Canadian Journal of Zoology*, 64(2),
407 pp.407-415.
- 408 Smolker, R.A., Mann, J., Smuts, B.B., 1993. Use of signature whistles during separations and
409 reunions by wild bottlenose dolphin mothers and infants. *Behavioral Ecology and*
410 *Sociobiology*, 33(6), pp.393-402.
- 411 Tokuda, I., Riede, T., Neubauer, J., Owren, M.J., Herzog, H., 2002. Nonlinear analysis of irregular
412 animal vocalizations. *The Journal of the Acoustical Society of America*, 111(6), pp.2908-2919.
- 413 Vergara, V., Barrett-Lennard, L.G., 2008. Vocal development in a beluga calf (*Delphinapterus*
414 *leucas*). *Aquatic Mammals*, 34(1), p.123.
- 415 Vergara, V., Michaud, R., Barrett-Lennard, L., 2010. What can captive whales tell us about their
416 wild counterparts? Identification, usage, and ontogeny of contact calls in belugas (*Delphinapterus*
417 *leucas*). *International Journal of Comparative Psychology*, 23(3).
- 418 Wang, D., Wang, K.X., Akamatsu, T., K., 1999. Study on whistling of the Chinese river dolphin
419 (*Lipotes vexillifer*). *Oceanologia et Limnologia Sinica*, 30, pp.353-360.

- 420 Wang, K., Wang, D., Akamatsu, T., Fujita, K., Shiraki, R., 2006. Estimated detection distance of
421 a baiji's (Chinese river dolphin, *Lipotes vexillifer*) whistles using a passive acoustic survey
422 method. *The Journal of the Acoustical Society of America*, 120(3), pp.1361-1365.
- 423 Wilden, I., Herzel, H., Peters, G., Tembrock, G., 1998. Subharmonics, biphonation, and
424 deterministic chaos in mammal vocalization. *Bioacoustics*, 9(3), pp.171-196.
- 425 Yurk, H., Barrett-Lennard, L., Ford, J.K.B. and Matkin, C.O., 2002. Cultural transmission within
426 maternal lineages: vocal clans in resident killer whales in southern Alaska. *Animal*
427 *Behaviour*, 63(6), pp.1103-1119.
- 428 Zwamborn, E.M., Whitehead, H., 2017. Repeated call sequences and behavioural context in long-
429 finned pilot whales off Cape Breton, Nova Scotia, Canada. *Bioacoustics*, 26(2), pp.169-183.
- 430