

A peer-reviewed version of this preprint was published in PeerJ on 6 November 2018.

[View the peer-reviewed version](https://peerj.com/articles/5709) (peerj.com/articles/5709), which is the preferred citable publication unless you specifically need to cite this preprint.

Abyzova GA, Nikitin MA, Popova OV, Pasternak AF. 2018. Genetic population structure of the pelagic mollusk *Limacina helicina* in the Kara Sea. PeerJ 6:e5709 <https://doi.org/10.7717/peerj.5709>

Genetic population structure of the pelagic mollusk *Limacina helicina* in the Kara Sea

Galina Anatolievna Abyzova ^{Corresp., 1}, Mikhail Aleksandrovich Nikitin ², Olga Vladimirovna Popova ², Anna Fedorovna Pasternak ¹

¹ Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia

² Belozersky Institute for Physico-Chemical Biology, Lomonosov Moscow State University, Moscow, Russia

Corresponding Author: Galina Anatolievna Abyzova
Email address: abyzova.ga@ocean.ru

Background. Pelagic pteropods *Limacina helicina* are widespread and can play an important role in the food webs and in biosedimentation in Arctic and Subarctic ecosystems. Previous publications have shown differences in the genetic structure of populations of *L. helicina* from populations found in the Pacific Ocean and Svalbard area. Currently, there are no data on the genetic structure of *L. helicina* populations in the seas of the Siberian Arctic. We assessed the genetic structure of *L. helicina* from the Kara Sea populations and compared them with samples from around Svalbard and the North Pacific. We also compared *L. helicina* from the different habitats within the Kara Sea.

Methods. We examined genetic differences in *L. helicina* from three different locations in the Kara Sea via analysis of a fragment of the mitochondrial gene COI. We also compared a subset of samples with *L. helicina* from previous studies to find connections between populations from the Atlantic and Pacific Oceans.

Results. 65 individual *L. helicina* from the Kara Sea were sequenced to produce 19 different haplotypes. This is comparable with numbers of haplotypes found in Svalbard and Pacific samples (24 and 25, respectively). Haplotypes from different locations sampled around Arctic and Subarctic were combined into two significantly different groups: H1 and H2. The H2 includes sequences from the Kara Sea and Svalbard, was present only in the Atlantic sector of the Arctic. The other genetic group, H1, is widespread and found throughout all *L. helicina* populations. Phi-st analyses also indicated significant genetic difference between the Atlantic and Pacific regions, but no differences between Svalbard and the Kara Sea.

Discussion. The obtained results support our hypothesis about genetic similarity of *L. helicina* populations from the Kara Sea and Svalbard: the majority of haplotypes belongs to the haplotype group H2, with the H1 group representing a minority of the haplotypes present. In contrast, in the Canadian Arctic and the Pacific Ocean only haplogroup H1 is found. The negative values of Fu's F_s indicate directed selection or expansion of the population. The reason for this pattern could be due to an isolation of the *Limacina helicina* population during the Pleistocene glaciation and a subsequent rapid expansion of this species after the last glacial maximum.

1 **Genetic population structure of the pelagic mollusk *Limacina***
2 ***helicina* in the Kara Sea**

3 Galina Anatolievna Abyzova¹, Mikhail Aleksandrovich Nikitin², Olga Vladimirovna Popova²,
4 Anna Fedorovna Pasternak¹

5 ¹ Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia

6 ² Belozersky Institute for Physico-Chemical Biology, Lomonosov Moscow State University,
7 Moscow, Russia

8 Corresponding Author:

9 Galina Abyzova¹

10 36, Nahimovskiy prospekt, Moscow, 117997, Russia

11 Email address: abyzova.ga@ocean.ru

12 Abstract

13 **Background.** Pelagic pteropods *Limacina helicina* are widespread and can play an important
14 role in the food webs and in biosedimentation in Arctic and Subarctic ecosystems. Previous
15 publications have shown differences in the genetic structure of populations of *L. helicina* from
16 populations found in the Pacific Ocean and Svalbard area. Currently, there are no data on the
17 genetic structure of *L. helicina* populations in the seas of the Siberian Arctic. We assessed the
18 genetic structure of *L. helicina* from the Kara Sea populations and compared them with samples
19 from around Svalbard and the North Pacific. We also compared *L. helicina* from the different
20 habitats within the Kara Sea.

21 **Methods.** We examined genetic differences in *L. helicina* from three different locations in the
22 Kara Sea via analysis of a fragment of the mitochondrial gene COI. We also compared a subset
23 of samples with *L. helicina* from previous studies to find connections between populations from
24 the Atlantic and Pacific Oceans.

25 **Results.** 65 individual *L. helicina* from the Kara Sea were sequenced to produce 19 different
26 haplotypes. This is comparable with numbers of haplotypes found in Svalbard and Pacific
27 samples (24 and 25, respectively). Haplotypes from different locations sampled around Arctic
28 and Subarctic were combined into two significantly different groups: H1 and H2. The H2
29 includes sequences from the Kara Sea and Svalbard, was present only in the Atlantic sector of
30 the Arctic. The other genetic group, H1, is widespread and found throughout all *L. helicina*
31 populations. Phi-st analyses also indicated significant genetic difference between the Atlantic
32 and Pacific regions, but no differences between Svalbard and the Kara Sea.

33 **Discussion.** The obtained results support our hypothesis about genetic similarity of *L. helicina*
34 populations from the Kara Sea and Svalbard: the majority of haplotypes belongs to the haplotype
35 group H2, with the H1 group representing a minority of the haplotypes present. In contrast, in the
36 Canadian Arctic and the Pacific Ocean only haplogroup H1 is found. The negative values of Fu's
37 Fs indicate directed selection or expansion of the population. The reason for this pattern could be
38 due to an isolation of the *Limacina helicina* population during the Pleistocene glaciation and a
39 subsequent rapid expansion of this species after the last glacial maximum.

40 Introduction

41 Pelagic pteropods *Limacina helicina* (Phipps, 1774) are widespread in marine Arctic and
42 Subarctic ecosystems, where their local abundance and biomass are comparable to or greater
43 than that of copepods (Bernard & Froneman, 2005; Hunt et al., 2008). Pteropods are able to form
44 locally dense aggregations in the water column (Percy & Fife, 1985). *Limacina helicina* is the
45 main food of many zooplankton organisms and predators of higher trophic levels, such as fish,
46 whales, and birds (Hunt et al., 2008), and play a key role in the food web and in
47 biosedimentation (Gilmer & Harbison, 1986; Noji et al., 1997; Bernard & Froneman, 2009;
48 Manno et al., 2009).

49 The body of *Limacina helicina* is covered by a fragile calcium carbonate shell that protects
50 them from predation. The aragonite composition of the shell makes these animals extremely
51 sensitive to ocean acidification, which is expected to increase due to anthropogenic CO₂
52 emissions into the atmosphere (Teniswood et al., 2016). Consequently, this species represents a
53 good model organism for ecological, physiological and biogeographical for how climate change
54 is affecting the Arctic Ocean (Comeau et al., 2009; Lischka et al., 2011).

55 Despite the important role of pteropods in Arctic ecosystems, little is known about the
56 genetic structure of *Limacina helicina* populations. A high diversity of haplotypes was found in
57 local populations from the fjords of Svalbard (Sromek et al., 2015), including haplotypes typical
58 of these pteropods in the Pacific Ocean (Shimizu et al., 2017). However, studies on the genetic
59 structure of *Limacina helicina* have not been carried out in the Siberian Arctic seas.

60 In the Kara Sea, pteropods are a common component of the pelagic community and their
61 spatial distribution is patchy (Arashkevich et al., 2010; Flint et al., 2015) similar to the other
62 areas (Percy & Fife, 1985). Within patches, their abundance reached one million ind. m⁻², and
63 they are the dominant consumers of suspended matter and phytoplankton (Drits et al., 2015).

64 The Kara Sea is a typical shelf Siberian Arctic Sea: the warm, salty water from the Barents
65 Sea enters from the south into the Kara Sea, and the cold Arctic water penetrates from the north
66 (Zatsepin et al., 2015). We expect that the genetic structure of the populations of *Limacina* in the
67 Kara Sea is similar to that in the Svalbard region, which is also influenced by the Barents Sea
68 and Arctic basin waters (Stiansen & Filin, 2007). In the Kara Sea, however, the effect of waters
69 of different origin, combined with the impact of a strong river run-off, creates a mosaic of
70 biotopes, where the genetic structure of populations can be different. We tested these hypotheses

71 by examining intraspecific diversity of *Limacina helicina* in the Kara Sea using a fragment of the
72 mitochondrial gene COI.

73 **Materials & Methods**

74 *Limacina helicina* were selected from zooplankton samples collected during the cruise #63
75 of the RV Akademik Mstislav Keldysh in the Kara Sea that took place September-October 2015.
76 Samples were collected at three different locations: station 5265 in the south of the Kara Sea and
77 two stations in the Voronin and St. Anna troughs, 5239 and 5212, respectively (Fig. 1, Table 1).
78 In the top 20 m of the water column at the station AMK 5265 temperature was 6 °C and salinity
79 was 31.4. At station AMK 5239 temperature was 1.2 °C and salinity was 30 (this station affected
80 by freshwater runoff and melting ice), while at station AMK 5212 in the St. Anna trough was 4.3
81 °C and salinity was 34.3.

82 Pteropods were preserved in 96% ethanol immediately after collection. DNA was isolated
83 from a piece of the pteropodia of large individuals (1-7 mm) or from the whole animal in case of
84 small individuals (0.1-0.7 mm) using the ExtraGene™ DNA Prep 100 kit (Isogen, Moscow,
85 Russia) per the manufacturer's protocol.

86 Fragments of mitochondrial cytochrome oxidase subunit gene (COI) were amplified using
87 Encyclo Plus PCR kit (Eurogen, Moscow, Russia) using two standard primers: LCO-1490 (5'-
88 GTCAACAAATCATAAAGATATTGG-3') and HCO-2198 (5'-
89 TAAACTTCAGGGTGACCAAAAATCA-3') (Folmer et al., 1994). PCR was conducted using
90 the following common PCR cycle settings: 5 min at 95°C, 40 cycles of 95°C for 30 s, followed
91 by annealing at 48°C for 45 s, 72°C for 1 min, and then a final elongation at 72°C for 5 min.
92 PCR products were analyzed with a 1% agarose gel electrophoresis, purified and sequenced
93 using Applied Biosystems® 3500 Genetic Analyzer. Subsequently sequences were aligned and
94 analyzed using MEGA 6.0 (Tamura et al., 2013). The 503 bp fragments of COI gene were used
95 for comparison with all other *L. helicina* samples from the Arctic and Pacific available from the
96 GenBank database (Table 1). Low quality contigs (contigs containing more than 3 Ns) were
97 excluded from analysis. The software Popart 1.7 (Leigh & Bryant, 2015) was then used for
98 comparative analysis and identification of differences between populations as well as for
99 construction of a TCS haplotype network (Clement et al., 2002). Furthermore, the program
100 DnaSP (Rozas et al., 2017) was used for an estimation of genetic diversity in populations.

101 Finally, the Arlequin 3.5 (Excoifer & Lischer,2010) software was used for AMOVA (Analysis of
102 Molecular Variance) analysis and verification of neutrality.

103 **Results**

104 We analyzed 73 specimens of *L. helicina* from the Kara Sea. COI sequences were obtained
105 from 65 of these samples, and 8 sequences were discarded due to poor quality contigs, leaving 57
106 individual sequences for further analysis. The highest Phi-st value found between the southern
107 and northern parts was 0.023 ($p < 0.5$). Due to the lack of significant differences in genetic
108 structure between the three different Kara Sea collection locations, the data from these stations
109 were combined for comparison with Svalbard and Pacific populations. In total, 179 *L. helicina*
110 sequences from the Arctic and Pacific were downloaded from GenBank (Table 1). These
111 sequences were regarded as three large geographical subgroups: the Kara Sea, Svalbard (data
112 from Sromek et al., 2015), and the Pacific (Jennings et al., 2010; Chichvarkin, 2016; Shimizu et
113 al., 2017). We also added the data from the Canadian Arctic (Hunt et al., 2010; Jennings et al.,
114 2010; Layton, Martel & Hebert, 2014) for the haplotype network construction. A total of 65
115 haplotypes was found from all sequences, which were combined in two large haplogroups, which
116 differ from each other by 2 nucleotide substitutions (Fig. 2B). Each haplogroup represents a
117 typical star-like haplonet with numerous branches. These patterns are in agreement with analysis
118 of Shimizu et al (2017) and so we adopted their names of haplogroups as H1 and H2.
119 Haplogroup H2 includes majority of sequences from the Kara Sea and Svalbard, while the H1 is
120 widespread at all research locations and found throughout all *L. helicina* populations (Table 1).

121 The samples from Kara Sea were represented by 19 haplotypes with two being widespread
122 (Fig. 2A). The remaining haplotypes are structured by their variations, differing by 1-3
123 nucleotide substitutions. The majority of the Kara Sea individuals are represented by the H2
124 haplogroup (79%). The greatest variability of haplotypes was found at the St. Anna Trough in
125 the north of the sea. The H2 haplogroup also the predominant haplogroup found in samples from
126 Svalbard fjords (84%), while only H1 haplogroups were found in the Pacific region (Fig. 2B).
127 The highest haplotype variety (Table 2) was reported around Svalbard ($H = 0.771$). The variety
128 of haplotypes in the Kara Sea is similar ($H = 0.672$), despite a smaller number of analyzed
129 individuals. The variability of haplotypes in Pacific is significantly lower ($H = 0.449$) as well as

130 nucleotide diversity (Table 2). The Tajima's D and Fu's Fs neutral evolution model tests showed
131 significant negative values (Table 2).

132 The haplotype network (Fig. 2B) shows similarity between the Kara Sea and Svalbard
133 populations. The ratio between the H2 and the H1 haplogroups in this region is also similar – the
134 majority of individuals belongs to the H2 haplogroup. All individuals from Canadian Arctic and
135 Pacific share the H1 haplogroup.

136 Pairwise comparison of Phi-st showed no significant differences between the Kara Sea and
137 the Svalbard populations ($\Phi\text{-st} = -0.00109$, $p = 0.4775$), however the samples from the Kara
138 Sea and Svalbard differed significantly from the Pacific (Table 3).

139 Discussion

140 The obtained results support our hypothesis that *L. helicina* populations from the Kara Sea
141 would be genetically similar to those near Svalbard.

142 The haplotype network is very similar for populations from the Kara Sea and those from
143 the North of Atlantic near Svalbard (Fig. 2B), and the ratio of haplotype group H2 haplotype
144 group H1 is also similar. The majority of haplotypes belongs to haplotype group H2, a minor part
145 to the group H1. In contrast, in the Canadian Arctic and the Pacific only haplogroup H1 is found.
146 The H1 group of haplotypes is widespread and occurs at all stations and populations (Table 1,
147 Fig. 3), and populations from the Pacific Ocean and the Canadian Arctic were almost identical
148 and were represented by the same sequence. This is explained by the main currents through the
149 Bering Strait and indicates the possible direction of distribution of plankton communities from
150 the Pacific Ocean (Nelson et al., 2009; Questel et al., 2016). Typical star-like haplonet and the
151 conducted Tajima's D and Fu's Fs tests can point to the rapid population expansion. The negative
152 values of Fu's Fs indicate the presence of a large number of low frequency haplotypes, usually
153 described for loci under directed selection or expansion of the population after a severe decline
154 (however, see Niwa et al., 2016, for alternative explanation of negative D and F in abundant
155 marine organisms). The reason for this pattern could be the rapid expansion of this species after
156 the last glacial maximum. Similar dispersal was observed for other Arctic species that have
157 survived in the refugia, then quickly spread to their current habitats after the deglaciation
158 (Hewitt, 2000; Weydmann et al., 2017).

159 According to the previous studies (Sromek, Lasota, & Wolowicz, 2015; Shimizu et al.,
160 2017), *Limacina helicina* were formerly widely distributed in the Arctic and the Pacific, but the
161 populations were isolated in the Northern Atlantic during the glaciation. Haplotype group H1
162 may have persisted in Pacific refuge, and H2 - in Atlantic. Subsequently, during the retreat of the
163 glacier about 131 ky BP, there was an increase in genetic diversity and distribution around
164 Svalbard. A similar spread of Pacific fauna was shown for other groups of organisms in the
165 Atlantic region (Laakkonen et al., 2013). The recent distribution of *L. helicina* haplotypes could
166 be explained in a similar way. When the ice sheets disappeared between the Pacific and Atlantic,
167 the Pacific population could have resettled in the Arctic. This hypothesis is supported by the
168 existence of a separated haplotype H1 along with haplotype H2 (Fig. 3). The northward currents
169 from the Pacific to the Arctic region are currently passing through the Bering Strait (Palumbi &
170 Kessing, 1991; Questel et al., 2016), which is reflected by the wide distribution of the haplotypes
171 H1 throughout the Arctic and by the absence of haplogroup H2 south of the Bering Strait in the
172 Pacific Ocean.

173 The absence of significant differences between the Kara Sea and Svalbard and the similarity
174 of proportions of different haplotypes in these regions is consistent with an ongoing or recent
175 exchange between these two populations, which coincides with the oceanography in this area
176 (Stiansen & Filin, 2007).

177 Frequency of occurrence of different haplotypes varies between locations of the Kara Sea
178 (near the Kara Strait, the St. Anna and Voronin Troughs), but these differences are not
179 significant. In the south (station AMK 5265 near the Kara Strait), the percentage H1 haplotype
180 (14%) is lower than in the north at St. Anna or Voronin Troughs (st. AMK 5239, 5212) (26%).
181 This is in accordance with the penetration of water of different origins into the sea: in the south-
182 west at station AMK 5265, the warm and salty water of the Barents Sea origin penetrates through
183 the Kara Strait, while the northern part of the Kara Sea is strongly influenced by the Arctic saline
184 and cold water (Zatsepin et al. 2015). Since these populations were not significantly different
185 genetically, the different environments are not isolating either population.

186 Conclusions

187 This study represents the first research on the genetic structure of *Limacina helicina* in the Kara
188 Sea. The comparison of our own data from the Kara Sea with the published data obtained in the

189 Svalbard area, northwest Pacific, and Canadian Arctic, allowed us to conclude that the
190 distribution of haplotypes in the Kara Sea is similar to that in Svalbard. Although no significant
191 differences between habitats within the Kara Sea were found, the proportion of haplotypes H2
192 was higher near the Kara Strait than in the northern troughs. The analysis of the available data
193 provides insight into the population structure of this pteropod species, indicating possible
194 direction of post-glacial distribution of *Limacina helicina* in the Arctic. However, many
195 questions regarding the genetics of this mollusk in the Arctic still remain unresolved, and in
196 future studies we hope to better understand how far the western population of *Limacina helicina*
197 penetrates and how the haplotypes are distributed over other Arctic seas.

198 Acknowledgements

199 For the help in the collection of *Limacina helicina* and discussion of this project we would like to
200 thank A.V. Drits, as well as V.V. Aleshin and C. Gross for extremely helpful comments on this
201 manuscript. We are particularly grateful for the laboratory assistance from the 2017 Invertebrate
202 Zoology bachelor students (Lomonosov Moscow State University). In addition, we gratefully
203 acknowledge the native English speaker Elizabeth Schmidt for constructive comments of the
204 manuscript. We thank Pieternella Luttkhuizen, Katharina Jörger and an anonymous reviewer for
205 important and valuable comments that significantly improved the paper.

206 References

- 207 1. Arashkevich EG, Flint MV, Nikishina AB, Pasternak AF, Timonin AG, Vasilieva JV,
208 Mosharov SA, Soloviev KA. 2010. The role of zooplankton in the transformation of the
209 organic matter in the Ob estuary, on the shelf, and in the deep regions of the Kara Sea.
210 *Oceanology*. 50(5):780-792.
- 211 2. Bernard KS, Froneman PW. 2005. Trophodynamics of selected mesozooplankton in the
212 west-Indian sector of the Polar Frontal Zone, Southern Ocean. *Polar Biology* 28(8):594-
213 606.
- 214 3. Bernard KS, Froneman PW. 2009. The sub-Antarctic euthecosome pteropod, *Limacina*
215 *retroversa*: Distribution patterns and trophic role. *Deep Sea Research Part I:*
216 *Oceanographic Research Papers* 56(4):582-598.

- 217 4. Chichvarkhin A. 2016. Shallow water sea slugs (Gastropoda: Heterobranchia) from the
218 northwestern coast of the Sea of Japan, north of Peter the Great Bay, Russia. *PeerJ*. 4,
219 e2774-e2774.
- 220 5. Clement M, Snell Q, Walker P, Posada D, Crandall K. 2002. TCS: Estimating gene
221 genealogies. Parallel and Distributed Processing Symposium, International Proceedings.
222 2: 184.
- 223 6. Comeau S, Gorsky G, Jeffree R, Teyssié JL, Gattuso JP. 2009. Impact of ocean
224 acidification on a key arctic- pelagic mollusc (*Limacina helicina*). *Biogeosciences*.
225 6:1877–1882.
- 226 7. Drits AV, Arashkevich EG, Nikishina AB, Sergeeva VM, Solovyev KA, Flint MV. 2015.
227 Mesozooplankton grazing impact on phytoplankton in the northern regions of the Kara
228 Sea in autumn. *Oceanology*. 55(4):595-605.
- 229 8. Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: A new series of programs to
230 perform population genetics analyses under Linux and Windows. *Molecular Ecology*
231 *Resources*. 10:564–567.
- 232 9. Flint MV, Poyarkov SG, Timonin AG, Soloviev KA. 2015. The structure of the
233 mesoplankton community in the area of the continental slope of the St. Anna trough
234 (Kara Sea). *Oceanology*. 55(4):583-594.
- 235 10. Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for
236 amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan
237 invertebrates. *Molecular Marine Biology and Biotechnology*. 3(5):294-299.
- 238 11. Gilmer RW, Harbison GR. 1986. Morphology and field behavior of pteropod molluscs:
239 feeding methods in the families Cavoliniidae, Limacinidae and Peraclididae (Gastropoda:
240 Thecosomata). *Marine Biology*. 91(1):47-57.
- 241 12. Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature*, 405(6789):907-
242 913.
- 243 13. Hunt B, Strugnell J, Bednarsek N, Linse K, Nelson RJ, Pakhomov E, Würzberg L. 2010.
244 Poles apart: the “bipolar” pteropod species *Limacina helicina* is genetically distinct
245 between the Arctic and Antarctic oceans. *PLoS One*. 5(3):e9835.

- 246 14. Hunt BPV, Pakhomov EA, Hosie GW, Siegel V, Ward P, Bernard K. 2008. Pteropods in
247 Southern Ocean ecosystems. *Prog Oceanogr.* 78:193–221.
- 248 15. Jennings RM, Bucklin A, Ossenbrügger H, Hopcroft RR. 2010. Species diversity of
249 planktonic gastropods (Pteropoda and Heteropoda) from six ocean regions based on DNA
250 barcode analysis. *Deep Sea Research Part II: Topical Studies in Oceanography.* 57(24):
251 2199-2210.
- 252 16. Laakkonen HM, Lajus DL, Strelkov P, Väinölä R. 2013. Phylogeography of amphi-
253 boreal fish: tracing the history of the Pacific herring *Clupea pallasii* in North-East
254 European seas. *BMC Evolutionary biology.* 13(1):67.
- 255 17. Layton KK, Martel AL, Hebert PD. 2014. Patterns of DNA barcode variation in
256 Canadian marine molluscs. *PLoS One.* 9(4):e95003.
- 257 18. Leigh JW, Bryant D. 2015. popart: full-feature software for haplotype network
258 construction. *Methods in Ecology and Evolution.* 6(9):1110-1116.
- 259 19. Lischka S, Büdenbender J, Boxhammer T, Riebesell U. 2011. Impact of ocean
260 acidification and elevated temperatures on early juveniles of the polar shelled pteropod
261 *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences.*
262 8(4):919.
- 263 20. Manno C, Tirelli V, Accornero A, Fonda Umani S. 2009. Importance of the contribution
264 of *Limacina helicina* faecal pellets to the carbon pump in Terra Nova Bay (Antarctica).
265 *Journal of Plankton Research.* 32(2):145-152.
- 266 21. Nelson RJ, Carmack EC, McLaughlin FA, Cooper GA. 2009. Penetration of Pacific
267 zooplankton into the western Arctic Ocean tracked with molecular population genetics.
268 *Marine Ecology Progress Series.* 381:129-138.
- 269 22. Niwa, H. S., Nashida, K., & Yanagimoto, T. 2016. Reproductive skew in Japanese
270 sardine inferred from DNA sequences. *ICES Journal of Marine Science,* 73(9), 2181-
271 2189.
- 272 23. Noji TT, Bathmann UV, Bodungen BV, Voss M, Antia A, Krumbholz M, Klein B,
273 Peeken I, Noji CIM, Rey F. 1997. Clearance of picoplankton-sized particles and

- 274 formation of rapidly sinking aggregates by the pteropod, *Limacina reiroversa*. *Journal of*
275 *Plankton Research*. 19(7):863-875.
- 276 24. Palumbi SR, Kessing BD. 1991. Population biology of the trans-arctic exchange: MtDNA
277 sequence similarity between Pacific and Atlantic sea urchins. *Evolution*. 45(8):1790-
278 1805.
- 279 25. Percy JA, Fife FJ. 1985. Energy distribution in an Arctic coastal macrozooplankton
280 community. *Arctic*. 38:39-42.
- 281 26. Questel JM, Blanco-Bercial L, Hopcroft RR, Bucklin A. 2016. Phylogeography and
282 connectivity of the Pseudocalanus (Copepoda: Calanoida) species complex in the eastern
283 North Pacific and the Pacific Arctic Region. *Journal of plankton research*. 38(3):610-
284 623.
- 285 27. Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-
286 Onsins SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA Sequence Polymorphism Analysis
287 of Large Datasets. *Mol. Biol. Evol.* 34:3299-3302. DOI: 10.1093/molbev/msx248.
- 288 28. Schlitzer, R. 2018. Ocean Data View, odv. awi. de.
- 289 29. Shimizu K, Kimoto K, Noshita K, Wakita M, Fujiki T, Sasaki T. 2017. Phylogeography
290 of the pelagic snail *Limacina helicina* (Gastropoda: Thecosomata) in the subarctic
291 western North Pacific. *Journal of Molluscan Studies*. 1-8.
- 292 30. Sromek L, Lasota R, Wolowicz M. 2015. Impact of glaciations on genetic diversity of
293 pelagic mollusks: Antarctic *Limacina antarctica* and Arctic *Limacina helicina*. *Marine*
294 *Ecology Progress Series*. 525:143-152.
- 295 31. Stiansen JE, Filin A. 2007. Joint PINRO/IMR report on the state of the Barents Sea
296 ecosystem in 2006 with expected situation and considerations for management.
- 297 32. Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S. 2013. MEGA6: molecular
298 evolutionary genetics analysis version 6.0. *Molecular biology and evolution*.
299 30(12):2725-2729.

- 300 33. Tenniswood CM, Roberts D, Howard WR, Bray SG, Bradby JE. 2016. Microstructural
301 shell strength of the Subantarctic pteropod *Limacina helicina antarctica*. *Polar Biology*.
302 39(9):1643-1652.
- 303 34. Weydmann A, Przyłucka A, Lubośny M, Walczyńska KS, Serrão EA, Pearson GA,
304 Burzyński A. 2017. Postglacial expansion of the Arctic keystone copepod *Calanus*
305 *glacialis*. *Marine Biodiversity*. 1-9.
- 306 35. Zatsëpin AG, Poyarkov SG, Kremenetskiy VV, Nedospasov AA, Shchuka SA, Baranov
307 VI, Kondrashov AA, Korzh AO. 2015. Hydrophysical features of deep water troughs in
308 the western Kara Sea. *Oceanology*. 55(4):472-484.

Figure 1

Location of the stations in the Kara Sea where *L. helicina* were collected.

Schlitzer, R., Ocean Data View, <https://odv.awi.de>, 2018.




Figure 2

TCS network of *Limacina helicina* haplotypes.

(A) haplotypes from Kara Sea (this study). St. Anna trough is marked in light green, Voronin through - in black, and southern part of Kara Sea - in dark green. **(B)** haplotypes across Northern hemisphere based on the current research and the GenBank data. Svalbard population is marked in dark blue, the Kara Sea in green, Pacific in orange, and the Canadian Arctic in blue. The H2 haplogroup is shown on the left and the H1 haplogroup on the right.

Notes: each haplotype is colored according to the location where it was collected. Haplotype circle sizes indicate frequency (according to the Table 1).




Figure 3

Haplotype distribution among the Arctic populations of *Limacina helicina*.

Orange - haplogroup H1, Blue - haplogroup H2. Schlitzer, R., Ocean Data View, <https://odv.awi.de>, 2018.




Table 1 (on next page)

Geographical location of *Limacina helicina* samples and compositions of haplotypes.

N - number of analyzed individuals, H1 and H2 - haplogroups.

Location	Coordinates/ References		N	Haplogroup	N from haplogroup	GenBank accession number
	N°	E°				
Kara Sea, South St. 5265	70°53	58°18	22	H1	3	MH379290 - MH379311
				H2	19	
Kara Sea, Voronina Trough, St. 5239	78°36	88°04	12	H1	2	MH379312 - MH379330
				H2	10	
Kara Sea, St. Anna Trough, St. 5212	76°43	70°59	23	H1	7	MH379266 - MH379289
				H2	16	
Svalbard	Sromek et al., 2015		68	H1	11	AB859527 - AB859593
				H2	57	
Pacific Ocean	Jennings et al., 2010 Chichvarkin, 2016 Shimizu et al., 2017		105	H1	105	FJ876923, KX871888, KX871889, LC185015 - LC185073, LC229727 - LC229769
				H2	0	
Canadian Arctic	Hunt et al., 2010 Layton et al., 2014 Jennings et al., 2010		6	H1	6	GQ861826 - GQ861828, HM862494, HM862496, FJ876924
				H2	0	

Table 2 (on next page)

Estimates of genetic diversity in populations of *Limacina helicina* from regions of Arctic and Pacific Ocean. Nucleotide and haplotype diversity, neutrality test.

N: number of individuals; **Ns**: number of sites; **k**: number of haplotypes; **S**: polymorphic sites; **H**: haplotype diversity; **π** : nucleotide diversity; **(Π)**: average number of nucleotide differences; **D**: Tajima's D; **Fs**: Fu's Fs neutrality test.

1

	N	Ns	k	S	H	π	Π	D	F_s
Kara Sea	57	500	19	26	0,672	0,00301754	1,509	-2,35896 (p < 0.001)	-17,725 (p < 0.0001)
Svalbard	68	503	24	25	0,771	0,00338705	1,704	-2,10848 (p < 0.01)	-24,253 (p < 0.0001)
Pacific	105	503	26	26	0,449	0,00124309	0,625	-2,60329 (p < 0.001)	-42,81 (p < 0.0001)

2

Table 3 (on next page)

Pairwise F_{st} values and associated p -values among *Limacina helicina* populations from the three different geographical areas.

1

Compared areas	Φ_{st}	p-value
Within Kara Sea		
St Anna – Voronin	0.01574	0.21622
St Anna – South	0.02292	0.15315
Voronin – South	-0.00263	0.42342
Between different seas		
Kara Sea – Svalbard	-0.00109	0.47748
Kara Sea – Pacific	0.63422	0.00000
Svalbard – Pacific	0.60013	0.00000

2