

A peer-reviewed version of this preprint was published in PeerJ on 25 March 2019.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.6648) (peerj.com/articles/6648), which is the preferred citable publication unless you specifically need to cite this preprint.

Liao H, Du T, Zhang Y, Shi L, Huai X, Zhou C, Deng J. 2019. Capacity for heat absorption by the wings of the butterfly *Tirumala limniace* (Cramer) PeerJ 7:e6648 <https://doi.org/10.7717/peerj.6648>

Capacity for heat absorption by the wings of the butterfly *Tirumala limniace* (Cramer)

Huaijian Liao^{1,2,3}, Ting Du^{1,2}, Yuqi Zhang⁴, Lei Shi^{1,2}, Xiyu Huai¹, Chengli Zhou^{1,2}, Jiang Deng¹

¹ Research Institute of Resources Insects, Chinese Academy of Forestry, Kunming, 650224, China

² Key Laboratory of Cultivating and Utilization of Resources Insects of State Forestry Administration, Kunming, 650224, China

³ Institute of Leisure Agriculture, Jiangsu Academy of Agricultural Sciences, Nanjing, 210014, Jiangsu, China

⁴ College of Life Science, Southwest Forestry University, Kunming, 650224, China

Corresponding Authors: Huaijian Liao, Lei Shi

Email address: huaixiyu_08@126.com, leishi@139.com

ABSTRACT

Butterflies can directly absorb heat from the sun via their wings to facilitate autonomous flight. However, how is the heat absorbed by the butterfly from sunlight stored and transmitted in the wing? The answer to this scientific question remains unclear. The butterfly *Tirumala limniace* (Cramer) is a typical heat absorption insect, and its wing surface color is only composed of light and dark color. Thus, in this study, we measured a number of wing traits relevant for heat absorption including the thoracic temperature at different light intensities and wing opening angles, the thoracic temperature of butterflies with only one right fore wing or one right hind wing, the spectral reflectance of the wing surfaces, the thoracic temperature of butterflies with the scales removed or not in light or dark areas, and the real-time changes in heat absorption by the wing surfaces with temperature. High intensity light (600–60000 lx) allowed the butterflies to absorb more heat and 60–90° was the optimal angle for heat absorption. The heat absorption capacity was stronger in the fore wings than the hind wings. Dark areas on the wing surfaces were heat absorption areas. The dark areas in the lower region of the fore wing surface and the inside region of the hind wing surface were heat storage areas. Heat was transferred from the heat storage areas to the wing base through the veins near the heat storage areas of the fore and hind wings.

INTRODUCTION

Sunlight is the most important source of energy and it supports the survival and reproduction of most creatures on Earth. Butterflies can directly utilize the heat from the sun to facilitate their autonomous flight. Previous studies have shown that butterflies must directly absorb the heat from sunlight to increase their thoracic temperature above that of their surroundings in order to allow autonomous flight to occur (*Kemp & Krockenberger, 2004; Barton et al., 2014; Bonebrake et al., 2014; Kleckova et al., 2014; Mattila 2015; Liao et al., 2017*). Autonomous flight is required for the reproductive activities of butterflies, including dispersal, courtship behavior, mating behavior, and oviposition behavior (*Shreeve, 1992; Bennett et al., 2012*). Therefore, heat absorption is an indispensable prerequisite for the reproductive success of butterflies (*Gibbs et al., 2010; Westerman et al., 2014; Gibbs et al., 2018*).

Butterflies primarily absorb heat via their wings (*Shanks et al., 2015; Han et al., 2016; Niu et al., 2016*). When adult butterflies require heat for autonomous flight, they adjust their body posture and either fully expand their wings or angle them to allow the optimal absorption of sunlight, where the heat absorbed is then transferred to the thoracic muscles to enable flight (*Heinrich, 1990; Huey et al., 2003*). Some studies indicate that the angle of the incoming sunlight and the wing opening angle can both significantly influence the rate and amount of heat absorption in butterflies (*Plattner, 2004; Shanks et al., 2015; Niu et al., 2016*). Indeed, heat absorption is important, but it has to be modulated via behavior to avoid overheating (*Kingsolver, 1985*).

Based on a study of a new mechanism of behavioral thermoregulation, Kingsolver (1985) suggested that *Pieris* butterflies use their wings as solar reflectors to reflect solar radiation onto the body and to increase their body temperature. Some studies also indicate that butterflies rely on “photonic crystal” structures in their wings to absorb heat (*Li et al., 2004; Han et al., 2013; Luohong, 2014; Han et al., 2015a*). Therefore, the transfer of heat in a butterfly not only depends on reflection from the wing surface, but it also is due to the internal tissue in the wing. Thus, what is the specific tissue involved with heat transfer in the wing? The answer to this question remains unclear. Previous studies of solar heat absorption by butterfly wings have focused on the forewings (*Devries, 2015; Wu et al., 2015*) whereas none have considered the hind wings. As a consequence, it is

not clear whether the fore wings or hind wings have a greater capacity for heat absorption.

Light intensity also significantly affects heat absorption by butterflies. In the butterfly *Heteronympha merope*, as the intensity of solar radiation increases, the thoracic temperature excess increases significantly regardless of wings being fully extended or closed (Barton *et al.*, 2014). Kingsolver (1983) found that short-term meteorological variations, such as intermittent cloudy periods, can rapidly reduce the body temperature excess and flight activity to zero in *Colias* butterflies. The light intensity also significantly influences the ability of the butterfly *Tirumala limniace* to absorb heat, where it was shown that the equilibrium thoracic temperature of adults was higher when exposed to 2240 lx compared with other light intensities and the time required to reach it was shorter. In addition, the thoracic temperature excess and rate of thoracic temperature increase were higher and achieved more quickly (Liao *et al.*, 2017), the sunlight intensity is about 2240 lx from 07:00am to 08:00am in summer in Yuanjiang county, Yunnan province, China.

Previous studies have shown that color significantly affects the absorption of heat in butterflies (Schmitz, 1994; Van Dyck *et al.*, 1997; Brashears *et al.*, 2016; Stuart-Fox *et al.*, 2017). Some studies indicate that more heat is absorbed when the wings of butterflies are darker, (Kingsolver, 1987; Schmitz, 1994; Berwaerts *et al.*, 2001; Trullas *et al.*, 2007). Some butterflies exhibit wing surface melanism to increase their ability to absorb heat (Guppy, 1986; Trullas *et al.*, 2007), but wing melanization is not directly functional to increasing heat absorption, as melanin is involved in numerous functions (Stuart-Fox *et al.*, 2017). Compared with the distal wing area, melanism on the wing surface near the body is more conducive to heat absorption by butterflies (Kingsolver, 1987; Brashears *et al.*, 2016). However, few studies have considered the specific wing surface areas that are used for heat absorption.

In this study, we aimed to understand the effects of the light intensity and wing opening angle on heat absorption, as well as clarifying the differences in the heat absorption capacity of the fore and hind wings, the distribution of heat absorption and non-heat absorption areas on the wing surfaces, and whether the wing has the ability to store heat temporarily and the location of the heat storage areas on the wing surfaces in order to identify the wing tissue that serves as the heat transfer channel. Elucidating the heat

absorption capacity of butterflies may provide a deeper understanding of the utilization of solar heat by insects.

Tirumala limniace (Cramer) is a typical butterfly that directly utilizes solar heat. In field, we have investigated that adults are highly flight active when on the sunny day, but on cloudy and rainy days, the butterflies do not fly even when the air temperature is high. The wings of the male and female butterflies have the same morphology and the wing surface only comprises black (black background) and light blue (light blue bands) areas (Chen *et al.*, 2008). According to the color depth, the wing surface can be divided into dark and light areas, while the wings of other butterflies in Yuanjiang County have more colors composition. Thus, the color composition of the wing surface is suitable for studying the heat absorption capacity of butterflies. At 243–2240 lx, high light intensity enabled adult butterflies to quickly absorb more heat than at low light intensities to elevate their thoracic temperature to above that of the environmental temperature and to take off earlier than at low light intensities (Liao *et al.*, 2017). However, the light intensity range considered in a previous study by Liao *et al.* (2017) was too narrow to determine the appropriate light intensity for heat absorption by the adult *T. limniace* butterfly. In China, *T. limniace* is mainly distributed in the dry hot valleys of south Yunnan Province, Guangxi Province, and Guangdong Province (Chen *et al.*, 2008). Yuanjiang County, Yunnan Province, China, which was the site of the artificial breeding ground, is also in the actual habitat range for this butterfly. In this study, according to the natural light intensity in Yuanjiang County, a light intensity range of 600–60000 lx was selected to study the heat absorption capacity of *T. limniace* butterflies in order to investigate the mechanism responsible for solar heat utilization by insects.

Thus, in the present study, we measured the absorption of heat by the butterfly *T. limniace* at different light intensities and wing opening angles in order to determine the most appropriate light intensity and wing opening angle for heat absorption. The thoracic temperature was measured in a butterfly with only one right fore wing or one right hind wing to elucidate the difference in the heat absorption capacity of the fore and hind wings. The spectral reflectance was measured for the wings to identify the distribution of the heat absorption and non-heat absorption areas on the fore and hind wings. In order to demonstrate that dark areas are responsible for heat absorption on butterfly wings, we investigated heat absorption in butterflies with the scales removed or present in light or

dark areas. To identify the wing surface areas involved with heat storage and transfer after heat absorption by butterfly wings, we monitored the real-time temperature changes on the wing surface during heat absorption in adult *T. limniace*. We also monitored the real-time temperature changes in the heat storage areas and non-heat storage areas, as well as the wing veins involved with heat transfer, to further determine the heat storage areas and the transfer channel on the wing surface. Based on the results, we obtained a clear understanding of the heat absorption capacity of the wings in the butterfly *T. limniace* to facilitate further research into the mechanism responsible for the utilization of solar heat by insects.

MATERIALS AND METHODS

Insects

T. limniace pupae were purchased from the Ornamental Insect Technology Development, Kunming Zhonglin Co. Ltd, Kunming, China. The pupae were placed on a cylindrical net (height: 65 cm, diameter: 50 cm) on top of a towel so they could readily spread their wings after emergence. Subsequently, the pupae were transferred to a room at $25 \pm 1^\circ\text{C}$ with a relative humidity of $75\% \pm 5\%$ under a 14:10 light:dark photoperiod. After the adults emerged, the females and males were reared separately in two different cages to prevent mating. Five-day-old unmated adults were used in this study. All the butterflies used in the experiment were living.

In China, *T. limniace* is mainly distributed in the dry hot valleys of south Yunnan Province, Guangxi Province, and Guangdong Province. Yuanjiang County, Yunnan Province, China, which was the artificial breeding ground, is also in their normal range habitat. This butterfly is important for ecological and environmental quality monitoring in south China. The optimal temperature range for survival and flight is $25\text{--}36^\circ\text{C}$. During the first and second day after emergence, adult *T. limniace* gradually unfurl their wings and exhibit low flight activity. Their flight activity increases significantly from 3 days and the peak flight activity occurs from 4 days to 7 days during the summer and autumn when the thermal absorption activity is also very high. Thus, unmated adults aged 5 days were used in this study.

Effects of light intensity and wing opening angle on the heat absorption capacity

We investigated the effects of the light intensity and wing opening angle on the heat absorption capacity of adult *T. limniace* butterflies. Yuanjiang County, Yunnan Province, China, is a suitable habitat for the butterfly *T. limniace*, and its population is mainly observed from July to October every year. Thus, the natural light intensity on September 11, 2017 in Yuanjiang County, Yunnan Province, China, was selected as the reference standard for the light intensity in this study because the skies were clear and cloudless on this day. Based on the natural light intensity levels between 07:00 to 20:00 h on September 11, 2017 (Figure 1), six light intensity levels were selected for this study, i.e., 600, 5000, 15,000, 25,000, 40,000, and 45,000 lx. The left and right wings form a specific angle when they are open where this angle ranges from 0° to 180°. Based on this angle range, 10 wing opening angles were tested in this study, i.e., 15°, 30°, 45°, 60°, 75°, 80°, 90°, 120°, 150°, and 180°. An orthogonal design was used in this experiment to study the effects of the light intensity and wing opening angle on heat absorption by the butterfly wings.

According to the specific angle required, a plane with the selected angle was cut with a knife along the middle line of a white plastic foam board (side length: 10 cm, thickness: 6 cm) and the adult *T. limniace* wings were fixed on this plane. Thus, a new unmated 5-day-old adult butterfly was fixed on the plane at a specific wing opening angle using double-sided adhesive tape. The plane with the affixed adult butterfly was placed in a dark room (length: 100 cm, width: 80 cm, height: 80 cm). The walls of the dark room were made of black-out cloth and the front wall of the dark room comprised a curtain made of awning cloth to facilitate the placement of butterflies. A 75 W halogen lamp (OSRAM-64841FLSP, Osram China Lighting Ltd, Shanghai, China) was hung 60 cm above the bottom of the dark room. The light intensity of the lamp was adjusted with a potentiometer (CHILFU, Shanghai Wenfu Electric Co. Ltd, Shanghai, China).

Before turning the lamp on, the butterfly fixed on the plane was maintained in the dark room for 15 min in order to stabilize its body temperature to avoid the influence of body temperature on heat absorption by the butterfly. After switching the lamp on, the thoracic

temperature was recorded every 10 s until the temperature was maintained at a set temperature for 90 s, thereby indicating that the thoracic temperature of the butterfly had reached equilibrium under the specific light intensity. The thoracic temperature was measured using a thermocouple thermometer (Thermometer GM1312, Shenzhen Jumaoyuan Science and Technology Co. Ltd, Shenzhen, China). A thermistor was inserted through the mesoscutellum and placed in or near the dorsal longitudinal muscles to record the thoracic temperature. During each run of the experiment, we recorded the equilibrated thoracic temperature, rate of thoracic temperature increase (Total change in temperature over time), ΔT as the difference in temperature between the thorax and the ambient temperature, and the time when the thoracic temperature reached equilibrium. Three female and three male adult butterflies were tested for each orthogonal treatment with the different light intensities and wing opening angles. In total, 180 female and 180 male butterflies were tested in this experiment.

Difference in the heat absorption capacity of the fore and hind wings

To determine the difference in the heat absorption capacity of the fore and hind wings, we measure the thoracic temperature in a butterfly with only one right fore wing or only one right hind wing under six light intensities, i.e., 600, 5000, 15,000, 25,000, 40,000 and 45,000 lx. The two hind wings and left fore wing were cut from a new unmated 5-day-old adult butterfly using surgical scissors so only the right fore wing remained (fore wing treatment). The two fore wings and left hind wing were cut from another new unmated 5-day-old adult butterfly using surgical scissors so only the right hind wing remained (hind wing treatment). The butterfly was then fixed on a horizontal plane (side length: 10 cm, thickness: 6 cm) without any angle, before it was placed and maintained in the dark room for 15 min so its body temperature could stabilize before turning the lamp on. After switching the lamp on, during each run of the experiment, we recorded the equilibrated thoracic temperature, the rate ($^{\circ}\text{C}/\text{min}$) of thoracic temperature increase, ΔT , and the time required to reach equilibrium. In this experiment, we tested six female and six male butterflies with only one right fore wing at each light intensity, and six female and six male butterflies with only one right hind wing at each light intensity. In total, we tested 36 female and 36 male butterflies in the fore wing treatment, and 36 female and 36 male butterflies in the hind wing treatment. We also measured the wing expanse for both the

216 fore and hind wings of the adult butterflies using 37 female and 34 male adult butterflies.

217 **Reflectance of light and dark areas**

218 To determine the distribution of the heat absorption and non-heat absorption areas on the
219 fore and hind wings, we measured the spectral reflectance from the wings of adult
220 butterflies. A higher spectral reflectance indicates weaker light absorption and heat
221 absorption, whereas lower spectral reflectance denotes stronger light absorption and heat
222 absorption. The wing of the *T. limniace* butterfly appears the same in both sexes, but they
223 differ because there is a scaly bag with a protruding ear in the middle of the Cu₂ chamber
224 in the male butterfly. The wing surface is black with many transparent azure markings
225 (Figure 2). Therefore, the wing surface can be divided into light areas and dark areas
226 according to the color shade on the wing surface.

227 The right fore and right hind wings were cut from a female butterfly and a male
228 butterfly. During the cutting process, the integrity of the wings was maintained as far as
229 possible and we prevented scale removal from the wings. The wings were then placed in
230 the dark room. The light intensity in the dark room was set to 5000 lx. The spectral
231 reflectance of the wings was measured using a spectrograph (SOC710 HS-Portable,
232 Surface Optics Corporation, San Diego, USA). As shown in Figure 2, the dorsal surfaces
233 of the fore and hind wings were divided into three parts comprising the border, middle,
234 and inside parts. The spectral reflectance was then calculated for the dark and light areas
235 in these three parts. The average spectral reflectance of all the light areas in each part was
236 taken as the spectral reflectance of the light areas in this part, and the average spectral
237 reflectance of all the dark areas in each part was taken as the spectral reflectance of the
238 dark areas in this part. In addition, the excised fore and hind wings were placed under a
239 VHX-1000 super-high magnification lens zoom 3D microscope (VHX-1000, Keyence
240 (China) Co. Ltd, Shanghai, China) to obtain images. The areas of the light areas, dark
241 areas, and the entire wing surface were measured with an X/Y measurement system under
242 the microscope. The ratios of the light areas and dark areas on the wing surfaces were
243 also determined.

244 **Heat absorption capacities of dark and light areas on the wings**

In order to demonstrate that the dark area was responsible for heat absorption via the butterfly wings, we investigated heat absorption in butterflies treated with four different scale removal treatments comprising the removal of scales from dark areas (DSR treatment), the removal of scales from light areas (LSR treatment), the removal of all scales from the wings (ASR treatment), and leaving the scales intact on the wings (IN treatment). Scales are the main tissues for heat absorption of butterfly wings. And the structure color of the wing surface is determined by the scale structure and arrangement. Thus, the presence or absence of scales can be used to determine the heat absorption ability of a certain area of wing surface. The scales were gently removed from the wing surfaces with a fine brush in each treatment. The treated butterflies were then placed in the dark room. The light intensity was 5000 lx. After switching the lamp on, during the course of each experiment, we recorded the equilibrated thoracic temperature, rate of thoracic temperature increase, ΔT , and the time required to reach equilibrium. Three female and three male butterflies were tested for each treatment. All of the butterflies used in this experiment were new and unmated 5-day-old adult *T. limniace*.

Heat storage and transfer in the wing

We monitored the real-time temperature changes on the wing surface in order to determine the wing surface areas involved with heat storage and transfer during heat absorption by the butterfly wings. The color and markings on the wing surfaces are the same in male and female butterflies, so we only measured the real-time temperature changes on the wing surfaces of male butterflies during heat absorption. A new unmated 5-day-old male butterfly was fixed to a square white plastic foam board (10 × 10 cm) using insect dissection needles. The wings were fully extended to facilitate real-time temperature monitoring on the wing surface. The butterfly was then placed in the dark room and the light intensity was set to 5,000 lx. The heat absorption rate under 5,000 lx was slower than that at light intensity levels of 15,000–25,000 lx, but it was too slow at 600 lx, so we selected 5,000 lx because it yielded a suitable heat absorption rate for real-time wing temperature monitoring. The real-time temperature changes on the wing surface were recorded using a thermal infrared imager (Flir A600s, FLIR Systems, Inc., Wilson, OR, USA) until the thoracic temperature reached equilibrium.

To determine if dark areas served as heat storage areas, we selected 6 points in the dark

area and 6 points in the light area and compared their real-time temperature differences. In addition, we selected evenly distributed points from border of wing surface to the wing base to identify heat transfer channels. The distributions of all points on the fore and hind wings are shown in Figure 3A. The distributions of the points on the wing vein on the wing surfaces are shown in Figure 3B. Five to seven points were selected for each heat transfer channel. The initial heat absorption rate of the areas as heat absorption is faster than that of the areas as heat storage. When the heat is transferred from the area as heat absorption to the heat storage area, the heat absorption rate of the storage area increases rapidly and exceeds that of the areas as heat absorption. Finally, the equilibrium temperature of the heat storage area is higher than that of the non-heat storage area.

Statistical analysis

The interaction effects of the light intensity and wing opening angle on heat absorption were analyzed using two-way ANOVA. We analyzed ΔT , the thoracic equilibrium temperature, rate of thoracic temperature increase, and the time required to reach equilibrium under different light intensities and wing opening angles simultaneously using one-way ANOVA. Duncan's multiple-range test was used to compare multiple means when significant differences were detected. Differences in ΔT , the thoracic equilibrium temperature, rate of thoracic temperature increase, the time required to reach equilibrium, and the wing expanse between the fore and hind wings were validated using the Student's *t*-test. The differences of spectral reflectance among different areas in male and female wings were analyzed by Kruskal-Wallis test. The differences in the areas of the light and dark wing surface areas were analyzed using chi-square test. Differences in ΔT , the thoracic equilibrium temperature, rate of thoracic temperature increase, and the time required to reach equilibrium in the SDR, SLR, SAL, and IN treatments were analyzed using one-way ANOVA. Duncan's multiple-range test was used to compare multiple means when significant differences were detected. The trends of real-time temperatures of the monitoring points in the dark areas, light areas, and wing veins were analyzed using time-series ExpDec2 of exponential models ($y = A1 \cdot \exp(-x/t1) + A2 \cdot \exp(-x/t2) + y0$). $A1$ and $A2$ is the pre-factor, and they can indicate an increasing trend of the y value; $t1$ and $t2$ are the relaxation time; $y0$ is the offset, it can reflect the final equilibrium temperature.

RESULTS

Interaction effects of light intensity and wing opening angle on heat absorption by adult butterflies

The wing opening angle and light intensity significantly affected the heat absorption capacity of adult *T. limniace* butterflies (Table 1). The equilibrium thoracic temperature, time required to reach equilibrium temperature, ΔT , and the rate of thoracic temperature increase were affected significantly by the wing opening angle and light intensity in both male and female butterflies (Table 1). The interaction effect of the light intensity and wing opening angle on the rate of thoracic temperature increase was significant in both male and female butterflies, but the interaction effects on the equilibrium thoracic temperature, time required to reach equilibrium, and ΔT were not significant (Table 1).

Effects of light intensity on the heat absorption capacity

The light intensity significantly affected the heat absorption capacity in adult *T. limniace* (Tables 2–5). In both the female and male butterflies, within the range of 600–60,000 lx, the equilibrium thoracic temperature became significantly higher as the light intensity increased under the same wing opening angle (Table 2). The time required to reach the equilibrium temperature became shorter as the light intensity increased (Table 3). ΔT became significantly higher as the light intensity increased (Table 4). The rate of thoracic temperature increase also became faster as the light intensity increased (Table 5).

However, we also found that when the light intensity was above 40,000 lx, the equilibrium temperature could rapidly exceed 70°C, which was much higher than the butterflies could tolerate, and thus exposure to a light intensity above 40,000 lx caused the death of the butterflies half an hour after the experiment finished. In summary, a higher light intensity could allow butterflies to absorb more heat (Tables 2–5).

Effects of wing opening angles on the heat absorption capacity

The wing opening angle also significantly influenced the heat absorption capacity of adult *T. limniace* (Tables 2–5). Compared with wing opening angles 15–45° and

120–180°, when male and female butterflies were at wing opening angles 60–90°, the equilibrium temperatures were higher (Table 2), the times required to reach the equilibrium temperature were much shorter (Table 3), the values of ΔT were much higher (Table 4), and the rates of thoracic temperature increase were faster (Table 5). However, the influence of wing opening angle was only seen at some light intensities, particularly the higher light intensities. Thus, the optimal wing opening angles for heat absorption ranged from 60° to 90°.

Heat absorption capacities of the fore wing and hind wing

The heat absorption capacities of both the fore wing and hind wing were significantly affected by the light intensity where the capacity of the fore wings was higher than that of the hind wings (Table 6). The Student's *t*-test was used to analyze the differences in the heat absorption capacities of the fore wing and hind wing in both female and male butterflies (Table 6). The results showed that the heat absorption capacity of the fore wing was much higher than that of the hind wing in both female and male butterflies (details see Table 6). The equilibrium temperatures and values of ΔT were positively higher for the fore wings than the hind wings under 600–40,000 lx (Table 6), while the times required to reach equilibrium temperature under 600–60,000 lx were significantly shorter in the fore wings than the hind wings, and the rates of thoracic temperature increase were also significantly faster in the fore wings than the hind wings (Table 6). In addition, we found that in both male and female butterflies, the expanse of the fore wing was significantly longer than that of the hind wing (male: $t_{66} = 24.717$, $P < 0.0001$; female: $t_{72} = 29.124$, $P < 0.0001$; Figure 4). In summary, within the range of 600–60,000 lx, the fore and hind wings could both absorb much more heat at a faster rate as the light intensity increased (Table 6). The heat absorption capacity of the fore wing was much higher than that of the hind wing (Table 6), and the expanse of the fore wing was also much larger than that of the hind wing (Figure 4).

Identification of heat absorption and non-heat absorption areas

The spectral reflectance of the wings indicated that the dark areas of the wings were used for heat absorption, whereas the light areas did not absorb heat (Figure 5). The spectral

reflectance values of the light areas on both the fore and hind wings were all much higher than those of the dark areas (Male fore wing: $Z=585.340$, $P<0.0001$; Male hind wing: $Z=589.787$, $P<0.0001$; Female fore wing: $Z=583.716$, $P<0.0001$; Female hind wing: $Z=574.677$, $P<0.0001$; Figure 5). The dark areas were significantly larger on both the fore and hind wings than the light areas (male fore wing: $X^2 = 20.841$, $P < 0.0001$; male hind wing: $X^2 = 10.562$, $P = 0.0012$; female fore wing: $X^2 = 16.237$, $P < 0.0001$; female hind wing: $X^2 = 7.242$, $P = 0.0071$). However, the light areas still accounted for about 20% of the fore wing area and about 29% of the hind wing area. In summary, the dark areas could be used to absorb heat, whereas the light areas may prevent the butterfly from absorbing too much heat.

Heat absorption capacities of dark and light areas on the wings

Scale removal treatment significantly impacted equilibrium thoracic temperature (Figure 6). Based on the multirange tests, that the effect was due to removal of the dark scales and all scales, while there was no difference in the response variables between intact butterflies and those with light scales removed. When the scales were removed from the light areas of the wings, the equilibrium thoracic temperature, ΔT , the time required to reach equilibrium temperature, and the rate of thoracic temperature increase were not significantly different from those in the butterflies without scale removal. However, their equilibrium thoracic temperatures (male: $F_{3,8} = 28.269$, $P < 0.0001$; female: $F_{3,8} = 21.132$, $P < 0.0001$) and the values of ΔT s (male: $F_{3,8} = 15.686$, $P < 0.0001$; female: $F_{3,8} = 29.17$, $P < 0.0001$) were significantly higher, and the time required to reach the equilibrium temperature (male: $F_{3,8} = 5.96$, $P = 0.0019$; female: $F_{3,8} = 29.170$, $P < 0.0001$) and the rates of thoracic temperature increase (male: $F_{3,8} = 34.839$, $P < 0.0001$; female: $F_{3,8} = 31.613$, $P < 0.0001$) were significantly shorter and faster, respectively, than those of butterflies with all scales removed from the wings or with scales removed from the dark areas (Figure 6). These results indicate that the dark areas of the wings are the areas where the butterflies absorb heat, whereas the light areas cannot absorb heat.

Heat storage and transfer in the wings

Real-time temperatures on the wing surface during heat absorption

The wing temperature increased significantly in *T. limniace* as the lighting time increased, and the temperatures of the wing base and thorax also increased (Figure 7). After the light was switched on, heat absorption by the wings began immediately. The thoracic temperature reached equilibrium after lighting for 480 s. The temperatures of the light areas on both the fore and hind wings were always lower than those of the dark areas during heat absorption in adult *T. limniace*.

As the lighting period continued, heat clearly started to accumulate in some areas of the wing surface and the temperature was significantly higher than that of other areas (Figure 7). The temperatures of the dark areas in the middle of wing cell 1A-Cu₂ on the hind wing were higher than those of the other wing areas after lighting for 240 s. The temperatures of the dark area in the mid-posterior region near the wing base were markedly higher in wing cells A-CU₃ and Cu₂-Cu₃ on the fore wing, and wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ on the hind wings compared with other wing areas during lighting from 300–540 s. These results indicate that the dark areas in the mid-posterior region near the wing base in wings cells A-CU₃ and Cu₂-Cu₃ on the fore wings, and wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ on the hind wings were the temporary heat storage areas on the wing surfaces (Figure 3 & Figure 7).

According to the real-time temperature changes on the wing surface, it appeared that heat was transferred from the heat storage areas to the wing base and thorax through veins Cu₂, Cu₃, Cu, and A on the fore wings, and veins 1A, Cu₂, Cu₁, Cu, M₁, M₃, M, R₂, and R on the hind wings during lighting for 0–540 s.

Real-time temperatures in the heat storage and non-storage areas on the wing surface during heat absorption

The temperature at each point on the fore and hind wing surfaces increased as the lighting period increased as the analysis of time-series ExpDec2 of exponential models (Figure 8 & Table 7). On the fore wing, the base of the wing (Sp7) heated at a conspicuously slow rate initially (A1=-7.40247), then heated at a higher rate than other points at later times (A2=-18.4286). The points in the dark area heated at a similar rate initially (Sp8: A1=-11.7911; Sp9: A1=-11.158; Sp10: A1=-10.0475; Sp11: A1=-11.7152; Sp12: A1=-11.173), then the points Sp8–Sp11 heated at a higher rate than the point Sp12 without heat storage (Sp8: A2=-10.7115; Sp9: A2=-11.5553; Sp10: A2=-11.7129; Sp11:

A2=-10.5312; Sp12: A2=-9.14257). The points in the light area always heated at a similar rate with the points in the dark area, but the temperatures were always lower than the points for heat storage (the values of y_0 s of the points Sp1–Sp6 was significantly lower than the points Sp8–Sp11: $t_8=2.864$, $P=0.021$). In summary, Sp8–Sp11 could temporarily store heat and then transfer it to Sp7, and the heat storage was greater closer to Sp7. Sp12 could absorb heat from the light source but it could not store the heat.

In the hind wing, the base of the wing (H-Sp7) heated at a conspicuously slow rate initially (A1=-6.07772), then heated at a higher rate than other points at later times (A2=-19.64). The points in the dark area heated at a similar rate initially (H-Sp8: A1=-10.7477; H-Sp9: A1=-13.0178; H-Sp10: A1=-10.9644; H-Sp11: A1=-12.9741; H-Sp12: A1=-13.2487), then the points for heat storage heated at a higher rate than the point Sp12 without heat storage (H-Sp8: A2=-13.9014; H-Sp9: A2=-10.1396; H-Sp10: A2=-11.233; H-Sp11: A2=-9.03949; H-Sp12: A2=-7.43888). We also found that the values of y_0 s of H-Sp1, H-Sp2 and H-Sp3 were similar with the y_0 s of the points H-Sp8–H-Sp11 ($t_5=20.683$, $P=0.525$); The possible reason is that these points were too close to the base of the wing and were affected by the heat of the base of the wing, but their y_0 values were still lower than the point H-Sp8 in the dark area close to them. The temperatures of other three points in the light area were always lower than the points H-Sp8–H-Sp11 (y_0 : $t_5=2.986$, $P=0.031$). In summary, H-Sp8–H-Sp11 could temporarily store heat and then transfer the heat to H-Sp7, and the heat storage was greater closer to H-Sp7. H-Sp12 could absorb heat from the light source, but it could not store the heat (Figure 8C–D).

Real-time temperatures in the heat transfer channel in the wing veins during heat absorption

Both in the fore and hind wings, the temperatures of all the veins increased as the lighting time increased as the analysis of time-series ExpDec2 of exponential models (Figure 9 & 10). The heat rate of the points in all veins failed to show a certain rule according to the distance from the base of the wing (Figure 9 & 10). But, in all vein channels, the closer points get to the base of the wing, the higher the final equilibrium temperature was (Figure 9 & 10), and the value of y_0 was higher (table 8). These indicated that with the heat transfer in the vein, the closer the points were to the wing base, the higher the

temperatures; conversely, the farther the points were from the wing base, the lower the temperatures (Figure 9 & 10). In thus, heat may mainly transfer along the veins from the points at the edge of the wing surface to the points at the base of the wing. The potential transfer channels were showed in Figure 3 and Figure 9.

DISCUSSION

Effects of the light intensity on heat absorption

We found that the light intensity significantly affected the heat absorption capacity of the butterfly *T. limniace*. Within the light intensity range from 600 lx to 60000 lx, as the light intensity increased, the thoracic equilibrium temperature and ΔT increased significantly, the time to reach the equilibrium temperature was significantly shorter, and the rate of thoracic temperature increase was significantly faster. Our previous studies also showed that compared with other light intensities (243, 860, and 1280 lx), the equilibrium thoracic temperature of adults exposed to 2240 lx was higher and the time required to reach it was shorter. In addition, the value of ΔT was higher and the rate of thoracic temperature increase was achieved more quickly (Liao et al., 2017). These results suggest that in an appropriate range, a high light intensity can allow adult *T. limniace* butterflies to rapidly absorb more heat to elevate their thoracic temperature compared with low light intensities. Similar results were obtained for the butterfly *Heteronympha merope* where its body temperature excess also increased as the level of solar radiation increased (Barton et al., 2014).

The heat absorbed by butterflies is used mainly for autonomous flight and it is directly involved with reproductive behavior (Shreeve, 1992; Kemp and Krockenberger, 2004; Bennett et al., 2012; Bonebrake et al., 2014; Kleckova et al., 2014; Mattila 2015; Shanks et al., 2015; Han et al., 2016; Niu et al., 2016). In *Bicyclus anynana*, active males increase their own likelihood of copulation and active females increase their likelihood of being courted (Westerman et al., 2014). The flight activity of *T. limniace* butterflies is higher under a high light intensity compared with those exposed to a low light intensity (Liao et al., 2017). The high thorax temperature recorded in *Junonia villida* is probably linked to its high flight speed (Nève and Hall, 2016). We found that a higher light intensity increased the heat absorbed by *T. limniace* adults from light sources. Thus,

within an appropriate light intensity range, a higher light intensity can allow *T. limniace* adults to absorb more heat to facilitate autonomous flight and increase the flight activity, thereby enhancing the likelihood of reproduction.

We also found that when the light intensity was above 40,000 lx, the equilibrium temperature could rapidly exceed 70°C, which was much higher than the butterflies could tolerate, and thus exposure to a light intensity above 40,000 lx caused the death of the butterflies half an hour after the experiment finished. In the summer and autumn season of Yuanjiang county, Yunan province, China, the peak of flight activity of butterfly *T. limniace* mainly appeared at 09:00-12:00 and 15:00-17:00 (Chen *et al.*, 2008). During these time periods, the sunlight intensity is lower than 40,000 lx, which allows the butterfly to absorb the right amount of heat for autonomous flight. More than 40,000 lx light intensity could make the butterflies to absorb more heat than the body can tolerate, burning the body and causing death. Some studies indicated that butterflies can adjust their basking posture to reduce heat absorption (Rawlins, 1980; Tsuji *et al.*, 1986), e.g. closing the wings and turning the body opposite the sunlight. In *T. limniace*, in the field we observed that when the sunlight was above 40,000 lx, the butterflies would fly into the woods and perch on branches to avoid the sunlight.

Effects of the wing opening angle on heat absorption

The wing opening angle also significantly influenced the heat absorption capacity of *T. limniace* adults. Some previous studies have shown that butterflies with fully open wings have the highest heat absorption capacity (Tsuji *et al.* 1986; Berwaerts *et al.*, 2001; De Keyser *et al.*, 2015). When the butterfly *Polyommatus icarus* fully opens its wings, the warming rates are maximized and its body temperature is also highest (De Keyser *et al.*, 2015). *Pararge aegeria* butterflies with fully opened wings are able to absorb more heat and cool faster compared with those with half-opened wings (Berwaerts *et al.*, 2001). Other studies suggest that the heat absorption capacity of butterflies is highest in the optimal range for the wing opening angle (Kingsolver and Moffat, 1982; Barton *et al.*, 2014). For example, in *Pieris*, normal and black base manipulation of butterflies can achieve maximum temperature excesses at wing angles of 30–40°, whereas the optimal wing angles for the highest temperature excess in black distal treatment butterflies are 60–90° (Kingsolver, 1987). Thus, regardless of whether the wings are normal or

melanized, heat absorption by butterflies requires the optimal wing opening angle. We found that the optimal wing opening angle range for heat absorption by *T. limniace* was 60–90°. In this range, the equilibrium temperatures were significantly higher, the times required to reach the equilibrium temperature were much shorter, the values of ΔT were higher, and the rates of thoracic temperature increase were significantly faster. Thus, an appropriate wing opening angle can enhance the heat absorption capacity of butterflies. In wild, when butterflies need heat to autonomous fly, adults *T. limniace* doesn't fully open their wings, however, they would make the wings maintained a certain angle. Thus, wing opening angles 60–90° were suitable for heat absorption of butterfly *T. limniace*.

Differences in heat absorption between the fore and hind wings

Our results showed that the heat absorption capacity of the fore wings was significantly higher than that of the hind wings. The equilibrium temperatures and values of ΔT s were significantly higher for the fore wings than the hind wings in males exposed to light at 600–40,000 lx and females exposed to light at 600 lx, 15000 lx and 40,000 lx. The times required to reach equilibrium temperature and the rates of thoracic temperature increase were significantly shorter and faster, respectively, in the fore wings than those in the hind wing for both males and females exposed to light at 600–60,000 lx. These results suggest that the fore wing is the major tissue used for heat absorption in butterflies. Thus, the wing size significantly limits the heat absorption capacity of butterflies (*Kammer and Bracchi, 1973; Berwaerts et al., 2001, 2002*). Some studies suggest that butterflies with larger wings can absorb more heat than those with smaller wings (*Heinrich, 1986; Kingsolver, 1987; Schmitz, 1994*). Thus, a study of 20 Australian butterflies showed that those with the highest wing loadings had the highest thorax temperature at take-off (*Nève and Hall, 2016*). In our study, the wing expanse of the fore wing was significantly larger than that of the hind wing in both male and female butterflies. Therefore, the fore wings allow *T. limniace* butterflies to absorb more heat than the hind wings under the same lighting conditions. However, in *Pararge aegeria*, butterflies with larger wings heated up more slowly compared with those with small wings (*Berwaerts et al., 2001*).

Identification of heat absorption and non-heat absorption areas

The spectral reflectance from light areas on both the fore and hind wings was much

higher than that from dark areas in *T. limniace* adults. Thus, the absorbance and thermal absorptivity of the dark areas was significantly greater than that of the light areas. Color can affect an animal's temperature because dark surfaces absorb more solar energy than light surfaces and the energy is converted into heat (Stuart-Fox *et al.*, 2017). Previous studies indicate that wing color can limit the heat absorption capacity, where darker coloration can absorb more heat from sunlight or other light sources than lighter coloration (Watt, 1968; Kingsolver, 1987, 1988; Berwaerts *et al.*, 2001). For example, darker *Pararge aegeria* butterflies have a much higher warming rate (Van Dyck *et al.*, 1997). We found that when the scales were removed from the light areas on the wings, the equilibrium thoracic temperature and value of ΔT s were significantly higher, and the time required to reach the equilibrium temperature and the rate of thoracic temperature increase were significantly shorter and faster, respectively, than those in butterflies with all the scales removed from the wings or with scales removed from the dark areas. Thus, the dark areas of the wings absorb heat in the butterfly *T. limniace* rather than the light areas.

Some studies suggest that melanization of the wing surfaces can promote the absorption of heat from sunlight in butterflies (Kingsolver and Wiernasz, 1991; Stoehr and Wojan, 2016). The butterfly *Parnassius phoebus* lives at high altitudes and high latitudes, and it uses melanization to obtain heat from solar radiation, where the color of its wings becomes darker as the altitude and latitude increase (Guppy, 1986). Melanization inside the wing surfaces can improve the heat absorption capacity of butterflies whereas melanization at the distal wing surfaces cannot affect heat absorption (Kammer and Bracchi, 1973; Kingsolver, 1987; Brashears *et al.*, 2016). Wasserthal (1975) showed that the most of the heat transferred from the wing to the body originates from the 15% of the wing surface nearest to the body in the butterflies *Papilio machaon*, *Apatura ilia*, and *Papilio troilus*. Thus, the dark areas of the wing surface near the body are the main heat absorption areas in butterflies. Our results are similar to those obtained in previous studies. In *T. limniace*, we found that the temperatures were markedly higher in the dark areas in the mid-posterior region near the wing base in wing cells A-CU₃ and Cu₂-Cu₃ in the fore wings, as well as in wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ in the hind wings compared with other wing areas during lighting for 300–540 s.

The color of a butterfly wing mainly comprises iridescence and structural colors

(Nijhout, 1991; Michielsen et al., 2010; Han et al., 2016; Siddique et al., 2017). The part of the wing with a structural color is used to absorb heat from the sun (Bosi et al., 2008). In *Pararge aegeria*, normal butterflies reach a higher equilibrium thoracic temperature than descaled butterflies, probably because the wings without scales absorb less radiation (Berwaerts et al., 2001). In the present study, we found that removing the scales from the dark areas of the wings significantly reduced the heat absorption capacity of butterflies, whereas removing the scales from the light areas did not affect normal heat absorption. Therefore, the dark areas on the dorsal wing surfaces in *T. limniace* may mainly comprise structural colors.

High reflectance can reduce the absorption of heat from sunlight (Stuart-Fox et al., 2017). Thus, light colors are not conducive to heat absorption by butterflies. We found that removing the scales from the light areas on the wing surfaces did not affect the absorption of heat by butterflies. In *Anartia fatima*, it was previously shown that blackening the white bands on the wing surface did not affect its equilibrium temperature, but adding white bands decreased the rate of heating (Brashears et al., 2016). Thus, the light areas on the wing surfaces may prevent butterflies from absorbing heat excessively quickly and increasing the body temperature to an intolerable level. We also found that light areas accounted for about 20% of the fore wing area and about 29% of the hind wing area. However, some studies indicate that light areas have high spectral reflectance and they could be used as information for intraspecies identification or for detecting the opposite sex in butterflies (Emmel, 1972; Taylor, 1973; Seymoure and Aiello, 2015). The role of the light areas on butterfly wings in heat absorption will be addressed in our future research.

Heat storage and transfer in the wing

In the butterfly *T. limniace*, the temperatures of the dark areas were markedly higher in the mid-posterior region near the wing base in wing cells A-Cu₃ and Cu₂-Cu₃ on the fore wings, as well as in wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ on the hind wings compared with the other wing areas during lighting for 300–540 s. Hence, the heat absorbed by the wings of the butterfly *T. limniace* needs to be temporarily stored in heat storage areas on the wing surface, before it is transferred to the wing base and thorax to elevate the thoracic temperature above the ambient level to trigger autonomous flight.

Thus, the dark areas in the mid-posterior region near the wing base in wing cells A-Cu₃ and Cu₂-Cu₃ in the fore wings, and in wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ in the hind wings were temporary heat storage areas on the wing surface. The heat was transferred from the heat storage areas to the wing base and thorax through the veins near the storage areas. During the heat transfer process, the temperatures of the wing base and thorax were the same as the heat storage areas and much higher than those of other areas of the wing surfaces on both the fore and hind wings, and the thoracic temperature reached equilibrium after lighting for 480 s. Thus, the channels for heat transfer were the veins near the storage areas.

Kingsolver (1985) suggested that the wings act as solar reflectors in *Pieris* butterflies to reflect solar radiation onto the body in order to increase its temperature. However, some studies have shown that the absorption of heat by butterfly wings depends mainly on the internal structure of the scale itself, where the structure called a “photonic crystal” can convert the absorbed light into heat for autonomous flight (Li *et al.*, 2004; Han *et al.*, 2013; Luohong, 2014; Han *et al.*, 2015a, b). In the butterfly *Trogonoptera brookiana*, the scales have longitudinal ridges that run through the scales and the surfaces of the scales comprises a set of raised longitudinal quasiparallel lamellae (ridges), where the spaces between adjacent ridges are filled with a netlike reticulum comprising pores (Han *et al.*, 2015a). This “photonic crystal” structure in the scales may help the wings to absorb heat. Therefore, the heat absorbed is not reflected to the wing base and thorax through the wing surface, but instead it is transmitted via the internal tissue in the wing. Our results showed that the heat was transferred from the heat storage areas to the wing base and thorax through veins Cu₂, Cu₃, Cu, and A in the fore wings, and veins 1A, Cu₂, Cu₁, Cu, M₁, M₃, M, R₂, and R in the hind wings during lighting for 0–540 s. This suggests that the heat absorbed by the wings is transferred from the heat storage areas on the wing surface to the wing base and thorax through wing veins.

CONCLUSIONS

In this study, we showed that in the optimal light intensity range, a higher light intensity could help butterflies to absorb more heat, and the optimal wing opening angles for heat absorption ranged from 60° to 90°. The heat absorption capacity of the fore wing was

much greater than that of the hind wing. The dark areas on the wing surfaces were the areas where heat was absorbed by the butterfly wing. The dark areas in the mid-posterior region near the wing base in wing cells A-CU₃ and Cu₂-Cu₃ in the fore wing, and in wing cells 1A-Cu₂, Cu₁-Cu₂, M₃-Cu₁, and R₂-M₁ in the hind wings were temporary heat storage areas on the wing surface. The heat was transferred from heat storage areas to the wing base and thorax through veins Cu₂, Cu₃, Cu, and A in the fore wings, and veins 1A, Cu₂, Cu₁, Cu, M₁, M₃, M, R₂, and R in the hind wings during lighting for 0–540s. In the future, we will investigate the heat storage mechanism in butterfly wings, as well the heat transfer mechanism in the wing veins and the role of light areas on the wings in heat absorption.

ACKNOWLEDGEMENTS

We would like to thank our reviewers for their comments, which have improved the manuscript tremendously. We also thank Chuanjing Liu for her assistance in original materials. We thank International Science Editing for language editing.

REFERENCES

- Barton M, Porter W, Kearney M. 2014.** Behavioural thermoregulation and the relative roles of convection and radiation in a basking butterfly. *Journal of Thermal Biology* 41: 65-71 DOI 10.1016/j.jtherbio.2014.02.004.
- Bennett VJ, Smith WP, Betts MG. 2012.** Evidence for mate guarding behavior in the Taylor's Checkerspot Butterfly. *Journal of Insect Behavior* 25: 183-196 DOI 10.1007/s10905-011-9289-1.
- Berwaerts K, Van Dyck H, Aerts P. 2002.** Does flight morphology relate to flight performance? An experimental test with the butterfly *Pararge aegeria*. *Functional Ecology* 16: 484-491 DOI 10.1046/j.1365-2435.2002.00650.x.
- Berwaerts K, Van Dyck H, Vints E, Matthysen E. 2001.** Effect of manipulated wing characteristics and basking posture on thermal properties of the butterfly *Pararge aegeria* (L.). *Journal of Zoology* 255: 261-267 DOI 10.1017/S0952836901001327.
- Bonebrake TC, Boggs CL, Stamberger JA, Deutsch CA, Ehrlich PR. 2014.** From global change to a butterfly flapping: biophysics and behaviour affect tropical climate change impacts. *Proceedings of the Royal Society B: Biological Sciences* 281: 20141264 DOI 10.1098/rspb.2014.1264.
- Bosi SG, Hayes J, Large MC, Poladian L. 2008.** Color, iridescence, and

- thermoregulation in Lepidoptera. *Applied Optics* 47: 5235-5241 DOI 10.1364/AO.47.005235.
- Brashears J, Aiello A, Seymoure BM. 2016.** Cool Bands: Wing bands decrease rate of heating, but not equilibrium temperature in *Anartia fatima*. *Journal of Thermal Biology* 56: 100-108 DOI 10.1016/j.jtherbio.2016.01.007.
- Chen XM, Zhou CL, Shi JY, Shi L, Yi CH. 2008.** Ornamental butterflies in China. Beijing: China Forestry Publishing House, Beijing, pp108–109.
- De Keyser R, Breuker CJ, Hails RS, Dennis RLH, Shreeve TG. 2015.** Why small is beautiful: wing colour is free from thermoregulatory constraint in the Small Lycaenid Butterfly, *Polyommatus icarus*. *Plos One* 10: e0122623 DOI 10.1371/journal.pone.0122623.
- Devries P. 2015.** Butterfly Reflections in Thirds, Springer International Publishing, DOI 10.1007/978-3-319-20457-4_11.
- Emmel TC. 1972.** Mate selection and balanced polymorphism in the tropical nymphalid butterfly, *Anartia fatima*. *Evolution* 26: 96-107 DOI 10.1111/j.1558-5646.1972.tb00177.x.
- Gibbs M, Van Dyck H, Breuker CJ. 2018.** Flight-induced transgenerational maternal effects influence butterfly offspring performance during times of drought. *Oecologia* 186: 383-391 DOI 10.1007/s00442-017-4030-1.
- Gibbs M, Van Dyck H, Karlsson B. 2010.** Reproductive plasticity, ovarian dynamics and maternal effects in response to temperature and flight in *Pararge aegeria*. *Journal of Insect Physiology* 56: 1275-1283 DOI 10.1016/j.jinsphys.2010.04.009.
- Guppy CS. 1986.** The adaptive significance of alpine melanism in the butterfly *Parnassius phoebus* F. (Lepidoptera: Papilionidae). *Oecologia* 70: 205-213 DOI 10.1007/BF00379241.
- Han Z, Li B, Mu Z, Yang M, Niu S. 2015a.** An ingenious super light trapping surface templated from butterfly wing scales. *Nanoscale Research Letters* 10: 1-8 DOI 10.1186/s11671-015-1052-7.
- Han Z, Li B, Mu Z, Yang M, Niu S, Zhang J, Ren L. 2015b.** Fabrication of the replica templated from butterfly wing scales with complex light trapping structures. *Applied Surface Science* 355: 290-297 DOI 10.1016/j.apsusc.2015.07.119.
- Han Z, Mu Z, Li B, Niu S, Zhang J, Ren L. 2016.** A high-transmission, multiple antireflective surface inspired from bilayer 3D ultrafine hierarchical structures in butterfly wing scales. *Small* 12: 713-720 DOI 10.1002/sml.201502454.
- Han Z, Niu S, Zhang L, Liu Z, Ren L. 2013.** Light Trapping effect in wing scales of

- 707 butterfly *Papilio peranthus* and its simulations. *Journal of Bionic Engineering* 10:
708 162-169 DOI 10.1016/S1672-6529(13)60211-5.
- 709 **Heinrich B. 1986.** Comparative thermoregulation of four montane butterflies of different
710 mass. *Physiological Zoology* 59: 616-626 DOI 10.1086/physzool.59.6.30158609.
- 711 **Heinrich B. 1990.** Is "reflectance" basking real? *Journal of Experimental Biology* 154:
712 31-43.
- 713 **Huey RB, Hertz PE, Sinervo B. 2003.** Behavioral drive versus behavioral inertia in
714 evolution: a null model approach. *The American Naturalist* 161: 357-366 DOI
715 10.1086/346135.
- 716 **Kammer AE, Bracchi J. 1973.** Role of the wings in the absorption of radiant energy by
717 a butterfly. *Comparative Biochemistry and Physiology Part A: Physiology* 45:
718 1057-1063 DOI 10.1016/0300-9629(73)90342-3.
- 719 **Kemp DJ, Krockenberger AK. 2004.** Behavioural thermoregulation in butterflies: the
720 interacting effects of body size and basking posture in *Hypolimnas bolina* (L.)
721 (Lepidoptera: Nymphalidae). *Australian Journal of Zoology* 52: 229-239 DOI
722 10.1071/ZO03043.
- 723 **Kingsolver J. 1985.** Thermal ecology of *Pieris* butterflies (Lepidoptera: Pieridae): a new
724 mechanism of behavioral thermoregulation. *Oecologia* 66: 540-545 DOI
725 10.1007/BF00379347.
- 726 **Kingsolver J, Moffat R. 1982.** Thermoregulation and the determinants of heat transfer in
727 *Colias* butterflies. *Oecologia* 53: 27-33 DOI 10.1007/BF00377132.
- 728 **Kingsolver JG. 1983.** Thermoregulation and flight in *Colias* butterflies: Elevational
729 patterns and mechanistic limitations. *Ecology* 64: 534-545 DOI 10.2307/1939973.
- 730 **Kingsolver JG. 1987.** Evolution and coadaptation of thermoregulatory behavior and wing
731 pigmentation pattern in *Pierid* butterflies. *Evolution* 41: 472-490 DOI
732 10.1111/j.1558-5646.1987.tb05819.x.
- 733 **Kingsolver JG. 1988.** Thermoregulation, flight, and the evolution of wing pattern in
734 *Pierid* butterflies: The topography of adaptive landscapes. *Integrative &*
735 *Comparative Biology* 28: 899-912 DOI 10.1093/icb/28.3.899.
- 736 **Kingsolver JG, Wiernasz DC. 1991.** Seasonal polyphenism in wing-melanin pattern and
737 thermoregulatory adaptation in *Pieris* butterflies. *The American Naturalist* DOI 137:
738 816-830 DOI 10.1086/285195.
- 739 **Kleckova I, Konvicka M, Klecka J. 2014.** Thermoregulation and microhabitat use in
740 mountain butterflies of the genus *Erebia*: Importance of fine-scale habitat
741 heterogeneity. *Journal of Thermal Biology* 41: 50-58 DOI

- 742 10.1016/j.jtherbio.2014.02.002.
- 743 **Li B, Li Q, Zhou J, Li L. 2004.** Photonic structures in butterfly *Thaumantis diores*.
744 *Chinese Science Bulletin* 49: 2545-2546.
- 745 **Liao H, Shi L, Liu W, Du T, Ma Y, Zhou C, Deng J. 2017.** Effects of light intensity on
746 the flight behaviour of adult *Tirumala limniace* (Cramer) (Lepidoptera:
747 Nymphalidae: Danainae). *Journal of Insect Behavior* 30: 139-154 DOI
748 10.1007/s10905-017-9602-8.
- 749 **Luohong S. 2015.** Butterfly Wings: Nature's Fluttering Kaleidoscope. New York:
750 Springer International Publishing, pp127-134 DOI 10.1007/978-3-319-03125-5_11.
- 751 **Mattila ALK. 2015.** Thermal biology of flight in a butterfly: genotype, flight metabolism,
752 and environmental conditions. *Ecology and Evolution* 5: 5539-5551 DOI
753 10.1002/ece3.1758.
- 754 **Michielsen K, De Raedt H, Stavenga DG. 2010.** Reflectivity of the gyroid biophotonic
755 crystals in the ventral wing scales of the Green Hairstreak butterfly, *Callophrys rubi*.
756 *Journal of the Royal Society Interface* 7: 765-771 DOI 10.1098/rsif.2009.0352.
- 757 **Nève G, Hall C. 2016.** Variation of thorax flight temperature among twenty Australian
758 butterflies (Lepidoptera: Papilionidae, Nymphalidae, Pieridae, Hesperidae,
759 Lycaenidae). *European Journal of Entomology* 113: 571-578 DOI
760 10.14411/eje.2016.077.
- 761 **Nijhout HF. 1991.** The development and evolution of butterfly wing patterns.
762 Washington D C, United States: Smithsonian Institution Scholarly.
- 763 **Niu SC, Li B, Ye JF, Mu ZZ. 2016.** Angle-dependent discoloration structures in wing
764 scales of *Morpho menelaus* butterfly. *SCIENCE CHINA Technological Sciences* 59:
765 749-755 DOI 10.1007/s11431-016-6007-4.
- 766 **Plattner L. 2004.** Optical properties of the scales of *Morpho rhetenor* butterflies:
767 theoretical and experimental investigation of the back-scattering of light in the
768 visible spectrum. *Journal of the Royal Society Interface* 1: 49-59 DOI
769 10.1098/rsif.2004.0006.
- 770 **Rawlins J E. 1980.** Thermoregulation by the black swallowtail butterfly, *Papilio*
771 *polyxenes* (Lepidoptera: Papilionidae). *Ecology* 61: 345-357 DOI 10.2307/1935193 .
- 772 **Schmitz H. 1994.** Thermal characterization of butterfly wings-1. Absorption in relation
773 to different color, surface structure and basking type. *Journal of Thermal Biology* 19:
774 403-412 DOI 10.1016/0306-4565(94)90039-6.
- 775 **Seymour BM, Aiello A. 2015.** Keeping the band together: evidence for false boundary
776 disruptive coloration in a butterfly. *Journal of Evolutionary Biology* 28: 1618-1624

- DOI 10.1111/jeb.12681.
- Shanks K, Senthilarasu S, French-Constant RH, Mallick TK. 2015.** White butterflies as solar photovoltaic concentrators. *Scientific Reports* 5: 12267 DOI 10.1038/srep12267.
- Shreeve TG. 1992.** Adult behavior. In: The ecology of butterflies in Britain (Ed. by Dennis RLH). Oxford, UK: Oxford University Press, pp22-45.
- Siddique RH, Donie YJ, Gomard G, Yalamanchili S, Merdzhanova T, Lemmer U, Holscher H. 2017.** Bioinspired phase-separated disordered nanostructures for thin photovoltaic absorbers. *Science Advances* 3: e1700232 DOI 10.1126/sciadv.1700232.
- Stoehr AM, Wojan EM. 2016.** Multiple cues influence multiple traits in the phenotypically plastic melanization of the cabbage white butterfly. *Oecologia* 182: 1-11 DOI 10.1007/s00442-016-3694-2.
- Stuart-Fox D, Newton E, Clusella-Trullas S. 2017.** Thermal consequences of colour and near-infrared reflectance. *Philosophical Transactions of the Royal Society B* 372: 20160345 DOI 10.1098/rstb.2016.0345.
- Taylor OR. 1973.** A non-genetic "Polymorphism" in *Anartia fatima* (Lepidoptera: Nymphalidae). *Evolution* 27: 161 DOI 10.1111/j.1558-5646.1973.tb05928.x.
- Trullas CT, van Wyk JH, Spotila JR. 2007.** Thermal melanism in ectotherms. *Journal of Thermal Biology* 32: 235-245 DOI 10.1016/j.jtherbio.2007.01.013.
- Tsuji J, Kingsolver JG, Watt WB. 1986.** Thermal physiological ecology of *Colias* butterflies in flight. *Oecologia* 69: 161-170 DOI 10.1007/BF00377616.
- Van Dyck H, Matthysen E, Dhondt AA. 1997.** The effect of wing colour on male behavioural strategies in the speckled wood butterfly. *Animal Behaviour* 5: 39-51 DOI 10.1006/anbe.1996.0276.
- Wasserthal LT. 1975.** The rôle of butterfly wings in regulation of body temperature. *Journal of Insect Physiology* 21: 1921-1930 DOI 10.1016/0022-1910(75)90224-3.
- Watt WB. 1968.** Adaptive significance of pigment polymorphisms in *Colias* butterflies. I. Variation of melanin pigment in relation to thermoregulation. *Evolution* 22: 437-458 DOI 10.1111/j.1558-5646.1968.tb03985.x.
- Westerman E, Drucker CB, Monteiro A. 2014.** Male and female mating behavior is dependent on social context in the butterfly *Bicyclus anynana*. *Journal of Insect Behavior* 27: 478-495 DOI 10.1007/s10905-014-9441-9.
- Wu L, Ren W, Song Y, Xin M, Niu S, Han Z. 2015.** High light absorption properties and optical structures in butterfly *Heliophorus ila* Lycaenidae wing scales. *Rsc Advances*

5: 46011-46016 DOI 10.1039/C5RA04135J.

Table 1 Effects of the light intensity and wing opening angle on heat absorption by adult *Tirumala limniace* butterflies.

Sex	Source	<i>d.f.</i>	Type III SS	Mean Square	<i>F</i>	<i>P</i>
Male	Equilibrium thoracic temperature	Wing opening angle	9	2154.7588	239.4176	6.1 <0.0001
		Light intensity	5	44467.6879	8893.5375	226.55 <0.0001
		Wing opening angle*Light intensity	45	443.29354	9.85097	0.25 1
	Time required to reach equilibrium temperature	Wing opening angle	9	161.94397	17.993775	6.91 <0.0001
		Light intensity	5	1165.495037	233.099007	89.46 <0.0001
		Wing opening angle*Light intensity	45	87.682895	1.948509	0.75 0.8659
	ΔT	Wing opening angle	9	2009.57726	223.28636	6.16 <0.0001
		Light intensity	5	40432.775	8086.555	223.02 <0.0001
		Wing opening angle*Light intensity	45	503.31054	11.18468	0.31 1
	Rate of thoracic temperature increase	Wing opening angle	9	2912.84731	323.6497	19.99 <0.0001
		Light intensity	5	11774.35845	2354.67169	145.42 <0.0001
		Wing opening angle*Light intensity	45	3145.22568	69.8939	4.32 <0.0001
	Equilibrium thoracic temperature	Wing opening angle	9	1651.61128	183.51236	2.59 0.0093
		Light intensity	5	40823.08502	8164.617	15.02 <0.0001
		Wing opening angle*Light intensity	45	730.02801	16.22284	0.23 1
Female	Time to reach equilibrium temperature	Wing opening angle	9	394.8589642	43.8732182	25.28 <0.0001
		Light intensity	5	975.0365768	195.0073154	112.36 <0.0001
		Wing opening angle*Light intensity	45	88.2546565	1.9612146	1.13 0.2964
	ΔT	Wing opening angle	9	1441.50025	160.16669	4.28 <0.0001
		Light intensity	5	41421.11186	8284.22237	221.13 <0.0001
		Wing opening angle*Light intensity	45	815.41275	18.12028	0.48 0.9967
	Rate of thoracic temperature increase	Wing opening angle	9	1873.31856	208.14651	11.76 <0.0001
		Light intensity	5	10673.83577	2134.76715	120.64 <0.0001
		Wing opening angle*Light intensity	45	1574.69478	34.99322	1.98 0.0018

Table 2 Effects of the light intensity and wing opening angle on the equilibrium thoracic temperature in adult *Tirumala limniace* butterflies.

Sex	Wing opening angle	600 lx	5000 lx	15000 lx	25000 lx	40000 lx	60000 lx
Male	15°	30.60±2.47d,A	47.47±4.78c,A	61.85±2.64b,A	64.06±2.33ab,BCDE	72.26±3.87ab,A	78.19±8.12a,A
	30°	32.62±2.01d,A	48.05±3.00c,A	61.75±2.44b,A	63.43±1.81b,CDE	69.34±2.23b,A	80.98±6.65a,A
	45°	34.22±2.72d,A	52.803±4.29c,A	63.45±2.99bc,A	67.96±2.55bc,ABC	77.26±4.81ab,A	87.41±9.68a,A
	60°	36.25±1.73d,A	54.39±5.48c,A	63.57±2.12bc,A	70.69±2.07ab,AB	80.81±6.56a,A	81.81±5.69a,A
	75°	36.28±1.53d,A	53.34±3.85c,A	67.06±2.05b,A	71.79±1.44b,A	77.52±4.41ab,A	86.84±5.13a,A
	80°	35.51±1.73e,A	48.82±3.57d,A	63.98±2.44c,A	67.63±1.51bc,ABC	76.42±4.39ab,A	81.78±6.66a,A
	90°	35.30±0.91e,A	49.27±4.10d,A	63.11±3.13c,A	67.27±3.19bc,ABCD	75.28±2.94ab,A	83.51±3.45a,A
	120°	32.87±1.87d,A	43.78±1.73c,A	61.23±2.08b,A	62.36±1.34b,CDE	72.93±1.17a,A	77.19±1.48a,A
	150°	30.31±2.60d,A	43.00±2.11c,A	59.17±1.15b,A	60.67±2.68b,DE	68.85±0.98a,A	73.48±3.16a,A
	180°	30.43±2.13d,A	41.35±4.96c,A	59.55±1.66b,A	59.47±1.10b,E	64.21±1.69b,A	74.19±4.27a,A
Female	15°	33.52±5.12d,A	45.10±5.42cd,A	57.63±7.47bc,A	64.89±1.50b,AB	83.30±5.27a,A	86.27±3.67a,A
	30°	32.07±4.44d,A	46.89±5.00cd,A	59.37±6.11bc,A	58.90±0.52bc,B	71.30±0.63ab,BC	77.09±7.57a,A
	45°	35.93±5.03d,A	50.09±5.39cd,A	62.10±5.85bc,A	64.56±0.49bc,AB	76.66±1.29ab,AB	84.23±8.49a,A
	60°	34.67±3.96d,A	48.99±6.29cd,A	60.34±6.50bc,A	66.43±0.69abc,A	74.06±2.34ab,AB	82.65±11.00a,A
	75°	35.58±3.65d,A	49.26±5.81c,A	60.35±6.09bc,A	65.64±1.50ab,A	73.80±1.58ab,AB	78.40±5.21a,A
	80°	34.63±3.72d,A	47.24±3.47cd,A	59.25±8.07bc,A	65.10±2.46ab,AB	72.25±2.24ab,BC	79.32±6.60a,A
	90°	35.73±3.51d,A	46.15±6.78cd,A	55.56±6.46bc,A	63.21±2.30ab,AB	69.86±3.54ab,BC	74.39±4.54a,A
	120°	31.95±4.10e,A	43.89±5.97de,A	54.64±4.56cd,A	62.38±1.96bc,AB	69.66±3.65ab,BC	75.95±3.63a,A
	150°	31.01±3.62d,A	43.53±6.31cd,A	52.33±7.15bc,A	58.90±3.08ab,B	62.78±2.66ab,BC	71.76±0.62a,A
	180°	30.80±3.33d,A	42.39±5.36cd,A	51.79±5.90bc,A	58.93±2.11ab,B	67.12±3.97a,C	71.81±4.26a,A

Different lowercase and capital letters indicate significant differences at $P < 0.05$ in the results obtained at different light intensities and wing opening angles, respectively.

Table 3 Effects of the light intensity and wing opening angle on the time required to reach equilibrium temperature in adult *Tirumala limniace* butterflies.

Sex	Wing opening angle	600 lx	5000 lx	15000 lx	25000 lx	40000 lx	60000 lx
Male	15°	11.97±1.36a,A	9.68±1.17ab,A	7.94±0.81b,A	4.44±0.27c,A	4.32±0.46c,A	3.89±0.35c,AB
	30°	10.60±1.72a,A	8.64±1.84ab,A	6.26±0.43bc,AB	4.12±0.42c,A	4.36±0.31c,A	4.67±0.77c,A
	45°	7.92±1.31a,A	6.35±0.50ab,A	4.73±0.20bc,BC	3.84±0.10c,AB	3.84±0.65c,A	3.48±0.71c,ABC
	60°	9.81±1.25a,A	6.22±1.11b,A	3.52±0.53c,C	3.18±0.08c,BC	3.18±0.69c,AB	2.68±0.95c,ABCD
	75°	8.98±2.04a,A	6.11±1.45ab,A	3.25±0.51bc,C	2.47±0.04c,C	2.08±0.54c,BC	1.60±0.40c,CD
	80°	10.89±1.49a,A	6.33±1.77b,A	3.70±0.29bc,C	2.81±0.18c,C	2.93±0.52c,ABC	2.56±0.69c,BCD
	90°	6.51±0.05a,A	5.85±0.58a,A	4.04±0.41b,BC	2.84±0.27c,C	1.38±0.10d,C	1.29±0.14d,D
	120°	11.62±1.54a,A	6.24±1.89b,A	4.67±1.00b,BC	2.64±0.48b,C	2.69±0.54b,ABC	2.64±0.73b,ABCD
	150°	9.23±0.56a,A	7.17±1.43a,A	3.77±0.71b,C	2.49±0.11b,C	2.83±0.40b,ABC	2.95±0.65b,ABCD
	180°	11.04±1.53a,A	5.83±1.65b,A	5.33±1.48b,BC	3.05±0.25b,BC	2.13±0.53b,BC	3.16±0.40b,ABCD
Female	15°	13.64±1.40a,A	10.25±0.94b,A	9.51±1.26b,A	4.72±0.52c,A	6.22±0.81c,A	5.03±0.37c,A
	30°	12.74±2.15a,A	10.71±1.52a,A	8.69±1.29ab,A	4.12±0.40c,AB	4.69±0.65bc,B	4.66±0.79bc,AB
	45°	8.01±0.41a,B	6.27±0.71b,B	4.84±0.21bc,B	3.21±0.24cd,BC	3.46±0.50cd,BC	2.79±0.94d,BC
	60°	8.18±1.14a,B	4.77±0.37b,B	4.23±0.93b,B	3.02±0.61b,BC	2.65±0.47b,C	3.18±1.15b,ABC
	75°	6.48±0.11a,B	4.73±0.30b,B	3.14±0.56c,B	2.16±0.21cd,C	2.34±0.09cd,C	1.46±0.28d,C
	80°	9.13±0.40a,B	6.91±0.57ab,B	4.93±1.68bc,B	2.92±0.32c,BC	2.76±0.52c,C	2.64±0.71c,C
	90°	8.31±1.10a,B	4.85±0.66b,B	3.47±0.66bc,B	2.83±0.20bc,BC	1.98±0.50c,C	1.43±0.11c,C
	120°	9.26±0.42a,B	5.37±0.48b,B	3.84±0.51c,B	3.21±0.21c,BC	3.55±0.17c,BC	2.57±0.49,C
	150°	9.32±0.46a,B	6.47±0.32b,B	3.47±0.35c,B	3.15±0.34cd,BC	2.96±0.33cd,C	2.33±0.14d,C
	180°	9.12±1.22a,B	6.67±1.31a,B	3.76±0.75b,B	3.55±0.67b,AB	3.33±0.43b,BC	2.36±0.35b,C

Different lowercase and capital letters indicate significant differences at $P < 0.05$ in the results obtained at different light intensities and wing opening angles, respectively.

Table 4 Effects of the light intensity and wing opening angle on the thoracic temperature excess (ΔT) in adult *Tirumala limniace* butterflies.

Sex	Wing opening						
	angle	600 lx	5000 lx	15000 lx	25000 lx	40000 lx	60000 lx
Male	15°	9.01±2.01b,B	21.41±4.01b,A	37.71±1.57a,BC	41.07±2.45a,BCD	48.96±3.53a,ABC	50.80±8.71a,A
	30°	10.89±0.95d,AB	22.78±2.84c,A	37.17±0.89b,BC	40.81±2.72b,BCD	45.79±1.62ab,BC	54.47±7.13a,A
	45°	12.72±2.23d,AB	27.52±3.91cd,A	39.25±1.60bc,AB	45.68±2.00ab,ABC	54.01±3.95ab,AB	60.34±9.89a,A
	60°	14.56±1.06d,A	28.97±5.05c,A	39.37±0.66bc,AB	47.58±1.70ab,AB	57.72±5.74a,A	55.18±6.36a,A
	75°	15.15±1.40e,A	27.84±3.42d,A	42.39±0.89c,A	49.25±1.10bc,A	54.77±3.86ab,AB	61.35±5.88a,A
	80°	14.33±1.09c,A	23.47±3.78c,A	40.12±1.01b,AB	45.40±1.36ab,ABCD	53.87±4.12a,AB	56.87±7.37a,A
	90°	14.07±0.14e,A	25.01±3.60d,A	38.89±1.73c,ABC	45.12±3.13bc,ABCD	52.65±2.55ab,AB	58.93±3.89a,A
	120°	11.65±1.70d,AB	18.68±2.47c,A	37.27±1.26b,BC	39.27±1.85b,CD	49.94±0.41a,ABC	52.34±1.58a,A
	150°	8.96±1.52d,B	18.52±1.62c,A	35.26±0.62b,C	38.98±2.36b,CD	45.20±1.22a,BC	48.16±3.48a,A
	180°	10.34±1.95c,AB	15.36±4.23c,A	36.91±0.49b,BC	38.22±2.28b,D	41.65±2.45ab,C	48.49±4.22a,A
Female	15°	10.24±3.12c,A	21.09±3.81c,A	33.06±5.95b,A	43.37±0.72b,A	60.16±4.40a,A	62.58±3.01a,A
	30°	9.03±2.63d,A	23.27±3.72c,A	34.91±4.65b,A	36.35±1.35b,A	48.04±1.25a,BC	53.51±5.89a,A
	45°	12.80±3.10e,A	25.18±3.54d,A	37.53±4.31c,A	42.57±0.93bc,A	53.81±0.51ab,AB	59.59±6.94a,A
	60°	11.70±1.87d,A	25.03±4.51cd,A	35.51±4.96bc,A	44.49±0.82ab,A	51.30±1.48a,BC	58.97±8.52a,A
	75°	12.75±2.00e,A	25.38±4.17d,A	35.78±4.52c,A	43.78±0.82bc,A	51.18±0.81ab,BC	55.04±3.50a,A
	80°	11.16±1.34e,A	23.37±1.52d,A	34.66±6.53c,A	43.84±1.72bc,A	49.71±1.48ab,BC	56.05±4.39a,A
	90°	12.66±2.06d,A	22.96±4.84cd,A	31.16±4.79bc,A	38.83±4.68ab,A	47.80±2.59a,BCD	50.24±2.28a,A
	120°	8.83±1.69e,A	19.94±4.02d,A	28.71±3.80c,A	41.82±1.33b,A	47.11±2.69ab,BCD	52.35±2.16a,A
	150°	9.21±1.90c,A	20.21±4.04b,A	27.40±5.23b,A	38.11±2.12a,A	40.48±2.06a,D	47.85±1.97a,A
	180°	7.62±1.42d,A	18.72±3.19c,A	28.16±4.02b,A	39.17±1.76a,A	45.71±3.01a,CD	47.28±2.82a,A

Different lowercase and capital letters indicate significant differences at $P < 0.05$ in the results obtained at different light intensities and wing opening angles, respectively.

Table 5 Effects of the light intensity and wing opening angle on the rate ($^{\circ}\text{C}/\text{min}$) of thoracic temperature increase in adult *Tirumala limniace* butterflies.

Sex	Wing opening angle	600 lx	5000 lx	15000 lx	25000 lx	40000 lx	60000 lx
Male	15 $^{\circ}$	0.78 \pm 0.19d,E	2.36 \pm 0.67d,A	4.86 \pm 0.59c,D	9.26 \pm 0.46b,D	11.42 \pm 0.42a,CD	12.87 \pm 1.09a,B
	30 $^{\circ}$	1.07 \pm 0.16e,CDE	2.94 \pm 0.79d,A	6.01 \pm 0.52c,CD	10.00 \pm 0.49b,D	10.55 \pm 0.39ab,D	11.82 \pm 0.54a,B
	45 $^{\circ}$	1.63 \pm 0.24d,ABC	4.46 \pm 0.92d,A	8.32 \pm 0.50c,BCD	11.88 \pm 0.33b,CD	14.58 \pm 1.66b,CD	17.72 \pm 1.14a,B
	60 $^{\circ}$	1.55 \pm 0.26d,BCD	4.96 \pm 1.30cd,A	11.65 \pm 1.58bc,AB	14.98 \pm 0.50b,BC	19.31 \pm 2.67ab,CD	23.93 \pm 4.76a,B
	75 $^{\circ}$	1.80 \pm 0.31d,AB	5.49 \pm 2.09d,A	13.71 \pm 2.12cd,A	19.98 \pm 0.61bc,A	29.28 \pm 5.61b,B	41.47 \pm 6.65a,A
	80 $^{\circ}$	1.36 \pm 0.20e,BCDE	4.50 \pm 1.78de,A	10.99 \pm 0.99cd,AB	16.37 \pm 1.45bc,AB	19.12 \pm 2.17ab,CD	24.63 \pm 4.40a,B
	90 $^{\circ}$	2.16 \pm 0.03e,A	4.49 \pm 1.12de,A	9.82 \pm 1.05cd,ABC	16.16 \pm 1.88c,AB	38.36 \pm 2.75b,A	46.26 \pm 3.92a,A
	120 $^{\circ}$	1.00 \pm 0.05d,DE	4.07 \pm 1.81cd,A	8.92 \pm 2.21bcd,BCD	15.88 \pm 2.78abc,BC	20.19 \pm 4.18ab,BCD	23.83 \pm 7.42a,B
	150 $^{\circ}$	0.96 \pm 0.12c,DE	2.92 \pm 0.87c,A	9.97 \pm 1.69b,ABC	15.68 \pm 1.13ab,BC	16.65 \pm 2.37a,CD	17.98 \pm 3.74a,B
Female	180 $^{\circ}$	0.94 \pm 0.17e,DE	3.00 \pm 1.23de,A	7.90 \pm 1.77cd,BCD	12.61 \pm 0.71bc,BCD	21.47 \pm 4.12a,BC	15.59 \pm 1.00ab,B
	15 $^{\circ}$	0.77 \pm 0.26c,B	2.14 \pm 0.55c,B	3.44 \pm 0.18c,C	9.40 \pm 1.02b,BC	10.00 \pm 1.53ab,C	12.60 \pm 1.25a,C
	30 $^{\circ}$	0.72 \pm 0.16d,B	2.33 \pm 0.60cd,B	4.03 \pm 0.06c,C	8.93 \pm 0.58b,C	10.63 \pm 1.38ab,BC	11.75 \pm 0.81a,C
	45 $^{\circ}$	1.57 \pm 0.30d,AB	4.27 \pm 1.13cd,AB	7.80 \pm 1.00bcd,B	13.45 \pm 1.24bc,BC	16.17 \pm 2.19b,BC	25.35 \pm 6.32a,BC
	60 $^{\circ}$	1.52 \pm 0.37c,AB	5.18 \pm 0.57c,A	8.82 \pm 1.17bc,AB	16.26 \pm 3.79ab,AB	20.79 \pm 4.24a,ABC	21.32 \pm 3.89a,C
	75 $^{\circ}$	1.97 \pm 0.31d,A	5.50 \pm 1.24cd,A	11.64 \pm 0.82c,A	20.61 \pm 1.99b,A	21.92 \pm 1.01b,AB	39.78 \pm 5.38a,A
	80 $^{\circ}$	1.23 \pm 0.18c,AB	3.47 \pm 0.53c,AB	7.77 \pm 1.35bc,B	15.36 \pm 1.65ab,ABC	19.67 \pm 4.50a,ABC	24.23 \pm 5.83a,BC
	90 $^{\circ}$	1.64 \pm 0.45e,AB	4.69 \pm 0.59d,AB	9.38 \pm 1.39c,AB	14.00 \pm 2.36b,BC	27.93 \pm 8.03a,A	35.74 \pm 3.70a,AB
	120 $^{\circ}$	0.97 \pm 0.23c,B	3.68 \pm 0.57c,AB	7.50 \pm 0.62bc,B	13.22 \pm 1.34b,BC	13.40 \pm 1.42b,BC	22.46 \pm 5.58a,BC
	150 $^{\circ}$	0.98 \pm 0.19d,B	3.13 \pm 0.65d,AB	7.92 \pm 1.22c,B	12.46 \pm 1.69b,BC	13.89 \pm 0.92b,BC	20.64 \pm 0.93a,C
	180 $^{\circ}$	0.86 \pm 0.21d,B	3.30 \pm 1.32cd,AB	7.92 \pm 1.48bc,B	12.14 \pm 2.95b,BC	14.15 \pm 1.72b,BC	20.87 \pm 2.81a,C

Different lowercase and capital letters indicate significant differences at $P < 0.05$ in the results obtained at different light intensities and wing opening angles, respectively.

889 **Table 6** Heat absorption capacities of the fore wing and hind wing in adult *Tirumala limniace*.

Light intensity	Equilibrium thoracic temperature (°C)		Time (min) required to reach equilibrium temperature		ΔT (°C)		Rate (°C/min) of thoracic temperature increase	
	Fore wing	Hind wing	Fore wing	Hind wing	Fore wing	Hind wing	Fore wing	Hind wing
Male		30.99±0.65		11.23±0.8	11.56±0.55	10.37±0.7		0.95±0.1
	600 lx	*	8	8.79±0.39*	3	ns	5	1.32±0.06*
		47.55±0.67			27.12±0.54	21.53±1.2	4.78±0.25*	2.69±0.2
	5000 lx	**	7	5.73±0.22*	8.43±0.94	**	0	*
								7
	15000			4.65±0.18*		38.96±0.87	29.70±2.3	8.45±0.42*
		60.63±0.73						3.83±0.4
	lx	**	6	*	7.93±0.52	**	1	*
								0
	25000			4.17±0.25*		44.49±0.34	40.80±1.3	10.85±0.59
Female		67.32±0.52						6.74±0.6
	lx	**	7	*	6.28±0.46	*	1	**
								9
	40000			3.51±0.10*		45.23±0.81	38.44±1.1	12.98±0.55
		68.18±1.02						6.59±0.4
	lx	**	5	*	5.96±0.41	**	2	**
								5
	60000			3.65±0.09*		51.98±2.77	45.97±0.5	14.25±0.68
		76.96±2.88						7.78±0.6
	lx	ns	2	*	6.12±0.52	ns	9	**
Female		31.38±0.46		11.44±0.7	13.28±0.81		1.41±0.08*	0.54±0.1
	600 lx	**	0	9.39±0.21*	8	**	5.90±0.90	*
								0
		47.72±1.31		12.23±1.3	27.49±1.01	21.96±2.5	3.43±0.13*	1.80±0.1
	5000 lx	ns	1	8.02±0.18*	4	ns	1	*
								3
	15000			4.98±0.13*		40.16±0.75	34.82±0.6	8.09±0.27*
		61.47±0.53						4.05±0.1
	lx	**	7	*	8.64±0.22	**	7	*
								7
Female	25000			4.03±0.17*		43.06±0.75	41.12±0.6	10.80±0.52
		65.92±0.62						5.26±0.3
	lx	ns	0	*	8.06±0.75	ns	0	**
								5
	40000			3.73±0.21*		50.43±0.56	44.18±1.1	13.72±0.71
		72.87±0.46						6.86±0.4
Female	lx	**	1	*	6.58±0.48	**	1	**
								3
	60000			3.74±0.17*		50.26±1.19	49.84±1.4	13.54±0.56
		75.57±1.05						8.12±0.4
	lx	ns	4	*	6.22±0.33	ns	4	**
								7

890 **, *, and ns indicate the significance levels of differences in the results (equilibrium thoracic temperature, time
891 required to reach equilibrium temperature, ΔT , and rate of thoracic temperature increase) according to the Student's
892 *t*-test between the fore and hind wings at $P < 0.01$, $P < 0.05$, and $P > 0.05$, respectively.

Table 7 The results of time-series ExpDec2 of exponential models analysis for trends of real-time temperatures of the monitoring points in the heat storage areas and non-heat storages in butterflies *Tirumala limniace*.

ExpDec2 Model: $y = A1 \cdot \exp(-x/t1) + A2 \cdot \exp(-x/t2) + y0$

Wing type	Area	Point	y0	A1	t1	A2	t2	r	F	P
Fore wing	Light area	Sp1	49.10929	-12.4412	27.33323	-9.97643	204.0954	0.99816	1.74E+08	<0.0001
		Sp2	49.57089	-11.0479	179.7561	-11.9106	24.73968	0.99842	1.96E+08	<0.0001
		Sp3	47.28997	-10.1232	24.05404	-10.6675	137.0862	0.99856	2.35E+08	<0.0001
		Sp4	48.98872	-10.933	140.0638	-11.428	23.81826	0.99847	2.13E+08	<0.0001
		Sp5	47.99227	-10.6245	129.5564	-10.7854	25.69463	0.99846	2.14E+08	<0.0001
		Sp6	48.50339	-11.4961	26.27294	-10.2551	137.9374	0.99842	2.09E+08	<0.0001
	Wing base	Sp7	52.29589	-7.40247	12.81485	-18.4286	217.0853	0.99897	2.03E+08	<0.0001
	The dark area for heat storage	Sp8	50.75721	-11.7911	228.043	-10.7115	24.1697	0.99702	1.09E+08	<0.0001
		Sp9	49.82083	-11.158	143.3776	-11.5553	21.9387	0.99868	2.56E+08	<0.0001
		Sp10	49.65997	-10.0475	166.3924	-11.7129	24.33015	0.9976	1.50E+08	<0.0001
		Sp11	49.53145	-11.7152	24.88213	-10.5312	138.5937	0.99852	2.28E+08	<0.0001
	Border in dark area	Sp12	47.91015	-11.173	24.91333	-9.14257	145.2361	0.99726	1.43E+08	<0.0001
Hind wing	Light area	H-Sp1	49.17633	-10.4439	25.37711	-11.6125	215.0843	0.99741	1.18E+08	<0.0001
		H-Sp2	50.26874	-14.1702	238.9041	-8.59803	22.34626	0.99829	1.59E+08	<0.0001
		H-Sp3	49.76717	-7.90298	21.82562	-15.0295	244.5349	0.99901	2.50E+08	<0.0001
		H-Sp4	48.85711	-11.3694	25.41496	-10.9769	161.4886	0.99852	2.08E+08	<0.0001
		H-Sp5	48.40098	-10.3218	159.2493	-11.6644	29.78131	0.99848	1.97E+08	<0.0001
		H-Sp6	48.54843	-12.3569	31.24274	-9.28688	151.8395	0.9983	1.89E+08	<0.0001
	Wing base	H-Sp7	52.09365	-6.07772	11.27207	-19.64	235.8899	0.99925	2.53E+08	<0.0001
	The dark area for heat storage	H-Sp8	51.53298	-10.8477	218.4568	-13.9014	28.09526	0.99883	2.47E+08	<0.0001
		H-Sp9	50.90148	-13.0178	30.27843	-10.1396	194.0823	0.99808	1.60E+08	<0.0001
		H-Sp10	49.46966	-10.9644	24.55755	-11.233	145.7819	0.99839	2.01E+08	<0.0001
		H-Sp11	49.06514	-12.9741	31.49138	-9.03949	155.2494	0.9979	1.53E+08	<0.0001
	Border in dark area	H-Sp12	48.04232	-13.2487	34.84247	-7.43888	185.9954	0.99798	1.74E+08	<0.0001

904 **Table 8** The results of time-series ExpDec2 of exponential model analysis for trends of real-time
 905 temperatures of the monitoring points in the heat transfer channels of butterflies *Tirumala limniace*.

906 **ExpDec2 Model:** $y = A1 \cdot \exp(-x/t1) + A2 \cdot \exp(-x/t2) + y0$

Wing type	Heat transfer channel	Point	y0	A1	t1	A2	t2	r	F	P
Fore wing	Vein A channel	A-1	51.65031	-7.84003	12.75891	-17.2268	207.2903	0.99856	1.60E+08	<0.0001
		A-2	50.5731	-11.4145	195.1354	-11.8801	24.10951	0.99828	1.81E+08	<0.0001
		A-3	49.41572	-11.5431	23.41794	-10.7117	148.3088	0.99858	2.39E+08	<0.0001
		A-4	49.06832	-10.6785	24.03947	-11.0778	133.0077	0.9985	2.25E+08	<0.0001
		A-5	47.6016	-10.7138	26.48365	-9.33575	144.2478	0.9974	1.45E+08	<0.0001
	Vein Cu-Cu3 channel	Cu-1	52.26994	-18.9101	211.9113	-6.97851	12.25504	0.99928	2.77E+08	<0.0001
		Cu-2	49.67318	-11.2014	21.28217	-10.1931	197.2287	0.99303	5.58E+07	<0.0001
		Cu-3	49.61431	-8.88343	204.3245	-11.9366	23.56786	0.99061	4.53E+07	<0.0001
		Cu3-4	49.38674	-11.2699	22.10509	-10.4084	150.2459	0.99763	1.55E+08	<0.0001
		Cu3-5	49.59448	-11.1955	22.80552	-10.8877	135.7532	0.99861	2.52E+08	<0.0001
		Cu3-6	49.06385	-11.857	25.909	-9.68216	144.6714	0.99801	1.80E+08	<0.0001
	Vein Cu-Cu2 channel	Cu-1	52.26994	-18.9101	211.9113	-6.97851	12.25504	0.99928	2.77E+08	<0.0001
		Cu-2	49.67318	-11.2014	21.28217	-10.1931	197.2287	0.99303	5.58E+07	<0.0001
		Cu-3	49.61431	-8.88343	204.3245	-11.9366	23.56786	0.99061	4.53E+07	<0.0001
		Cu-4	48.88576	-11.0924	23.04441	-9.71285	160.9608	0.9957	9.08E+07	<0.0001
		Cu2-5	48.67759	-9.81106	147.3665	-11.1861	26.23936	0.9984	2.25E+08	<0.0001
		Cu2-6	47.66185	-9.15046	138.1042	-11.0852	25.98615	0.99752	1.55E+08	<0.0001
Hind wing	Vein A channel	1A-1	52.1982	-24.3625	250.3032	-2.16858	17.72072	0.99953	2.62E+08	<0.0001
		1A-2	50.81314	-14.4636	236.0123	-8.44482	17.70421	0.99773	1.26E+08	<0.0001
		1A-3	50.97916	-12.6379	26.86096	-11.4056	233.0519	0.99857	2.06E+08	<0.0001
		1A-4	49.37377	-10.3706	212.6243	-12.156	29.12618	0.99859	2.17E+08	<0.0001
		1A-5	47.77004	-12.3352	38.22446	-7.54459	313.1892	0.9978	1.71E+08	<0.0001
	Vein Cu-Cu2 channel	Cu-1	52.06394	-4.74979	11.58034	-21.4292	239.0546	0.99951	3.26E+08	<0.0001
		Cu-2	50.40223	-9.85654	22.84531	-12.667	238.1548	0.99708	1.05E+08	<0.0001
		Cu-3	50.69774	-12.9494	23.92048	-10.4392	219.3316	0.99674	1.04E+08	<0.0001
		Cu2-4	50.08737	-13.0645	30.0606	-9.55433	198.0817	0.99787	1.51E+08	<0.0001
		Cu2-5	47.87697	-12.3765	33.8684	-7.79342	191.6719	0.99745	1.42E+08	<0.0001
	Vein Cu-Cu1 channel	Cu-1	52.06394	-4.74979	11.58034	-21.4292	239.0546	0.99951	3.26E+08	<0.0001
		Cu-2	50.40223	-9.85654	22.84531	-12.667	238.1548	0.99708	1.05E+08	<0.0001
		Cu-3	50.45283	-12.7617	23.48126	-10.4624	201.8555	0.99669	1.02E+08	<0.0001
		Cu-4	50.10416	-12.5917	28.82723	-9.83279	178.2873	0.99704	1.10E+08	<0.0001
		Cu1-5	49.79113	-13.0948	31.51791	-9.12978	159.7369	0.99766	1.39E+08	<0.0001
		Cu1-6	48.74334	-13.6109	37.68257	-7.2046	208.5855	0.9979	1.65E+08	<0.0001

Vein Cu-M-M ₃ channel	Cu-1	49.43767	-11.7424	28.44878	-10.0544	141.542	0.99771	1.46E+08	<0.0001
	Cu-2	48.1947	-11.6759	29.62228	-8.88782	150.9702	0.99744	1.41E+08	<0.0001
	Cu-3	52.06394	-4.74979	11.58034	-21.4292	239.0546	0.99951	3.26E+08	<0.0001
	Cu-4	50.40223	-9.85654	22.84531	-12.667	238.1548	0.99708	1.05E+08	<0.0001
	M-5	50.45283	-12.7617	23.48126	-10.4624	201.8555	0.99669	1.02E+08	<0.0001
	M ₃ -6	49.65581	-12.0016	27.14308	-10.1177	170.5792	0.9977	1.44E+08	<0.0001
	M ₃ -7	49.23426	-10.7417	184.3841	-11.596	26.7489	0.99845	2.00E+08	<0.0001
Vein R-M-M ₁ channel	R-1	52.24029	-20.6188	232.3034	-5.47058	11.76307	0.99934	2.63E+08	<0.0001
	R-2	50.9655	-8.43454	13.48904	-15.4319	215.4063	0.99783	1.24E+08	<0.0001
	R-3	49.8506	-10.9777	207.8621	-10.8279	24.80101	0.99655	9.79E+07	<0.0001
	M-4	49.69803	-10.7669	184.4267	-11.2514	23.31607	0.99702	1.16E+08	<0.0001
	M ₁ -5	49.16783	-11.2546	26.17524	-10.719	143.3344	0.99783	1.50E+08	<0.0001
	M ₁ -6	48.86012	-10.2357	135.2823	-10.939	26.87515	0.99784	1.61E+08	<0.0001
	M ₁ -7	48.2474	-11.5281	29.4377	-8.72021	176.2904	0.99685	1.19E+08	<0.0001
Vein R-R ₂ channel	R-1	52.24029	-20.6188	232.3034	-5.47058	11.76307	0.99934	2.63E+08	<0.0001
	R-2	50.9655	-8.43454	13.48904	-15.4319	215.4063	0.99783	1.24E+08	<0.0001
	R-3	49.8506	-10.9777	207.8621	-10.8279	24.80101	0.99655	9.79E+07	<0.0001
	R ₂ -4	49.27588	-11.1911	23.79768	-10.8977	151.1524	0.99836	2.04E+08	<0.0001
	R ₂ -5	48.42841	-10.3834	27.89841	-10.5232	146.3984	0.99825	1.89E+08	<0.0001




Figure 1 Natural light intensity on September 11, 2017 in Yuanjiang County, Yunnan Province, China.




Figure 2 The wing surface was divided into border, middle, and inside parts to obtain spectral reflectance measurements from the wings of adult *Tirumala limniace* butterflies. A: Male, B: female.




Figure 3 Distribution of the potential heat storage areas (A) and potential heat transfer channels (B) on the wing surface in adult *Tirumala limniace* butterflies. In A, the light gray area represents the heat storage area and the blue points are points used for real-time temperature monitoring in the heat storage areas (fore wing: Sp8–Sp11, hind wing: H-Sp8–H-Sp11), light areas (fore wing: Sp1–Sp6, hind wing: H-Sp1–H-Sp6), wing base (fore wing: Sp7, hind wing: H-Sp7) and non-storage areas (fore wing: Sp12, hind wing: H-Sp12). In B, the red points are the points used for real-time temperature monitoring during heat transfer in the wing veins. The heat transfer channels are the vein Cu-Cu₂ channel, vein Cu-Cu₃ channel, and vein A channel in the fore wing, and the vein R-R₂ channel, vein R-M-M₁ channel, vein Cu-M-M₃ channel, vein Cu-Cu₁ channel, vein Cu-Cu₂ channel, and vein 1A channel in the hind wing.




Figure 4 Expanse of the fore and hind wings in adult *Tirumala limniace* butterflies. * indicates a significant difference at $P < 0.05$ in the wing expanse between the fore and hind wings.




Figure 5 Spectral reflectance of the male fore wing (A), male hind wing (B), female fore wing (C) and female hind wing (D) in adult *Tirumala limniace* butterflies.




Figure 6 Equilibrium thoracic temperature (A), time required to reach the equilibrium temperature (B), thoracic temperature excess (ΔT) (C) and rate of thoracic temperature increase (D) in adult *Tirumala limniace* under four scale removal treatments. DSR treatment: scales removed from dark areas; LSR treatment: scales removed from light areas; ASR treatment: all scales removed from the wing surface; and IN treatment: scales on the wings were left intact. Different lowercase and capital letters indicate significant differences at $P < 0.05$ in the results obtained for males and females among the four scale removal treatments.




Figure 7 Real-time temperature on the wing surface during heat absorption by adult *Tirumala limniace* butterflies. The numbers below each subgraph indicate the monitoring time in seconds.




Figure 8 Real-time temperature at the monitoring points in the heat storage areas and non-heat storage areas during heat absorption by the wing surfaces in adult *Tirumala limniace* butterflies. A: Fore wing; B: real-time temperatures at monitoring points on the fore wing during lighting for 500–540 s; C: hind wing; D: real-time temperatures at monitoring points on the hind wing during lighting for 500–540s.




Figure 9 Real-time temperatures at monitoring points on the heat transfer channels on the fore wing during heat absorption by adult *Tirumala limniace* butterflies. A: Vein A channel; B: vein Cu-Cu₃ channel; C: vein Cu-Cu₂ channel.




Figure 10 Real-time temperatures at monitoring points on the heat transfer channels on the hind wing during heat absorption by adult *Tirumala limniace* butterflies. A: vein 1A channel; B: vein Cu-Cu₂ channel; C: vein Cu-M-M₃ channel; D: vein R-R₂ channel; E: vein Cu-Cu₁ channel; F: vein R-M-M₁ channel.