

## **Hypothesis: Avian flight originated in arboreal archosaurs gliding on membranous wings**

The oldest feathers known to date have been found in archosaurs capable to fly. However, some of them (scansoriopterygids) flied by the use of a membrane rather than feathers. We therefore propose a new mechanism for the origin of avian flight by the use of membranous wings on both the forelimbs and the hindlimbs. It complements the Beebe's (1915) prediction of the tetrapteryx stage. Paleontological and embryological evidence suggest that feathers are a modification of reptilian scales. Scansoriopterygids were covered mostly by down-like feathers which seemingly acted as a thermal isolation rather than were adapted for flight. Certain early birds, including scansoriopterygids, possessed elongate shafted tail feathers, which were probably used principally for display and resembled elongate scales. We suppose that display is the primary function of early feathers, which were preadapted for the thermal isolation and also the flight. The body of theropods was covered mostly by typical reptilian scales, yet some ornithischian dinosaurs possessed filamentous integumentary structures which might have had a comparable display function. However, it is doubtful that these structures in dinosaurs were homologous with avian feathers. Early birds probably shared with theropod dinosaurs an incipient endothermy, which was inherited from their common ancestor from the Triassic Period.

# 1 HYPOTHESIS: AVIAN FLIGHT ORIGINATED IN ARBOREAL ARCHOSAURS

## 2 GLIDING ON MEMBRANOUS WINGS

3  
4 PIOTR BAJDEK<sup>1,\*</sup>, TOMASZ SULEJ<sup>1</sup>

5  
6 P. Bajdek ([p.bajdek@twarda.pan.pl](mailto:p.bajdek@twarda.pan.pl)), T. Sulej ([sulej@twarda.pan.pl](mailto:sulej@twarda.pan.pl)), Institute of Paleobiology,  
7 Polish Academy of Sciences, Twarda 51/55, 00-818 Warsaw, Poland<sup>1</sup>

8  
9 \*Corresponding author

10  
11 The oldest feathers known to date have been found in archosaurs capable to fly. However, some  
12 of them (scansoriopterygids) flied by the use of a membrane rather than feathers. We therefore  
13 propose a new mechanism for the origin of avian flight by the use of membranous wings on both  
14 the forelimbs and the hindlimbs. It complements the Beebe's (1915) prediction of the tetrapteryx  
15 stage. Paleontological and embryological evidence suggest that feathers are a modification of  
16 reptilian scales. Scansoriopterygids were covered mostly by down-like feathers which seemingly  
17 acted as a thermal isolation rather than were adapted for flight. Certain early birds, including  
18 scansoriopterygids, possessed elongate shafted tail feathers, which were probably used  
19 principally for display and resembled elongate scales. We suppose that display is the primary  
20 function of early feathers, which were preadapted for the thermal isolation and also the flight. The  
21 body of theropods was covered mostly by typical reptilian scales, yet some ornithischian  
22 dinosaurs possessed filamentous integumentary structures which might have had a comparable

display function. However, it is doubtful that these structures in dinosaurs were homologous with avian feathers. Early birds probably shared with theropod dinosaurs an incipient endothermy, which was inherited from their common ancestor from the Triassic Period.

**Keywords:** origins of avian flight, function of feathers, Scansoriopterygidae

## Introduction

The origin and primary function of feathers has been debated for a long time (Holmgren 1955; Prum 1999). Currently, the prevailing view point on the origin of avian feathers is that they originated from filamentous integumentary structures found in some dinosaur fossils, which might constitute a thermal isolation in endothermic cursorial theropods (Prum 1999; Makovicky and Zanno 2011; Chatterjee 2015). On the contrary, others suggest that feathers are a modification of reptilian scales (Dzik et al. 2010), whereas dinosaurs and early birds are thought to have been ectothermic (Ruben and Jones 2000; Feduccia 2012) and birds most likely not to be descendants of dinosaurs and instead of that to have evolved from arboreal forms of archosaurs (Czerkas and Feduccia 2014). Certain authors offer rather several alternatives for the origin of birds (James and Pourtless 2009) and the origin of feathers (Feduccia 2012). Most of the recent papers on the assembly of the avian body plan ignore non-dinosaur hypotheses for the origin of birds (e.g., Bhullar et al. 2012; Puttick et al. 2014; O'Connor and Zhou 2015; Nesbitt et al. 2017; Sullivan et al. 2017; Cau 2018) and archosaur species possessing unequivocal flight feathers are described as theropod dinosaurs (e.g., Hu et al. 2018; Rauhut et al. 2018).

However, only a mix of these two orthodox viewpoints can be sustained by biological data in the light of paleontological finds of the last decade. Herein, we discuss the available evidence on the origin of birds, archosaurian integumentary structures and thermal physiology. We propose a new mechanism for the origin of avian flight by the use of membranous wings at the very early stage, which complements the Beebe's (1915) prediction of the tetrapteryx stage. Also, we provide a corrected reconstruction of the scansoriopterygid *Yi qi* (Xu et al. 2015) showing that it was capable of flight only if its membranous wings were also present on the hindlimbs.

### **The origin of avian flight**

Important facts for the debate on the origin of avian flight may be provided by the archosaurian family Scansoriopterygidae as it encloses some of the earliest known flying maniraptorans (late Middle Jurassic–early Late Jurassic). Czerkas and Feduccia (2014) considered *Scansoriopteryx* an arboreal bird capable of parachuting or gliding, which would have represented the four-winged “tetrapteryx” stage of the avian evolution, as predicted by Beebe (1915). Interestingly, Czerkas and Feduccia (2014) noted that the body of *Scansoriopteryx* was covered solely by down-like feathers which were composed of individual strands and there were no indications of pennaceous feathers on the forelimb. The lack of preserved wing feathers in *Epidexipteryx* is also peculiar, but it had shafted elongate tail feathers (Zhang et al. 2008). The Early Cretaceous *Zhongornis haoae* may be closely related to scansoriopterygids but no feathers are preserved on the only known specimen (O'Connor and Sullivan 2014).

The paradox of the apparent lack of flight feathers in scansoriopterygids was explained by the description of putative patches of membranous tissue and a long rod-like bone extending from each wrist in the scansoriopterygid *Yi*, which also had abundant down-like feathers (Xu et al. 2015). Each of the known flying tetrapods, including both extinct and extant groups, involves a membrane for flight, as for example bats, flying squirrels and flying lizards do (McGuire and Dudley 2011; Xu et al. 2015). It is an intriguing question whether birds are truly an exception to this rule. Notably, structures such as the contour feathers are far more complex than the membrane, so the membrane seems to be a simpler solution for the development of the ability of flight. In fact, a membrane (propatagium) is present in the forewings of extant birds (Brown et al. 1994) and contributes to the avian flight producing a significant part of the lift (Brown and Cogley 1996). Also, a membrane is present as a rudiment in the hindlimbs of certain baby birds (see Beebe 1915, fig. 10). One can therefore imagine that birds originated from archosaurs which used a membrane as the principal flight apparatus. If this is true, in the early avian evolution the membrane, still present in scansoriopterygids, was subsequently covered with contour feathers resulting in typical tetrapteryx forms such as *Anchiornis* (Hu et al. 2009) and *Microraptor* (Xu et al. 2003) (Fig. 1). In flying objects the center of gravity must be located forward relatively to the center of lift (wings) for good stability and maneuverability (Dizdarevic and Dizdarevic 2005). For this reason, scansoriopterygids probably represented the tetrapteryx body plan (Fig. 2), just like many other primitive birds did before shortening of the tail and reinforcement of the forewings (Beebe 1915).

The fossil feathers *Praeornis* from the Callovian–Oxfordian of Kazakhstan have a rachis and also three vanes (Dzik et al. 2010). The morphology of a three-vaned feather can be far more

88 parsimoniously derived from a reptilian scale than from a simple filament, for example as in the  
89 scenario of such a transformation proposed by Dzik et al. (2010). Feathers are likely to be  
90 modified scales because in the embryological development the feather sac represents the tip of a  
91 scale (Duerden 1922; Holmgren 1955). Given the presence of the scale-like feathers on the tail of  
92 scansoriopterygids (Zhang et al. 2008), feathers hypothetically have first appeared on the tail of  
93 early birds. The down-like feathers composed mostly of individual strands, which covered the  
94 body of scansoriopterygids (Czerkas and Feduccia 2014) and probably acted as a thermal  
95 isolation, would have been simplified from more complex vaned feathers like those present on  
96 their tail. The principal function of the elongate tail feathers in *Epidexipteryx* might have been the  
97 display, as suggested by Zhang et al. (2008). Elongate tail feathers of various types were common  
98 in birds from the Cretaceous Period, such as for example the dromaeosaurid *Microraptor*, the  
99 basal bird *Jeholornis*, the confuciusornithiform *Confuciusornis*, and the enantiornithine  
100 *Eopengornis* (Sullivan et al. 2017). The flight feathers would also have evolved as a modification  
101 of the shafted display feathers. At some stage of the evolution of birds, the display tail feathers  
102 covered other body parts including the wings but it is uncertain when they acquired the  
103 aerodynamic value. The reason of the replacement of the membrane by feathers requires more  
104 study but one possibility is the role of sexual selection.

105 *Praeornis* is a pennaceous feather and it is coeval with scansoriopterygids which apparently  
106 lacked feathers of this type. Comparable in age are also birds such as *Anchiornis* (Hu et al. 2009)  
107 and *Caihong* (Hu et al. 2018), which were equipped with flight feathers on both their forelimbs  
108 and hindlimbs. Therefore, scansoriopterygids can be ruled out as ancestors of the main lineage of

birds. The common ancestor of scansoriopterygids (e.g. *Yi*) and more derived birds of the Late Jurassic (e.g. *Anchiornis*) likely differed from scansoriopterygids in having a longer tail (Fig. 1).

# **The origin of birds and the integumentary structures of archosaurs**

It can be pointed out that the late Middle Jurassic–early Late Jurassic birds were already diversified including forms of a variety of flight apparatus types (a membrane, flight feathers). Thus, birds – here, defined as archosaurs equipped with feathers – would have evolved by the Early to Middle Jurassic. Some authors claim that scansoriopterygids and *Zhongornis* were not “birds” (Fostowicz-Frelik and O’Connor 2017) as in computerized cladistic analyses they are not recovered as members of the clade Aves (see Cau 2018). In fact, this is consistent with the viewpoint presented herein because we do not consider scansoriopterygids as direct ancestors of the main lineage of birds (Fig. 1). The bird-like maniraptorans and ornithomimosaurs of the Cretaceous age, which include common flightless forms, reveal little about bird origins no matter they are considered as theropod dinosaurs or, alternatively, as non-dinosaur birds (Feduccia 2012). This standpoint may be inconsistent with the cladistic approach as certain maniraptorans of the Cretaceous Period, such as for example oviraptorosaurs, are cladistically recovered as more basal than the Jurassic scansoriopterygids (see Sullivan et al. 2017; Cau 2018). The Late Jurassic (Oxfordian) *Limusaurus* might possibly represent a secondarily flightless bird as well, due to its supposedly non-dinosaur hand anatomy and digestive physiology (Bajdek 2018). Secondarily flightless forms would have appeared at very early stages of the bird evolution and such bird forms can be easily mistaken for theropod dinosaurs (see Maryńska et al. 2002).

Remarkably, filamentous structures found in some theropod fossils can be easily mistaken for down feathers of flightless birds. It is debated whether these filaments represent a kind of integumentary structures of theropods or collagen fibers of decomposed skin (Feduccia et al. 2005; Smithwick et al. 2017). However, fossils show that the skin of dinosaurs was covered by reptilian scales, as seen for example in compsognathids (Chiappe and Gröhlich 2010) and tyrannosaurids (Bell et al. 2017). Filamentous integumentary structures have been found in some ornithischian dinosaurs, such as the heterodontosaurid *Tianyulong* (Zheng et al. 2009) and the ceratopsian *Psittacosaurus* (see Mayr et al. 2002). Since these structures were present mainly on the tail whereas the whole body was covered by typical reptilian scales, it seems that their primary function was the display rather than the thermal isolation.

Elongate integumentary appendages of *Longisquama* from the Late Triassic are also thought by some researchers to be probably homologous with avian feathers (Jones et al. 2000). However, the function of the appendages and also the phylogenetic position of *Longisquama* within diapsids are uncertain (Buchwitz and Voigt 2012). Because the limbs of *Longisquama* were short and the enigmatic appendages were placed along the dorsum, its body plan does not conform to that expected for the direct ancestor of birds.

If it is assumed that the enigmatic integumentary structures present in certain ornithischian dinosaurs and *Longisquama* were modified scales, and that avian feathers are modified scales too, one would say that they were homologous. On the other hand, all these structures most likely evolved from typical reptilian scales independently. Whereas true avian feathers are very complex structures and have probably evolved just once, simple thin elongate scales such as those



encountered among ornithischians could have appeared multiple times in different lineages of reptiles (see James and Pourtless 2009).

Prum (1999) discussed several possible functions of filamentous integumentary structures in archosaurs, such as the thermal insulation, heat shielding, water repellency, communication, crypsis, and defense. Since the single-stranded feather-like structures which covered the body of scansoriopterygids are unlikely to have acted as a flight apparatus, it can be expected that they possessed an adaptive value of another kind (Czerkas and Feduccia 2014). For example, an incipient endothermy in dinosaurs and early birds has been suggested by researchers for a long time (e.g., Bakker 1975) but this issue is still a subject of controversy (see Feduccia 2012). The study of theropod and early avian gut contents, feces and gastric pellets suggests that both these groups were likely characterized by moderately high metabolic rates and hence likely an incipient endothermy (Bajdek 2018). Yet, the typical metabolic rates in theropods would have been a little lower than those of extant birds and mammals, which was suggested, among other clues, on the basis of the lack of respiratory turbinates (Ruben et al. 1998) and just moderately rapid digestive processes in theropods (Bajdek 2018). Remarkably, the evidence for an incipient endothermy in early birds is comparable to that for some Late Permian theriodont therapsids, which (a) were likely characterized by a relatively short food retention time in the gastrointestinal tract, as revealed by the study of fossil feces, and moreover (b) possibly had insulation in the form of hairs (Bajdek et al. 2016).

The arboreal and cursorial hypotheses for the origin of avian flight are still a subject of debate (Segre and Banet 2018). The model presented herein suggests that birds originated from arboreal forms of archosaurs that could spread the hindlimbs sideways for tree climbing and flight (Fig. 1).

Currently, the vast majority of researchers supposes that birds evolved from theropod dinosaurs (see Makovicky and Zanno 2011). However, several alternatives for the ancestral group of archosaurs have been proposed, including theropods, dinosauromorphs, crocodylomorphs, and “early archosaurs” (see James and Pourtless 2009). The dinosaur origin of birds is problematic due to the partially closed acetabulum in early birds such as *Anchiornis* and *Scansoriopteryx* (Czerkas and Feduccia 2014). The theropod hindlimbs had a fully upright stance and were clearly adapted for a cursorial locomotion (Burnham et al. 2010; Czerkas and Feduccia 2014). For this reason, in this paper we have chosen to use a broader term “archosaur” to call the arboreal ancestor of birds (see also Bajdek 2018).

## Conclusions

We propose that birds originated from tree-climbing archosaurs and probably passed through the stage of arboreal forms which glided on membranous wings. The membrane was present most likely on both the forelimbs and the hindlimbs. At some stage of the early bird evolution the membrane got covered with feathers resulting in “classical” tetrapteryx bird forms, which were predicted by William Beebe in 1915. Feathers probably evolved from reptilian scales and display is hypothetically the primary function of feathers, which were preadapted for the thermal isolation and flight.

## Author contributions

196 Piotr Bajdek conceived the idea of this work and wrote the manuscript.

197 Tomasz Sulej provided additional ideas and prepared the graphics.

198

## 199 **Acknowledgments**

200

201 We are grateful to our colleagues who kindly commented on early drafts of the manuscript: Alan  
202 Feduccia (University of North Carolina at Chapel Hill), Jerzy Dzik (Polish Academy of Sciences  
203 and University of Warsaw), and Mateusz Tałanda (University of Warsaw).

204

## 205 **References**

206

207 Bajdek P. 2018. Comparative digestive physiology of archosaurs with notes on bird origins.  
208 *PeerJ Preprints* 6: e26902v2 doi: 10.7287/peerj.preprints.26902v2

209

210 Bajdek, P., Qvarnström, M., Owocki, K., Sulej, T., Sennikov, A.G., Golubev, V.K., and  
211 Niedźwiedzki, G. 2016. Microbiota and food residues including possible evidence of pre-  
212 mammalian hair in Upper Permian coprolites from Russia. *Lethaia* 49: 455–477.

213

214 Bakker, R.T. 1975. Dinosaur renaissance. *Scientific American* 232: 58–78.

215

216 Beebe, W.A. 1915. Tetrapteryx stage in the evolution of birds. *Zoologica* 2: 39–52.

217

- 218 Bell, P.R., Campione, N.E., Persons IV, W.S., Currie, P.J., Larson, P.L., Tanke, D.H., and Bakker,  
219 R.T. 2017 Tyrannosauroid integument reveals conflicting patterns of gigantism and feather  
220 evolution. *Biology Letters* 13: 20170092. doi: 10.1098/rsbl.2017.0092  
221
- 222 Bhullar, B.A.S., Marugán-Lobón, J., Racimo, F., Bever, G.S., Rowe, T.B., Norell, M.A., and  
223 Abzhanov, A. 2012. Birds have paedomorphic dinosaur skulls. *Nature* 487:223. doi:  
224 10.1038/nature11146  
225
- 226 Brown, R.E. and Cogley, A.C. 1996. Contributions of the Propatagium to Avian Flight. *The*  
227 *Journal of Experimental Zoology* 276: 112–124.  
228
- 229 Brown, R.E., Baumel, J.J., and Klemm, R.D. 1994. Anatomy of the propatagium: The great  
230 horned owl (*Bubo virginianus*). *Journal of Morphology* 219: 205–224.  
231
- 232 Buchwitz, M. and Voigt, S. 2012. The dorsal appendages of the Triassic reptile *Longisquama*  
233 *insignis*: reconsideration of a controversial integument type. *Paläontologische Zeitschrift* 86:  
234 313–331.  
235
- 236 Burnham, D.A., Feduccia, A., Martin, L.D., and Falk, A.R. 2011. Tree climbing – a fundamental  
237 avian adaptation. *Journal of Systematic Palaeontology* 9: 103–107.

238

239 Cau, A. 2018. The assembly of the avian body plan: a 160-million-year long process. *Bollettino*  
240 *della Società Paleontologica Italiana* 57: 1–25.

241

242 Chatterjee, S. 2015. *The Rise of Birds. 225 Million Years of Evolution*. Johns Hopkins University  
243 Press.

244

245 Chiappe, L.M. and Göhlich, U.B. 2010. Anatomy of *Juravenator starki* (Theropoda:  
246 Coelurosauria) from the Late Jurassic of Germany. *Neues Jahrbuch für Geologie und*  
247 *Paläontologie Abhandlungen* 258: 257–296.

248

249 Czerkas, S. 1994. The history and interpretation of sauropod skin impressions. *Gaia* 10: 173–182.

250

251 Czerkas, S.A. and Feduccia, A. 2014. Jurassic archosaur is a non-dinosaurian bird. *Journal of*  
252 *Ornithology* 155: 841–851.

253

254 Dizdarevic, F. and Dizdarevic, M. 2005. Tailed flying wing aircraft. United States Patent: US  
255 6,923,403 B1.

256

257 Duerden, J.E. 1922. The origin of feathers from the scales of reptiles. *Journal of the Department*  
258 *of Agriculture* July 1922: 67–75.

259

- 260 Dzik, J., Sulej, T., and Niedźwiedzki G. 2010. Possible link connecting reptilian scales with avian  
261 feathers from the early Late Jurassic of Kazakstan. *Historical Biology* 22: 394–402.  
262
- 263 Feduccia, A. 2002. Birds are dinosaurs: simple answer to a complex problem. *The Auk* 119:  
264 1187–1201.  
265
- 266 Feduccia, A. 2012. *Riddle of the Feathered Dragons: Hidden Birds of China*. Yale University  
267 Press.  
268
- 269 Feduccia, A., Lingham-Soliar, T., and Hinchliffe, J.R. 2005. Do Feathered Dinosaurs Exist?  
270 Testing the Hypothesis on Neontological and Paleontological Evidence. *Journal of Morphology*  
271 266: 125–166.  
272
- 273 Fostowicz-Frelik, Ł. and O'Connor, J.K. 2017. The saga of birds. *Acta Palaeontologica Polonica*  
274 62: 844.  
275
- 276 Holmgren, N. 1955. Studies on the phylogeny of birds. *Acta Zoologica* 36: 243–328.  
277
- 278 Hu, D., Hou, L., Zhang, L., and Xu, X. 2009. A pre-*Archaeopteryx* troodontid theropod from  
279 China with long feathers on the metatarsus. *Nature* 461: 640–643.  
280

- 281 Hu, D., Clarke, J.A., Eliason, C.M., Qiu, R., Li, Q., Shawkey, M.D., Zhao, C., D’Alba, L., Jiang,  
282 J., and Xu, X. 2018. A bony-crested Jurassic dinosaur with evidence of iridescent plumage  
283 highlights complexity in early paravian evolution. *Nature Communications* 9: 217. doi:  
284 10.1038/s41467-017-02515-y
- 285
- 286 James, F.C. and Pourtless, J.A. 2009. Cladistics and the origin of birds: a review and two new  
287 analyses. *Ornithological Monographs* 66: 1–78.
- 288
- 289 Jones, T.D, Ruben, J.A, Martin, L.D., Kurochkin, E.N., Feduccia, A., Maderson, P.F.A.,  
290 Hillenius, W.J., Geist, N.R., Alifanov, V. 2000. Nonavian Feathers in a Late Triassic Archosaur.  
291 *Science* 288: 2202–2205.
- 292
- 293 Makovicky, P.J. and Zanno, L.E. 2011. Theropod diversity and the refinement of avian  
294 characteristics. In: Dyke, G. and Kaiser, G. (Eds.), *Living dinosaurs: the evolutionary history of*  
295 *modern birds*. John Wiley & Sons, pp. 9–29.
- 296
- 297 Maryańska, T., Osmólska, H., and Wolsan, M. 2002. Avialan status for Oviraptorosauria. *Acta*  
298 *Palaeontologica Polonica* 47: 97–116.
- 299
- 300 Mayr, G., Peters, D.S., Plodowski, G., and Vogel, O. 2002. Bristle-like integumentary structures  
301 at the tail of the horned dinosaur *Psittacosaurus*. *Naturwissenschaften* 89: 361–365.
- 302

- 303 McGuire, J.A. and Dudley, R. 2011. The Biology of Gliding in Flying Lizards (Genus *Draco*) and  
304 their Fossil and Extant Analogs. *Integrative and Comparative Biology* 51: 983–990.  
305
- 306 Nesbitt, S.J., Butler, R.J., Ezcurra, M.D, Barrett, P.M., Stocker, M.R., Angielczyk, K.D., Smith,  
307 R.M.H., Sidor, C.A., Niedźwiedzki, G., Sennikov, A.G., and Charig, A.J. 2017. The earliest bird-  
308 line archosaurs and the assembly of the dinosaur body plan. *Nature* 544: 484–487.  
309
- 310 O’Connor, J.K. and Sullivan, C. 2014. Reinterpretation of the Early Cretaceous maniraptoran  
311 (Dinosauria: Theropoda) *Zhongornis haoae* as a scansoriopterygid-like non-avian, and  
312 morphological resemblances between scansoriopterygids and basal oviraptorosaurs. *Vertebrata*  
313 *Palasiatica* 52: 3–30.  
314
- 315 O’Connor, J. and Zhou, Z. 2015. Early evolution of the biological bird: perspectives from new  
316 fossil discoveries in China. *Journal Fur Ornithologie*. doi: 10.1007/s10336-015-1222-5  
317
- 318 Prum, R.O. 1999. Development and Evolutionary Origin of Feathers. *Journal of Experimental*  
319 *Zoology* 285: 291–306.  
320
- 321 Puttick, M.N., Thomas, G.H., and Benton, M.J. 2014. High rates of evolution preceded the origin  
322 of birds. *Evolution* 2014: 1–14. doi: 10.1111/evo.12363  
323



324 Rauhut, O.W.M., Foth, C., and Tischlinger, H. 2018. The oldest *Archaeopteryx* (Theropoda:  
325 Avialiae): a new specimen from the Kimmeridgian/Tithonian boundary of Schamhaupten,  
326 Bavaria. *PeerJ* 6: e4191. doi: 10.7717/peerj.4191

327

328 Ruben, J.A. and Jones, T.D. 2000. Selective factors associated with the origin of fur and feathers.  
329 *American Zoologist* 40: 585–596.

330

331 Ruben, J.A., Jones, T.D., and Geist, N.R. 1998. Respiratory physiology of the dinosaurs.  
332 *BioEssays* 20: 852–859.

333

334 Segre, P.S. and Banet, A.I. 2018. The origin of avian flight: finding common ground. *Biological*  
335 *Journal of the Linnean Society*. doi: 10.1093/biolinnean/bly116

336

337 Smithwick, F.M., Mayr, G., Saitta, E.T., Benton, M.J., and Vinther, J. 2017. On the purported  
338 presence of fossilized collagen fibres in an ichthyosaur and a theropod dinosaur. *Palaeontology*  
339 60: 409–422.

340

341 Sullivan, C., Xu, X., and O'Connor, J.K. 2017. Complexities and novelties in the early evolution  
342 of avian flight, as seen in the Mesozoic Yanliao and Jehol Biotas of Northeast China.  
343 *Palaeoworld* 26: 212–229.

344


Xu, X., Zhou, Z., Wang, X., Kuang, X., Zhang, F., and Du, X. 2003. Four-winged dinosaurs from China. *Nature* 421: 335–340.

Xu, X., Zheng, X., Sullivan, C., Wang, X., Xing, L., Wang, Y., Zhang, X., O'Connor, J.K., Zhang, F., and Pan, Y. 2015. A bizarre Jurassic maniraptoran theropod with preserved evidence of membranous wings. *Nature* 521: 70–73.

Zhang, F., Zhou, Z., Xu, X., Wang, X., and Sullivan, C. 2008. A bizarre Jurassic maniraptoran from China with elongate ribbon-like feathers. *Nature* 455: 1105–1108.

Zheng, X.T., You, H.L., Xu, X., and Dong, Z.M. 2009. An Early Cretaceous heterodontosaurid dinosaur with filamentous integumentary structures. *Nature* 458: 333–336.

367




369 **Fig. 1. Origin and early evolution of birds.** Birds originated from tree-climbing archosaurs and  
 370 passed through the stage of arboreal forms which glided on membranous wings. It is uncertain at  
 371 which stage the tail feathers have first appeared.

372

373

374



**Fig. 2. Reconstructions of *Yi qi* showing the aerodynamic apparatus.** Model proposed by Xu et al. (2015) (A) and a new model proposed herein (B). Xu et al. (2015) provided three different alternatives for the construction of the front wing of *Yi* and only that one of the greatest extension of the membrane is reproduced (model A). Approximate positions of the center of mass (blue dots) and the center of lift (orange dots) are marked. Note that *Yi* was capable to fly only if the membrane was extended also on the hindlimbs, as the center of lift was located backward relatively to the center of mass (model B).