

**A peer-reviewed version of this preprint was published in PeerJ on 12 November 2018.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.5925) (peerj.com/articles/5925), which is the preferred citable publication unless you specifically need to cite this preprint.

Holwerda FM, Díez Díaz V, Blanco A, Montie R, Reumer JWF. 2018. Late Cretaceous sauropod tooth morphotypes may provide supporting evidence for faunal connections between North Africa and Southern Europe. PeerJ 6:e5925 <https://doi.org/10.7717/peerj.5925>

# Could Late Cretaceous sauropod tooth morphotypes provide supporting evidence for faunal connections between North Africa and Southern Europe?

Femke M Holwerda<sup>Corresp., 1, 2, 3</sup>, Verónica Díez Díaz<sup>4, 5</sup>, Alejandro Blanco<sup>2, 6</sup>, Roel Montie<sup>1, 2, 3</sup>, Jelle WF Reumer<sup>1</sup>

<sup>1</sup> Faculty of Geosciences, Utrecht University, Utrecht, Netherlands

<sup>2</sup> Vertebrate Palaeontology, SNSB-Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Bavaria, Germany

<sup>3</sup> Department of Earth and Environmental Sciences and GeoBioCenter, Ludwig-Maximilians-Universität München, Munich, Bavaria, Germany

<sup>4</sup> Leibniz-Institut für Evolutions-und Biodiversitätsforschung, Museum für Naturkunde, Berlin, Germany

<sup>5</sup> Humboldt Universität Berlin, Berlin, Germany

<sup>6</sup> Facultade de Ciencias, Centro de Investigacións Científicas Avanzadas (CICA), Universidade da Coruña, Coruña, Spain

Corresponding Author: Femke M Holwerda  
Email address: f.holwerda@lrz.uni-muenchen.de

The Cretaceous Kem Kem beds of Morocco and equivalent beds in Algeria have produced a rich fossil assemblage, yielding, amongst others, isolated sauropod teeth, which can be used in species diversity studies. These Albian-Cenomanian (~113 – 93.9 Ma) strata rarely yield sauropod body fossils, therefore, isolated teeth can help to elucidate the faunal assemblages from North Africa, and their relations with those of contemporaneous beds and geographically close assemblages. Eighteen isolated sauropod teeth from three localities (Erfoud and Taouz, Morocco, and Algeria) are studied here, to assess whether the teeth can be ascribed to a specific clade, and whether different tooth morphotypes can be found in the samples. Two general morphotypes are found, based on enamel wrinkling and general tooth morphology. Morphotype I, with mainly rugose enamel wrinkling, pronounced carinae, lemon-shaped to (sub)cylindrical cross-section and mesiodistal tapering towards an apical tip, shows affinities to titanosauriforms and titanosaurs. Morphotype II, characterized by more smooth enamel, cylindrical cross-section, rectangular teeth with no apical tapering and both labial and lingual wear facets, shows similarities to rebbachisaurids. Moreover, similarities are found between these northwest African tooth morphotypes, and tooth morphotypes from titanosaurs and rebbachisaurids from both contemporaneous finds from north and central Africa, as well as from the latest Cretaceous (Campanian-Maastrichtian, 83.6Ma – 66.0Ma) of the Ibero-Armorican Island. These results support previous hypotheses from earlier studies on faunal exchange and continental connections between North Africa and Southern Europe in the Cretaceous.

**Could Late Cretaceous sauropod tooth morphotypes provide supporting evidence for faunal connections between North Africa and Southern Europe?**

Femke M. Holwerda<sup>\*1,2,3</sup>, Verónica Díez Díaz<sup>4,5</sup>, Alejandro Blanco<sup>2,6</sup>, Roel Montie<sup>1,2,3</sup> and Jelle W.F. Reumer<sup>1</sup>

<sup>1</sup> Faculty of Geosciences, Utrecht University, Princetonlaan 8a, 3584 CB Utrecht, The Netherlands;

<sup>2</sup> Staatliche Naturwissenschaftliche Sammlungen Bayerns (SNSB), Bayerische; Staatssammlung für Paläontologie und Geologie, Richard Wagnerstrasse 10, 80333 Munich, Germany;

<sup>3</sup> Department of Earth and Environmental Sciences and GeoBioCenter, Ludwig Maximilians Universität, 80333 Munich, Germany;

<sup>4</sup> Museum für Naturkunde, Leibniz-Institut für Evolutions-und Biodiversitätsforschung, Invalidenstraße 43, 10115 Berlin, Germany;

<sup>5</sup> Humboldt Universität, Berlin, Germany;

<sup>6</sup> Centro de Investigaci3n Científicas Avanzadas (CICA), Facultade de Ciencias, Universidade da Coruña, 15071 A Coruña, Spain

\*Corresponding author: [f.m.holwerda@gmail.com](mailto:f.m.holwerda@gmail.com)

**ABSTRACT**

The Cretaceous Kem Kem beds of Morocco and equivalent beds in Algeria have produced a rich fossil assemblage, yielding, amongst others, isolated sauropod teeth, which can be used in

species diversity studies. These Albian-Cenomanian (~113 – 93.9 Ma) strata rarely yield sauropod body fossils, therefore, isolated teeth can help to elucidate the faunal assemblages from North Africa, and their relations with those of contemporaneous beds and geographically close assemblages. Eighteen isolated sauropod teeth from three localities (Erfoud and Taouz, Morocco, and Algeria) are studied here, to assess whether the teeth can be ascribed to a specific clade, and whether different tooth morphotypes can be found in the samples. Two general morphotypes are found, based on enamel wrinkling and general tooth morphology. Morphotype I, with mainly rugose enamel wrinkling, pronounced carinae, lemon-shaped to (sub)cylindrical cross-section and mesiodistal tapering towards an apical tip, shows affinities to titanosauriforms and titanosaurs. Morphotype II, characterized by more smooth enamel, cylindrical cross-section, rectangular teeth with no apical tapering and both labial and lingual wear facets, shows similarities to rebbachisaurids. Moreover, similarities are found between these northwest African tooth morphotypes, and tooth morphotypes from titanosaurs and rebbachisaurids from both contemporaneous finds from north and central Africa, as well as from the latest Cretaceous (Campanian-Maastrichtian, 83.6Ma – 66.0Ma) of the Ibero-Armorican Island. These results support previous hypotheses from earlier studies on faunal exchange and continental connections between North Africa and southern Europe in the Cretaceous.

## INTRODUCTION

The early Late Cretaceous of northwestern Africa is well-known for its rich vertebrate fauna, many taxa having been described in particular from the Albian–Cenomanian (~113 – 93.9 Ma) Kem Kem beds of Morocco, and the Albian–Cenomanian equivalent continental intercalaire of Algeria. The Moroccan Kem Kem beds include aquatic fauna such as sharks, lungfish,

47 coelacanths, bony fish, amphibians, turtles, crocodylomorphs, as well as terrestrial vertebrates  
 48 such as squamates, pterosaurs, sauropods, and an abundance of theropods (Lavocat, 1954;  
 49 Russell, 1996; Sereno et al., 1996; Wellnhofer & Buffetaut, 1999; Cavin et al., 2010; Richter,  
 50 Mudroch & Buckley, 2013; Läng et al., 2013; Mannion & Barrett, 2013). Despite this large  
 51 diversity, most fossil material consists of isolated elements from theropods and chondrichthyans  
 52 (e.g. *Spinosaurus*, *Carcharodontosaurus*, *Onchopristis*, Läng et al., 2013; C. Underwood  
 53 pers.comm. 2018). Läng et al. (2013) attributed this to the deltaic palaeoenvironment being  
 54 unsuitable for the setting of stable terrestrial vegetation. Because of this, the herbivorous fauna  
 55 has not received much attention thus far, and sauropod material is rare (C. Underwood, pers.  
 56 comm. 2018, but see McGowan & Dyke, 2009). Studies of sauropod material from this region  
 57 thus far found *Rebbachisaurus garasbae*, and other rebbachisaurids (Lavocat, 1954; de  
 58 Lapparent & Gorce, 1960; Russell, 1996; Mannion & Barrett, 2013; Wilson & Allain, 2015) as  
 59 well as several titanosauriform remains, and also a possible titanosaurian (De Broin, Grenot &  
 60 Vernet, 1971; Kellner & Mader, 1997; Mannion & Barrett, 2013; Lamanna & Hasegawa, 2014;  
 61 Ibrahim et al., 2016). De Lapparent & Gorce (1960) also mentioned brachiosaurid finds,  
 62 however, these remains are now considered to be rebbachisaurid or titanosauriform (Mannion,  
 63 2009; Mannion & Barrett, 2013).  
 64 Sauropod body fossils are restricted to mostly isolated elements, or, if associated, material is not  
 65 as numerous as with theropod material (see e.g. Mahler, 2005; Novas, Dalla Vecchia & Pais,  
 66 2005; Cau & Maganuco, 2009). Sauropod teeth, however, are preserved in relative abundance.  
 67 One isolated sauropod tooth had already been reported on by Kellner & Mader, 1997. Sauropod  
 68 teeth are commonly preserved in the fossil record due to their hardness, resilience against  
 69 weathering, and due to their high tooth replacement rates (see e.g. Calvo, 1994; Erickson, 1996;

García & Cerda, 2010). Studying isolated teeth has previously been applied to assessing theropod species diversity in North Africa (Richter, Mudroch & Buckley, 2013). Sauropod teeth can be used for a similar purpose as well, as morphological classifications based on shape, size and position of wear facets (Calvo, 1994; Salgado & Calvo, 1997; Chure et al., 2010; Mocho et al., 2016; Carballido et al., 2017), and enamel wrinkling patterns (Carballido & Pol, 2010; Díez Díaz, Suberbiola & Sanz, 2012; Díez Díaz, Tortosa & Le Loeuff, 2013; Holwerda, Pol & Rauhut, 2015) have classified tooth assemblages into morphotypes or even down to family or genus level (e.g. *Amygdalodon*, *Patagosaurus*).

Mannion & Barrett (2013) suggested that the Cretaceous North African titanosauriforms may not be closely related to southern African forms, as the lineages were cut off from each other by the trans-Saharan seaway. Moreover, close relations are suggested between Cretaceous North African sauropods and Italian sauropods (Zarcone et al., 2010; Dal Sasso et al., 2016), and Iberian sauropods (Sallam et al., 2018; Díez Díaz et al., 2018). More specifically, close relations between Egyptian and European sauropods (Sallam et al., 2018) and between Tunisian and European sauropods (Fanti et al., 2015) have been found. These studies proposed faunal exchanges during the Late Cretaceous between northern Africa and southern Europe. Several migratory routes have been suggested, such as the ‘Apulian route’ during the Early Cretaceous (Dalla Vecchia, 2002; Canudo et al., 2009). Continental connections would have been made possible by peri-Adriatic carbonate platforms in the Mediterranean, connecting North Africa with Adria, throughout the Cretaceous, making migration possible between the northern African and southern European islands and peninsulas (Zarcone et al., 2010). Indeed, these carbonate platforms contain numerous tetrapod footprints, including those of sauropods (Zarcone et al., 2010). The hypothesis of a faunal exchange during the Cretaceous is not new; Late Cretaceous

abelisaurid theropods and titanosaurian sauropods from France are found to have Gondwanan affinities, indicating migration from Gondwana to Europe, an event which could already have taken place in the Early Cretaceous (Buffetaut, Mechin & Mechin-Salessy, 1988; Buffetaut, 1989). Next to sauropods, Early Cretaceous abelisaurid and carcharodontosaurid theropods were found with Gondwanan affinities, as well as other terrestrial fauna, such as amphibians, snakes, and ziphodont crocodyliforms (Le Loeuff, 1991; Vullo et al., 2005; Vullo, Neraudeau & Lenglet, 2007; Pereda-Suberbiola, 2009). Ösi, Apesteguía & Kowalewski, (2010) found Santonian theropods from the Mediterranean region to have both Gondwanan and North American affinities. Dalla Vecchia & Cau (2011) added a notosuchian from the Late Cretaceous of Italy, and Rabi & Sebök, (2015) a sebecosuchian to this faunal assemblage with Gondwanan affinities. Reviewing undescribed North African Cretaceous sauropod material could add information on both the biogeographical patterns of the Euro-Gondwanan area, as well as on sauropod species diversity in northwestern Africa.

Here, we present a morphological and quantitative analysis of a sauropod tooth assemblage from the Cenomanian of Morocco and Algeria. Teeth are categorized into two morphotypes, which are then compared to contemporaneous Cretaceous sauropod tooth morphotypes, including sauropod teeth from Africa and southern Europe.

# Institutional abbreviations:

BSPG: Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany

FAM: Fox-Amphoux-Métisson, France

MB.R.: Museum für Naturkunde, Berlin, Germany

116 MCCM-HUE: Museo de las Ciencias de Castilla-la Mancha, Spain

117 MHN-AIX-PV: Natural History Museum Aix-en-Provence, France

118 MPCA-Pv: Museo Provincial “Carlos Ameghino”, colección de paleovertebrados, Río Negro,

119 Argentina

120 PIMUZ: Palaeontological Institute and Museum, University of Zürich, Switzerland

121

## 122 **GEOLOGICAL SETTING**

123 Fourteen of the teeth studied here are from the Kem Kem beds of Morocco, and four were

124 supposedly found in the Late Cretaceous Continental Intercalaire of Algeria. Four of the

125 Moroccan sample are labeled as originating from ‘Taouz, Algeria,’ and four are labeled ‘Kem

126 Kem Morocco’. Taouz, Algeria, might actually mean Taouz, Morocco, which is the southern part

127 of the Kem Kem beds, and ten samples are labeled as originating from Erfoud, Morocco, which

128 is the more northern part of the Kem Kem beds (Figure 1). The Kem Kem area is located in the

129 south-east of Morocco (Figure 1). Here, the Kem Kem beds form an escarpment around the

130 eastern end of the Anti-Atlas, from near Goulmima in the northwest (C. Underwood pers. comm.

131 2018; see Figure 1) past Erfoud in the north (Wellnhofer & Buffetaut, 1999; Cavin & Forey,

132 2004) and along the east, paralleling the Algerian border (C. Underwood pers. comm. 2018).

133 The Kem Kem ends to the west, south of Taouz, thereby in total stretching to about 300 km of

134 outcrop (C. Underwood pers. comm. 2018; see Figure 1). The Kem Kem is usually mentioned to

135 be Cenomanian in age, as it has been found to match ammonites from the Lower Cenomanian of

136 Bahariya, Egypt (Le Loeuff et al., 2012). However, the age range could span over the Albian–

137 Cenomanian (C. Underwood, pers. comm. 2018), which is closer to the age given to the Algerian

138 Cretaceous Continental Intercalaire (Lefranc & Guiraud, 1990; Le Loeuff et al., 2012) and to the



139 fossil-rich Cretaceous ‘Continental Intercalaire’ beds of Tunisia (Fanti et al., 2016) as well as  
 140 sauropod bonebeds from Niger (Sereno & Wilson, 2005). The Kem Kem beds are considered to  
 141 be made up of two formations (see Figure 1): the fossil-rich lower Ifezouane Formation and the  
 142 upper Aoufous Formation, rich in ichnofossils (Cavin et al., 2010; Belvedere et al., 2013); also  
 143 named the lower sandy unit (a braided fluvial system) and the upper marly unit (a coastal  
 144 lagoon), respectively (Cavin et al., 2010, Belvedere et al., 2013, Mannion & Barrett, 2013;  
 145 Ibrahim et al., 2014). Practically all fossil vertebrates originate from the lower Ifezouane  
 146 Formation (Cavin et al., 2010, C. Underwood pers. comm. 2018). The Continental Intercalaire of  
 147 Algeria is less studied than the Kem Kem, and the age ranges from Barremian to Turonian.  
 148 However, most authors set the age of the beds close to the Moroccan border (where our Algerian  
 149 specimens are supposedly from), to Albian-Cenomanian-Turonian, with the Cenomanian layers  
 150 being the most fossil-rich (Läng et al., 2013; Benyoucef et al., 2015; Meister et al., 2017). As  
 151 said before, the labeling, however, of ‘Taouz, Algeria’ is most likely not correct, as Taouz is  
 152 situated in Morocco, south of Erfoud, close to the border with Algeria, where many fossils from  
 153 the southern Kem Kem exposures, towards Ouzina, are collected over a broad expanse, usually  
 154 in mines around Bagaa (M. Dale, C. Underwood, pers. comm. 2018; Figure 1). Indeed, Taouz,  
 155 Morocco, is indicated as fossil locality in other Kem Kem fossil vertebrate studies (e.g.  
 156 Wellnhofer & Buffetaut, 1999; Cavin et al., 2010; Forey, López-Arbarello & MacLeod, 2011;  
 157 Richter, Mudroch & Buckley, 2013). As the Kem Kem outcrops at present run parallel to the  
 158 border, the labelling of ‘Algeria’ is probably still be correct, as the ‘Kem Kem’ beds extend out  
 159 across the border (Alloul et al., 2018), however, the specific provenance of these teeth is unclear.  
 160 Next to the labelling, the colour of the fossils likely confirms the provenance on the labels, as  
 161 fossils from Bagaa (Taouz) are chocolate brown in colour, and fossils from north of Erfoud show

a range of colours, usually shades of beige and black (C. Underwood, pers. comm. 2018), matching the provenance on the collection reference (see Description). As most of the fossils retrieved from the Kem Kem and equivalent beds from Algeria are found via mining, the provenance is unfortunately usually unclear and only traceable to a regional provenance (Forey & Cavin, 2007; Rodrigues et al., 2011; C. Underwood, pers.comm).

## MATERIALS AND METHODS

In this study, eighteen isolated sauropod teeth from the Kem Kem beds and Continental Intercalaire are studied for shape, size, position of wear facets (where applicable) and enamel wrinkling. The four teeth from around Taouz (Morocco) are BSPG 1993 IX 331A, BSPG 1993 IX 331B, BSPG 1993 IX 331C, and BSPG 1993 IX 313A, see Figure 2A-D. The ten teeth from Erfoud (Morocco) are labeled PIMUZ A/III 0823, and are given the additional labeling of a, b, c, etc., for convenience, see Figure 3A-J. The four Algerian specimens are BSPG 1993 IX 2A, BSPG 1993 IX 2B, BSPG 1993 IX 2C, and BSPG 1993 IX 2D, see Figure 2E-H. Measurements were taken with a caliper to mm scale. For imaging, the teeth were photographed using normal and macro settings. Scanning Electron Microscopy (SEM) pictures were taken of the Munich sample at the Zoologisches Institut in Munich to obtain a detailed view of the enamel wrinkling patterns. SEM images were not possible to obtain for the Zürich sample. The specimens were examined using a LEO 1430VP SEM. In this study, the proposed dental orientations of Smith and Dodson (2003) are followed. The Slenderness Index (SI, sensu Upchurch, 1998) was measured for each crown tooth by dividing the apicobasal length by the mesiodistal width in the middle of the crown. The Compression Index (CI, sensu Díez Díaz, Tortosa & Le Loeuff, 2013)

was measured for each tooth by dividing the labiolingual width by the mesiodistal width in the middle of the crown. The abrasion stages of the functional dentition proposed by Saegusa & Tomida (2011) and Wiersma & Sander (2016) will be assessed for each tooth in order to assess possible tooth position in the toothrow. The angles of the wear facets were measured with respect to the labiolingual axes of the teeth. See Table 1 for all tooth measurements.

The studied sample was compared with teeth from other sauropod taxa in a quantitative approach by multivariate analysis. The comparative tooth sample is measured by first hand observations (i.e., the Lo Hueco, Massecaps, *Ampelosaurus*, Fox-Amphoux-Métisson sample, *Patagosaurus*, *Lapparentosaurus*) as well as from literature, where SI and CI were reported or could be confidently measured or estimated, and a minimum sample size of three tooth specimens was reached (Upchurch & Barrett, 2000; Barrett et al., 2002; Carpenter & Tidwell, 2005; Apesteguía, 2007; Freire, Medeiros & Lindoso, 2007; Díaz et al., 2012a,b; Díez Díaz, Tortosa & Le Loeuff, 2013; Díez Díaz, Ortega & Sanz, 2014; Holwerda, Pol & Rauhut, 2015; França et al., 2016; Averianov & Sues, 2017). Taxa from different periods and palaeobiogeographic origins were included where possible. A total of 102 teeth were grouped by taxon, if possible, or sorted by morphotype, finally creating 17 different groups. Both damaged/worn as well as unworn teeth were used, due to paucity of sample size. See Supplementary Table for this data. Differences in SI and CI ratios were tested amongst groups through statistical analyses. The shape of the cross section and the number, angle and size of the wear facets were not considered for this purpose because these features may be more related to other functional factors (i.e., tooth position, stage of tooth wear) rather than a taxonomic factor. Due to the small sample size, the non-normal distribution and the non-homoscedastic variances amongst sample groups, the non-parametric

multivariate one-way PERMANOVA test was performed, followed by post-hoc tests assessing differences for each pair of groups (Hammer & Harper, 2006). The analysis was implemented using PAST v3.20 (Hammer, Harper & Ryan, 2001). Finally, the tooth groups were depicted in a dispersion plot, together with additional taxa for which sample size was too small to be included in the statistical analyses (see Discussion).

## RESULTS

### Morphological description

#### Moroccan sample: Taouz

##### BSPG 1993 IX 331A (Figure 2A)

The crown of this chocolate- to reddish-brown coloured, apicobasally elongated tooth is more or less cylindrical. It tapers towards the apex, both mesiodistally and labiolingually. The labial side is strongly convex, the lingual side is straight to concave. The convexity increases towards the apex as the distal 1/3rd bends more strongly towards the lingual side. The tooth has an almost circular cross section at the base of the crown, becoming slightly more flattened in the labiolingual direction apically. It has a SI of 4.31 and a CI of 0.85 (see Table 1). Two distinct wear facets are present on the lingual and the apical side of the tooth (F2 abrasion stage). The lingual wear facet has an angle of almost 90 degrees with respect to the labiolingual axis. As no mesial and distal wear facets are present this tooth was probably located anteriorly in the upper jaw (premaxilla). A polished surface is found on the labial side of the crown. Either damage or wear is present on the mesial and distal edges on the carinae, exposing the dentine. The enamel

wrinkling pattern is more pronounced on the labial side than on the lingual side of the tooth (see Figure 4A-B). On the labial as well as the lingual side, the pattern is more pronounced in the middle of the tooth, and fades out slightly toward the apex and the base. The labial enamel wrinkling pattern consists of frequently anastomosing, sinuous grooves and crests of varying width with a general apicobasal orientation. Grooves and crests are discontinuous; crests are often interrupted by pits and islets. The crests are rounded to triangular in shape. The distribution of crests and grooves is roughly equal. Compared to the other teeth, excepting BSPG 1993 IX 331B, the crests protrude sharply, and the grooves are relatively deep. On the lingual side, more pits are present, and the grooves and crests appear slightly less rounded in shape, but retain their apicobasal orientation. The grooves appear more shallow on the lingual side.

#### BSPG 1993 IX 331B (Figure 2B)

This unworn tooth has the same colour and enamel texture as BSPG 1993 IX 331A. It curves towards the lingual side, with the labial side slightly more convex on the upper half, resulting in a labiolingually tapering apex. The tooth crown is convex toward the distal side, tapering to a mesiodistally narrow apex. The tooth is generally distally inclined, however the apex curves slightly towards the mesial side. It has an oval cross section at the base, which becomes “lemon-like” (sensu Díez Díaz et al., 2013) at the apex due to the presence of pronounced carinae on the mesial and distal edges. Its SI is 2.61 and the CI 0.77, see Table 1. The carina on the distal side is slightly more pronounced and continues further basally than the one on the mesial side. The apex contains a polished surface on the mesial side.

The wrinkling is similar to BSPG 1993 IX 331A, however, due to the unworn state of BSPG 1993 IX 331B, it is more pronounced. The enamel wrinkling pattern consists of sharply

protruding, angular, narrow and discontinuous grooves (see Figure 4C-D). The wrinkling on the lingual side is more pronounced than the labial side, and also appears slightly more rounded. Although it is fully developed, the crown lacks wear facets, and the root does not seem to have any resorption, as in the F1 abrasion stage proposed by Wiersma & Sander (2017). This, together with the coarse enamel pattern of the crown (Figure 4C-D), with no signs of abrasion by occlusion, indicates that this was probably a recently erupted (unused) tooth. Due to this we cannot hypothesize the placement of this tooth in the jaw. This specimen differs from the other teeth in this study by the strong curvature of the crown, and its low SI ratio. This can be attributed to a distal (posterior) placement of the tooth in the jaw, as for example is seen in the nearly complete tooth row of *Giraffatitan* (MB.R.2181.21), *Sarmientosaurus* (Martinez et al., 2016), and *Abydosaurus* (Chure et al., 2010). In addition, and because of its size, it probably belonged to a juvenile individual.

#### BSPG 1993 IX 331C (Figure 2C)

This chocolate to reddish-brown coloured tooth differs from the first two Kem Kem teeth, in that it is rather straight both labially and lingually, shows little tapering and has a distinctive enamel wrinkling pattern. The apex shows tapering due to labial and lingual wear facets, and only a slight apical mesiodistal tapering is present. A slight curvature towards the lingual side is present. The base and middle of the tooth is oval in cross section; the apex shows a slightly more elliptical shape. It has a SI of 4.11 and a CI of 0.83, see Table 1.

Four wear facets are present on the tooth, one on each of the labial, lingual, mesial, and distal surfaces, respectively. The lingual wear facet is angled at around 60 degrees with respect to the labiolingual axis. The labial wear facet is angled at almost 90 degrees with respect to the

labiolingual axis. Both the mesial and distal wear facets are more pronounced on the lingual side, and are positioned basal to the lingual wear facet. They appear almost parallel to the tooth's main axis. The number and development of the wear facets indicate that this tooth is between the F4 and F5 abrasion stages, with an important tooth-to-tooth contact. The food-to-tooth contact seems to not be as important, as the crown enamel is not as worn as in other specimens of this sample (see e.g. BSPG 1993 IX 313A). Due to the higher development of the lingual wear facet and the placement of the mesial and distal ones, this tooth was probably located in the maxilla. The enamel of BSPG 1993 IX 331C is ornamented with thick mesiodistally oriented sharply protruding ridges and relatively wide grooves. At about two thirds of the apicobasal height, the grooves slope towards the base from the mesial and distal edges at an angle of about 45 degrees, and meet in the midpoint. Towards the base, the ridges become more horizontally positioned. Some grooves seem to be connected, forming a chevron-like morphology. This wrinkling pattern is also seen in PIMUZ A/III 0823j. Some pits are visible in the apex, probably due to a diet with some grit content.

The four wear facets of BSPG 1993 IX 331C are not seen in similar shapes in any of the other teeth. Moreover, the deep grooves of the enamel wrinkling on the labial and lingual sides of the tooth do not resemble any taphonomic patterns as described by King et al. (1999), but this does not rule out taphonomic processes completely. However, the peculiar pattern is also seen in PIMUZ A/III 0832j, therefore, it is probably natural.

#### BSPG 1993 IX 313A (Figure 2D)

The upper half of this chocolate-brown coloured, worn tooth crown is inclined towards the lingual side, with the labial side curving convexly, and the lingual side curving concavely. The

apex tapers mesiodistally, and the mesial side of the tooth inclines distally, creating a convex mesial apical end. The tooth has an oval cross-section at the base, becoming more “lemon-like” (sensu Díez Díaz et al., 2013) towards the apex due to the presence of protruding carinae on the mesial and distal edges. The carina on the distal edge continues further towards the base than the carina on the mesial edge. Its SI and CI are 3.35 and 0.76, respectively, see Table 1. The apex contains one wear facet on the lingual side, angled at around 75 degrees with respect to the labiolingual axis of the tooth. The presence of only one wear facet, the occlusal one, groups this tooth within the F2 abrasion stage. Due to this, this tooth was probably placed anteriorly in the upper jaw (premaxilla). The enamel appears smooth, except for thin apicobasally-oriented discontinuous grooves, (see Figure 4E-F).

# Moroccan sample: Erfoud

## PIMUZ A/III 0823a (Figure 3A)

This reddish-brown worn tooth crown is apicobasally straight in lingual and labial view. Only at the apex, in labial view, does the tip taper very slightly to the distal side. In both mesial and distal views, the tooth is seen to curve towards the lingual side, showing a moderate convexity on the labial side, and an equally moderate concavity on the lingual side; also seen in BSPG 1993 IX 331A. The width at the base is only slightly higher than at the middle of the crown, showing a moderate mesiodistal tapering towards the apex. The cross-section of the tooth is oval to elliptical at the base, to oval to possibly lemon-shaped at the apex. The carinae are not pronounced on this tooth, therefore the apical lemon-shaped cross-section is not clear. The



carinae on both sides of the tooth appear worn. This tooth has a SI value of 4.92 and a CI value of 0.58, see Table 1.

Labiolingual tapering towards the apex is caused by the presence of a round to oval lingual wear facet (meaning a F2 abrasion stage). This could mean that this tooth could have been a premaxillary one.

The enamel wrinkling pattern is not well-preserved, possibly due to abrasion of the tooth. Only faintly reticulate wrinkling is seen on mostly the labial side of the tooth. This pattern could be a worn enamel wrinkling type similar to that of the more unworn BSPG 1993 IX 331A and BSPG 1993 IX 331B. No protruding ridges or deep prominent grooves are present in the enamel wrinkling pattern of this tooth.

### PIMUZ A/III 0823b (Figure 3B)

This worn, grey to beige coloured tooth displays a unique morphology amongst the tooth sample, as it is a relatively apicobasally short, mesiodistally wide tooth, with a D-shaped cross-section, representing more a general non-neosauropod eusauropod tooth shape (though this feature also exists in basal titanosaurs); Barrett and Upchurch, 2005; Holwerda et al., 2015). In labial and lingual view, the tooth is apicobasally straight at its lower half, showing a similar mesiodistal width at the base as at the middle. The tooth then tapers mesiodistally at its upper half, and also tapers slightly towards the distal side. Moreover, in lingual view, the upper half of the crown, towards the apex, is slightly mesiodistally constricted, giving the tooth a pear-shaped appearance. In mesial and distal view, the tooth curves towards the lingual side, creating a convex labial and concave lingual side. The cross-section is oval at the base, to D-shaped at the apex. The SI is 2.38 and the CI 0.69, see Table 1.

Both carinae are worn, exposing dentine on both sides, and a prominent oval wear facet is present on the lingual side (F2 abrasion stage). This tooth could have been a premaxillary one. Another worn surface is present on the apex of the labial surface, where the enamel is worn away to expose the dentine. The enamel wrinkling consists of rugose, protruding, continuous anastomosing ridges, which are flattened by wear and/or abrasion throughout. Between these ridges, discontinuous grooves and pits are visible.

### PIMUZ A/III 0823c (Figure 3C)

The crown of this well-preserved, unworn tooth is black, whilst the root is cream-coloured to beige. The crown is more or less apicobasally straight in labial and lingual view, however a faint convexity is seen on the medial side, showing a curvature of the entire tooth, from root to apex, towards the distal side. Towards the apex the crown tapers sharply, creating a triangular apical tip. In mesial and distal view, there is only a faint convexity visible on the labial side, whereas the lingual side remains apicobasally straight. The cross-section at the base is oval, and the cross-section at the apex is lemon-shaped (sensu Díez Díaz, Tortosa & Le Loeuff, 2013). It presents a SI of 3.78 and a CI of 0.67, See Table 1.

As no wear facets are present, this tooth has a F1 abrasion stage. No more information about its placement in the jaw can be deduced. Both carinae are well-preserved as sharply protruding ridges, where the distal carina is more pronounced than the mesial, and shows a faint sinusoidal curvature. Furthermore, the distal carina shows ridges inclining from the apicobasally oriented wrinkling towards the carina, creating denticle-like structures, or pseudodenticles (Bonaparte, Riga, Apesteguia 2006). The enamel wrinkling consists of rugose, discontinuous, sinusoidal,

rounded protrusions and islets, which are intersected by shallow but broad anastomosing grooves and pits. The enamel is most pronounced on the lingual side, whilst the labial side shows a polished surface on the upper half of the crown, where the wrinkling is worn away. This tooth was probably used before it was lost, despite the lack of wear facets, because of its labial polished surface (indicating food-to-tooth contact).

#### PIMUZ A/III 0823d (Figure 3D)

This reddish to chocolate-brown, worn and polished tooth crown is apicobasally straight in labial and lingual view. In labial view, the tooth is more rectangular in shape, whilst in lingual view, the mesiodistal width increases from the base towards the middle, after which it tapers towards the apex. The apex of the crown inclines slightly towards the distal side, especially in lingual view. In labial view, two grooves are present, the distal of which is slightly displaced towards the middle of the tooth. In mesial and distal view, the labial side is mildly convex, and the lingual side is more apicobasally straight. The cross-section from base to apex remains similar, being oval to elliptical, without protruding carinae on mesial and distal sides. The SI is 2.33 and the CI 0.67, see Table 1.

Symmetrical, oval wear facets are present on the lingual and labial apices. The presence of both labial and lingual wear facets in the same tooth is an interesting feature that has not been considered in the five abrasion stages (Wiersma & Sander, 2017). García and Cerda (2010) described several apical wear patterns from a sample of titanosaurian teeth from Argentina, and the morphology of the crown and wear facets of PIMUZ A/III 0823d is highly similar to the tooth MPCA-Pv-55 (fig. 5.4). The apex of these teeth (with both labial and lingual apical wear facets) is usually straight and perpendicular to the apicobasal axis of the crown, and the

inclination and development of both wear facets is normally different. García & Cerda (2010) suggested that the presence of both a labial and a lingual wear facet on the same tooth could be related to the position of the teeth on the tooth row, the occlusal pattern, and the replacement of the opposite teeth (see García & Cerda, 2010, fig. 7, for a more graphic explanation of this hypothesis). The absence of mesial and distal wear facets suggests that it was probably anteriorly located in the dentary or (pre)maxilla (it is not possible to be more accurate, as both wear facets are equally developed). The mesial and distal sides of the apex do not taper, which further adds to the rectangular appearance of the tooth. The mesial and distal carinae are worn down to smooth, polished ridges, however, the dentine is not exposed.

The enamel wrinkling pattern is mostly polished away by (food-to-tooth or taphonomic) abrasion, and only at the base of the labial and lingual sides a faint wrinkling is present. This consists of rounded, anastomosing protrusions, giving a pebbly textural shape, with wide but shallow grooves and pits in between. At the base of the carinae, the wrinkling becomes slightly more rugose, showing angular protrusions angling at ~30 degrees with respect to the apicobasal axis, towards the carinae, with deep wide grooves running parallel to these.

#### PIMUZ A/III 0823e (Figure 3E)

The crown of this greyish-brown tooth tapers gently towards the apex in labial and lingual view, and inclines slightly towards the distal side. In labial and lingual view, the tooth crown is apicobasally straight. In mesial and distal view, the tooth is labially strongly convex, but lingually only mildly concave. The apex tapers to a point labiolingually. The cross-section at the base is elliptical to cylindrical, whilst the apex is more D-shaped to elliptical. It has a SI of 4.11

and a CI of 0.78, see Table 1. Both carinae are worn, showing marginal grooves where dentine almost appears through the enamel.

A large oval wear facet is present on the lingual apical side. However, the labial side also shows a less pronounced wear facet, symmetrically placed to that of the lingual side. This tooth was probably placed anteriorly in the upper part of the mouth (possibly premaxilla), as can be deduced from the more developed lingual wear facet and the absence of mesial and distal facets. The poor development of the labial wear facet indicates that its lower new opposite tooth probably erupted only shortly prior to its loss. We consider a F2 abrasion stage to be appropriate for this tooth, due to the poor development of the labial wear facet, and its placement in the premaxilla (because of the absence of mesial and distal wear facets).

The enamel wrinkling pattern is finely reticulate, with a symmetrical distribution of small rounded protrusions surrounded by thin grooves. However, at the base of the crown, the enamel shows deep continuous grooves interspersed with pits, whilst the protrusions are worn away. Part of the enamel at the root is damaged or gone.

#### PIMUZ A/III 0823f (Figure 3F)

This black, smooth, tooth crown shows traces of red sediment. In labial and lingual views, the lower half of the tooth crown is straight, after which it shows a strong inclination towards the distal side, giving the mesial apical half a convex, and the distal apical half a straight to concave, shape. In mesial and distal views, the tooth is rather apicobasally straight, only a faint labial convexity and lingual concavity is seen. The cross-section at the base is round to elliptical, and more cylindrical to a very faint lemon-shape (due to the carinae) at the apex. Apically, the tooth tapers to a sharp tip which is inclined distally. The SI is 4.75 and the CI is 0.88. Both mesial and

distal carinae show a prominent but smooth ridge protruding towards mesial and distal sides on the apical half of the crown, the lower half of the crown does not show carinae.

As no apical wear facets are present we consider this tooth to show a F1 abrasion stage. Due to this, the placement of this tooth in the snout cannot be hypothesized. The enamel is completely smooth, suggesting, together with the lack of clear wear facets, that this tooth was relatively unused and unworn.

#### PIMUZ A/III 0823g (Figure 3G)

This grey to reddish brown tooth crown follows the same general morphology as PIMUZ A/III 0823f, in that the lower half of the crown is apicobasally straight in both labial and lingual views, as well as in mesial and distal views. The upper half of the crown bends towards the distal side, providing the mesial side with a convex curvature. The apex tapers to a sharp tip in labial and lingual view, as well as in mesial and distal view. Both mesial and distal carinae are present, which are prominently but smoothly protruding at the apical half of the crown. The cross-section is elliptical at the base to strongly lemon-shaped at the apex. This tooth has a SI value of 4.83 and a CI value of 0.83, see Table 1.

An apicobasally-elongated, mesiolabially-oriented wear facet, as well as an equally elongated, distolingual wear facet is present, giving the apical half of the crown a more constricted shape, and creating a faint buttress at the base of the mesial and distal constriction. Finally, a faint, small, round apical wear facet is present. The presence of these three wear facets place this tooth in a F4 abrasion stage, as the apex is already gently rounded. This tooth probably was placed in the maxilla. The slight displacement of both mesial and distal wear facets indicate that this tooth

was rotated. The enamel is completely smooth, however, apicobasal striations in the enamel are present on the labial apical side of the tooth.

#### PIMUZ A/III 0823h (Figure 3H)

The smallest tooth of the sample, this reddish-brown crown shows a similar morphology to PIMUZ A/III 0823f and PIMUZ A/III 0823g. The crown is straight in labiolingual and mesiodistal views, with only the mesial apical end inclining towards the distal side in labiolingual view. The apex tapers to a sharp tip. The cross-section is elliptical to cylindrical at the base and at the apex. The SI is 4.33 and the CI is 0.75, see Table 1. The tooth is completely smooth and nearly unworn; only a small, circular, apical wear facet is present (F1-F2 abrasion stage). This information is not sufficient to propose its likely placement in the snout. Due to its small size and unworn state, this tooth might originate from a juvenile.

#### PIMUZ A/III 0823i (Figure 3I)

This rectangular, red tooth is the second smallest of the Erfoud sample. It has a rectangular shape, showing almost no tapering in labial and lingual view, and only a slight apical tapering in mesial and distal view. In mesial and distal view, the labial side is mildly convex and the lingual side is mildly concave. The cross-section is oval at the base to oval to lemon-shaped at the apex. It has a SI of 3.82 and a CI of 0.73, see Table 1. The labial apical side shows a high-angled, oval wear facet, whilst a low-angled elongated wear facet is present on the lingual side, which stretches to halfway down the apicobasal length of the crown. This tooth presents the same condition as PIMUZ A/III 0823d, with both labial and lingual apical wear facets. As the apical labial wear facet is more developed, this tooth may have

been placed in the lower jaw. The absence of mesial and distal wear facets suggests that it was probably anteriorly located in the dentary. The enamel of the tooth is completely smooth.

#### PIMUZ A/III 0823j (Figure 3J)

This dark brown to black tooth tapers gently towards the apex from the base, whilst being slightly mesiodistally constricted at around the middle. The labial side is convex, whilst the lingual side remains apicobasally straight. Both mesial and distal carinae are present as smooth ridges which do not protrude. The cross-section is elliptical, both at the base as well as at the apex. It presents a SI value of 3.78 and a CI value of 1.11, see Table 1.

The upper half of the tooth crown is polished. However, only on the lingual side, a low angled wear facet is present. Both at the mesial and distal apical sides, a low-angled, oval, polished surface is present as well, indicating tooth overlapping (F4 abrasion stage, as the crown apex is rounded). The location and development of these three wear facets indicate that this tooth was probably located in the maxilla.

The enamel wrinkling pattern is only visible at the base of the tooth, as, similar to BSPG 1993 IX 331C, there is a mesiodistally-positioned, chevron-like pattern of deep wide grooves and ridges, which disperse towards the carinae.

#### **Algerian sample**

#### BSPG 1993 IX 2A (Figure 2E)

The tooth is covered in black enamel, and has a reddish-brown base. The crown is apicobasally straight. The mesial and distal sides taper strongly towards the apex, whereas the labial and lingual sides taper slightly. The mesial side, however, shows a slight convexity, and curves



towards the distal side, which is straight apically. The tooth is mesiodistally widest at the middle, after which it tapers towards the apex. Carinae are present on the mesial and distal edges of the upper third of the crown, as smooth protruding ridges. The carina on the distal side shows a slightly sinusoidal curvature at about halfway of the apicobasal length of the tooth, which is also seen in PIMUZ A/III 0823c. The distal carina reaches slightly further basally. The cross-section at the base is oval, becoming lemon-like at the apex due to the carinae (sensu Díez Díaz, Tortosa & Le Loeuff, 2013). The SI is 4.35 and the CI 0.78, (see Table 1).

A wear facet is present on the lingual side, angled at around 50 degrees with respect to the labiolingual axis. The presence of only one wear facet, the occlusal one, groups this tooth within the F2 abrasion stage. As this facet is located lingually, and there are no mesial and distal wear facets, this tooth was probably placed in the maxilla.

The apical part of the labial side also shows a polished surface, as in PIMUZ A/III 0823c. The enamel wrinkling pattern on both the labial and lingual sides consists of apicobasally-oriented grooves and ridges, which are less broad and more sinuous on the labial side than on the lingual side (see Figure 4G-H). While unworn, was probably similar to the crown enamel of BSPG 1993 IX 2B (Figure 4I-J), but it has been worn by tooth-to-food contact (like BSPG 1993 IX 313A, Figure 4E-F). The enamel on the labial side is slightly thicker than on the lingual side.

BSPG 1993 IX 2B (see Figure 2F)

This worn tooth is grey to beige in colour. The tooth is curved lingually, with the labial side convex, and the lingual side concave. This curvature increases towards the apex. A very slight mesiodistal tapering can be seen at the apical part of the crown, however the tooth appears rectangular when observed from the labial and lingual side. Labiolingual tapering appears due to

the presence of a labial and lingual wear facet intersecting at the apex. The lingual side of the tooth is relatively flat in the mesiodistal direction when compared to the labial side. This gives the tooth an oval cross section at the base, becoming D-shaped toward the centre, and more cylindrical to lemon-shaped apically due to the presence of carinae. Carinae are present on the mesial and distal edges, although the distal carina is more pronounced and continues further basally. It has a SI of 3.65 and a CI of 0.71, (see Table 1).

Two distinct wear facets are present on BSPG 1993 IX 2B, one on the labial and the other on the lingual side. The labial wear facet is larger (almost 15 mm in apicobasal length), and angled toward the mesial side. The labial wear facet is angled at around 72 degrees with respect to the labiolingual axis of the crown. The lingual wear facet is smaller. It cuts the labiolingual axis at almost 90 degrees. This tooth presents the same condition as PIMUZ A/III 0823d and PIMUZ A/III 0823i, with both labial and lingual apical wear facets. As the apical labial wear facet is more developed in BSPG 1993 IX 2B, this tooth may have been placed in the lower jaw. The absence of mesial and distal wear facets suggests that it was probably anteriorly located in the dentary.

The enamel wrinkling pattern consists mainly of apicobasally-oriented, sinuous grooves and ridges. The ridges are sharply protruding and are triangular in shape (see Figure 4I-J). The enamel wrinkling pattern of BSPG 1993 IX 2B appears quite similar to BSPG 1993 IX 331A, BSPG 1993 IX 331B, and the labial side of BSPG 1993 IX 2A. The enamel wrinkling of BSPG 1993 IX 331A and of BSPG 1993 IX 331B is slightly more pronounced than the enamel wrinkling of BSPG 1993 IX 2B, a difference perhaps caused by greater wear on the latter. A noteworthy difference between the enamel wrinkling patterns of this tooth and all other teeth from this

sample is that the enamel wrinkling is more pronounced on what seems to be the lingual side, instead of the labial side.

#### BSPG 1993 IX 2C (Figure 2G)

This grey to beige coloured tooth is badly preserved and largely covered with reddish sediment. The crown is fairly straight apicobasally, and tapers apically (mesiodistally as well as labiolingually). The mesial apical side shows a curvature towards the distal side, as seen in BSPG 1993 IX 2A. From the base to the middle, the width of the crown expands slightly labiolingually, after which it tapers to the apex. This expansion mainly seems to occur on the convex (probably labial) side, and is also seen in BSPG 1993 IX 2A. The distal edge shows faint traces of a carina but that cannot currently be accurately determined. The cross-section at the base is slightly oval, becoming almost circular at the middle of the crown, and then becoming more lemon-like apically due to the presence of carinae. The SI is 4.09 and the CI is 0.87, (see Table 1).

A possible wear facet can be seen on the labial side, cutting the labiolingual axis of the crown at a low angle (35 degrees), so it was probably at a F2 abrasion stage. The absence of mesial and distal wear facets, and the labial position of the one present, indicate that this tooth probably belonged to the anterior part of the dentary. Because of the damaged state of this tooth, no SEM pictures were taken.

#### BSPG 1993 IX 2D (Figure 2H)

This reddish-brown coloured crown tapers mesiodistally towards the apex, as well as labiolingually, resulting in a sharp apical tip. The labial side is convex, whilst the lingual side is

more concave. The upper third of the crown shows stronger mesiodistal tapering on the mesial side. Distinctly protruding carinae are present on the mesial and distal edges of the crown. The mesial carina is slightly more distinct due to the curvature of the mesial apical part, however, it only runs halfway along the tooth towards the base, whereas the distal carina continues further basally, to the upper 3/4th of the tooth. The tooth has an oval cross section at the base, becoming strongly lemon-shaped apically due to the distinct carinae. It presents a SI of 3.62 and a CI of 0.71, (see Table 1).

Wear facets are not present (F1 abrasion stage), so its position on the snout cannot be deduced. The enamel is smooth except for some pits (see Figure 4K-L). This feature is interesting, not only because this tooth is the one that has the most worn crown enamel, but also for the presence of pits on it. The individual probably fed on low vegetation, with a significant quantity of grit. This tooth had clearly been used for feeding, but occlusion features are not present, so it was probably located anteriorly in the snout, with its opposite tooth unerupted throughout its functional life.

## DISCUSSION

### *Systematic discussion and comparisons*

The tooth sample from northwestern Africa shares the presence of mesial and distal margins extending parallel to each other along almost the entire length of the crown with neosauropod (diplodocoid and titanosauriform) sauropods, together with the absence of a mesiodistal expansion at the base of the crown (Calvo, 1994; Upchurch, 1995, 1998; Salgado & Calvo, 1997; Wilson & Sereno, 1998; Upchurch & Barrett, 2000; Barrett et al., 2002; Wilson, 2002;

Upchurch, Barrett & Dodson, 2004). These teeth also share with diplodocoids and titanosaurs the loss of some plesiomorphic features of Sauropoda, which are retained in some, but not all, basal titanosauriforms (e.g. *Giraffatitan*, *Euhelopus*), such as the presence of a lingual concavity with a median ridge, and labial grooves (Upchurch, 1998; Barrett et al., 2002).

The general crown outline is similar in all the teeth of the sample: parallel-sided crowns showing slight labiolingual compression with mesial and distal carinae, which express a higher amount of protrusion in BSPG 1993 IX 331B, BSPG 1993 IX 2A, BSPG 1993 IX 2D, PIMUZ 0823c and PIMUZ A/III 0823h. The labiolingual compression (seen in apical view) is more conspicuous in the Algerian teeth, though PIMUZ A/III 0823a from Erfoud, Morocco, also shows this amount of compression. BSPG 1993 IX 331B and BSPG 1993 IX 2D, PIMUZ A/III 0823d, PIMUZ A/III 0823h, PIMUZ A/III 0823i present a different crown morphology from the rest of the sample (see below). However, this difference in the crown morphology between these teeth and the rest of the sample could be due to different positions in the tooth row, as occurs in most eusauropods and other sauropodomorphs (e.g. Carballido & Pol, 2010; Chure et al., 2010; Holwerda et al., 2015; Martinez et al., 2016; Mocho et al., 2016, Carballido et al., 2017; Wiersma & Sander, 2017). Lingual buttresses seen in titanosauriforms from the Lower Cretaceous of Japan (Barret et al., 2002), South Korea (Lim, Martin & Baek, 2001), in *Giraffatitan* (Janensch, 1935), and in *Astrodon* (Carpenter & Tidwell, 2005), or circular bosses as seen in *Euhelopus* (Barrett & Wang, 2007; Poropat & Kear, 2013) a possible macronarian from the Upper Jurassic of Portugal (Mocho et al., 2017), and an unknown (possible euhelopodid) titanosauriform from the Barremian of Spain (Canudo et al., 2002) are not seen in any of our Moroccan or Algerian tooth samples. Neither do the Algerian and Moroccan tooth samples show mesial and distal buttresses

such as in an isolated tooth from the Santonian of Hungary (Ösi, Csiki-Sava & Prondvai, 2017). Moreover, diplodocoid pencil-like or needle-like teeth as in Limaysaurus (Calvo & Salgado, 1995; Salgado et al., 2004) are not found in this sample either. In addition, two further enamel types can be found in the sample: rugosely or reticulately wrinkled (BSPG 1993 IX 331A, 331B, 331C, PIMUZ A/III 0823a,b,c,e, BSPG 1993 IX 2B and 2C) or smooth (BSPG 1993 IX 313A, PIMUZ A/III 0823d,f,g,h,i, BSPG 1993 IX 2A, and 2D) enamel. The differences between more rugose and more reticulate wrinkling in the enamel ornamentation could be due to the wear of the tooth and the diet of the individual animal.

Finally, although the CI is similar in both morphotypes, the SI range is wider in the Moroccan Erfoud sample, whilst remaining within a similar range between the Moroccan Taouz and the Algerian sample.

After the morphological descriptions and similarities, two main morphotypes could be distinguished, based on shared features of shape, expression of wear facets, carinae and enamel wrinkling. One morphotype, however, may harbour further submorphotypes, which are also discussed, however, without further sampling this could not be proven valid. It should also be noted that the morphotypes proposed here might not always reflect biological morphotypes. The morphotypes are here discussed and compared to other sauropods, including, but not restricted to, biogeographically similar and contemporaneous sauropods (e.g. of Spain, France, northwest and central Africa).

***Morphotype I: BSPG 1993 IX 331A, BSPG 1993 IX 331B, BSPG 1993 IX 313A, BSPG 1993 IX 2A, BSPG 1993 IX 2C, BSPG 1993 2D; PIMUZ A/III 0823a, PIMUZ A/III 0823b, PIMUZ***

*A/III 0823c, PIMUZ A/III 0823e, PIMUZ A/III 0823f, PIMUZ A/III 0823g, PIMUZ A/III 0823h*

### **General**

This is the most abundant morphotype in the sample. It consists of teeth with high SI (2,2-4,9) and CI (0,6-0,9), rugose enamel wrinkling (but see discussion on this), prominent mesial and distal carinae, together with a labial convexity, and a slightly distal inclination of the apex, and a subcylindrical to lemon-shaped cross-section. There are many intra-group specific morphological differences within this morphotype, however as teeth both differ morphologically within one toothrow, as well as between upper and lower toothrows (Sereno & Wilson, 2005; Wilson, 2005; Zaher et al., 2011; Holwerda, Pol & Rauhut, 2015; Martínez et al., 2016; Mocho et al., 2016; Wiersma & Sander, 2016), enamel wrinkling as well as the presence of carinae are taken as the main drivers for comparisons (see Carballido & Pol, 2010; Holwerda, Pol & Rauhut 2015). Moreover, enamel wrinkling can be demonstrated to change over ontogeny, with indications that juveniles have smooth, or smoother enamel wrinkling in comparison with adult animals (Fiorillo, 1991, 1998; Díaz et al., 2012; Díaz, Suberbiola & Sanz, 2012; Díez Díaz, Ortega & Sanz, 2014; Holwerda, Pol & Rauhut, 2015). As stated above, this abundant Morphotype I arguably still hosts several (sub)morphotypes, which are also discussed below.

### **Discussion**

BSPG 1993 IX 331A and BSPG 1993 IX 331B (Figure 2A, 2B) show a different SI, and different CI. However, they share a similar morphology in terms of expression of carinae, labial convexity, and enamel wrinkling pattern, and most likely belong to the same morphotype. Similarly, PIMUZ A/III 0823a shares the same apicobasal elongation of BSPG 1993 IX 331A, as

well as a worn-down version of the rugose and highly sinuous enamel wrinkling. This reticulate, worn-down version of the rugose type of enamel wrinkling is further found in BSPG IX 1993 313A, BSPG IX 1993 2B, as well as in PIMUZ A/III 0823b and PIMUZ A/III 0823e. BSPG 1993 331A resembles a large Campanian-Maastrichtian titanosaur tooth from Río Negro, Argentina (Garcia, 2013), in having a similar crown-root transition, as well as having high CI and SI ratios (0.73 vs 0.85 and 3.73 vs 4.36 for the tooth described by Garcia, (2013) and BSPG 1993 IX 331A respectively). However, the enamel wrinkling patterns between these two teeth show great differences; the enamel of BSPG 1993 IX 331A shows highly sinuous patterns not visible on the Argentine tooth. The tooth also resembles the cylindrical morphotype with circular cross-section of the southeastern French Campanian-Maastrichtian Fox-Amphoux-Métisson morphotype (FAM 03.06, 03.11, and 04.17; Díaz et al., 2012b, fig. 9). The SI ratios are high in both the Algerian/Moroccan and the French sample (~4, excepting BSPG 1993 IX 331B). These teeth also display a similar labial convexity, which becomes stronger towards the apex in both the Algerian/Moroccan and the French sample. The enamel wrinkling differs between these two, however, as the Kem Kem teeth show a much more pronounced enamel wrinkling. The apicobasal elongation, slight labial convexity, and presence on most of these teeth of an apically based labial and lingual wear facet, also resembles that of the Late Cretaceous Mongolian *Nemegtosaurus* (Wilson, 2005). Finally, the moderate tapering (or lack thereof) resembles that of teeth of the Late Cretaceous *Alamosaurus* from the USA (Kues, Lehman & Rigby, 1980), although the enamel wrinkling on *Alamosaurus* is smoother than in the Moroccan/Algerian sample, though apicobasal striations are present on both.

Even though the reticulate enamel wrinkling could be a worn-down version of the rugose wrinkling described for this morphotype, BSPG 1993 IX 313A, PIMUZ A/III 0823b and PIMUZ



686 A/III 0823e (Figure 2D, Figure 3B, 3E) slightly deviate from the general shape of this  
 687 morphotype in displaying a mesiodistally constricted apex in labial and lingual view. A  
 688 difference in order of tooth row might explain this, as teeth from the mesial side of the toothrow  
 689 are larger and more robust than teeth from the distal end of the toothrow. This is not seen in  
 690 titanosaurs (García & Cerda, 2010), however, it is observed in titanosauriforms (e.g. *Giraffatitan*  
 691 MB.R.2181.21; *Abydosaurus*, Chure et al., 2010). This shape is further seen in more robust  
 692 (lower SI, higher mesiodistal and labiolingual width) titanosauriform morphotypes, such as the  
 693 Cenomanian titanosauriform indet. tooth from France, described by Vullo, Neraudeau & Lenglet,  
 694 (2007), *Astrodon* (Carpenter & Tidwell, 2005), the more robust tooth form of the Late  
 695 Cretaceous Asian *Mongolosaurus* (Mannion, 2011), *Europatitan* from the Barremian-Aptian of  
 696 Spain (Torcida Fernández-Baldor et al., 2017), the posterior maxillary teeth of the Cenomanian-  
 697 Turonian Patagonian *Sarmientasaurus* (Martínez et al., 2016), *Ligabuesaurus* from the Aptian-  
 698 Albian of Neuquen, Argentina, (Bonaparte, González Riga & Apesteguía, 2006) an unnamed  
 699 titanosaur tooth sample from the Upper Cretaceous Bissekty Fm., Uzbekistan (Averianov &  
 700 Sues, 2016), and lastly, the robust types of the Massecaps and *Ampelosaurus* morphotypes from  
 701 the Campanian-Maastrichtian of southeastern France (Díez Díaz, Tortosa & Le Loeuff, 2013).  
 702 Finally, BSPG 1993 IX 313A, with its strong apical labial convexity, is morphologically similar  
 703 to both the small tooth type of the Cenomanian-Campanian Chinese *Huabeisaurus* (D’Emic et  
 704 al., 2013), as well as the small Campanian-Maastrichtian *Atsinganosaurus* tooth type from  
 705 France by sharing a labiolingual compression (CI: 0.76), a high-angled apical wear facet, and  
 706 lemon-shaped cross section due to the presence of apical carinae (Díez Díaz, Tortosa & Le  
 707 Loeuff, 2013; fig. 3, MHN-AIX-PV.1999.22). It also matches *Huabeisaurus* and  
 708 *Atsinganosaurus* in having smooth enamel. One hypothesis for this smooth enamel could be the

tooth-to-food contact, that could have worn a previously more coarse enamel, like the ones of BSPG 1993 IX 331A and BSPG 1993 IX 331B (Figure 4A-D). However, the enamel of BSPG 1993 IX 313A does not present the conspicuous pits that appear on the enamel of BSPG 1993 IX 2D (Figure 4K-L) from the Algerian sample. This could be due to a diet with a greater quantity of grit in the Algerian individual, as happens in many sauropods, especially the ones that fed in the lower levels of the trees (Fiorillo, 1998; Upchurch & Barrett, 2000; García & Cerda, 2010). The individual of BSPG 1993 IX 313A may have fed on soft vegetation, as has been suggested for *Diplodocus* (Fiorillo, 1998).

Also included in Morphotype I are BSPG 1993 IX 2A, BSPG 1993 IX 2C, PIMUZ A/III 0823c (Figure 2E, 2G, Figure 3C). These teeth deviate from the general cylindrical and general apical tapering shape of Morphotype I in that they all display a mesiodistally slender base, after which the width increases towards the middle, and then tapers towards the rounded apical tip. This is also arguably seen in *Rinconosaurus* (Calvo & González Riga, 2003), as well as in an unnamed titanosaur tooth sample from the Upper Cretaceous Bissekty Formation., Uzbekistan (Averianov & Sues, 2016), although the latter teeth show a lingual median ridge, which the Moroccan and Algerian samples do not. BSPG 1993 IX 2A and BSPG 1993 IX 2C display a similar low angled lingual wear facet (~50 degrees relative to the labiolingual tooth axis), and both BSPG 1993 IX 2A and PIMUZ A/III 0823c show a similar labial polished surface. BSPG 1993 IX 2C is too worn, however the carinae of the well-preserved BSPG 1993 IX 2A and PIMUZ A/III 0823c are prominently present, as mesially and distally offset ridges. Moreover, the distal carina is more developed than the mesial one in both, showing a sinusoidal curvature in distal view, as seen to a lesser extent in *Huabeisaurus* (D'Emic et al., 2013). Note, however, that the carinae are not as prominent as in *Euhelopus* (Barrett & Wang, 2007; Poropat & Kear, 2013). Furthermore, the

distal carina of PIMUZ A/III 0823c shows pseudodenticles, a feature seen in the Late Jurassic South African *Giraffatitan* (Janensch, 1935), the Lower Cretaceous Patagonian *Ligabuesaurus* (Bonaparte, Riga & Apestegui 2006), an unnamed titanosauriform from the Early Cretaceous of Denmark (Bonde, 2012), the Late Jurassic French *Vouivria* (Mannion, Allain & Moine, 2017), and the Early Cretaceous African *Malawisaurus* (Gomani, 2005).

Furthermore, BSPG 1993 IX 2A resembles the morphotype B of Lo Hueco, Spain (MCCM-HUE 2687, (Díez Díaz, Ortega & Sanz, 2014; fig. 5). Both morphotypes are cylindrical, have a high SI ( $>4.3$ ), a strong apical distal or medial inclination of the tooth (giving the tooth a far from straight outline) with a high-angled wear facet. Moreover, the enamel wrinkling of BSPG 1993 IX 2A matches that of the Lo Hueco morphotype, in that both morphotypes show coarse, but not rugose, discontinuous wrinkling, with smooth longitudinal ridges, although BSPG 1993 IX 2A shows more pronounced enamel wrinkling than Lo Hueco morphotype B.

PIMUZ A/III 0823c also resembles Lo Hueco morphotype A, in the general morphology of the crown and its length, its labiolingual compression, and its SI and CI values (see e.g. HUE-685, (Díez Díaz, Ortega & Sanz, 2014, fig. 2).

It could be argued that, due to the differences of BSPG 1993 IX 2A, BSPG 1993 IX 2C, PIMUZ A/III 0823c from the rest of Morphotype I, these teeth could be gathered in a separate morphotype, however, as the sample size is low, with only three teeth, further analysis to help distinguish any (sub)morphotypes is not possible and lies outside of the scope of the current study, until a higher tooth sample can be examined. Therefore, although these teeth are discussed as potentially separate, the most parsimonious conclusion is to leave them in Morphotype I.

Finally, BSPG 1993 2D, PIMUZ A/III 0823f, PIMUZ A/III 0823g, PIMUZ A/III 0823h are assigned to Morphotype I, although the enamel wrinkling differs from the rugose type (both the

worn and unworn expression of it). BSPG 1993 IX 2D, PIMUZ A/III 0823f, PIMUZ A/III 0823g and PIMUZ A/III 0823h (Figure 2H, Figure 3F, 3G, 3H) all share a smooth enamel type, a strong distal displacement of the apex in labial and lingual view, an acutely tapering, sharp apical tip, and a distally inclined base of 45 degrees. One tooth (PIMUZ A/III 0823h) is likely to be a juvenile tooth. This is interesting, as thus far teeth of juvenile sauropods show a difference in expression of enamel wrinkling, as found by Fiorillo (1991, 1998), Díez Díaz et al. (2012), Díaz, Suberbiola & Sanz, (2012), Díez Díaz, Ortega & Sanz, (2014), and Holwerda, Pol & Rauhut (2015) in several titanosaurian teeth from the Ibero-Armorican Island. The smoothness of enamel wrinkling in juvenile teeth, therefore, could express differently in adults, further supporting the most parsimonious conclusion of including these last teeth in Morphotype I. Barrett et al., (2016) described some juvenile (hatchling or even embryonic) sauropod teeth of titanosauriforms/camarasaurids from the Berriasian of southern France, which match the apical morphology of this Moroccan tooth. However, the Moroccan tooth lacks the basal mesial constriction and the lingual buttress of juvenile teeth described by Barrett et al. (2016). Juvenile teeth are also described by García & Cerda, (2010) for sauropod embryos from Auca Mahuevo, however these small teeth show a more acutely tapering apical tip than the Moroccan sample. PIMUZ A/III 0823f and PIMUZ A/III 0823h resemble the Spanish Lo Hueco morphotype A in the general morphology of their crowns and their slight apical asymmetry (which could be due to wear). PIMUZ A/III 0823f also reflects the Lo Hueco morphotype A in that it has a similar length, labiolingual compression, and SI and CI values (Díez Díaz, Ortega & Sanz, 2014). As PIMUZ A/III 0823h probably belonged to a juvenile specimen, the size of this tooth is smaller when compared to the other ones, and this changes the SI and CI values. Juvenile teeth need to

be used with caution when included in comparative studies, especially when using quantitative variables such as the SI and CI (see Statistical Analysis).

PIMUZ A/III 0823g is the only tooth with smooth enamel that shows striations on the surface, providing the cross-section with a slightly rhomboid shape, however the strongly protruding carinae also still give a lemon-shaped cross section. Striations are seen on teeth of *Huabeisaurus* (D'Emic et al., 2013), brachiosaurid teeth from South Korea (Lim et al., 2001), smooth enamel type titanosaur teeth from the Bissekty Fm., Uzbekistan (Averianov & Sues, 2016), and in combination with smooth enamel, in *Atsinganosaurus* (though some teeth do display gentle wrinkling on the middle, see Díez Díaz, Tortosa & Le Loeuff, (2013), as well as in *Demandasaurus* (Torcida Fernández-Baldor et al., 2011), although the carinae are less conspicuous on the latter taxon. Smooth enamel, in combination with a sharp apex, is further seen in the Albian *Karongasaurus* from Malawi (Gomani, 2005). Smooth enamel exists on *Limaysaurus* (Salgado et al., 2004), however this tooth is more needle-shaped than the teeth of the Moroccan/Algerian sample. As these teeth are not worn, the apical morphology differs from the other teeth of Morphotype I, which are all mostly worn apically. However, as worn versus unworn teeth cannot be properly morphologically distinguished with isolated teeth, these are still assigned to Morphotype I.

Given the resemblance of Morphotype I to predominantly Cretaceous sauropods, and moreover, given the size and shape range, this morphotype is difficult to assign to any particular group. The size and shape range suggests a titanosauriform origin, as derived titanosaurs usually show a more conservative morphological range (García & Cerda, 2010). However, given the similarity between some teeth of Morphotype I with those of titanosaurs, it is more likely that both non-titanosaurian titanosauriforms as well as titanosaurs are present in the tooth sample of

Morphotype I, which reflects sauropod postcranial diversity of the Cretaceous of northwest Africa recorded by previous studies, as both titanosauriforms (Mannion & Barrett, 2013; Fanti, Cau & Hassine, 2014; Lamanna & Hasegawa, 2014) as well as derived titanosaurs (Ibrahim et al., 2016) have been reported to be present. Even though some of the teeth resemble more specific groups (e.g. *Euhelopus*, *Huabeisaurus*) the general morphotype cannot be assigned to any more specific group. No affinity with other groups (e.g. diplodocids, rebbachisaurids) is found for Morphotype I either.

**Morphotype II: BSPG 1993 IX 331C, BSPG 1993 IX 2B, PIMUZ A/III 0823d, PIMUZ A/III 0823i, PIMUZ A/III 0823j**

### **General**

A smaller tooth sample from both the Moroccan and Algerian sample deviates morphologically from Morphotype I. These teeth show a more rectangular morphology, a lack of apical tapering, and distinct labial and lingual wear facets.

### **Discussion**

BSPG 1993 IX 331C and PIMUZ A/III 0823j (Figure 2C, Figure 3J) both show a distinctive enamel wrinkling, not seen in the other teeth of the sample. It displays a ‘chevron’ pattern on the labial and lingual surfaces, with rugose striations anastomosing towards the carinae. This pattern is arguably present on the labial side of the enamel of another Moroccan Kem Kem sauropod tooth, tentatively (but not conclusively) assigned to rebbachisauridae by Kellner & Mader (1997). This latter tooth, however, also shows striations (Kellner & Mader, 1997), which are not seen on BSPG 1993 IX 331C, and PIMUZ A/III 0823j. As Carballido & Pol (2010) and Holwerda, Pol & Rauhut (2015) showed, enamel wrinkling on sauropod teeth can be an

autapomorphic diagnostic feature. As both teeth are worn, however, the other morphological features besides enamel wrinkling must be taken into account. The teeth do share morphological similarities in crown shape with the tooth described by Kellner & Mader (1997), which is apicobasally straight. The rectangular morphology of these teeth, caused by a lack of mesiodistal tapering towards the apex, together with a slightly higher CI due to higher labiolingual compression than the other teeth from the Moroccan/Algerian sample, showing an oval to cylindrical cross-section, is shared with BSPG 1993 IX 2B, as well as PIMUZ A/III 0823d and PIMUZ A/III 0823i (Figure 2F, Figure 3D, 3I). These five teeth further share labial and lingual wear facets, which are symmetrically placed, though in PIMUZ A/III 0823i, as well as in BSPG 1993 IX 2B, one high-angled and low-angled wear facet is seen, which is characteristic of the Aptian-Albian *Nigersaurus* (Serenó & Wilson, 2005; p.166, fig. 5.7; Sereno et al., 2007, Fig. 2). The size of PIMUZ A/III 0823i matches that of *Nigersaurus*, however, the other teeth are much larger than the isolated *Nigersaurus* tooth described by Sereno and Wilson (2005). The enamel asymmetry, a diagnostic feature for *Nigersaurus*, could not be accurately measured on the Moroccan and Algerian tooth sample. To a lesser degree, the labial and lingual wear facet symmetry is seen in both the Brazilian Aptian *Tapuisaurus* (Zaher et al., 2011), as well as the Late Cretaceous *Maxakalisaurus* (Kellner et al., 2006; França et al., 2016). However, these teeth show a higher SI than the Moroccan/Algerian sample, as well as stronger labial convexity. The morphology of Morphotype II also does not closely match the conical, peg-shaped teeth of *Demandsaurus* (Torcida Fernández-Baldor et al., 2011), a rebbachisaurid from the Early Cretaceous of Spain. However, the enamel is relatively smooth on both Morphotype II and *Demandsaurus*, and moreover, the teeth of Morphotype II are unworn, whereas the *Demandsaurus* teeth are relatively unworn.



The morphological features on BSPG 1993 IX 2B further show a set of derived titanosaurian characteristics, with a D-shaped cross-section in the middle of the tooth, as seen in *Mongolosaurus* (Mannion, 2011), but also seen in a titanosauriform tooth from Argentina (García and Cerda, 2010, MPCA-Pv-55, fig. 5). As previously stated, differences in tooth shape can be explained by their relative placement in the toothrow, as seen in *Mongolosaurus* (having both a cylindrical as well as a D-shaped tooth; Barrett et al., 2002; Mannion, 2011) and *Nemegtosaurus* (Wilson, 2005). Both *Mongolosaurus* as well as *Nemegtosaurus*, however, show a distinct tapering of the apex, which BSPG 1993 IX 2B does not show. Furthermore, differences in shape between upper and lower toothrows is also observed in diplodocoids, such as *Diplodocus* and *Nigersaurus* (Serenó & Wilson, 2005). The labial wear facet in particular is characteristic for diplodocids and dicraeosaurids (Serenó & Wilson 2005), and while diplodocids are not known from the Cretaceous, rebbachisaurids and dicraeosaurids are (e.g. Sereno & Wilson, 2005; Gallina et al., 2014). As rebbachisaurids such as *Rebbachisaurus garasbae* are found in the Kem Kem beds of Morocco (Lavocat, 1954; Mannion & Barrett, 2013; Wilson & Allain, 2015), the Aptian-Albian of Niger (*Nigersaurus* and rebbachisaurids, Sereno et al., 1996; Wilson & Sereno, 2005; Sereno et al., 2007), and also the Aptian-Albian of Tunisia (Fanti, Cau & Hassine, 2014; Fanti et al., 2015), and tentatively in the late Barremian-early Aptian of Spain (Pereda Suberbiola et al., 2003; Torcida Fernández-Baldor et al., 2011), they are both biogeographically close, as well as originating from beds near-contemporaneous to those of our tooth sample. Morphotype II therefore is tentatively assigned to Rebbachisauridae indet.

### ***Quantitative analysis***



Though quantitative analysis has been applied to theropod teeth in several previous studies (e.g., Hendrickx & Mateus, in review; Samman et al., 2005; Fanti & Therrien, 2007; Ösi, Apesteguía & Kowalewski, 2010; Hendrickx, Mateus & Araújo, 2015), a statistical approach to sauropod tooth diversity has thus far only been applied through using one variable - the SI - by Chure et al. (2010), and the number of wear facets on the apex (Averianov & Sues, 2016). Quantitative analyses on sauropod teeth with two or more variables is therefore an area of study that has not received much attention. Here, we conduct an analysis of sauropod teeth based on multivariate statistical tests. The one-way PERMANOVA revealed significant differences between some of the tested groups ( $F=14.46$ ,  $p=0.0001$ ). Specific comparisons for each pair of groups are listed in Table 2. Morphotype I shows significant differences in SI and CI ratios with the D-shaped Morphotype from Fox-Amphoux, *Patagosaurus*, *Euhelopus*, the Shiramine morphotype, and *Maxakalisaurus* (Table 2). Differences with some of these groups are congruent with the qualitative approach described above. On the other hand, the SI and CI values of Morphotype I did not significantly differ from those of *Atsinganosaurus*, *Lirainosaurus*, *Astrodon*, the cylindrical Morphotype from Fox-Amphoux, both tooth morphotypes from Lo Hueco, and the titanosaurian teeth from Masecaps, Neuquén, Uzbekistan, and the Bauru Formation, Brazil. (see Table 2). This is in agreement with the qualitative discussion of Morphotype I, where the robust tooth type showed morphological similarities with *Astrodon*, and the elongated cylindrical type was shown to be morphologically comparable to the Neuquén and Uzbekistan tooth types. It also reinforces the morphological similarities found between Morphotype I with Ibero-Armorican titanosaurs.

Concerning Morphotype II, the SI and CI ratios differ significantly from *Atsinganosaurus*, *Patagosaurus*, *Euhelopus*, and *Maxakalisaurus* (see Table 2) confirming that even though the

wear facets of Morphotype II match those of *Maxakalisaurus*, there is no further morphological overlap. Unfortunately, the available sample size of *Nigersaurus* and other rebbachisaurids was not large enough to statistically test their similarities with confidence (see Supplementary Table). However, the very small sample size of this study is not well-suited to this type of approach, therefore the results of this analysis must be taken with caution, and any relationships found between sauropod taxa of our sample must be regarded as tentative. The sauropod relationships suggested by the qualitative and quantitative analysis might not be supported if the sample size is increased. For example, most of the tooth groups of the studied sample were not distinguished after the Bonferroni correction on the post-hoc tests. This might mean that titanosaurian teeth do not possess any clear diagnostic features, or, more likely, that only two variables - the SI and the CI - are not enough to distinguish between titanosaurs, as most taxa and morphotypes strongly group together (see Figure 5). In Figure 5, most teeth from the Moroccan/Algerian samples cluster together, and there is also overlap with other sauropod taxa and/or tooth morphotypes. The teeth with high SI and high CI are *Maxakalisaurus*, *Rapetosaurus*, *Petrobasaurus* and *Limaysaurus*, whilst teeth with low SI and low CI are *Patagosaurus* and *Jobaria* (see Figure 5). The taxon with low SI but high CI is *Lapparentosaurus* (Figure 5). From the central cluster, Morphotype I and II overlap mostly with the Shiramine teeth, the Uzbekistan teeth, *Astrodon*, and the Franco-Iberian teeth (i.e. the Lo Hueco, Masecaps, Fox-Amphoux-Métissons and *Atsinganosaurus* types; Figure 5). Titanosauriform teeth are known to show a large range in morphology, and, moreover, they show some convergence in morphology with more basal sauropod teeth (Garcia & Cerda 2010). However, it is possible, after the qualitative and quantitative analyses here, that titanosaurian teeth show a similar convergence. Finally, as mentioned previously, the large range of size and shape within one tooththrow could also cause

taxa to accrete in Figure 5, though postcranial morphology would cause more morphological scatter. To resolve these issues, a larger sample size, not only of the case study at hand, but also of other (Cretaceous) sauropod teeth, could possibly aid in obtaining a better resolution of future quantitative analyses, however this is beyond the scope of the present study.

# ***Euro-Gondwanan Cretaceous sauropod diversity and palaeobiogeographical implications***

To summarize, the described tooth sample from the Albian–Cenomanian Kem Kem beds of Morocco and the Continental Intercalaire of Algeria shows a predominantly non-titanosaurian titanosauriform/titanosaurian assemblage, with only one morphotype (II, with four teeth) showing tentative rebbacchisaurid affinities. When compared with the biogeographically nearest tooth assemblages, namely titanosaurian teeth found in the Ibero-Armorican Island, as well as northwest and central Africa, several probable palaeobiogeographical patterns can be assessed.

Morphotype I shows morphological affinities with titanosauriforms. Titanosaurian sauropods in the Campanian Kem Kem beds of Morocco have been reported previously (Ibrahim et al., 2016). This titanosaurian predominance is also seen in the later stages of the Cretaceous of Egypt (Lamanna et al., 2017; Sallam et al., 2018), as well as Spain and France (see e.g. Le Loeuff, 1995, 2005; Sanz et al., 1999; Garcia et al., 2010; Díaz et al., 2016; Vila, Sellés & Brusatte, 2016). As fourteen teeth from our sample share several similarities with the cylindrical morphotype from Fox-Amphoux-Métisson, with the morphotype A from Lo Hueco and with the D-shaped morphotype from Fox-Amphoux-Métisson, as well as with *Atsinganosaurus* and the morphotype B from Lo Hueco (Díaz et al., 2012; Díaz, Suberbiola & Sanz, 2012; Díez Díaz,

Tortosa & Le Loeuff, 2013; Díez Díaz, Ortega & Sanz, 2014), this might suggest a possible close affinity between the Cenomanian North African titanosaurian faunas and those from the Campanian-Maastrichtian of southern Europe.

Morphotype II shows affinities with rebbachisaurids. Rebbachisaurid presence has already been noted from the Early Cretaceous of Niger (Serenó et al., 1999; Sereno & Wilson, 2005), Tunisia (Fanti et al., 2013, 2015; Fanti, Cau & Hassine, 2014), and Morocco (Lavocat, 1954, Kellner & Mader, 1997, Mannion & Barrett, 2013, Wilson & Allain, 2015), as well as the United Kingdom (Mannion, Upchurch & Hutt, 2011) and Spain (Pereda Suberbiola et al., 2003; Torcida Fernández-Baldor et al., 2011). Again, the North African - southern European connection is a tentative explanation for the dispersion of rebbachisaurids between Gondwana and Europe in the Cretaceous (see Figure 6); in addition, rebbachisaurids also seem to be a relatively diverse clade within North and Central Africa during the end of the Aptian-Cenomanian (e.g. Mannion & Barrett, 2013; Fanti et al., 2014; Wilson and Allain, 2015).

In previous studies, faunal connections have been demonstrated to exist between North Africa and Italy, such as temporary continental connections during the Barremian (125 Ma, the so-called Apulian route), or more permanent connections caused by carbonate platforms in the peri-Adriatic (Gheerbrant & Rage, 2006; Canudo et al., 2009; Zarcone et al., 2010; Torcida Fernández-Baldor et al., 2011). These last authors suggested that this route could have allowed for the divergence of the rebbachisaurids *Demandasaurus* and *Nigersaurus*. Moreover, Fanti et al., (2016) point to a rebbachisaurid dispersal event leading from Gondwana to a European lineage in the Early Cretaceous. Perhaps the hypothesized continental connection also allowed

961 titanosaurs to migrate between Laurasia and Gondwana. Furthermore, next to European  
 962 sauropods, also theropods, crocodyliforms, amphibians, and snakes and even batoids from  
 963 southern Europe (France, Spain, Italy, Croatia) have been found to show Gondwanan affinities  
 964 (Soler-Gijón & López-Martínez, 1998; Gardner, Evans, & Sigogneau-Russell, 2003; Pereda-  
 965 Suberbiola et al. 2009, 2015; Sweetman & Gardner, 2013; Csiki-Sava et al., 2015; Blanco et al.,  
 966 2016, 2017; Dal Sasso et al., 2016; Blanco, in press).  
 967 The discovery of titanosaurian teeth with similar morphologies from the Cenomanian of Algeria  
 968 and Morocco, and from the late Campanian-early Maastrichtian of the Ibero-Armorican Island,  
 969 could indicate that some of these European titanosaurian faunas had a Gondwanan origin.  
 970 However, this hypothesis needs to be taken with some caution until more postcranial remains  
 971 (with associated cranial specimens) are found and described from the Early-Late Cretaceous of  
 972 North Africa and southwestern Europe, as our statistical analysis does not show a high support of  
 973 any definite grouping between the North African and southern European tooth morphotypes.  
 974 Csiki-Sava et al. (2015) suggested that European titanosaurs do not seem to have a southern  
 975 influence, and palaeobiogeographical analyses, as well as this study, show both a Gondwanan  
 976 (South American) and North American input in European Cretaceous fauna (e.g. Upchurch,  
 977 2008; Ezcurra & Agnolín, 2011), which might cautiously be supported by our qualitative and  
 978 quantitative analyses as well. However, more recent studies have added more information about  
 979 the North African-southern European connection between these sauropod faunas in the Late  
 980 Cretaceous (see e.g. Sallam et al., 2018; Díez Díaz et al., 2018). In Figure 6, a tentative  
 981 palaeobiogeographical construction is shown using the Algerian and Moroccan morphotypes and  
 982 their morphological counterparts from northwest and central Africa, as well as southern Europe,  
 983 and possible migratory routes based on previous hypotheses.

984

## 985 CONCLUSIONS

986 A sample of eighteen teeth from the Cretaceous Kem Kem beds from Erfoud and Taouz,  
 987 Morocco, and from the Continental Intercalaire, Algeria, has been studied. The overwhelming  
 988 majority of this sample shows titanosauriform/titanosaur affinities, with a smaller group showing  
 989 rebbachisaurid affinities. This is congruent with the results of the statistical analyses. However,  
 990 the small size of the comparative samples and the relative scarcity of this type of analysis in  
 991 studies thus far, begs for caution in interpreting results until further study is possible.  
 992 On the other hand, similarities between tooth samples from northwestern and central Africa, and  
 993 southwestern Europe, do strongly hint at a possible sauropod faunal exchange through  
 994 continental connections in the early Late Cretaceous between North and Central Africa, and  
 995 between North Africa and southwestern Europe. These results support previous hypotheses from  
 996 earlier studies on faunal exchange and continental connections.

997

## 998 ACKNOWLEDGEMENTS

999 Oliver Rauhut (BSPG Munich, Germany), Christian Klug and Torsten Scheyer (PIMUZ Zürich,  
 1000 Switzerland) are profoundly thanked for kindly allowing us to use material from their  
 1001 collections. We are indebted to Enrico Schwabe (ZSM Munich, Germany) for his kind assistance  
 1002 in SEM imaging. Many thanks also to Daniela Schwarz and Jens Koch (Museum für Naturkunde  
 1003 Berlin, Germany) for allowing us to use their *Giraffatitan* specimen. Jeff Liston kindly checked  
 1004 the English. The comments and recommendations of editor Kenneth de Baets, and reviewers  
 1005 Paul Barrett, Attila Ösi, and one anonymous reviewer, greatly improved this manuscript. This  
 1006 work was part of the BSc thesis of RM at Utrecht University (the Netherlands).

# REFERENCES

- Alloul, T., Rage, J.-C., Hamdidouche, R., Jalil, N.-E., 2018. First report on Cretaceous vertebrates from the Algerian Kem Kem beds. A new procoelous salamander from the Cenomanian, with remarks on African Caudata. *Cretaceous Research* 84, 384–388.
- Apesteguía S. 2007. The sauropod diversity of the La Amarga Formation (Barremian), Neuquén (Argentina). *Gondwana Research* 12:533–546. DOI: 10.1016/j.gr.2007.04.007.
- Averianov A, Sues H-D. 2017. Sauropod teeth from the Upper Cretaceous Bissekty Formation of Uzbekistan. *Historical Biology* 29:641–653.
- Barrett PM, Hasegawa Y, Manabe M, Isaji S, Matsuoka H. 2002. Sauropod dinosaurs from the Lower Cretaceous of eastern Asia: taxonomic and biogeographical implications. *Palaeontology* 45:1197–1217. DOI: 10.1111/1475-4983.00282.
- Barrett PM, Pouech J, Mazin J-M, Jones FM. 2016. Teeth of Embryonic or Hatchling Sauropods from the Berriasian (Early Cretaceous) of Cherves-de-Cognac, France. *Acta Palaeontologica Polonica* 61:591–596. DOI: 10.4202/app.00257.2016.
- Barrett PM, Wang X-L. 2007. Basal titanosauriform (Dinosauria, Sauropoda) teeth from the Lower Cretaceous Yixian Formation of Liaoning Province, China. *Palaeoworld* 16:265–271. DOI: 10.1016/j.palwor.2007.07.001.
- Belvedere M, Jalil N-E, Breda A, Gattolin G, Bourget H, Khaldoune F, Dyke GJ. 2013. Vertebrate footprints from the Kem Kem beds (Morocco): A novel ichnological approach

- 1029 to faunal reconstruction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 383–
- 1030 384:52–58. DOI: 10.1016/j.palaeo.2013.04.026.
- 1031 Benyoucef M, Läng E, Cavin L, Mebarki K, Adaci M, Bensalah M. 2015. Overabundance of
- 1032 piscivorous dinosaurs (Theropoda: Spinosauridae) in the mid-Cretaceous of North Africa:
- 1033 The Algerian dilemma. *Cretaceous Research* 55:44–55.
- 1034 Blanco A. in press. *Igdabatis marmii* sp. nov. (Myliobatiformes) from the lower Maastrichtian
- 1035 (Upper Cretaceous) of northeastern Spain: an Ibero-Armorican origin for a Gondwanan
- 1036 batoid. *Journal of Systematic Palaeontology*. DOI: 10.1080/14772019.2018.1472673.
- 1037 Blanco A, Bolet A, Blain HA, Fondevilla V, Marmi J, 2016. Late Cretaceous (Maastrichtian)
- 1038 amphibians and squamates from northeastern Iberia. *Cretaceous Research*, 57:624–638.
- 1039 DOI:10.1016/j.cretres.2015.07.005.
- 1040 Blanco A, Szabó M, Blanco À, Marmi J. 2017. Late Cretaceous (Maastrichtian) Chondrichthyes
- 1041 and Osteichthyes from northeastern Iberia. *Palaeogeography, Palaeoclimatology,*
- 1042 *Palaeoecology*, 465, 278–294. DOI: 10.1016/j.palaeo.2016.10.039
- 1043 Bonaparte JF, González Riga BJ, Apesteguía S. 2006. *Ligabuesaurus leanzai* gen. et sp. nov.
- 1044 (Dinosauria, Sauropoda), a new titanosaur from the Lohan Cura Formation (Aptian, Lower
- 1045 Cretaceous) of Neuquén, Patagonia, Argentina. *Cretaceous Research* 27:364–376.
- 1046 Bonde N. 2012. Danish dinosaurs: a review. *Bernissart Dinosaurs and Early Cretaceous*
- 1047 *Terrestrial Ecosystems* 464:435–451.
- 1048 Buffetaut E. 1989. Archosaurian reptiles with Gondwanan affinities in the Upper Cretaceous of
- 1049 Europe. *Terra nova* 1:69–74.
- 1050 Buffetaut E, Mechin P, Mechin-Salessy A. 1988. Un dinosaure théropode d’affinités
- 1051 gondwaniennes dans le Crétacé supérieur de Provence. *Comptes rendus de l’Académie des*



- 1052 sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la
- 1053 Terre 306:153–158.
- 1054 Calvo JO. 1994. Jaw mechanics in sauropod dinosaurs. *Gaia* 10:183–193.
- 1055 Calvo JO, González Riga BJ. 2003. *Rinconsaurus caudamirus* gen. et sp. nov., a new
- 1056 titanosaurid (Dinosauria, Sauropoda) from the Late Cretaceous of Patagonia, Argentina.
- 1057 *Revista geológica de Chile* 30:333–353.
- 1058 Canudo JJ, Barco JL, Pereda-Suberbiola X, Ruiz-Omeñaca JJ, Salgado L, Fernández-Balador FT,
- 1059 Gasulla JM. 2009. What Iberian dinosaurs reveal about the bridge said to exist between
- 1060 Gondwana and Laurasia in the Early Cretaceous. *Bulletin de la Société Géologique de*
- 1061 *France* 180:5–11. DOI: 10.2113/gssgfbull.180.1.5.
- 1062 Carballido JL, Holwerda FM, Pol D, Rauhut OW. 2017. An Early Jurassic sauropod tooth from
- 1063 Patagonia (Cañadon Asfalto Formation: implications for sauropod diversity.. *Publicación*
- 1064 *Electrónica de la Asociación Paleontológica Argentina* 17.
- 1065 Carballido JL, Pol D. 2010. The dentition of *Amygdalodon patagonicus* (Dinosauria:
- 1066 Sauropoda) and the dental evolution in basal sauropods. *Comptes Rendus Palevol* 9:83–93.
- 1067 Carpenter K, Tidwell V. 2005. Reassessment of the Early Cretaceous sauropod *Astrodon*
- 1068 *johnsoni* Leidy 1865 (Titanosauriformes). In: *Thunder-lizards*. Indiana University Press,
- 1069 Bloomington and Indianapolis, 78–114.
- 1070 Cau A, Maganuco S. 2009. A new theropod dinosaur, represented by a single unusual caudal
- 1071 vertebra, from the Kem Kem Beds (Cretaceous) of Morocco. *Atti della Società italiana di*
- 1072 *scienze naturali e del museo civico di storia naturale di Milano* 150:239–257.
- 1073 Cavin L, Forey PL. 2004. New mawsoniid coelacanth (Sarcopterygii: Actinistia) remains from
- 1074 the Cretaceous of the Kem Kem beds, Southern Morocco. *Mesozoic fishes* 3:493–506.

- 1075 Cavin L, Tong H, Boudad L, Meister C, Piuze A, Tabouelle J, Aarab M, Amiot R, Buffetaut E,  
1076 Dyke G, others. 2010. Vertebrate assemblages from the early Late Cretaceous of  
1077 southeastern Morocco: an overview. *Journal of African Earth Sciences* 57:391–412.
- 1078 Chure D, Britt B, Whitlock J, Wilson J. 2010. First complete sauropod dinosaur skull  
1079 from the Cretaceous of the Americas and the evolution of sauropod dentition.  
1080 *Naturwissenschaften* 97:379–391. DOI: 10.1007/s00114-010-0650-6.
- 1081 Chure D., Britt BB., Whitlock JA., Wilson JA. 2010. First complete sauropod dinosaur skull  
1082 from the Cretaceous of the Americas and the evolution of sauropod dentition.  
1083 *Naturwissenschaften* 97:379–391.
- 1084 Csiki-Sava Z, Buffetaut E, Ősi A, Pereda-Suberbiola X, Brusatte SL. 2015. Island life in the  
1085 Cretaceous - faunal composition, biogeography, evolution, and extinction of land-living  
1086 vertebrates on the Late Cretaceous European archipelago. *ZooKeys* 469:1–161. DOI:  
1087 10.3897/zookeys.469.8439.
- 1088
- 1089 Dal Sasso C, Pierangelini G, Famiani F, Cau A, Nicosia U. 2016. First sauropod bones from Italy  
1090 offer new insights on the radiation of Titanosauria between Africa and Europe.  
1091 *Cretaceous Research* 64:88–109.
- 1092 Dalla Vecchia FM. 2002. Cretaceous dinosaurs in the Adriatic–Dinaric carbonate platform (Italy  
1093 and Croatia): paleoenvironmental implications and paleogeographical hypotheses.  
1094 *Memorie della Società Geologica Italiana* 57:89–100.
- 1095 Dalla Vecchia FM, Cau A. 2011. The first record of a notosuchian crocodyliiform from Italy.  
1096 *Rivista Italiana di Paleontologia e Stratigrafia (Research In Paleontology and*  
1097 *Stratigraphy)* 117.

- 1098 De Broin F, Grenot C, Vernet R. 1971. Sur la découverte d'un nouveau gisement de Vertébrés  
1099 dans le Continental Intercalaire saharien: la Gara Samani (Algérie). *Comptes Rendus de*  
1100 *l'Académie des Sciences de Paris* 272:1219–1221.
- 1101 D'Emic MD, Mannion PD, Upchurch P, Benson RBJ, Pang Q, Zhengwu C. 2013. Osteology of  
1102 *Huabeisaurus allocotus* (Sauropoda: Titanosauriformes) from the Upper Cretaceous of  
1103 China. *PLoS ONE* 8:e69375. DOI: 10.1371/journal.pone.0069375.
- 1104 Díez Díaz VD, Garcia G, Knoll F, Suberbiola X., Valentin X. 2012a. New cranial remains of  
1105 titanosaurian sauropod dinosaurs from the Late Cretaceous of Fox-Amphoux-Métisson  
1106 (Var, SE France). *Proceedings of the Geologists' Association* 123:626–637.
- 1107 Díez Díaz, V., Garcia, G., Pereda Suberbiola, X., Jentgen, B., Stein, K., Godefroit, P. and  
1108 Valentin, X. 2018. The titanosaurian dinosaur *Atsinganosaurus velauciensis*  
1109 (Sauropoda) from the Upper Cretaceous of southern France: new material, phylogenetic  
1110 affinities, and palaeobiogeographical implications. *Cretaceous Research* 91: 429–456.
- 1111  
1112 Díez Díaz VD, Mocho P, Páramo A, Escaso F, Marcos-Fernández F, Sanz JL, Ortega F. 2016. A  
1113 new titanosaur (Dinosauria, Sauropoda) from the Upper Cretaceous of Lo Hueco  
1114 (Cuenca, Spain). *Cretaceous Research* 68:49–60.
- 1115 Díez Díaz V, Ortega F, Sanz JL. 2014. Titanosaurian teeth from the Upper Cretaceous of “Lo  
1116 Hueco” (Cuenca, Spain). *Cretaceous Research*:285–291.
- 1117 Díez Díaz V, Pereda Suberbiola X, Sanz JL. 2012b. Juvenile and adult teeth of the titanosaurian  
1118 dinosaur *Lirainosaurus* (Sauropoda) from the Late Cretaceous of Iberia. *Geobios*  
1119 45:265–274.

- 1120 Díez Díaz V, Tortosa T, Le Loeuff J. 2013. Sauropod diversity in the Late Cretaceous of  
1121 southwestern Europe: The lessons of odontology. *Annales de Paléontologie* 99:119–  
1122 129. DOI: 10.1016/j.annpal.2012.12.002.
- 1123 Erickson GM. 1996. Incremental lines of von Ebner in dinosaurs and the assessment of tooth  
1124 replacement rates using growth line counts. *Proceedings of the National Academy of*  
1125 *Sciences of the United States of America* 93:14623–14627.
- 1126 Ezcurra MD, Agnolín FL. 2011. A New Global Palaeobiogeographical Model for the Late  
1127 Mesozoic and Early Tertiary. *Systematic Biology*.
- 1128 Fanti F, Cau A, Cantelli L, Hassine M, Auditore M. 2015. New information on *Tataouinea*  
1129 *hannibalis* from the Early Cretaceous of Tunisia and implications for the tempo and  
1130 mode of rebbachisaurid sauropod evolution. *PLOS ONE* 10:e0123475. DOI:  
1131 10.1371/journal.pone.0123475.
- 1132 Fanti F, Cau A, Hassine M. 2014. Evidence of titanosauriforms and rebbachisaurids (Dinosauria:  
1133 Sauropoda) from the Early Cretaceous of Tunisia. *Journal of African Earth Sciences*  
1134 90:1–8.
- 1135 Fanti F, Cau A, Hassine M, Contessi M. 2013. A new sauropod dinosaur from the Early  
1136 Cretaceous of Tunisia with extreme avian-like pneumatization. *Nature Communications*  
1137 4. DOI: 10.1038/ncomms3080.
- 1138 Fanti F, Cau A, Panzarin L, Cantelli L. 2016. Evidence of iguanodontian dinosaurs from the  
1139 Lower Cretaceous of Tunisia. *Cretaceous Research* 60:267–274.
- 1140 Fanti F, Therrien F. 2007. Theropod tooth assemblages from the Late Cretaceous Maevarano  
1141 Formation and the possible presence of dromaeosaurids in Madagascar. *Acta*  
1142 *Palaeontologica Polonica* 52:155.

- 1143 Fiorillo AR. 1991. Dental microwear on the teeth of *Camarasaurus* and *Diplodocus*:  
1144 implications for sauropod paleoecology. In: *Fifth Symposium on Mesozoic Terrestrial*  
1145 *Ecosystems and Biota*. Contributions from the Palaeontological Museum University of  
1146 Oslo, 24.
- 1147 Fiorillo AR. 1998. Dental micro wear patterns of the sauropod dinosaurs *Camarasaurus* and  
1148 *Diplodocus*: evidence for resource partitioning in the Late Jurassic of North America.  
1149 *Historical Biology* 13:1–16.
- 1150 Forey PL, Cavin L. 2007. A new species of *Cladocycclus* (Teleostei: Ichthyodectiformes) from  
1151 the Cenomanian of Morocco. *Palaeontologia Electronica* 10:1–10.
- 1152 Forey PL, López-Arbarello A, MacLeod N. 2011. A new species of *Lepidotes* (Actinopterygii:  
1153 Semiontiformes) from the Cenomanian (Upper Cretaceous) of Morocco. *Palaeontologia*  
1154 *electronica* 14:1–12.
- 1155 França MA, Júlio C de A, Riff D, Hsiou AS, Langer MC. 2016. New lower jaw and teeth  
1156 referred to *Maxakalisaurus topai* (Titanosauria: Aeolosaurini) and their implications for  
1157 the phylogeny of titanosaurid sauropods. *PeerJ* 4:e2054.
- 1158 Freire PC, Medeiros MA, Lindoso RM. 2007. Sauropod teeth diversity in the Laje do Coringa  
1159 fossiliferous site, Eocenomanian of northeastern Brazil. *Paleontologia: Cenários de*  
1160 *Vida* 1:523e532.
- 1161 Gallina PA, Apesteguía S, Haluza A, Canale JI. 2014. A diplodocid sauropod survivor from the  
1162 Early Cretaceous of South America. *PLoS ONE* 9:e97128. DOI:  
1163 10.1371/journal.pone.0097128.
- 1164 Garcia G, Amico S, Fournier F, Thouand E, Valentin X. 2010. A new titanosaur genus  
1165 (Dinosauria, Sauropoda) from the Late Cretaceous of southern France and its

- 1166 paleobiogeographic implications. *Bulletin de la Société géologique de France* 181:269–
- 1167 277.
- 1168 García Ra, Cerda IA 2010. Dentition and histology in titanosaurian dinosaur embryos from
- 1169 Upper Cretaceous of Patagonia, Argentina. *Palaeontology* 53:335–346.
- 1170 García RA, Cerda IA. 2010. Dentición de titanosaurios (Dinosauria, Sauropoda) del Cretácico
- 1171 Superior de la provincia de Río Negro, Argentina: morfología, inserción y reemplazo.
- 1172 *Ameghiniana* 47:45–60.
- 1173 Garcia RA, Salgado L. 2012. The Titanosaur Sauropods from the Late Campanian—Early
- 1174 Maastrichtian Allen Formation of Salitral Moreno, Río Negro, Argentina. *Acta*
- 1175 *Palaeontologica Polonica* 58:269–284.
- 1176 Gardner JD, Evans SE, Sigogneau–Russell D. 2003. New albanerpetontid amphibians from the
- 1177 Early Cretaceous of Morocco and Middle Jurassic of England. *Acta Palaeontologica*
- 1178 *Polonica* 48: 301–319.
- 1179 Gomani EM. 2005. Sauropod dinosaurs from the early Cretaceous of Malawi, Africa.
- 1180 *Palaeontologia Electronica* 8:27A.
- 1181 Hammer Ø, Harper DAT, Ryan PD. 2001. Past: Paleontological Statistics Software Package for
- 1182 education and data analysis. *Palaeontologia Electronica* 4:1–9.
- 1183 Hendrickx C, Mateus O. 2016. Morphofunctional analysis of the quadrate of Spinosauridae
- 1184 (Dinosauria: Theropoda) and the first definitive evidence of two cohabiting spinosaurids
- 1185 in the Cenomanian of North Africa. *PLoS ONE*.
- 1186 Hendrickx C, Mateus O, Araújo R. 2015. A proposed terminology of theropod teeth (Dinosauria,
- 1187 Saurischia). *Journal of Vertebrate Paleontology* 35:e982797.

- 1188 Holwerda FM, Pol D, Rauhut OWM. 2015. Using dental enamel wrinkling to define sauropod  
1189 tooth morphotypes from the Cañadón Asfalto Formation, Patagonia, Argentina. *PLOS*  
1190 *ONE* 10:e0118100. DOI: 10.1371/journal.pone.0118100.
- 1191 Ibrahim N, Dal Sasso C, Maganuco S, Fabbri M, Martill DM, Gorscak E, Lamanna MA. 2016.  
1192 Evidence of a derived titanosaurian (Dinosauria, Sauropoda) in the “Kem Kem beds” of  
1193 Morocco, with comments on sauropod paleoecology in the Cretaceous of Africa. In:  
1194 *Cretaceous Period: Biotic Diversity and Biogeography*. New Mexico Museum of Natural  
1195 History & Science,.
- 1196 Ibrahim N, Varricchio DJ, Sereno PC, Wilson JA, Dutheil DB, Martill DM, Baidder L, Zouhri S.  
1197 2014. Dinosaur footprints and other ichnofauna from the Cretaceous Kem Kem Beds of  
1198 Morocco. *PLoS ONE* 9:e90751. DOI: 10.1371/journal.pone.0090751.
- 1199 Kellner A, Campos D, Azevedo S, Trotta M, Henriques D, Craik M, Silva H. 2006. On a new  
1200 titanosaur sauropod from the Bauru Group, Late Cretaceous of Brazil. *Boletim do Museu*  
1201 *Nacional* 74:1–31.
- 1202 Kellner AW, Mader BJ. 1997. Archosaur teeth from the Cretaceous of Morocco. *Journal of*  
1203 *Paleontology* 71:525–527.
- 1204 Kues BS, Lehman T, Rigby JK. 1980. The teeth of *Alamosaurus sanjuanensis*, a Late Cretaceous  
1205 sauropod. *Journal of Paleontology* 54:864–869.
- 1206 Lamanna MC, Hasegawa Y. 2014. New titanosauriform sauropod dinosaur material from the  
1207 Cenomanian of Morocco: implications for paleoecology and sauropod diversity in the  
1208 Late Cretaceous of north Africa. *Bulletin of Gunma Museum of Natural History* 18:1–9.

- 1209 Läng E, Boudad L, Maio L, Samankassou E, Tabouelle J, Tong H, Cavin L. 2013. Unbalanced  
1210 food web in a Late Cretaceous dinosaur assemblage. *Palaeogeography,*  
1211 *Palaeoclimatology, Palaeoecology* 381–382:26–32. DOI: 10.1016/j.palaeo.2013.04.011.
- 1212 Lamanna, M.C., Gorscak, E., Díez Díaz, V., Schwarz, D., El-Dawoudi, I. 2017. Reassessment Of  
1213 A Partial Titanosaurian Sauropod Dinosaur Skeleton From The Upper Cretaceous  
1214 (Campanian) Quseir Formation Of The Kharga Oasis, Egypt. XV Annual Meeting of the  
1215 European Association of Vertebrate Palaeontologists. *Zitteliana*: 50.
- 1216 de Lapparent AF, Gorce F. 1960. The dinosaurs of the “Continental Intercalaire” of the central  
1217 Sahara. *Memoirs of the Geological Society of France XXXIX*.
- 1218 Lavocat R. 1954. Reconnaissance géologique dans les Hammadas des confins algéro-marocains  
1219 du Sud. *Notes et Mémoires du Service Géologique du Maroc* 116:1–147.
- 1220 Le Loeuff J. 1991. The Campano-Maastrichtian vertebrate faunas from southern Europe and  
1221 their relationships with other faunas in the world; palaeobiogeographical implications.  
1222 *Cretaceous Research* 12:93–114. DOI: 10.1016/S0195-6671(05)80019-9.
- 1223 Le Loeuff J. 1995. *Ampelosaurus atacis* (nov. gen., nov. sp.), a new titanosaurid (Dinosauria,  
1224 Sauropoda) from the Late Cretaceous of the Upper Aude Valley (France). *Comptes*  
1225 *Rendus de l’Academie des Sciences-Serie II-Sciences de la Terre et des Planetes*  
1226 321:693–700.
- 1227 Le Loeuff J. 2005. Osteology of *Ampelosaurus atacis* (Titanosauria) from southern France.  
1228 *Thunder-Lizards, The Sauropodomorph Dinosaurs*:115–137.
- 1229 Le Loeuff J, Läng E, Cavin L, Buffetaut E. 2012. Between Tendaguru and Bahariya: on the age  
1230 of the Early Cretaceous dinosaur sites from the Continental Intercalaire and other African  
1231 formations. *Journal of Stratigraphy* 36:e502.



- 1232 Lefranc JP, Guiraud R. 1990. The Continental Intercalaire of northwestern Sahara and its  
1233 equivalents in the neighbouring regions. *Journal of African Earth Sciences (and the*  
1234 *Middle East)* 10:27–77.
- 1235 Lim J-D, Martin LD, Baek K-S. 2001. The first discovery of a brachiosaurid from the Asian  
1236 continent. *Naturwissenschaften* 88:82–84.
- 1237 Mahler L. 2005. Record of Abelisauridae (Dinosauria: Theropoda) from the Cenomanian of  
1238 Morocco. *Journal of Vertebrate Paleontology* 25:236–239.
- 1239 Mannion PD. 2009. A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of Wight,  
1240 England. *Cretaceous Research* 30:521–526.
- 1241 Mannion PD. 2011. A reassessment of *Mongolosaurus haplodon* Gilmore, 1933, a titanosaurian  
1242 sauropod dinosaur from the Early Cretaceous of Inner Mongolia, People’s Republic of  
1243 China. *Journal of Systematic Palaeontology* 9:355–378. DOI:  
1244 10.1080/14772019.2010.527379.
- 1245 Mannion PD, Allain R, Moine O. 2017. The earliest known titanosauriform sauropod dinosaur  
1246 and the evolution of Brachiosauridae. *PeerJ* 5:e3217.
- 1247 Mannion PD, Barrett PM. 2013. Additions to the sauropod dinosaur fauna of the Cenomanian  
1248 (early Late Cretaceous) Kem Kem beds of Morocco: Palaeobiogeographical implications  
1249 of the mid-Cretaceous African sauropod fossil record. *Cretaceous Research* 45:49–59.  
1250 DOI: 10.1016/j.cretres.2013.07.007.
- 1251 Mannion PD, Upchurch P, Hutt S. 2011. New rebbachisaurid (Dinosauria: Sauropoda) material  
1252 from the Wessex Formation (Barremian, Early Cretaceous), Isle of Wight, United  
1253 Kingdom. *Cretaceous Research* 32:774–780.

- 1254 Martínez RDF, Lamanna MC, Novas FE, Ridgely RC, Casal GA, Martínez JE, Vita JR, Witmer  
1255 LM. 2016. A basal lithostrotian titanosaur (Dinosauria: Sauropoda) with a complete skull:  
1256 implications for the evolution and paleobiology of Titanosauria. *PLOS ONE*  
1257 11:e0151661. DOI: 10.1371/journal.pone.0151661.
- 1258 McGowan AJ, Dyke GJ. 2009. A surfeit of theropods in the Moroccan Late Cretaceous?  
1259 Comparing diversity estimates from field data and fossil shops. *Geology* 37:843–846.  
1260 DOI: 10.1130/G30188A.1.
- 1261 Meister C, Piuze A, Cavin L, Boudad L, Bacchia F, Ettachfini EM, Benyoucef M. 2017. Late  
1262 Cretaceous (Cenomanian-Turonian) ammonites from southern Morocco and south  
1263 western Algeria. *Arabian Journal of Geosciences* 10:1.
- 1264 Mocho P, Royo-Torres R, Malafaia E, Escaso F, Silva B, Ortega F. 2016. *Turiasauria*-like teeth  
1265 from the Upper Jurassic of the Lusitanian Basin, Portugal. *Historical Biology* 28:861–  
1266 880. DOI: 10.1080/08912963.2015.1049948.
- 1267 Novas FE, Dalla Vecchia F, Pais DF. 2005. Theropod pedal unguals from the Late  
1268 Cretaceous (Cenomanian) of Morocco, Africa. *Revista del Museo Argentino de Ciencias*  
1269 *Naturales* 7:167–175.
- 1270 Ösi A, Csiki-Sava Z, Prondvai E. 2017. A Sauropod Tooth from the Santonian of Hungary and  
1271 the European Late Cretaceous ‘Sauropod Hiatus.’ *Scientific reports* 7:3261.
- 1272 Ösi A, Apesteguía S, Kowalewski M. 2010. Non-avian theropod dinosaurs from the early Late  
1273 Cretaceous of Central Europe. *Cretaceous Research* 31:304–320.
- 1274 Pereda-Suberbiola X., Corral JC., Astibia H., Badiola A., Bardet N., Berreteaga A., Buffetaut E.,  
1275 Buscalioni AD., Cappetta H., Cavin L. 2015. Late Cretaceous continental and marine

- 1276 vertebrate assemblages of the Laño Quarry (Basque-Cantabrian Region, Iberian
- 1277 Peninsula): an update. *Journal of Iberian Geology* 41:101.
- 1278 Pereda Suberbiola X, Torcida F, Izquierdo LA, Huerta P, Montero D, Pérez G. 2003. First
- 1279 rebbachisaurid dinosaur (Sauropoda, Diplodocoidea) from the early Cretaceous of Spain:
- 1280 paleobiogeographical implications. *Bulletin de la Societe Geologique de France* 174:471–
- 1281 479. DOI: 10.2113/174.5.471.
- 1282 Pereda-Suberbiola X. 2009. Biogeographical affinities of Late Cretaceous continental tetrapods
- 1283 of Europe: a review. *Bulletin de la Société géologique de France* 180:57–71.
- 1284 Poropat SF, Kear BP. 2013. Photographic atlas and three-dimensional reconstruction of the
- 1285 holotype skull of *Euhelopus zdanskyi* with description of additional cranial elements.
- 1286 *PLoS ONE* 8:e79932. DOI: 10.1371/journal.pone.0079932.
- 1287 Rabi M, Sebök N. 2015. A revised Eurogondwana model: Late Cretaceous notosuchian
- 1288 crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal links
- 1289 between Europe and Gondwana during most of the Cretaceous. *Gondwana Research*
- 1290 28:1197–1211.
- 1291 Richter U, Mudroch A, Buckley LG. 2013. Isolated theropod teeth from the Kem Kem Beds
- 1292 (Early Cenomanian) near Taouz, Morocco. *Paläontologische Zeitschrift* 87:291–309.
- 1293 DOI: 10.1007/s12542-012-0153-1.
- 1294 Rodrigues T, Kellner AW, Mader BJ, Russell DA. 2011. New pterosaur specimens from the
- 1295 Kem Kem beds (Upper Cretaceous, Cenomanian) of Morocco. *Rivista Italiana di*
- 1296 *Paleontologia e Stratigrafia (Research In Paleontology and Stratigraphy)* 117.
- 1297 Russell DA. 1996. Isolated dinosaur bones from the Middle Cretaceous of the Tafilalt, Morocco.
- 1298 *Bulletin du Muséum National d'Histoire Naturelle* 4:349–402.

- 1299 Saegusa H, Tomida Y. 2011. Titanosauriform teeth from the Cretaceous of Japan. *Anais da*  
1300 *Academia Brasileira de Ciências* 83:247–265.
- 1301 Salgado L, Calvo JO. 1997. Evolution of titanosaurid sauropods. II: the cranial evidence.  
1302 *Ameghiniana* 34:33–48.
- 1303 Salgado L, Garrido A, Cocca SE, Cocca JR. 2004. Lower Cretaceous rebbachisaurid sauropods  
1304 from Cerro Aguada del León (Lohan Cura Formation), Neuquén Province, northwestern  
1305 Patagonia, Argentina. *Journal of Vertebrate Paleontology* 24:903–912.
- 1306 Sallam HM, Gorscak E, O'Connor PM, El-Dawoudi IA, El-Sayed S, Saber S, Kora MA, Sertich  
1307 JJ, Seiffert ER, Lamanna MC. 2018. New Egyptian sauropod reveals Late Cretaceous  
1308 dinosaur dispersal between Europe and Africa. *Nature Ecology & Evolution*:1.
- 1309 Samman T, Powell GL, Currie PJ, Hills LV. 2005. Morphometry of the teeth of western North  
1310 American tyrannosaurids and its applicability to quantitative classification. *Acta*  
1311 *Palaeontologica Polonica* 50:757.
- 1312 Sanz J, Powell J, Le Loeuff J, Martinez R, Pereda Suberbiola X. 1999. Sauropod remains from  
1313 the Upper Cretaceous of Laño (northcentral Spain). Titanosaur phylogenetic  
1314 relationships. *Estudios del Museo de Ciencias Naturales de Alava* 14:235–255.
- 1315 Sereno PC, Beck AL, Dutheil DB, Larsson HC, Lyon GH, Moussa B, Sadleir RW, Sidor CA,  
1316 Varricchio DJ, Wilson GP. 1999. Cretaceous sauropods from the Sahara and the uneven  
1317 rate of skeletal evolution among dinosaurs. *Science* 286:1342–1347.
- 1318 Sereno PC, Dutheil DB, Iarochene M, Larsson HCE, Lyon GH, Magwene PM, Sidor CA,  
1319 Varricchio DJ, Wilson JA. 1996. Predatory Dinosaurs from the Sahara and Late  
1320 Cretaceous Faunal Differentiation. *Science* 272:986–991. DOI:  
1321 10.1126/science.272.5264.986.

- 1322 Sereno PC, Wilson JA. 2005. Structure and evolution of a sauropod tooth battery. In: Curry  
1323 Rogers K, Wilson JA eds. *The Sauropods: Evolution and Paleobiology*. Berkeley, CA:  
1324 University of California Press, 157–177.
- 1325 Sereno PC, Wilson JA, Witmer LM, Whitlock JA, Maga A, Ide O, Rowe TA. 2007. Structural  
1326 extremes in a Cretaceous dinosaur. *PLoS ONE* 2:e1230. DOI:  
1327 10.1371/journal.pone.0001230.
- 1328 Soler-Gijón R, López-Martínez N. 1998. Sharks and rays (chondrichthyes) from the  
1329 Upper Cretaceous red beds of the south-central Pyrenees (Lleida, Spain): indices of  
1330 an India-Eurasia connection. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 141, 1–12.
- 1331 Sweetman SC, Gardner JD. 2013. A new albanerpetontid amphibian from the Barremian (Early  
1332 Cretaceous) Wessex Formation of the Isle of Wight, southern England. *Acta*  
1333 *Palaeontologica Polonica* 58 (2): 295–324.
- 1334 Torcida Fernández-Baldor FT, Canudo JI, Huerta P, Moreno-Azanza M, Montero D. 2017.  
1335 *Europatitan eastwoodi*, a new sauropod from the lower Cretaceous of Iberia in the initial  
1336 radiation of somphospondylans in Laurasia. *PeerJ* 5:e3409.
- 1337 Torcida Fernández-Baldor F, Canudo JI, Huerta P, Montero D, Pereda Suberbiola X. 2011.  
1338 *Demandasaurus darwini*, a new rebbachisaurid sauropod from the Early Cretaceous of  
1339 the Iberian Peninsula. *Acta Palaeontologica Polonica* 56:535–552. DOI:  
1340 10.4202/app.2010.0003.
- 1341 Upchurch P. 1995. The evolutionary history of sauropod dinosaurs. *Philosophical Transactions*  
1342 *of the Royal Society of London. Series B: Biological Sciences* 349:365–390.
- 1343 Upchurch P. 1998. The phylogenetic relationships of sauropod dinosaurs. *Zoological Journal of*  
1344 *the Linnean Society* 124:43–103.

- 1345 Upchurch P. 2008. Gondwanan break-up: legacies of a lost world? *Trends in ecology &*  
1346 *evolution* 23:229–236.
- 1347 Upchurch P, Barrett PM. 2000. The evolution of sauropod feeding mechanisms. In: Sues H-D ed.  
1348 *Evolution of herbivory in terrestrial vertebrates: perspectives from the fossil record*.  
1349 Cambridge: Cambridge University Press, 79–122.
- 1350 Upchurch P, Barrett PM, Dodson P. 2004. Sauropoda. In: Weishampel DB, Dodson P, Osmólska  
1351 H eds. *The Dinosauria. Second edition*. Berkeley, CA: University of California Press,  
1352 259–322.
- 1353 Vila B, Sellés AG, Brusatte SL. 2016. Diversity and faunal changes in the latest Cretaceous  
1354 dinosaur communities of southwestern Europe. *Cretaceous Research* 57:552–564. DOI:  
1355 10.1016/j.cretres.2015.07.003.
- 1356 Vullo R, Néraudeau D, Allain R, Cappetta H. 2005. Un nouveau gisement à microrestes de  
1357 vertébrés continentaux et littoraux dans le Cénomaniens inférieur de Fouras (Charente-  
1358 Maritime, Sud-Ouest de la France). *Comptes Rendus Palevol* 4:95–107.
- 1359 Vullo R, Neraudeau D, Lenglet T. 2007. Dinosaur teeth from the Cenomanian of Charentes,  
1360 western France: evidence for a mixed Laurasian-Gondwanan assemblage. *Journal of*  
1361 *Vertebrate Paleontology* 27:931–943.
- 1362 Wellnhofer P, Buffetaut E. 1999. Pterosaur remains from the Cretaceous of Morocco.  
1363 *Paläontologische Zeitschrift* 73:133.
- 1364 Wiersma K, Sander PM. 2016. The dentition of a well-preserved specimen of *Camarasaurus* sp.:  
1365 implications for function, tooth replacement, soft part reconstruction, and food intake.  
1366 *PalZ*:1–17. DOI: 10.1007/s12542-016-0332-6.

- 1367 Wilson JA. 2002. Sauropod dinosaur phylogeny: critique and cladistic analysis. *Zoological*  
1368 *Journal of the Linnean Society* 136:215–275.
- 1369 Wilson JA. 2005. Redescription of the Mongolian sauropod *Nemegtosaurus mongoliensis*  
1370 Nowinski (Dinosauria: Saurischia) and comments on Late Cretaceous sauropod diversity.  
1371 *Journal of Systematic Palaeontology* 3:283–318.
- 1372 Wilson JA, Allain R. 2015. Osteology of *Rebbachisaurus garasbae* Lavocat, 1954, a diplodocoid  
1373 (Dinosauria, Sauropoda) from the early Late Cretaceous-aged Kem Kem beds of  
1374 southeastern Morocco. *Journal of Vertebrate Paleontology* 35:e1000701.
- 1375 Wilson JA, Sereno PC. 1998. Early evolution and higher-level phylogeny of sauropod dinosaurs.  
1376 *Journal of Vertebrate Paleontology* 18:1–79.
- 1377 Zaher H, Pol D, Carvalho AB, Nascimento PM, Riccomini C, Larson P, Juarez-Valieri R,  
1378 Pires-Domingues R, da Silva NJ, de Almeida CD. 2011. A complete skull of an Early  
1379 Cretaceous sauropod and the evolution of advanced titanosaurs. *PLoS ONE* 6:e16663.
- 1380 Zarcone G, Petti FM, Cillari A, Di Stefano P, Guzzetta D, Nicosia U. 2010. A possible bridge  
1381 between Adria and Africa: new palaeobiogeographic and stratigraphic constraints on the  
1382 Mesozoic palaeogeography of the Central Mediterranean area. *Earth-Science Reviews*  
1383 103:154–162.

1384

## 1385 Figure and Table legends

1386

- 1387 Figure 1: Geological setting of Kem Kem beds, Morocco, and continental intercalaire, Algeria.  
1388 Taouz and Erfoud are indicated. The localities of Lo Hueco, Spain, and Fox-Amphoux-  
1389 Métißons and Masecaps are also portrayed.

A: stratigraphical column of Kem Kem beds (after Ibrahim et al., 2014). B: stratigraphical column of continental intercalaire, Algeria, (after Forey and Cavin, 2010).

Figure 2: Images of BSPG 1993 IX 331A (A), BSPG 1993 IX 331B (B), BSPG 1993 IX 331C (C), BSPG 1993 IX 313A (D), BSPG 1993 IX 2A (E), BSPG 1993 IX 2B (F), BSPG 1993 IX 2C (G), and BSPG 1993 IX 2D (H).

In basal view, apical view, labial view, lingual view, distal view, and mesial view. The scale bar equals 1 cm.

Figure 3: Images of PIMUZ A/III 0823a (A), PIMUZ A/III 0823b (B) PIMUZ A/III 0823c (C), PIMUZ A/III 0823d (D), PIMUZ A/III 0823e (E), PIMUZ A/III 0823f (F), PIMUZ A/III 0823g(G), PIMUZ A/III 0823h (H), PIMUZ A/III 0823i (I), PIMUZ A/III 0823j (J).

In apical view, basal view, labial view, lingual view, distal view, and mesial view. The scale bar equals 1 cm.

Figure 4: SEM pictures of enamel wrinkling. BSPG 1993 IX 331A in labial (A) and lingual (B) view, BSPG 1993 IX 331B in labial (C) and lingual (D) view, BSPG 1993 IX 313A in labial (E) and lingual (F) view, BSPG 1993 IX 2A in labial (G) and lingual (H) view, BSPG 1993 IX 2B in labial (I) and lingual (J) view, and BSPG 1993 IX 2D in labial (K) and lingual (L) view. The scale bar equals 500µm.

Figure 5: Dispersion plot of Cretaceous tooth morphotypes, with Jurassic outgroups. See coloured boxes in legend for sauropod groups.



1413

1414 Figure 6: Palaeobiogeographical reconstruction of northwest Africa and southern Europe during  
1415 the Cenomanian (black line) and Campanian-Maastrichtian (grey line), with possible migration  
1416 routes (dotted green line). Adapted from Csiki-Sava et al., (2015).

1417 Tooth morphotype I next to tooth morphotypes A and B from Lo Hueco (from Díez Díaz, Ortega  
1418 & Sanz, 2014) Fox-Amphoux-Métissons cylindrical type and D-shaped morphotype (from Díez  
1419 et al., 2012), the Masecaps titanosaur and *Atsinganosaurus* (from Díez Díaz, Tortosa & Le  
1420 Loeuff, 2013). Tooth morphotype II next to *Nigersaurus* (from Sereno et al., 2007), and  
1421 *Demandasaurus* (From Torcida Fernández-Baldor et al., 2011).

1422

1423 Table 1: Measurements, wear stage (after Saegusa & Tomida, 2011) and enamel morphology of  
1424 each tooth. Measurements in mm.

1425

1426 Table 2: One-way PERMANOVA comparisons for each pair of sauropod groups.

1427

1428 Supplementary Table 1: All tooth samples for the quantitative analysis with SI and CI from pers.  
1429 obs. or literature.


1430

1431

# Figure 1

Geological setting of Kem Kem beds, Morocco, and Continental Intercalaire, Algeria.

Taouz and Erfoud are indicated. localities of Lo Hueco, Spain, and France (FAM, AIX, Masecaps) are also portrayed. A: Map of Europe and Africa. B: Close-up of North Africa and Southern Europe. C: stratigraphical column of Kem Kem beds (after Belvedere et al., 2013; Ibrahim et al., 2014). D: stratigraphical column of Continental Intercalaire, Algeria, (after Forey, López-Arbarello & MacLeod, 2011).



# Figure 2

Images of BSPG 1993 IX 331A (A), BSPG 1993 IX 331B (B), BSPG 1993 IX 331C (C), BSPG 1993 IX 313A (D), BSPG 1993 IX 2A (E), BSPG 1993 IX 2B (F), BSPG 1993 IX 2C (G), and BSPG 1993 IX 2D (H).

In basal view, apical view, labial view, lingual view, distal view, and mesial view. The scale bar equals 1 cm.  
*Images taken by RM.*



1cm

# Figure 3

Images of PIMUZ A/III 0823a (A), PIMUZ A/III 0823b (B) PIMUZ A/III 0823c (C), PIMUZ A/III 0823d (D), PIMUZ A/III 0823e (E), PIMUZ A/III 0823f (F), PIMUZ A/III 0823g(G), PIMUZ A/III 0823h (H), PIMUZ A/III 0823i (I), PIMUZ A/III 0823j (J).

In apical view, basal view, labial view, lingual view, distal view, and mesial view. The scale bar equals 1 cm. *Images taken by FH.*






# Figure 4

SEM pictures of enamel wrinkling.

BSPG 1993 IX 331A in labial (A) and lingual (B) view, BSPG 1993 IX 331B in labial (C) and lingual (D) view, BSPG 1993 IX 313A in labial (E) and lingual (F) view, BSPG 1993 IX 2A in labial (G) and lingual (H) view, BSPG 1993 IX 2B in labial (I) and lingual (J) view, and BSPG 1993 IX 2D in labial (K) and lingual (L) view. The scale bar equals 500 $\mu$ m. *Images taken by FH and RM.*









# Figure 5

Dispersion plot of Cretaceous sauropod tooth morphotypes, with Jurassic outgroups.


See coloured boxes in legend for sauropod groups. A: All groups highlighted. B: Morphotype I and II highlighted.



# Figure 6

Palaeobiogeographical reconstruction of northwest Africa and southern Europe using sauropod tooth morphotypes.

A: northwest Africa and southern Europe during the Cenomanian (black line) and Campanian-Maastrichtian (grey line), with possible migration routes (dotted green line), adapted from Csiki-Sava et al., (2015). B: Tooth morphotype I next to tooth morphotypes A and B from Lo Hueco (from Díez Díaz, Ortega & Sanz, 2014) Fox-Amphoux-Métisçons cylindrical type and D-shaped morphotype (from Díez et al., 2012), the Masecaps titanosaur and *Atsinganosaurus* (from Díez Díaz, Tortosa & Le Loeuff, 2013). C: Tooth morphotype II next to *Nigersaurus* (from Sereno et al., 2007), and *Demandasaurus* (From Torcida Fernández-Baldor et al., 2011).



# **Table 1**(on next page)

Measurements, wear stage (after Saegusa & Tomida, 2011) and enamel morphology of each tooth.

Measurements in mm.

Specimen nr	Apico-basal length	Abrasion stage	Mesio-distal base	Mesio-distal middle	Mesio-distal apex	Labio-lingual base	Labio-lingual middle	Labio-lingual apex	SI	CI	Abrasion stage	Number and location of wear facets	Hypothetical placement in the snout	Apical third cross-section	Enamel ornamentation
BSPG 1993 IX 331A	60	53	12	12,3	6,2	10	10,5	4	4,3089	0,8536	F2	2 (ap, ln)	Premaxilla	Elliptical	Anastomosed
BSPG 1993 IX 331B	48	24	10,8	11,1	4,5	8,5	8,5	3	2,1621	0,7657	F1	0	-	Elliptical	Anastomosed
BSPG 1993 IX 331C	37	37	10	9	5,5	7,5	7,5	2	4,1111	0,8333	F4-F5	4 (lb, ln, m, d)	Maxilla	Elliptical	(worn) scratches and apical pits
BSPG 1993 IX 313A	27,5	27,5	8,2	8,2	4,5	6	6,2	1	3,3536	0,7560	F2	1 (ln)	Premaxilla	Elliptical	Longitudinal ridges (but worn)
BSPG 1993 IX 2A	51	50	10,5	11,5	5,5	8,5	9	2	4,3478	0,7826	F2	1 (ln)	Maxilla	Lemon shaped	Longitudinal ridges (but worn)
BSPG 1993 IX 2B	47,5	47,5	13	13	11,5	9	9,2	3	3,6538	0,7076	F2-F3	2 (lb, ln)	Anterior dentary	Lemon shaped	Longitudinal ridges
BSPG 1993 IX 2C	47	47	10	11,5	5	8	10	2	4,0869	0,8695	F2	1 (lb)	Anterior dentary	Lemon shaped	No (because of preservation)
BSPG 1993 IX 2D	38	38	9,5	10,5	1,5	6,5	7,5	1,5	3,6190	0,7142	F1	0	Anterior (?)	Lemon shaped	(worn) w/pits and scratches
PIMUZ A/III 0823 a	66	59	15	12	7	8	7	4	4,9166	0,5833	F2	1 (ln)	Premaxilla	Elliptical	Anastomosed (but worn)
PIMUZ A/III 0823 b	39	31	14,5	13	7	8	9	7	2,3846	0,6923	F2	1 (ln)	Premaxilla	D-shaped	Anastomosed
PIMUZ A/III 0823 c	57	34	11	9	5	6	6	3	3,7777	0,6666	F1	0	-	Lemon shaped	Anastomosed w/protrusions
PIMUZ A/III 0823 d	24	21	7	9	6	6	6	4	2,3333	0,6666	F2-F3	2 (lb, ln)	Anterior	Elliptical (worn)	Anastomosed w/protrusions
PIMUZ A/III 0823 e	43	37	10	9	8	9	7	4	4,1111	0,7777	F2	2 (lb, ln)	Premaxilla	Lemon shaped	Reticulate w/ protrusions
PIMUZ A/III 0823 f	42	38	9	8	4	6	7	3	4,75	0,875	F1	0	-	Elliptical	Smooth
PIMUZ A/III 0823 g	31	29	7	6	3	6	5	3	4,8333	0,8333	F4	3 (ln, m, d)	Maxilla	Lemon shaped (faint)	Smooth (w/ apicobasal striations)
PIMUZ A/III 0823 h	26	26	5	6	5	4	4,5	3	4,3333	0,75	F1-F2	1 (ap)	-	Elliptical	Smooth
PIMUZ A/III 0823 i	21	21	5	5,5	3	3	4	2	3,8181	0,7272	F2-F3	2 (lb, ln)	Anterior dentary	Lemon shaped	Smooth
PIMUZ A/III 0823 j	34	34	11	9	8	9	10	4	3,7777	1,1111	F4	3 (ln, m, d)	Maxilla	Elliptical (worn)	Chevron-like

## Table 2 (on next page)

One-way PERMANOVA comparisons for each pair of sauropod groups.

\*non-neosauropod eusauropod; \*\*non-titanosaurian titanosauriform; \*\*\*titanosaur; ?unknown or resolved in this study

	<i>Morph_I</i>	<i>Morph_II</i>	<i>Atsinganosaurus</i>	<i>Fox-Amphoux C-Morph</i>	<i>Fox Amphoux D-Morph</i>	<i>Lirainosaurus</i>	<i>LH Morph A</i>	<i>LH Morph B</i>	<i>Massecaps titanosaur</i>	<i>Patagosaurus</i>	<i>Euhelopus</i>	<i>Shiramine form</i>	<i>Astrodon</i>	<i>Neuquen titano indet</i>	<i>Uzbekistan Neosauropod</i>	<i>Maxakalisaurus</i>	<i>Bauru titanosaur</i>
<i>Morph_I</i> <sup>?</sup>		0,397 5	0,191 7	0,258 4	<b>0,038</b> 3	0,751 9	0,472 9	0,355 6	0,412 7	<b>0,000</b> 1	<b>0,000</b> 2	<b>0,009</b> 6	0,640 8	0,077 4	0,262 3	<b>0,000</b> 6	0,415 2
<i>Morph_II</i> <sup>?</sup>			<b>0,022</b> 8	0,075 2	0,180 9	0,862 3	0,875 8	0,115 6	1 1	<b>0,000</b> 8	<b>0,003</b> 8	0,113 1	0,856 8	0,370 2	0,090 9	<b>0,008</b> 7	0,763 2
<i>Atsinganosaurus</i> <sup>***</sup>				0,838 3	<b>0,008</b> 3	0,388 8	<b>0,042</b> 7	0,697 1	<b>0,034</b> 7	<b>0,000</b> 6	<b>0,002</b> 2	<b>0,009</b> 3	0,139 9	<b>0,049</b> 5	0,686 4	<b>0,016</b> 5	<b>0,035</b> 7
<i>FoxAmphoux C-Morph</i> <sup>***</sup>					<b>0,017</b> 2	0,189 2	0,053 2	0,645 8	0,095 4	<b>0,005</b> 4	<b>0,012</b> 2	<b>0,026</b> 8	0,142 3	0,106 5	0,600 5	<b>0,017</b> 6	0,102 8
<i>FoxAmphoux D-Morph</i> <sup>***</sup>						0,147 4	0,079 2	<b>0,014</b> 3	0,144 9	<b>0,000</b> 9	<b>0,007</b> 4	0,358 1	0,111 2	0,985 5	<b>0,005</b> 7	<b>0,008</b> 2	0,154 5
<i>Lirainosaurus</i> <sup>***</sup>							0,666 2	0,236 2	0,503 2	<b>0,022</b> 4	<b>0,037</b> 4	0,136 4	0,933 8	0,436 3	0,374 3	<b>0,05</b> 9	0,598 9
<i>LoHueco Morph A</i> <sup>***</sup>								0,093 2	0,711 9	<b>0,001</b> 5	<b>0,001</b> 9	<b>0,036</b> 6	0,908 1	0,321 3	0,091 3	<b>0,007</b> 9	0,660 7
<i>LoHueco Morph B</i> <sup>***</sup>									0,070 7	<b>0,001</b> 7	<b>0,002</b> 4	<b>0,009</b> 9	0,189 1	0,076 5	0,951 5	<b>0,009</b> 5	0,121 2
<i>Massecaps titanosaur</i> <sup>***</sup>										<b>0,005</b> 9	<b>0,011</b> 7	0,109 9	0,749 1	0,461 8	0,093 6	<b>0,019</b> 8	0,503 2
<i>Patagosaurus</i> <sup>*</sup>											<b>0,023</b> 3	<b>0,010</b> 5	<b>0,002</b> 1	<b>0,004</b> 7	<b>0,000</b> 2	<b>0,001</b> 2	<b>0,006</b> 1
<i>Euhelopus</i> <sup>**</sup>												0,190 2	<b>0,005</b> 1	0,081 8	<b>0,000</b> 2	<b>0,002</b> 2	<b>0,012</b> 8
<i>Shiramine form</i> <sup>**</sup>													0,143 3	0,563 9	<b>0,001</b> 4	<b>0,008</b> 6	0,170 7
<i>Astrodon</i> <sup>**</sup>														0,329 7	0,188 5	<b>0,007</b> 7	0,652 7
<i>Neuquen titanosaur</i> <sup>***</sup>															<b>0,021</b> 9	<b>0,007</b> 2	0,47 2

<i>Uzbekistan Neosauropod?</i>																<b>0,000 8</b>	0,096 8
<i>Maxakalisaurus***</i>																	<b>0,017 4</b>
<i>Bauru titanosaur***</i>																	

1