# A peer-reviewed version of this preprint was published in PeerJ on 2 April 2019.

<u>View the peer-reviewed version</u> (peerj.com/articles/6667), which is the preferred citable publication unless you specifically need to cite this preprint.

Villaseñor-Derbez JC, Fitzgerald S. 2019. Spatial variation in allometric growth of invasive lionfish has management implications. PeerJ 7:e6667 https://doi.org/10.7717/peerj.6667

# Spatial variation in allometric growth of invasive lionfish has management implications

Juan Carlos Villaseñor-Derbez Corresp., 1 , Sean Fitzgerald 1

<sup>1</sup> Bren School of Environmental Science and Management, University of California, Santa Barbara, Santa Barbara, CA, United States

Corresponding Author: Juan Carlos Villaseñor-Derbez Email address: jvillasenor@bren.ucsb.edu

Lionfish (*Pterois volitans / miles*) are an invasive species in the Western Atlantic and the Caribbean. Improving management of invasive lionfish populations requires accurate total biomass estimates, which depend on accurate estimates of allometric growth. Sedentary species like lionfish often exhibit high levels of spatial variation in life history characteristics. We review 17 published length-weight relationships for lionfish taken throughout their invasive range and found substantial regional differences in allometric growth parameters. The spatial pattern we observed is consistent with findings from other studies focusing on genetics or age-at-length. We show that the use of *ex situ* parameters can result in up to a threefold under- or overestimation of total weight, but using parameters from nearby regions reduces this error. These findings can have major implications for management in terms of predicting effects on local ecosystems, evaluating the effectiveness of removal programs, or estimating biomass available for harvest.

# Spatial variation in allometric growth of invasive lionfish has management

- implications
- **Juan Carlos Villaseñor-Derbez<sup>1</sup> and Sean Fitzgerald**
- $_{\rm 5}$   $^{-1}$  Bren School of Environmental Sciences and Management, University of California
- 6 Santa Barbara, Santa Barbara, California, USA
- 7 Corresponding author:
- 8 Juan Carlos Villaseñor-Derbez<sup>1</sup>
- 9 Email address: jvillasenor@ucsb.edu

## 10 ABSTRACT

Lionfish (Pterois volitans / miles) are an invasive species in the Western Atlantic and the Caribbean. 11 Improving management of invasive lionfish populations requires accurate total biomass estimates, which 12 depend on accurate estimates of allometric growth. Sedentary species like lionfish often exhibit high 13 levels of spatial variation in life history characteristics. We review 17 published length-weight relationships 14 for lionfish taken throughout their invasive range and found substantial regional differences in allometric 15 growth parameters. The spatial pattern we observed is consistent with findings from other studies 16 focusing on genetics or age-at-length. We show that the use of ex situ parameters can result in up to a 17 threefold under- or overestimation of total weight, but using parameters from nearby regions reduces 18 this error. These findings can have major implications for management in terms of predicting effects on 19 local ecosystems, evaluating the effectiveness of removal programs, or estimating biomass available for 20 harvest. 21

# 22 INTRODUCTION

Lionfish (Pterois volitans/miles complex) are an invasive species in the western Atlantic and Caribbean 23 Sea, likely introduced through liberation of aquarium-kept organisms (Betancur-R et al., 2011). They 24 are the first invasive marine vertebrates established along the North Atlantic Caribbean coasts (Schofield, 25 2009, 2010; Sabido-Itza et al., 2016) and their presence has been labeled as a major marine invasion 26 because they threaten local biodiversity, spread rapidly, and are difficult to manage (Hixon et al., 2016). 27 Lionfish have established invasive populations in coral reefs, estuaries, mangroves, hard-bottomed areas, 28 and mesophotic reefs (Barbour et al., 2010; Jud et al., 2011; Muñoz et al., 2011; Claydon et al., 2012; 29 Andradi-Brown et al., 2017; Gress et al., 2017). 30 A substantial amount of research describes lionfish impacts throughout its invaded range. A meta-31 analysis by Peake et al. (2018) showed that invasive lionfish prey on at least 167 different species across 32 the tropical and temperate North Atlantic. Their feeding behavior and high consumption rates can reduce 33 recruitment and population sizes of native reef-fish species, and can further endanger reef fish (Green et al. 34 (2012); Rocha et al. (2015); but see Hackerott et al. (2017)). For example, field experiments by Albins 35 and Hixon (2008) showed that lionfish establishment led to reduced recruitment of native fishes by nearly 36 80% over a five-week period in Florida. Green et al. (2012) reported that prey fish biomass declined by 37 65% over two years as lionfish biomass increased along Bahamian coral reefs. Their trophic impacts can 38 be minimized if local lionfish biomass is controlled by culling (Arias-Gonzalez et al., 2011). 39 Governments and non-profit organizations have sought to reduce lionfish densities through removal 40 programs and incentivizing its consumption (Chin et al., 2016). In some cases, these have shown to 41 significantly reduce –but not quite eliminate– lionfish abundances at local scales (de Leon et al., 2013; 42 Sandel et al., 2015). Complete eradication of lionfish through fishing is unlikely because of their rapid 43

- <sup>44</sup> recovery rates and ongoing recruitment to shallow-water areas from persistent populations in mesophotic
- ecosystems (Barbour et al., 2011; Andradi-Brown et al., 2017). However, promoting lionfish consumption

might create a level of demand capable of incentivizing a stable fishery while controlling shallow-water
 populations, thus creating alternative livelihoods and avoiding further impacts to local biota.

The feasibility of establishing fisheries through lionfish removal programs has been extensively

<sup>49</sup> evaluated through field observations and empirical modeling (Barbour et al., 2011; Morris et al., 2011; de

<sup>50</sup> Leon et al., 2013; Johnston and Purkis, 2015; Sandel et al., 2015; Usseglio et al., 2017). Determining the

<sup>51</sup> feasibility of such initiatives requires modeling the change in biomass in response to changes in fishing <sup>52</sup> mortality (*i.e.* culling). A common way to model this is via length-structured population models, where

<sup>52</sup> mortality (*i.e.* culling). A common way to model this is via length-structured population models, where <sup>53</sup> fish lengths are converted to weight to calculate total biomass (Barbour et al., 2011; Côté et al., 2014;

fish lengths are converted to weight to calculate total biomass (Barbour et al., 2011; Côté et al., 2014;
 Andradi-Brown et al., 2017). The allometric length-weight relationship is thus an essential component of

<sup>55</sup> these models, but this relationship can vary across regions as a response to biotic and abiotic conditions

<sup>56</sup> (Johnson and Swenarton, 2016).

Outcomes of previous studies suggest lionfish are likely to exhibit spatial heterogeneity in the length-57 weight relationship, which we summarize in two main causes. First, culling programs are effective in 58 reducing local adult populations largely because lionfish exhibit high levels of site fidelity and small home 59 ranges (Fishelson, 1997; Kochzius and Blohm, 2005; Jud and Layman, 2012; Côté et al., 2014). It is 60 know that fish with sedentary behavior are likely to exhibit high levels of spatial variation in important life 61 history characteristics such as growth or natural mortality rates (Gunderson et al., 2008; Hutchinson, 2008; 62 Wilson et al., 2012; Guan et al., 2013). Second, genetic analysis of lionfish suggests biological differences 63 due to the existence of two genetically distinct invasive subpopulations between the northwest Atlantic 64 and the Caribbean (Betancur-R et al., 2011). Site-specific parameters are necessary to accurately estimate 65 biomass when allometric relationships are spatially variable, and this variability is increasingly important 66 when estimating the potential effectiveness of lionfish culling programs (Barbour et al., 2011; Morris et al., 67 2011; Côté et al., 2014; Johnston and Purkis, 2015). However, the region-wide differences in allometric 68 growth parameters has remained unexplored for lionfish, despite the large number of site-specific studies 69 reporting the length-weight relationship. 70 Here, we compare previously published length-weight relationships for lionfish populations in North 71

<sup>72</sup> Carolina, Northern and Southern Gulf of Mexico, the Southern Mexican Caribbean, Bahamas, Little

<sup>73</sup> Cayman, Jamaica, Bonaire, Puerto Rico, and Costa Rica (Barbour et al., 2011; Darling et al., 2011; de

Leon et al., 2013; Fogg et al., 2013; Dahl and Patterson, 2014; Edwards et al., 2014; Toledo-Hernández,

<sup>75</sup> 2014; Sandel et al., 2015; Aguilar-Perera and Quijano-Puerto, 2016; Sabido-Itza et al., 2016; Sabido-Itzá

et al., 2016; Chin et al., 2016). We also collected lionfish length and weight data in the central Mexican

77 Caribbean and report the first allometric growth equation for this region. The objective of this paper is to

describe the spatial pattern of length-weight relationships of lionfish across the Caribbean and Western

79 Atlantic and to discuss implications of these spatial differences.



**Figure 1.** Locations where allometric growth parameters of lionfish (*Pterois spp*) have been reported. Circle sizes indicate sample size from each study, colors indicate the *b* coefficient from Eq. 1.

## 80 METHODS

We reviewed 12 published studies and obtained 17 length-weight relationships for the North Atlantic 81 (n = 1), Gulf of Mexico (n = 7), and Caribbean (n = 9), Table 2, Fig 3). We collected information on 82 sampling methods, sex differentiation, location, and depth ranges from each study when available. Only 83 two studies reported parameters for each gender (Aguilar-Perera and Quijano-Puerto, 2016; Fogg et al., 84 2013), so we assumed both genders were included in a study if gender was unspecified. Reviewed studies 85 presented information for organisms obtained at depths between 0.5 m and 57 m. Three studies explicitly 86 stated that their organisms were sampled with pole spears (Dahl and Patterson, 2014; Aguilar-Perera and 87 Quijano-Puerto, 2016; Chin et al., 2016; Sabido-Itzá et al., 2016), and five studies mentioned that some of 88 their organisms were obtained with pole spears (or other type of harpoon) but also hand-held nets or fish 89 traps (Barbour et al., 2011; Fogg et al., 2013; Edwards et al., 2014; Toledo-Hernández, 2014; Sandel et al., 90 2015; Sabido-Itza et al., 2016; Sabido-Itzá et al., 2016), and two studies did not specify how organisms 91 were sampled (Darling et al., 2011; de Leon et al., 2013). Fogg et al. (2013) use spineless weight in their 92 calculations, so their parameters likely underestimated total weight. Since no spineless to total weight 93 conversions were available, these parameters were taken as reported. 94 We also used data from Villaseñor-Derbez and Herrera-Pérez (2014), who collected organisms from 95 10 sampling sites along the central Mexican Caribbean coast in 2010 (Supplementary Table 1). Sampling

<sup>96</sup> 10 sampling sites along the central Mexican Caribbean coast in 2010 (Supplementary Table 1). Sampling <sup>97</sup> locations included wall and carpet reefs at depths between 5.7 m and 38.1 m. All observed lionfish (n <sup>98</sup> = 109) were collected using hand nets and numbered collection bottles. The use of hand nets prevented <sup>99</sup> any weight loss due to bleeding and allowed better representation of small sizes by eliminating gear <sup>100</sup> selectivity. Organisms were euthanized via pithing and Total Length (TL; mm) and Total Weight (TW; g) <sup>101</sup> were recorded.

The weight-at-length relationship for lionfish in the central Mexican Caribbean was calculated with the allometric growth function:

$$TW = aTL^b \tag{1}$$

104

Where *a* is the ponderal index and *b* is the scaling exponent or allometric parameter.

<sup>105</sup> Transforming this equation via base-10 logarithms we obtain:

$$log_{10}(TW) = b \times log_{10}(TL) + log_{10}(a)$$
<sup>(2)</sup>

<sup>106</sup> This can be simplified and re-written as:

$$Y = bX + c \tag{3}$$

Where  $Y = log_{10}(TW)$ ,  $X = log_{10}(TL)$ , and  $c = log_{10}(a)$ . The coefficients (*c* and *b*) were estimated with an Ordinary Least Squares Regression and heteroskedastic-robust standard error correction (Zeileis, 2004). When the b = 3, it is said that the organism exhibits a perfect isometric growth, so the *b* coefficient was tested against the null hypothesis of isometric growth (*i.e.*  $H_0 : b = 3$ ). Coefficients were tested with a two-tailed Student's t, and the significance of the regression was corroborated with an F-test.

Some of the reviewed studies inconsistently defined *a* as either the ponderal index from Eq. 1 or the y-intercept (*c*) from Eq. 3. Other studies incorrectly reported parameters as mm-to-g conversions when they were in fact cm-to-g conversions. We standardized each study by converting coefficients and report all parameters as TL(mm) to TW (gr) conversions. Locations where allometric studies have been performed are shown in Figure 1 and summarized in Table 2.

We obtained a total of 18 parameter pairs by combining length-weight parameters extracted from 117 the literature and the additional pair calculated here. We used the central Mexican Caribbean as a case 118 study of how the use of *ex situ* parameters influences the accuracy of weight estimates for lionfish. We 119 estimated TW from the TL observations we collected in the central Mexican Caribbean (n = 109, with 120  $TL \in (34, 310)$ ) using each of the 18 parameter pairs and divided predicted weights by known observed 121 weights to obtain a simple measure of over- or underestimation. Difference in mean weight ratios across 122 the different parameter pairs were tested with a one-way analysis of variance (ANOVA) and Tukey's test 123 was used for post-hoc tests. All analyses were performed in R version 3.5.1 (R Core Team, 2018). Raw 124 data and code used in this work are available on github. 125

#### 126 RESULTS

The length-weight relationship for organisms from the central Mexican Caribbean resulted in the coefficient values  $a = 3.2056297 \times 10^{-6}$ , b = 3.2347391 and c = -5.4940866 ( $R^2 = 0.977$ , F(df = 1; 107) = 6928.67, p < 0.001). The allometric factor (b) was significantly different from b = 3(t(107) = 6.04; p < 0.001) indicating that lionfish present allometric growth. The length-weight coefficients estimated in this study were within the range identified by studies in other regions (Table 2). Figure 2 shows the relationship between TL and TW for this region, and model fit statistics are presented in Table 1.

	$log_{10}(TW)$			
с	-5.494 (0.083)***			
b	3.235 (0.039)***			
F Statistic	6928.67*** (df = 1; 107)			
Observations	109			
Adjusted R <sup>2</sup>	0.976			
Residual Std. Error	0.096 (df = 107)			
Note:	*p<0.1; **p<0.05; ***p<0.01			

**Table 1.** Coefficients of the linear model fit to Eq 3. Numbers in parenthesees represent heteroskedastic-robust standard errors.

There were significant differences in our predicted weights for the central Mexican Caribbean when using the different pairs of parameters (F(df = 17; 1944) = 61.55; p < 0.001). The lowest weight

estimates resulted from using the allometric parameters from Banco Chinchorro in the Caribbean, with



**Figure 2.** Length-weight relationship for 109 lionfish sampled in the central Mexican Caribbean. Points indicate samples, dashed black line indicates curve of best fit, marginal plots represent the density distribution of each variable.

<sup>137</sup> mean  $\pm$  SD of 40.37  $\pm$  58.74 gr (Sabido-Itzá et al., 2016), and the highest weight estimates came from <sup>138</sup> the Northern Atlantic with 73.76  $\pm$  96.11 gr (Barbour et al., 2011). To put this in context, true observed <sup>139</sup> weights were 52.56  $\pm$  76.58 gr. These correspond to predicted-to-observed weights ratios of 0.80  $\pm$  0.19 <sup>140</sup> and 1.76  $\pm$  0.50 (mean  $\pm$  SD), respectively.

The calculated ratio of predicted-to-observed weight ranged from 0.36 to 3.51, indicating that ex situ 141 parameters can result in major under- and overestimation. Tukey's post-hoc test suggests that weight ratios 142 for the central Mexican Caribbean were not different from those obtained with parameters from Little 143 Cayman, the Bahamas, and some sites in the Gulf of Mexico (Tukey's HSD p > 0.05). Weight estimates 144 using parameters from the Gulf of Mexico and North-Western Atlantic were higher on average than those 145 from the Caribbean (Fig 3). The average ( $\pm$  SD) predicted-to-observed weight ratios from these three 146 regions were  $1.24 \pm 0.309$ ,  $1.76 \pm 0.496$ , and  $1.17 \pm 0.398$ , respectively. Predicted-to-observed weight 147 ratios are presented in Figure 4. Spineless weight parameters from Fogg et al. (2013) still produced 148 predicted-to-observed weight ratios > 1. 149

**Table 2.** Summary of 18 allometric growth parameters available for lionfish in the invaded range from peer-reviewed literature and this study. All parameters have been adjusted to convert from millimeters to grams. n = Sample size, Sex specifies whether data was presented for Females (F), Males (M), or both genders combined (B), a = scaling parameter for Eq. 1 (presented in  $\times 10^{-5}$ ), c = y-intercept for Eq. 3, b = exponent or slope for Eq. 1 or Eq. 3, respectively. The Fit column contains the reported  $R^2$  of the model fit.

Region	Sex	n	a	b	с	Fit	Reference
Caribbean	В	458	3.6	2.81	-4.44	-	Sandel et al., 2015
Caribbean	В	419	2.8	2.85	-4.56	0.8715	Chin et al., 2016
Caribbean	В	1450	2.3	2.89	-4.64	0.96	de Leon et al., 2013
Caribbean	В	1887	0.3	3.24	-5.52	0.97	Edwards et al., 2014
Caribbean	В	-	0.25	3.29	-5.60	-	Darling et al., 2011
Caribbean	В	2143	0.52	3.18	-5.28	0.9907	Sabido-Itza et al., 2016
Caribbean	В	227	0.8	3.11	-5.10	0.958	Toledo-Hernández et al., 2014
Caribbean	В	449	0.23	3.25	-5.64	0.97	Sabido-Itza et al., 2016b
Caribbean	В	368	0.32	3.19	-5.50	0.98	Sabido-Itza et al., 2016b
Caribbean	В	109	0.32	3.23	-5.49	0.9766	This study
GoM	В	934	0.21	3.34	-5.68	0.98	Dahl & Patterson, 2014
GoM	В	472	0.29	3.30	-5.54	0.95	Aguilar-Perera & Quijano-Puerto, 2016
GoM	F	67	0.12	3.47	-5.93	0.95	Aguilar-Perera & Quijano-Puerto, 2016
GoM	Μ	59	0.42	3.23	-5.38	0.95	Aguilar-Perera & Quijano-Puerto, 2016
GoM	В	582	0.14	3.43	-5.86	0.99	Fogg et al., 2013
GoM	М	119	0.27	3.31	-5.57	0.97	Fogg et al., 2013
GoM	F	115	0.68	3.14	-5.17	0.94	Fogg et al., 2013
North Atlantic	В	774	2.9	2.89	-4.54	-	Barbour et al., 2011





**Figure 3.** Length-weight relationships (n = 18) for 12 studies and this study. Colors indicate studies from which the parameters were extracted. Dotted, dashed and solid lines show models for males, females, and combined sexes, respectively. The dashed black line represents the relationship estimated in this study.



**Figure 4.** Violin plot of predicted-to-observed weight ratios for 18 pairs of allometric parameters. Blue circles indicate median values and Like letters indicate values that do not differ significantly.

8/11

### 150 DISCUSSION

Our results suggest that lionfish exhibit highly variable allometric relationships across the invaded range, and that this variation is related to space. Moreover, we shot that the use of *ex situ* parameters may lead to highly biased weight estimates. Our comparison of observed weights to those predicted with locally-informed parameters and *ex situ* parameters showed that weight can be overestimated by more than a three-fold, and highlights the need to use local information. Here we discuss the implications of our findings and highlight potential future research directions.

We detected substantial differences in weight-at-length between organisms from the Caribbean, Gulf 157 of Mexico, and North-Western Atlantic. Groupings of predicted-to-observed weight ratios aligned with 158 the spatial distribution of the examined studies, suggesting that these differences are mediated by space. 159 These regional allometric differences mirror similar patterns in age-at-length of lionfish across both their 160 invaded and native regions (Pusack et al., 2016). Variation may be driven by genetics or by organisms 161 exposure to distinct environmental conditions. For example, Betancur-R et al. (2011) used mitochondrial 162 DNA to demonstrate the existence of two distinct population groups, identified as the "Caribbean group" 163 and "Northern Group", and Fogg et al. (2015) alternatively suggested that age-at-length differences may 164 be climate-driven. Differences in weight-at-length could also reflect differential energy input or usage, or 165 a combination of both. Future research is needed to determine which processes are at work here. 166

Differences in length-weight relationships have traditionally been highlighted as potential pitfalls to 167 fishery management. For example, Wilson et al. (2012) show that small-scale variations in length-at-age 168 and fishing mortality in other Scorpaeniformes translate to differential landings, effort, and catch per 169 unit effort in the live fish fishery of California, and that these differences must be taken into account in 170 management plans. The lionfish case poses the opposite scenario, where the manager desires to eradicate 171 the species. To accurately gauge both the effectiveness of lionfish removal efforts and the resources 172 needed to successfully manage an invasion, we must acknowledge and understand regional biological 173 differences in important variables such as allometric growth parameters. 174

175 The results presented here have major implications for management. For example, Edwards et al. (2014) simulated a lionfish culling program under two scenarios, one using length-at-age and length-to-176 weight parameters from North Carolina and one using parameters from Little Cayman. Their results show 177 that using different parameters caused up to a four-year difference in the time required for the simulated 178 lionfish population to recover to 90% of its initial biomass after removals ceased. Here, we show that 179 180 using one set of length-weight parameters versus another for a given length can result in more than a threefold under- or overestimation of total weight. These spatially-driven differences become especially 181 important when allocating resources for lionfish removal programs, incentivizing lionfish fisheries as a 182 source of alternative livelihoods, or estimating ecosystem impacts. Research efforts focused on invasive 183 lionfish populations need to use parameters calculated for their region to the extent possible, or at least 184 use reasonable sets of different parameters that provide upper and lower bounds in their results. 185

### **ACKNOWLEDGEMENTS**

The authors would like to thank thank Nils Van Der Haar and Michael Doodey from Dive Aventuras as
 well as Guillermo Lotz-Cador who provided help to collect samples.

<sup>189</sup> Conflict of Interest: The authors declare that they have no conflict of interest.

### 190 REFERENCES

Aguilar-Perera, A. and Quijano-Puerto, L. (2016). Relations between fish length to weight, and otolith length and weight, of the lionfish pterois volitans in the parque nacional arrecife alacranes, southern

- <sup>193</sup> gulf of mexico. *Rev. biol. mar. oceanogr.*, 51(2):469–474.
- Albins, M. and Hixon, M. (2008). Invasive indo-pacific lionfish pterois volitans reduce recruitment of
   atlantic coral-reef fishes. *Mar. Ecol. Prog. Ser.*, 367:233–238.
- Andradi-Brown, D. A., Grey, R., Hendrix, A., Hitchner, D., Hunt, C. L., Gress, E., Madej, K., Parry, R. L.,
- <sup>197</sup> Régnier-McKellar, C., Jones, O. P., a Arteaga, M., Izaguirre, A. P., Rogers, A. D., and Exton, D. A.
- <sup>198</sup> (2017). Depth-dependent effects of culling-do mesophotic lionfish populations undermine current
- <sup>199</sup> management? *R Soc Open Sci*, 4(5):170027.
- Arias-Gonzalez, J. E., Gonzalez-Gandara, C., Luis Cabrera, J., and Christensen, V. (2011). Predicted

- impact of the invasive lionfish pterois volitans on the food web of a caribbean coral reef. *Environ Res*, 111(7):917–925. Barbour A Montgomery M Adamson A Diaz-Ferguson E and Silliman B (2010) Mangrove use
- Barbour, A., Montgomery, M., Adamson, A., Diaz-Ferguson, E., and Silliman, B. (2010). Mangrove use
   by the invasive lionfish pterois volitans. *Mar. Ecol. Prog. Ser.*, 401:291–294.
- Barbour, A. B., Allen, M. S., Frazer, T. K., and Sherman, K. D. (2011). Evaluating the potential efficacy
   of invasive lionfish (pterois volitans) removals. *PLoS ONE*, 6(5):e19666.
- Betancur-R, R., Hines, A., Acero, A., Orti, G., Wilbur, A., and Freshwater, D. (2011). Reconstructing the
   lionfish invasion: insights into greater caribbean biogeography. *J Biogeography*, 38:1281–1293.
- Chin, D. A., Aiken, K. A., and Buddo, D. (2016). Lionfish population density in discovery bay, jamaica.
   *International Journal of Scientific & Engineering Research*, 7(12):1327 1331.
- Claydon, J., Calosso, M., and Traiger, S. (2012). Progression of invasive lionfish in seagrass, mangrove
   and reef habitats. *Mar. Ecol. Prog. Ser.*, 448:119–129.
- Côté, I. M., Akins, L., Underwood, E., Curtis-Quick, J., and Green, S. J. (2014). Setting the record
   straight on invasive lionfish control: Culling works. *PeerJ*.
- <sup>215</sup> Dahl, K. A. and Patterson, W. F. (2014). Habitat-specific density and diet of rapidly expanding invasive <sup>216</sup> red lionfish, pterois volitans, populations in the northern gulf of mexico. *PLoS ONE*, 9(8):e105852.
- 217 Darling, E. S., Green, S. J., O'Leary, J. K., and Côté, I. M. (2011). Indo-pacific lionfish are larger and
- more abundant on invaded reefs: a comparison of kenyan and bahamian lionfish populations. *Biol Invasions*, 13(9):2045–2051.
- de Leon, R., Vane, K., Bertuol, P., Chamberland, V. C., Simal, F., Imms, E., and Vermeij, M. J. A.
- (2013). Effectiveness of lionfish removal efforts in the southern caribbean. *Endanger Species Res*, 22(2):175–182.
- Edwards, M. A., Frazer, T. K., and Jacoby, C. A. (2014). Age and growth of invasive lionfish (pterois spp.) in the caribbean sea, with implications for management. *BMS*, 90(4):953–966.
- Fishelson, L. (1997). Experiments and observations on food consumption, growth and starvation in dendrochirus brachypterus and pterois volitans (pteroinae, scorpaenidae). *Environmental Biology of Fishes*, 50(4):391–403.
- Fogg, A. Q., Evans, J. T., Ingram JR, G. W., Peterson, M. S., and Brown-Peterson, N. J. (2015). Comparing age and growth patterns of invasive lionfish among three ecoregions of the northern gulf of mexico. In
- age and growth patterns of invasive lionfish among three ecoregions of the northern gulf of mexico. In
   GCFI, G. and Institute, C. F., editors, *Proceedings of the 68 th Gulf and Caribbean Fisheries Institute*,
- Panama City. Gulf and Caribbean Fisheries Institute.
- Fogg, A. Q., Hoffmayer, E. R., Driggers, W. B., Campbell, M. D., Pellegrin, G. J., and Stein, W. (2013).
- Distribution and length frequency of invasive lionfish (pterois sp.) in the northern gulf of mexico. *GCR*,
   234
- Green, S. J., Akins, J. L., Maljković, A., and Côté, I. M. (2012). Invasive lionfish drive atlantic coral reef fish declines. *PLoS ONE*, 7(3):e32596.
- Gress, E., Andradi-Brown, D. A., Woodall, L., Schofield, P. J., Stanley, K., and Rogers, A. D. (2017).
   Lionfish (pterois spp.) invade the upper-bathyal zone in the western atlantic. *PeerJ*, 5:e3683.
- Guan, W., Cao, J., Chen, Y., and Cieri, M. (2013). Impacts of population and fishery spatial structures on fishery stock assessment. *Can. J. Fish. Aquat. Sci.*, 70(8):1178–1189.
- Gunderson, D. R., Parma, A. M., Hilborn, R., Cope, J. M., Fluharty, D. L., Miller, M. L., Vetter, R. D.,
- Heppell, S. S., and Greene, H. G. (2008). The challenge of managing nearshore rocky reef resources.
- 243 *Fisheries*, 33(4):172–179.
- Hackerott, S., Valdivia, A., Cox, C. E., Silbiger, N. J., and Bruno, J. F. (2017). Invasive lionfish had no
   measurable effect on prey fish community structure across the belizean barrier reef. *PeerJ*, 5:e3270.
- Hixon, M., Green, S., Albins, M., Akins, J., and Morris, J. (2016). Lionfish: a major marine invasion.
   *Mar. Ecol. Prog. Ser.*, 558:161–165.
- Hutchinson, W. F. (2008). The dangers of ignoring stock complexity in fishery management: the case of
   the north sea cod. *Biol Lett*, 4(6):693–695.
- Johnson, E. G. and Swenarton, M. K. (2016). Age, growth and population structure of invasive lionfish
- (pterois volitans/miles) in northeast florida using a length-based, age-structured population model.
   *PeerJ*, 4:e2730.
- Johnston, M. and Purkis, S. (2015). A coordinated and sustained international strategy is required to turn the tide on the atlantic lionfish invasion. *Mar. Ecol. Prog. Ser.*, 533:219–235.
- Jud, Z., Layman, C., Lee, J., and Arrington, D. (2011). Recent invasion of a florida (usa) estuarine system

- <sup>256</sup> by lionfish pterois volitans / p. miles. *Aquat. Biol.*, 13(1):21–26.
- Jud, Z. R. and Layman, C. A. (2012). Site fidelity and movement patterns of invasive lionfish, pterois
- spp., in a florida estuary. *Journal of Experimental Marine Biology and Ecology*, 414-415:69–74.
- Kochzius, M. and Blohm, D. (2005). Genetic population structure of the lionfish pterois miles (scorpaenidae, pteroinae) in the gulf of aqaba and northern red sea. *Gene*, 347(2):295–301.
- Morris, J. A., Shertzer, K. W., and Rice, J. A. (2011). A stage-based matrix population model of invasive
- lionfish with implications for control. *Biol Invasions*, 13(1):7–12.
- <sup>263</sup> Muñoz, R., Currin, C., and Whitfield, P. (2011). Diet of invasive lionfish on hard bottom reefs of the <sup>264</sup> southeast usa: insights from stomach contents and stable isotopes. *Mar. Ecol. Prog. Ser.*, 432:181–193.
- Peake, J., Bogdanoff, A. K., Layman, C. A., Castillo, B., Reale-Munroe, K., Chapman, J., Dahl, K.,
- Patterson III, W. F., Eddy, C., Ellis, R. D., Faletti, M., Higgs, N., Johnston, M. A., Muñoz, R. C.,
- Sandel, V., Villaseñor-Derbez, J. C., and Morris, J. A. (2018). Feeding ecology of invasive lionfish
- (pterois volitans and pterois miles) in the temperate and tropical western atlantic. *Biol Invasions*.
- Pusack, T. J., Benkwitt, C. E., Cure, K., and Kindinger, T. L. (2016). Invasive red lionfish (pterois volitans) grow faster in the atlantic ocean than in their native pacific range. *Environ. Biol. Fishes*, 99(6-7):571–579.
- 99(0-7):371-379.
- R Core Team (2018). *R: A Language and Environment for Statistical Computing*. R Foundation for
   Statistical Computing, Vienna, Austria.
- Rocha, L. A., Rocha, C. R., Baldwin, C. C., Weigt, L. A., and McField, M. (2015). Invasive lionfish
   preying on critically endangered reef fish. *Coral Reefs*, 34(3):803–806.
- 276 Sabido-Itzá, M., Aguilar-Perera, A., and Medina-Quej, A. (2016). Length-weight and length-length
- relations, and relative condition factor of red lionfish, pterois volitans (actinopterygii: Scorpaeniformes: Scorpaenidae), from two natural protected areas in the mexican caribbean. *Acta Icth et Piscat*, 46(4):270–285
- <sup>279</sup> 46(4):279–285.
- Sabido-Itza, M., Medina-Quej, A., De Jesus-Navarrete, A., Gomez-Poot, J., and Garcia-Rivas, M. (2016).
   Uso de la estructura de tallas como evidencia del establecimiento poblacional del pez le?n pterois
   volitans (scorpaeniformes: Scorpaenidae) en el sur del caribe mexicano. *RBT*, 64(1):353.
- <sup>283</sup> Sandel, V., Martínez-Fernández, D., Wangpraseurt, D., and Sierra, L. (2015). Ecology and management
- of the invasive lionfish pterois volitans/miles complex (perciformes: Scorpaenidae) in southern costa
- rica. *Rev Biol Trop*, 63(1):213–221.
- Schofield, P. (2009). Geographic extent and chronology of the invasion of non-native lionfish (pterois
- volitans [linnaeus 1758] and p. miles [bennett 1828]) in the western north atlantic and caribbean sea.
   AI, 4(3):473–479.
- Schofield, P. (2010). Update on geographic spread of invasive lionfishes (pterois volitans [linnaeus, 1758]
- and p. miles [bennett, 1828]) in the western north atlantic ocean, caribbean sea and gulf of mexico. *AI*,
   5(Supplement 1):S117–S122.
- Toledo-Hernández, C. (2014). Population ecology and genetics of the invasive lionfish in puerto rico. *AI*, 9(2):227–237.
- <sup>294</sup> Usseglio, P., Selwyn, J. D., Downey-Wall, A. M., and Hogan, J. D. (2017). Effectiveness of removals of <sup>295</sup> the invasive lionfish: how many dives are needed to deplete a reef? *PeerJ*, 5:e3043.
- <sup>296</sup> Villaseñor-Derbez, J. C. and Herrera-Pérez, R. (2014). Brief description of prey selectivity and ontogenetic
- changes in the diet of the invasive lionfish pterois volitans (actinopterygii, scorpaenidae) in the mexican caribbean. *PANAMJAS*, 9(2):131–135.
- <sup>299</sup> Wilson, J. R., Kay, M. C., Colgate, J., Qi, R., and Lenihan, H. S. (2012). Small-scale spatial variation in <sup>300</sup> population dynamics and fishermen response in a coastal marine fishery. *PLoS ONE*, 7(12):e52837.
- <sup>301</sup> Zeileis, A. (2004). Econometric computing with hc and hac covariance matrix estimators. *J Stat Softw*, 11(10).