

**A peer-reviewed version of this preprint was published in PeerJ on 9 April 2019.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.6458) (peerj.com/articles/6458), which is the preferred citable publication unless you specifically need to cite this preprint.

Wang C, Guo H, Han D, Yu Z, Zheng Y, Jiang H, Gao Y, Yuan B, Zhang J. 2019. Pituitary tissue-specific miR-7a-5p regulates FSH expression in rat anterior adenohypophyseal cells. PeerJ 7:e6458  
<https://doi.org/10.7717/peerj.6458>

# Pituitary tissue-specific miR-7a-5p regulates FSH expression in rat anterior adenohypophyseal cells

**Chang-Jiang Wang**<sup>1</sup>, **Hai-Xiang Guo**<sup>1</sup>, **Dong-Xu Han**<sup>1</sup>, **Ze-Wen Yu**<sup>1</sup>, **Yi Zheng**<sup>1</sup>, **Hao Jiang**<sup>1</sup>, **Yan Gao**<sup>1</sup>, **Bao Yuan**<sup>Corresp., 1</sup>, **Jia-Bao Zhang**<sup>Corresp. 1</sup>

<sup>1</sup> Department of Laboratory Animals, College of Animal Sciences, Jilin University, Changchun, Jilin, China

Corresponding Authors: Bao Yuan, Jia-Bao Zhang  
Email address: yuan\_bao@jlu.edu.cn, zjb@jlu.edu.cn

Follicle-stimulating hormone (FSH), which is synthesized and secreted by the anterior pituitary gland, plays an important role in regulating reproductive processes. In this study, using the TargetScan program, we predicted that miRNAs regulate FSH secretion. Dual-luciferase reporter assays were performed and identified miR-7a-5p. MiR-7a-5p has been reported to regulate diverse cellular functions. However, it is unclear whether miR-7a-5p binds to mRNAs and regulates reproductive functions. Therefore, we constructed a suspension of rat anterior pituitary cells and cultured them under adaptive conditions, transfected miR-7a-5p mimics or inhibitor into the cell suspension and detected expression of the FSHb gene. The results demonstrated that miR-7a-5p downregulated FSHb expression levels, while treatment with miR-7a-5p inhibitors upregulated FSHb expression levels relative to those of negative control groups, as shown by quantitative PCR analysis. The results were confirmed with a subsequent experiment showing that FSH secretion was reduced after treatment with mimics and increased in the inhibitor groups, as shown by ELISA. Our results indicated that miR-7a-5p downregulates FSHb expression levels, resulting in decreased FSH synthesis and secretion, which demonstrates the important role of miRNAs in the regulation of FSH and animal reproduction.

# **Pituitary tissue-specific miR-7a-5p regulates FSH expression in rat anterior adenohypophyseal cells**

Chang-Jiang Wang#, Hai-Xiang Guo#, Dong-Xu Han, Ze-Wen Yu, Yi Zheng, Hao Jiang, Yan Gao, Bao Yuan\*, Jia-Bao Zhang\*

Department of Laboratory Animals, College of Animal Sciences, Jilin University, Changchun 130062, Jilin, P.R. China

# These authors contributed equally to this work.

\* Corresponding author:

Jia-Bao Zhang, Department of Laboratory Animals, College of Animal Sciences, Jilin University, 5333 Xi'an Road, Changchun 130062 (P.R. China), Tel./Fax +86 431 87836551, E-mail: zjb515@126.com

Bao Yuan, Department of Laboratory Animals, College of Animal Sciences, Jilin University, 5333 Xi'an Road, Changchun 130062, (P.R. China), Tel./Fax +86 431 87836536, E-mail: yuan\_bao@jlu.edu.cn

# ABSTRACT

Follicle-stimulating hormone (FSH), which is synthesized and secreted by the anterior pituitary gland, plays an important role in regulating reproductive processes. In this study, using the TargetScan program, we predicted that miRNAs regulate FSH secretion. Dual-luciferase reporter assays were performed and identified miR-7a-5p. MiR-7a-5p reportedly regulates diverse cellular functions. However, whether miR-7a-5p binds to mRNAs and regulates reproductive functions is unclear. Therefore, we created a suspension of rat anterior pituitary cells and cultured them under adaptive conditions. The cell suspension was transfected with miR-7a-5p mimics or inhibitors, and the expression of the FSHb gene was detected. MiR-7a-5p downregulated FSHb expression levels, while compared to the negative control, miR-7a-5p inhibitors upregulated FSHb expression levels, as shown by quantitative PCR analysis. The results were confirmed with a subsequent experiment showing that FSH secretion was reduced after treatment with mimics and increased after treatment with inhibitors, as shown by ELISA. Our results indicated that miR-7a-5p downregulates FSHb expression levels, resulting in decreased FSH synthesis and secretion, demonstrating the important role of miRNAs in FSH regulation and animal reproduction.

## INTRODUCTION

The pituitary gland, which is located in the sella turcica at the base of the crania and is composed of the adenohypophysis and neurohypophysis(Hong et al. 2016), is involved in complex feedback mechanisms. This gland secretes several specific regulatory hormones to control peripheral endocrine organs, such as the thyroid, adrenal gland and gonads, after receiving information from the brain via the hypothalamus(Treier & Rosenfeld 1996). Consequently, the pituitary gland can regulate vital processes such as metabolism(Rodriguez-Pacheco et al. 2013), growth(Vyshnevs'ka & Bol'shova 2013), reproduction(Musumeci et al. 2015) and behavior(Shin et al. 2013). As the most important internal secretion gland in mammals, the pituitary plays a role throughout an animal's life by releasing seven types of hormones(Yuan et al. 2015).

Gonadotropin follicle-stimulating hormone (FSH) plays a critical role in modulating the reproductive health of both sexes. FSH stimulates spermatogenesis and androgen production in males and regulates the cyclic recruitment of follicles, follicle development, and ovulation triggering in females(Hunzicker-Dunn & Maizels 2006; Rull et al. 2018; Walker & Cheng 2005). FSH, which is synthesized and secreted by the anterior pituitary gland(Sheng et al. 2018), together with luteinizing hormone (LH), human chorionic gonadotropin (HCG) and thyroid-stimulating hormone (TSH), comprises a class of the glycoprotein hormone family(Telikicherla et al. 2011). FSH consists of a common  $\alpha$  subunit shared by all glycoprotein hormones and a unique  $\beta$  subunit that determines its unique biological function(Lamminen et al. 2005). The  $\alpha$  subunit combines with hormone-specific  $\beta$  subunits to form heterodimers, and the extracellular domain of the  $\beta$  subunit has important binding sites for the membrane-bound receptor(Fan & Hendrickson 2005).

MicroRNAs (miRNAs) are genome-encoded small noncoding RNAs that are 22 nucleotides long in their mature form(Bartel 2009). MiRNAs can regulate gene expression at the posttranscriptional level by recognizing the 3'UTR sequence of target messenger RNAs(Ambros 2004). MiRNAs participate in various physiological processes and have a significant impact on hormone regulation. For example, miR-8 regulates multiple peptide hormones and may contribute

to *Drosophila* growth(Lee et al. 2015). In 2013, Setyowati Karolina D found that miR-25 and miR-92a overexpression suppressed insulin I expression in rats(Setyowati Karolina et al. 2013). Furthermore, miR-325-3p is involved in suppressing LH translation and secretion(Nemoto et al. 2012), and miR-136-3p binds directly to LHR mRNA to downregulate this molecule(Kitahara et al. 2013). MiR-133b promotes estradiol synthesis and levels by targeting Foxl2(Dai et al. 2013), and miR-132 stimulates estradiol synthesis by repressing Nurr1 translation(Wu et al. 2015b). However, miRNAs and their regulatory roles in the pituitary and whether they affect FSH expression are unclear.

In this study, we investigated and established a link between miR-7a-5p and FSH secretion based on previous research(Han et al. 2017). We verified the complementary sequence region between miR-7a-5p and the FSHb 3'UTR through dual-luciferase reporter assays. Moreover, we measured FSHb expression and secretion after transfection of miR-7a-5p mimics/inhibitors to determine whether miR-7a-5p affects FSH secretion.

## 2. MATERIALS AND METHODS

### 2.1. Animals and ethics

This study was performed with the approval of the Jilin University College of Animal Science. Healthy 8- to 9-week-old sexually mature male Sprague Dawley (SD) rats were used in this study. SD rats were raised in a comfortable room with a 12 h dark/12 h light cycle and free access to food and water.

We cared for the animals and sacrificed the rats in strict accordance with animal welfare laws and regulations and animal welfare ethics requirements. This experiment was approved by the Institutional Animal Care and Use Committee of Jilin University (201705026).

### 2.2. Pituitary extraction and cell culture

SD rats were euthanized by anesthesia and cervical dislocation. Then, we removed the heads with ophthalmic scissors and placed them on gauze disinfected with alcohol-soaked cotton. Next, we separated the rat skin with scissors to expose the rat skull. We opened the skull with tweezers

and removed the brain to collect the pituitary gland. Next, the tissues were placed in precooled phosphate-buffered saline (PBS) containing 0.3% bovine serum albumin (BSA) and 1% penicillin/streptomycin. We also separated the neurohypophysis from the pituitary. To ensure aseptic operations during the experiment, we autoclaved all experimental instruments and performed the operations near an alcohol-burning lamp.

After separation, we used PBS (0.3% BSA and 0.1% penicillin/streptomycin) to clean the pituitary glands and wash the blood from the tissue. Next, we placed the samples in 1 ml Dulbecco's Modified Eagle's Medium-F12 (DMEM-F12) containing 2.5% collagenase type I and cut the pituitary into pieces with ophthalmic scissors. We placed the pituitary fragments in a temperature-controlled incubator containing 5% CO<sub>2</sub> at 37°C for 90 min. After a 90 min incubation, we diluted pituitary cells with PBS (0.3% BSA and 0.1% penicillin/streptomycin) and then filtered the mixture through a 200 mesh (75 µm) cell sieve, which allowed pituitary cells to pass, while cell clusters and unhomogenized tissue could not pass through. The collected cell solution was centrifuged at 200×g for 10 min. After centrifugation, we carefully discarded the supernatant and diluted the cell precipitate with DMEM-F12 containing 15% fetal bovine serum (FBS). Finally, the diluted cell suspension was transferred to 6-well plates and cultured in a temperature-controlled incubator with 5% CO<sub>2</sub> at 37°C. Pituitary cells were monitored for the next few days.

### 2.3. Transfection of miR-7a-5p mimics, inhibitors or siRNA

After cells were cultured in 6-well plates for 4 days and in 24-well plates for 1 day, transfection of miR-7a-5p mimics or inhibitors was carried out as FSHb expression and FSH secretion peaked when pituitary cells were cultured for approximately 6 days. We used 30 µl buffer, 3 µl reagent and 2.5 µl mimics/inhibitors in this transfection. First, we added buffer, reagent and mimics, inhibitors or siRNA to a centrifuge tube. Then, we used a vortex oscillator to centrifuge the sample briefly to ensure homogeneous mixing. Next, we transfected the mixture into 24-well plates filled with pituitary cell suspension. Finally, we placed the 24-well plates into a 5% CO<sub>2</sub> incubator for 24 h to provide adequate time for the reaction. All mimics, inhibitors and

siRNAs were purchased from RiboBio (Gouzhou, China), and the transfection method and operation steps were performed in strict accordance with the manufacturer's recommended protocol. The mimic sequence was the double-stranded sequence of the miRNA mature sequence (sense strand: 5'-UGGAAGACUAGUGAUUUUGUUGU-3') and its complementary sequence (antisense strand: 3'-ACCUUCUGAUCACUAAAACAACA-5'). The inhibitor was the single-stranded sequence of the reverse complement of the miRNA mature sequence, which underwent full-chain methylated modification (5'-mAmCmAmAmCmAmAmAmAmUmCmAmCmUmAmGmUmCmUmUmCmCmA-3').

### 3.4. RNA isolation and qRT-PCR

After transfection, we harvested rat primary anterior pituitary cells that were transfected with miR-7a-5p mimics or inhibitors in 24-well plates. We used 350  $\mu$ l or 600  $\mu$ l cell lysate RL containing 1%  $\beta$ -mercaptoethanol to cause cell rupture and then extracted RNA according to the instructions of an RNAprep Pure Cell/Bacteria Kit. We subsequently measured the concentration and purity of the RNA with a NanoDrop ND-2000 spectrophotometer (Beijing, China) to verify the operational accuracy and RNA quality with a standard. Total RNA extraction was performed on a clean bench, and the samples were maintained at low temperatures to prevent RNA degradation. Next, we obtained cDNA using a FastQuant RT Kit and acquired raw data via q-PCR with SuperReal PreMix Plus (SYBR Green). These reagent kits were purchased from Tiangen (Beijing, China). The mRNA and miRNA primers used in RT-PCR and qRT-PCR are listed in Table 1.

### 2.5. Detection of cell apoptosis by flow cytometry

Flow cytometry was applied to detect rat adenohypophysis cell apoptosis to evaluate the effect of transfection on cells after 24 h. At an appropriate time point, we used trypsin to digest adhesive cells and transferred them into 10-ml reaction tubes. Then, the cells were centrifuged at 200 $\times$ g for 5 min and harvested by cell sedimentation. Next, we resuspended cells in 500  $\mu$ l 1x working fluid by diluting 5 $\times$  binding buffer with double-distilled water. Then, Annexin V-fluorescein isothiocyanate (FITC) and propidium iodide (PI) were added to sample tubes and parameter



regulation tubes according to an Annexin V-FITC/PI Apoptosis Kit (Multi Sciences, Hangzhou, China). Finally, we analyzed cell apoptosis via flow cytometry for 1 h.

## 2.6. Construction of the reporter plasmids

The pmiR-FSHb-3'UTR-wild-type (WT) plasmid and pmiR-FSHb-3'UTR-MUT plasmid were constructed to verify the target sites of the FSHb 3'UTR and miR-7a-5p. The PCR primers and mutant sequence of the target site are listed in File S1.

## 2.7. Detection of FSH secretion

After transfection with miR-7a-5p mimics or inhibitors for 24 h, we cultured pituitary cells with serum-free medium instead of DMEM-F12 (15% FBS) because FBS may contain other hormones that influence the results. After 24 h, we collected the culture medium and measured the secretion level of FSH in the culture medium via a Rat FSH enzyme-linked immunosorbent assay (ELISA) Kit (Meilian Biotech Co., Ltd., Shanghai, China).

## 2.8. Statistical analysis

At least three replicates were performed for each experiment. One-way ANOVA and Chi-square tests were performed to evaluate the statistical significance of differences.  $P < 0.05$  was considered statistically significant.

# 3. RESULTS

## 3.1 Prediction and verification of the complementary region between miR-7a-5p and the FSHb 3'UTR

First, information on the complementary sequence between miR-7a-5p and the FSHb 3'UTR was acquired via the TargetScan program (<http://www.targetscan.org/>) (Fig. 1A). Then, to further confirm that miR-7a-5p targets the FSHb 3'UTR, we successfully mutated the target complementary sequence TCTTCCA to AGAAGGT and constructed a FSHb-3'UTR-WT plasmid and FSHb-3'UTR mutated (MT) plasmid (Fig. 1B). Finally, the constructed plasmids were cotransfected with miR-7a-5p mimics into 293T cells. As expected, the luciferase activity was reduced by 36% when we cotransfected the pmiR-FSHb-3'UTR WT plasmid and miR-7a-5p

mimics into 293T cells. In contrast, cotransfection of the FSHb-3'UTR MUT plasmid and miR-7a-5p mimics resulted in a 7% reduction in luciferase activity (Fig. 1C). Therefore, we concluded that miR-7a-5p may regulate FSHb expression by directly targeting the FSHb gene.

### 3.2 Expression level of miR-7a-5p in different developmental stages and various rat tissues

To ascertain whether miR-7a-5p expression shows discrepancies, we measured miR-7a-5p levels in different developmental stages and rat tissues. We selected 2-week-old rats as the nonsexually mature group and 4-month-old rats as the sexually mature group. Then, we detected the expression levels of miR-7a-5p by RNA isolation and qRT-PCR and normalized them to the levels in immature animals (by setting this value to 1). MiR-7a-5p expression was downregulated in mature rats compared to that in nonsexually mature rats (Fig. 2A). Next, we collected seven tissues from mature rats and detected the expression level of miR-7a-5p by qRT-PCR. Relative miR-7a-5p expression was extremely high (1378-fold) in the pituitary gland compared to that in the heart. Furthermore, miR-7a-5p was poorly expressed in the heart, liver, spleen, lung, and kidney, although it was slightly expressed in the brain (Fig. 2B). These findings suggested that miR-7a-5p is much more highly expressed in the rat pituitary gland than in other tissues.

### 3.3 Efficiency of transfection and subsequent impact

To detect the efficiency of transfection, we transfected the NC mimic with fluorescent markers into pituitary cells. We detected red fluorescence labeling in the cells. The results showed that the NC mimic was successfully transfected into pituitary cells. The transfection efficiency was approximately 70–80%, as shown in Fig. 3A. Flow cytometry was performed to assess the damage from transfection and certain reagents. No significant differences in cell apoptosis were observed among the four groups, indicating that the impact of transfection was negligible (Fig. 3B). We examined the expression levels of miR-7a-5p after transfection of miR-7a-5p mimics or inhibitors for 24 h to verify whether the mimics and inhibitors were transfected into pituitary cells. The expression levels of miR-7a-5p in cells transfected with mimics were increased, while the expression levels of miR-7a-5p in cells transfected with inhibitors were lower than those in cells transfected with controls (Fig. 3C). In other words, transfection was successful and reliable.

### 3.4. Effects of miR-7a-5p overexpression/blockade on FSH secretion

We examined FSHb mRNA levels and FSH hormone secretion after the transfection of miR-7a-5p mimics, inhibitors and siRNA into rat primary anterior pituitary cells for 24 h to further verify that miR-7a-5p affects FSHb expression and regulates animal reproduction. As a positive control, rat primary pituitary cells were transfected with FSHb siRNA, and we examined the levels of FSHb by quantitative RT-PCR and FSH secretion by ELISA 24 h after transfection. As expected, FSHb levels and FSH secretion significantly decreased ( $P<0.001$ ,  $P=0.002$ ) (Fig. 4A-B). The expression levels of FSHb decreased by 0.60-fold ( $P=0.002$ ) after transfection with miR-7a-5p mimics compared to those after transfection with the negative control. In contrast, after transfection with a miR-7a-5p inhibitor, FSHb levels increased by 1.5-fold ( $P<0.001$ ) (Fig. 4C). We subsequently measured FSH secretion levels. As expected, FSH secretion showed the same trend as that of FSHb expression. After transfection with miR-7a-5p mimics, the FSH concentration was substantially decreased. Moreover, the FSH concentration increased after transfection with the miR-7a-5p inhibitor (Fig. 4D).

These results indicated that miR-7a-5p can decrease FSHb expression and reduce FSH hormone secretion. Our findings provide additional evidence showing that miRNAs regulate FSH, demonstrating their potential role in the pituitary.

## 4. DISCUSSION

MiRNAs have crucial roles in multiple fundamental biological processes, such as cell proliferation(Hu et al. 2018), apoptosis(Ren et al. 2018), metastasis(Sun et al. 2018), migration(Liu et al. 2015; Ying et al. 2016), differentiation(Chen et al. 2018), and cell adhesion(Wu et al. 2015a). In addition, some diseases and cancers are associated with aberrant expression of miRNAs and subsequently, their target genes(Tuna et al. 2016). For example, miR-9-3p, miR-330-3p-3p, and miR-345-5p were significantly overexpressed in sera from patients with prostate cancer compared to those in sera from individuals without cancer, while patients who were in remission after androgen deprivation therapy (ADT) appeared to have significantly decreased miR-345-5p

levels(Tinay et al. 2018). Consequently, some changes in miRNA expression can be used as diagnostic and potential clinical biomarkers in cancer and other diseases(Min et al. 2018; Quan et al. 2018; Tuna et al. 2016). In the normal pituitary and pituitary adenomas, miRNAs affect cell proliferation, organ maturity and hormone secretion. In 2018, Lu B et al. found that miRNA-16 expression influences the proliferation and angiogenesis of pituitary tumors(Lu et al. 2018), and low expression of miR-23b and miR-130b may facilitate pituitary carcinogenesis(Leone et al. 2014). Moreover, miRNAs are functional components and have potential roles in regulating hormone secretion in the pituitary, such as the secretion of growth hormone (GH)(Hao & Waxman 2018; Qi et al. 2015), LH(Menon et al. 2015), TSH(Vadstrup 2006), and FSH(Han et al. 2017; Sheng et al. 2018). In this study, miR-7a-5p was highly expressed in the rat pituitary gland. This finding indicates that miR-7a-5p plays a potentially vital role in the secretion of hormones and the regulation of sequential production in animals.

According to many studies, the miR-7 family plays different roles in different cancers and has pathological significance in cancer. MiR-7-5p acts as a tumor suppressor in pancreatic ductal adenocarcinoma and suppresses cell proliferation, migration and invasion by targeting SOX18(Zhu et al. 2018); in MCF-10A mammary epithelial cells, this miRNA suppresses oncogenes by mediating the signaling of hepatocyte growth factor(Jeong et al. 2017). However, miR-7 overexpression in NFs dramatically increased cancer cell coculture growth rates and migratory activity(Shen et al. 2017). Moreover, the miR-7 family is involved in diverse cellular functions. MiRNA-7a plays a role in Müller glial differentiation via blockade of Notch3 expression(Baba et al. 2015) and alleviates the maintenance of neuropathic pain by regulating neuronal excitability(Sakai et al. 2013). In addition, miR-7 promotes *Drosophila* wing growth by controlling the Notch signaling pathway(Aparicio et al. 2015). Importantly, miR-7 may slow Parkinson's disease (PD) progression and regulate proliferation and the mTOR pathway(Titze-de-Almeida & Titze-de-Almeida 2018; Wang et al. 2013). In 2017, Ahmed K et al. found that normal pituitary development depended on the participation of miR-7a2 and that genetic deletion of miR-7a2 caused infertility(Ahmed et al. 2017). Moreover, the lack of miR-200b and miR-429

exerted the same biological effects, anovulation and infertility(Hasuwa et al. 2013). Accordingly, as a single member of the miR-7 family, miR-7a-5p may regulate pituitary development and reproduction. Based on our results, miR-7a-5p overexpression attenuates FSHb expression and decreases FSH secretion, contributing to the mechanism underlying FSH regulation by miRNAs.

In the past decade, miRNAs have been found in various tissues and organs. However, most mature miRNAs exhibit tissue-specific expression patterns with a precise timing trend that crucially contribute to cell identity and function(Choudhury et al. 2013; Landgraf et al. 2007). In 2002, Lagos-Quintana M et al. examined nine different mice and identified 34 highly tissue-specific novel miRNAs(Lagos-Quintana et al. 2002). In 2014, in a study of the pig genome, Martini P et al. predicted species-specific and conserved miRNAs and identified many tissue-specific miRNAs in different tissues(Martini et al. 2014). Moreover, some muscle-specific miRNAs, such as miR-1, miR-133a, miR-133b and miR-206, were validated in 2014(Takeuchi et al. 2014). In our study, miR-7a-5p was highly expressed in the brain and pituitary, consistent with the results of an in situ hybridization study(Herzer et al. 2012). These data indicate that miR-7a-5p has a potential regulatory function in the pituitary gland and brain.

Gonadotropin FSH, one of the major hormones secreted by the anterior pituitary gland, has a critical role in regulating reproduction(Ulloa-Aguirre et al. 1995). Therefore, elucidating the mechanisms involved in FSH regulation is important. Although many studies have reported that miRNAs can inhibit the secretion of FSH, there are many other factors that influence the secretion of FSH to ensure the growth and development of animals, such as follistatin(Meriggiola et al. 1994), hormones(Dumesic et al. 2009) and single nucleotide polymorphisms(Dai et al. 2009). For example, triiodothyronine differentially modulates FSH synthesis and secretion in male rats, and the Bu-shen-zhu-yun decoction promotes FSH synthesis and secretion. Gonadotropin releasing hormone (GnRH) is a major regulator of FSH secretion, and differential GnRH pulse frequencies and amplitudes affect FSH secretion levels and patterns(Belchetz et al. 1978; Savoy-Moore & Swartz 1987). Nevertheless, little is known regarding the association of miRNAs with FSH secretion. In 2013, several miRNAs were shown to target the FSHb mRNA 3'UTR after pituitary

cells were treated with 100 nM GnRH(Ye et al. 2013). Additionally, the activation of FSH expression is dependent on miR-132/212(Lannes et al. 2015). In our previous study, miR-186, miR-433 and miR-21-3p were confirmed to regulate FSHb expression and FSH secretion by directly targeting the FSHb 3'UTR(Han et al. 2017; Sheng et al. 2018). Furthermore, in this study, miR-7a-5p had the same effect on FSHb. This study will help improve our understanding of the regulatory functions of miRNAs in the pituitary, enriching our knowledge regarding the mechanism underlying FSH regulation.

## CONCLUSION

Overall, our study demonstrated a role for miR-7a-5p in suppressing FSHb expression and decreasing FSH secretion. These findings provide additional evidence that miRNAs may regulate FSH secretion by directly targeting FSHb.

## SUPPLEMENTARY MATERIAL

S1 File. Construction of the pmiR-FSHb-3'UTR-MUT reporter plasmid.

## FIGURE LEGENDS

### Table 1. Primers used for RT-qPCR.

### Fig. 1. Prediction and verification of the complementary region between miR-7a-5p and the FSHb 3'UTR.

(A) The complementary base pairing region of miR-7a-5p and FSHb predicted through the TargetScan program is shown in red. (B) A sequencing map shows the mutation of the target sequence from TCTTCC to AGAAGG. (C) Relative luciferase activity was examined after cotransfection of the plasmid with the miR-7a-5p NC/mimic into 293T cells for 48 h. As a negative control, the luciferase activity of cells cotransfected with the FSHb-3'UTR wild-type plasmid and the NC group was set to 1. At least three replicates were performed for each experiment. Mean values and standard deviations (SDs) of the data are shown. One-way ANOVA and the Chi-square test were applied to analyze statistical significance.  $P < 0.05$  was considered significant, and different letters (a and b) indicate significant differences between groups.

### Fig. 2. Expression levels of miR-7a-5p in different developmental stages and in various rat tissues.

(A) MiR-7a-5p expression was measured in immature and mature rat pituitary. (B) MiR-7a-5p expression in pituitary and other tissues. *U6* was used as an internal standard in this study. At least three replicates were performed for each experiment. Mean values and standard deviations (SDs) of the data are shown. One-way ANOVA and the Chi-square test were performed to assess statistical significance. Different letters (a and b) indicate significant differences between groups ( $P < 0.05$ ).

### Fig. 3. Efficiency of transfection and subsequent impact.

(A) Fluorescence labeling was detected after transfection with the NC mimic carrying fluorescence markers. (B) The percentage of apoptotic pituitary cells after transfection with miR-7a-5p NC/mimic/I-NC/inhibitor in rat pituitary cells. (C) The relative expression of miR-7a-5p after transfection with miR-7a-5p mimic/inhibitor in rat pituitary cells. At least three replicates of each experiment were performed. Mean values and standard deviations (SDs) are shown. Data management and analysis were performed by using SPSS 19.0. Different letters (a and b) indicate significant differences ( $P < 0.05$ ).

### Fig. 4. Effects of miR-7a-5p overexpression/blockade on FSH secretion.

(A) FSHb relative expression after transfection with siRNA in rat pituitary cells. (B) The FSH concentration was measured via ELISA after transfection with siRNA for 24 h in rat pituitary cells. (C) FSHb relative expression after transfection with miR-7a-5p NC/mimic/I-NC/inhibitor in rat pituitary cells. (D) FSH concentrations of supernatants transfected with miR-7a-5p NC, mimic, I-



NC, and inhibitor in rat pituitary cells. At least three replicates were performed for each experiment. Each transfection experiment was performed in rat pituitary cells. Mean values and standard deviations (SDs) are shown. One-way ANOVA and the Chi-square test were applied to evaluate the statistical significance of the differences. Different letters indicate significant differences ( $P < 0.05$ ).

# References

- Ahmed K, LaPierre MP, Gasser E, Denzler R, Yang Y, Rulicke T, Kero J, Latreille M, and Stoffel M. 2017. Loss of microRNA-7a2 induces hypogonadotropic hypogonadism and infertility. *J Clin Invest* 127:1061-1074. 10.1172/JCI90031
- Ambros V. 2004. The functions of animal microRNAs. *Nature* 431:350-355. 10.1038/nature02871
- Aparicio R, Simoes Da Silva CJ, and Busturia A. 2015. MicroRNA miR-7 contributes to the control of Drosophila wing growth. *Dev Dyn* 244:21-30. 10.1002/dvdy.24214
- Baba Y, Aihara Y, and Watanabe S. 2015. MicroRNA-7a regulates Muller glia differentiation by attenuating Notch3 expression. *Exp Eye Res* 138:59-65. 10.1016/j.exer.2015.06.022
- Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* 136:215-233. 10.1016/j.cell.2009.01.002
- Belchetz PE, Plant TM, Nakai Y, Keogh EJ, and Knobil E. 1978. Hypophysial responses to continuous and intermittent delivery of hypophthalamic gonadotropin-releasing hormone. *Science* 202:631-633.
- Chen WJ, Chen YH, Hsu YJ, Lin KH, and Yeh YH. 2018. MicroRNA-132 targeting PTEN contributes to cilostazol-promoted vascular smooth muscle cell differentiation. *Atherosclerosis* 274:1-7. 10.1016/j.atherosclerosis.2018.04.030
- Choudhury NR, de Lima Alves F, de Andres-Aguayo L, Graf T, Caceres JF, Rappsilber J, and Michlewski G. 2013. Tissue-specific control of brain-enriched miR-7 biogenesis. *Genes Dev* 27:24-38. 10.1101/gad.199190.112
- Dai A, Sun H, Fang T, Zhang Q, Wu S, Jiang Y, Ding L, Yan G, and Hu Y. 2013. MicroRNA-133b stimulates ovarian estradiol synthesis by targeting Foxl2. *Febs Letters* 587:2474-2482. 10.1016/j.febslet.2013.06.023
- Dai L, Zhao Z, Zhao R, Xiao S, Jiang H, Yue X, Li X, Gao Y, Liu J, and Zhang J. 2009. Effects of novel single nucleotide polymorphisms of the FSH beta-subunit gene on semen quality and fertility in bulls. *Anim Reprod Sci* 114:14-22. 10.1016/j.anireprosci.2008.08.021
- Dumesic DA, Lesnick TG, Stassart JP, Ball GD, Wong A, and Abbott DH. 2009. Intrafollicular antimullerian hormone levels predict follicle responsiveness to follicle-stimulating hormone (FSH) in normoandrogenic ovulatory women undergoing gonadotropin releasing-hormone analog/recombinant human FSH therapy for in vitro fertilization and embryo transfer. *Fertil Steril* 92:217-221. 10.1016/j.fertnstert.2008.04.047
- Fan QR, and Hendrickson WA. 2005. Structure of human follicle-stimulating hormone in complex with its receptor. *Nature* 433:269-277. 10.1038/nature03206
- Han DX, Sun XL, Xu MQ, Chen CZ, Jiang H, Gao Y, Yuan B, and Zhang JB. 2017. Roles of differential expression



- of microRNA-21-3p and microRNA-433 in FSH regulation in rat anterior pituitary cells. *Oncotarget* 8:36553-36565. 10.18632/oncotarget.16615
- Hao P, and Waxman DJ. 2018. Functional Roles of Sex-Biased, Growth Hormone-Regulated MicroRNAs miR-1948 and miR-802 in Young Adult Mouse Liver. *Endocrinology* 159:1377-1392. 10.1210/en.2017-03109
- Hasuwa H, Ueda J, Ikawa M, and Okabe M. 2013. miR-200b and miR-429 function in mouse ovulation and are essential for female fertility. *Science* 341:71-73. 10.1126/science.1237999
- Herzer S, Silahdaroglu A, and Meister B. 2012. Locked nucleic acid-based in situ hybridisation reveals miR-7a as a hypothalamus-enriched microRNA with a distinct expression pattern. *Journal Of Neuroendocrinology* 24:1492-1504. 10.1111/j.1365-2826.2012.02358.x
- Hong GK, Payne SC, and Jane JA, Jr. 2016. Anatomy, Physiology, and Laboratory Evaluation of the Pituitary Gland. *Otolaryngol Clin North Am* 49:21-32. 10.1016/j.otc.2015.09.002
- Hu X, Miao J, Zhang M, Wang X, Wang Z, Han J, Tong D, and Huang C. 2018. miRNA-103a-3p Promotes Human Gastric Cancer Cell Proliferation by Targeting and Suppressing ATF7 in vitro. *Molecules and Cells* 41:390-400. 10.14348/molcells.2018.2078
- Hunzicker-Dunn M, and Maizels ET. 2006. FSH signaling pathways in immature granulosa cells that regulate target gene expression: branching out from protein kinase A. *Cell Signal* 18:1351-1359. 10.1016/j.cellsig.2006.02.011
- Jeong D, Ham J, Park S, Lee S, Lee H, Kang HS, and Kim SJ. 2017. MicroRNA-7-5p mediates the signaling of hepatocyte growth factor to suppress oncogenes in the MCF-10A mammary epithelial cell. *Sci Rep* 7:15425. 10.1038/s41598-017-15846-z
- Kitahara Y, Nakamura K, Kogure K, and Minegishi T. 2013. Role of microRNA-136-3p on the expression of luteinizing hormone-human chorionic gonadotropin receptor mRNA in rat ovaries. *Biology of Reproduction* 89:114. 10.1095/biolreprod.113.109207
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, and Tuschl T. 2002. Identification of tissue-specific microRNAs from mouse. *Current Biology* 12:735-739.
- Lamminen T, Jokinen P, Jiang M, Pakarinen P, Simonsen H, and Huhtaniemi I. 2005. Human FSH beta subunit gene is highly conserved. *Molecular Human Reproduction* 11:601-605. 10.1093/molehr/gah198
- Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M, Lin C, Socci ND, Hermida L, Fulci V, Chiaretti S, Foa R, Schliwka J, Fuchs U, Novosel A, Muller RU, Schermer B, Bissels U, Inman J, Phan Q, Chien M, Weir DB, Choksi R, De Vita G, Frezzetti D, Trompeter HI, Hornung V, Teng G, Hartmann G, Palkovits M, Di Lauro R, Wernet P, Macino G, Rogler CE, Nagle JW, Ju J, Papavasiliou FN, Benzing T, Lichter P, Tam W, Brownstein MJ, Bosio A, Borkhardt A, Russo JJ, Sander C, Zavolan M, and Tuschl T. 2007. A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell* 129:1401-1414. 10.1016/j.cell.2007.04.040
- Lannes J, L'Hote D, Garrel G, Laverriere JN, Cohen-Tannoudji J, and Querat B. 2015. Rapid communication: A microRNA-132/212 pathway mediates GnRH activation of FSH expression. *Molecular Endocrinology* 29:364-372. 10.1210/me.2014-1390
- Lee GJ, Jun JW, and Hyun S. 2015. MicroRNA miR-8 regulates multiple growth factor hormones produced from Drosophila fat cells. *Insect Molecular Biology* 24:311-318. 10.1111/imb.12156
- Leone V, Langella C, D'Angelo D, Mussnich P, Wierinckx A, Terracciano L, Raverot G, Lachuer J, Rotondi S, Jaffrain-Rea ML, Trouillas J, and Fusco A. 2014. Mir-23b and miR-130b expression is downregulated in

- pituitary adenomas. *Molecular and Cellular Endocrinology* 390:1-7. 10.1016/j.mce.2014.03.002
- Liu Y, Yang K, Shi H, Xu J, Zhang D, Wu Y, Zhou S, and Sun X. 2015. MiR-21 modulates human airway smooth muscle cell proliferation and migration in asthma through regulation of PTEN expression. *Exp Lung Res* 41:535-545. 10.3109/01902148.2015.1090501
- Lu B, Liu GL, Yu F, Li WJ, Xiang XX, and Xiao HZ. 2018. MicroRNA16/VEGFR2/p38/NFkappaB signaling pathway regulates cell growth of human pituitary neoplasms. *Oncology Reports* 39:1235-1244. 10.3892/or.2018.6227
- Martini P, Sales G, Brugiolo M, Gandaglia A, Naso F, De Pitta C, Spina M, Gerosa G, Chemello F, Romualdi C, Cagnin S, and Lanfranchi G. 2014. Tissue-specific expression and regulatory networks of pig microRNAome. *Plos One* 9:e89755. 10.1371/journal.pone.0089755
- Menon B, Gulappa T, and Menon KM. 2015. miR-122 Regulates LH Receptor Expression by Activating Sterol Response Element Binding Protein in Rat Ovaries. *Endocrinology* 156:3370-3380. 10.1210/en.2015-1121
- Meriggiola MC, Dahl KD, Mather JP, and Bremner WJ. 1994. Follistatin decreases activin-stimulated FSH secretion with no effect on GnRH-stimulated FSH secretion in prepubertal male monkeys. *Endocrinology* 134:1967-1970. 10.1210/endo.134.4.8137764
- Min N, Sakthi Vale PD, Wong AA, Tan NWH, Chong CY, Chen CJ, Wang RYL, and Chu JJH. 2018. Circulating Salivary miRNA hsa-miR-221 as Clinically Validated Diagnostic Marker for Hand, Foot, and Mouth Disease in Pediatric Patients. *Ebiomedicine*. 10.1016/j.ebiom.2018.05.006
- Musumeci G, Castorina S, Castrogiovanni P, Loreto C, Leonardi R, Aiello FC, Magro G, and Imbesi R. 2015. A journey through the pituitary gland: Development, structure and function, with emphasis on embryo-foetal and later development. *Acta Histochem* 117:355-366. 10.1016/j.acthis.2015.02.008
- Nemoto T, Mano A, and Shibasaki T. 2012. Increased expression of miR-325-3p by urocortin 2 and its involvement in stress-induced suppression of LH secretion in rat pituitary. *Am J Physiol Endocrinol Metab* 302:E781-787. 10.1152/ajpendo.00616.2011
- Qi QE, Xi QY, Ye RS, Chen T, Cheng X, Li CY, Zhu XT, Shu G, Wang LN, Jiang QY, and Zhang YL. 2015. Alteration of the miRNA expression profile in male porcine anterior pituitary cells in response to GHRH and CST and analysis of the potential roles for miRNAs in regulating GH. *Growth Horm IGF Res* 25:66-74. 10.1016/j.ghir.2014.12.002
- Quan X, Ji Y, Zhang C, Guo X, Zhang Y, Jia S, Ma W, Fan Y, and Wang C. 2018. Circulating MiR-146a May be a Potential Biomarker of Coronary Heart Disease in Patients with Subclinical Hypothyroidism. *Cell Physiol Biochem* 45:226-236. 10.1159/000486769
- Ren R, Chen SD, Fan J, Zhang G, and Li JB. 2018. miRNA-138 regulates MLK3/JNK/MAPK pathway to protect BV-2 cells from H2O2-induced apoptosis. *Bratisl Lek Listy* 119:284-288. 10.4149/BLL\_2018\_053
- Rodriguez-Pacheco F, Novelle MG, Vazquez MJ, Garcia-Escobar E, Soriguer F, Rojo-Martinez G, Garcia-Fuentes E, Malagon MM, and Dieguez C. 2013. Resistin regulates pituitary lipid metabolism and inflammation in vivo and in vitro. *Mediators Inflamm* 2013:479739. 10.1155/2013/479739
- Rull K, Grigorova M, Ehrenberg A, Vaas P, Sekavin A, Nommemees D, Adler M, Hanson E, Juhanson P, and Laan M. 2018. FSHB -211 G>T is a major genetic modulator of reproductive physiology and health in childbearing age women. *Human Reproduction*. 10.1093/humrep/dey057
- Sakai A, Saitow F, Miyake N, Miyake K, Shimada T, and Suzuki H. 2013. miR-7a alleviates the maintenance of neuropathic pain through regulation of neuronal excitability. *Brain* 136:2738-2750. 10.1093/brain/awt191


- 447 Savoy-Moore RT, and Swartz KH. 1987. Several GnRH stimulation frequencies differentially release FSH and LH  
448 from isolated, perfused rat anterior pituitary cells. *Adv Exp Med Biol* 219:641-645.
- 449 Setyowati Karolina D, Sepramaniam S, Tan HZ, Armugam A, and Jeyaseelan K. 2013. miR-25 and miR-92a regulate  
450 insulin I biosynthesis in rats. *RNA Biol* 10:1365-1378. 10.4161/rna.25557
- 451 Shen Z, Qin X, Yan M, Li R, Chen G, Zhang J, and Chen W. 2017. Cancer-associated fibroblasts promote cancer cell  
452 growth through a miR-7-RASSF2-PAR-4 axis in the tumor microenvironment. *Oncotarget* 8:1290-1303.  
453 10.18632/oncotarget.13609
- 454 Sheng X, Bao Y, Zhang JS, Li M, Li YN, Xu QN, Zhang SH, and Li CT. 2018. Research Progress on InDel Genetic  
455 Marker in Forensic Science. *Fa Yi Xue Za Zhi* 34:420-427. 10.12116/j.issn.1004-5619.2018.04.016
- 456 Shin MS, Chang H, Namkoong C, Kang GM, Kim HK, Gil SY, Yu JH, Park KH, and Kim MS. 2013. Hypothalamic  
457 and pituitary clusterin modulates neurohormonal responses to stress. *Neuroendocrinology* 98:233-241.  
458 10.1159/000355625
- 459 Sun X, Dai G, Yu L, Hu Q, Chen J, and Guo W. 2018. miR-143-3p inhibits the proliferation, migration and invasion  
460 in osteosarcoma by targeting FOSL2. *Sci Rep* 8:606. 10.1038/s41598-017-18739-3
- 461 Takeuchi J, Sakamoto A, and Takizawa T. 2014. Sevoflurane anesthesia persistently downregulates muscle-specific  
462 microRNAs in rat plasma. *International Journal of Molecular Medicine* 34:291-298.  
463 10.3892/ijmm.2014.1739
- 464 Telikicherla D, Ambekar A, Palapetta SM, Dwivedi SB, Raju R, Sharma J, Prasad Ts K, Ramachandra Y, Mohan SS,  
465 Maharudraiah J, Mukherjee S, and Pandey A. 2011. A comprehensive curated resource for follicle  
466 stimulating hormone signaling. *BMC Res Notes* 4:408. 10.1186/1756-0500-4-408
- 467 Tinay I, Tan M, Gui B, Werner L, Kibel AS, and Jia L. 2018. Functional roles and potential clinical application of  
468 miRNA-345-5p in prostate cancer. *Prostate*. 10.1002/pros.23650
- 469 Titze-de-Almeida R, and Titze-de-Almeida SS. 2018. miR-7 replacement therapy in Parkinson's disease. *Curr Gene*  
470 *Ther*. 10.2174/1566523218666180430121323
- 471 Treier M, and Rosenfeld MG. 1996. The hypothalamic-pituitary axis: co-development of two organs. *Current Opinion*  
472 *In Cell Biology* 8:833-843.
- 473 Tuna M, Machado AS, and Calin GA. 2016. Genetic and epigenetic alterations of microRNAs and implications for  
474 human cancers and other diseases. *Genes Chromosomes Cancer* 55:193-214. 10.1002/gcc.22332
- 475 Ulloa-Aguirre A, Midgley AR, Jr., Beitins IZ, and Padmanabhan V. 1995. Follicle-stimulating isohormones:  
476 characterization and physiological relevance. *Endocrine Reviews* 16:765-787. 10.1210/edrv-16-6-765
- 477 Vadstrup S. 2006. The adaptation of TSH secretion to autonomy in non-toxic goiter may be based-on active regulation  
478 of set-point and sensitivity of central TSH-receptors, perhaps by the microRNA (MIR) gene. *Med Hypotheses*  
479 67:588-591. 10.1016/j.mehy.2005.12.049
- 480 Vyshnevs'ka OA, and Bol'shova OV. 2013. [Influence of replacement growth hormone therapy (hGH) on pituitary-  
481 thyroid and pituitary-adrenal systems in prepubertal children with GH deficiency]. *Lik Sprava*:81-87.
- 482 Walker WH, and Cheng J. 2005. FSH and testosterone signaling in Sertoli cells. *Reproduction* 130:15-28.  
483 10.1530/rep.1.00358
- 484 Wang Y, Liu J, Liu C, Naji A, and Stoffers DA. 2013. MicroRNA-7 regulates the mTOR pathway and proliferation  
485 in adult pancreatic beta-cells. *Diabetes* 62:887-895. 10.2337/db12-0451
- 486 Wu D, Cerutti C, Lopez-Ramirez MA, Pryce G, King-Robson J, Simpson JE, van der Pol SM, Hirst MC, de Vries  
487 HE, Sharrack B, Baker D, Male DK, Michael GJ, and Romero IA. 2015a. Brain endothelial miR-146a

- negatively modulates T-cell adhesion through repressing multiple targets to inhibit NF-kappaB activation. *J Cereb Blood Flow Metab* 35:412-423. 10.1038/jcbfm.2014.207
- Wu S, Sun H, Zhang Q, Jiang Y, Fang T, Cui I, Yan G, and Hu Y. 2015b. MicroRNA-132 promotes estradiol synthesis in ovarian granulosa cells via translational repression of Nurr1. *Reprod Biol Endocrinol* 13:94. 10.1186/s12958-015-0095-z
- Ye RS, Xi QY, Qi Q, Cheng X, Chen T, Li H, Kallon S, Shu G, Wang SB, Jiang QY, and Zhang YL. 2013. Differentially expressed miRNAs after GnRH treatment and their potential roles in FSH regulation in porcine anterior pituitary cell. *Plos One* 8:e57156. 10.1371/journal.pone.0057156
- Ying J, Liu Y, Yang R, Zhang Y, and Xu J. 2016. miR-100 Inhibits the Growth and Migration of Burn-Denatured Fibroblasts. *Med Sci Monit* 22:697-701.
- Yuan B, Han DX, Dai LS, Gao Y, Ding Y, Yu XF, Chen J, Jiang H, Chen CZ, and Zhang JB. 2015. A comprehensive expression profile of micrornas in rat's pituitary. *International Journal of Clinical and Experimental Medicine* 8:13289-13295.
- Zhu W, Wang Y, Zhang D, Yu X, and Leng X. 2018. MiR-7-5p functions as a tumor suppressor by targeting SOX18 in pancreatic ductal adenocarcinoma. *Biochem Biophys Res Commun* 497:963-970. 10.1016/j.bbrc.2018.02.005

# Figure 1

Prediction and verification of the complementary region between miR-7a-5p and the FSHb 3'UTR.



(A) The complementary base pairing region of miR-7a-5p and FSHb predicted through TargetScan program is shown in red. (B) A sequencing map shows the mutation of the target sequence from TCTTCC to AGAAGG. (C) The relative luciferase activity was examined after co-transfection of plasmid with the miR-7a-5p NC/mimic into 293T cell for 48 h. As a negative control, the luciferase activity of cells co-transfected with FSHb-3'UTR wild-type plasmid and the NC group was set to 1. At least three replicates of each experiment were performed. Mean values and standard deviations (SDs) of data are shown. One-way ANOVA and Chi-square test were applied to analyze statistical significance.  $P < 0.05$  was considered significant, and different letters (a and b) indicate significant differences between groups.



## Figure 2 (on next page)

The expression level of miR-7a-5p in different developmental stages and in various rat tissues

(A) MiR-7a-5p expression was measured in immature and mature rats. (B) MiR-7a-5p expression in pituitary and other tissues. *U6* was used as an internal standard in this study. At least three replicates of each experiment were performed. Mean values and standard deviations (SDs) of the data are shown. One-way ANOVA and the Chi-square test were performed to assess statistical significance. Different letters (a and b) indicate significant differences between groups ( $P < 0.05$ ).






# Figure 3

The efficiency of and impact after transfection.

(A) Fluorescence labeling was detected after transfection with mimic NC carrying fluorescence markers. (B) The percentage of apoptotic pituitary cells after transfection with miR-7a-5p NC/mimic/I-NC/ inhibitor. (C) The relative expression of miR-7a-5p after transfection with miR-7a-5p mimic/inhibitor. At least three replicates of each experiment were performed. Mean values and standard deviations (SDs) are shown. Data management and analysis was performed by SPSS 19.0. Different letters (a and b) indicate significant differences ( $P < 0.05$ ).






## Figure 4 (on next page)


Effect of the overexpression/blockade of miR-7a-5p on FSH secretion

(A) FSHb relative expression after transfection with siRNA. (B) The FSH concentration was measured via ELISA after transfection with siRNA for 24 h. (C) The FSHb relative expression after transfection with miR-7a-5p NC/mimic/I-NC/inhibitor. (D) The FSH concentration of supernatant transfected with miR-7a-5p NC, mimic, I-NC, and inhibitor. At least three replicates of each experiment were performed. Mean values and standard deviations (SDs) are shown. One-way ANOVA and Chi-square test were applied to evaluate the statistical significance of the differences. The different letters indicate significant differences ( $P < 0.05$ ).


A




B



C



D



# **Table 1**(on next page)

Primers used for RT-qPCR

# 1 Table 1. Primers used for RT-qPCR

**Table 1. Primers used for RT-qPCR**

primers name	sequence(5'-3')
U6 RT	CGCTTCACGAATTTGCGTGTCAT
miR-7a-5p RT	CTCAACTGGTGTCTCGTGGAGTCGGCAATTCAGTTGAGAGCCCAA A
U6 F	GCTTCGGCAGCACATATACTAAAAT
U6 R	CGCTTCACGAATTTGCGTGTCAT
miR-7a-5p F	ACACTCCAGCTGGGTGGAAGACTAGTGATTT
universal reverse	CTCAAGTGTCGTGGAGTCGGCAA
GAPDH F	GGAAACCCATCACCATCTTC
GAPDH R	GTGGTTCACACCCATCACAA
FSHb F	ATACCACTTGGTGTGAGGGC
FSHb R	TAGAGGGAGTCTGAGTGGCG

2

3