

**A peer-reviewed version of this preprint was published in PeerJ on 5 December 2019.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.8117) (peerj.com/articles/8117), which is the preferred citable publication unless you specifically need to cite this preprint.

Olszyński RM, Szczepocka E, Żelazna-Wieczorek J. 2019. Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin. PeerJ 7:e8117  
<https://doi.org/10.7717/peerj.8117>

# Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin

Rafał M. Olszyński<sup>Corresp., Equal first author, 1</sup>, Ewelina Szczepocka<sup>1</sup>, Joanna Żelazna-Wieczorek<sup>Corresp. Equal first author, 1</sup>

<sup>1</sup> Department of Algology and Mycology/Faculty of Biology and Environmental Protection, University of Lodz, Łódź, Polska

Corresponding Authors: Rafał M. Olszyński, Joanna Żelazna-Wieczorek  
Email address: rafal.olszynski@biol.uni.lodz.pl, joanna.zelazna@biol.uni.lodz.pl

**Background.** The ecological state of surface waters is typically assessed by a multi-aspect approach based on a determination of its chemical and physical parameters, by hydromorphology and the use of indicator organisms such as benthic diatoms. By assigning ecological indicator values, it is possible to create diatom indices which serve as the basic tool in assessing the ecological status of surface waters. These ecological indicator values are set according to classification systems, such as the Van Dam et al. 1994 system, which classifies species of diatoms according to seven different ecological factors. However, recent studies on the autecology of diatoms have shown the need to verify and establish new ecological indicator values. To this end, aquatic ecosystems are good environments to observe the range of tolerance of benthic diatoms to environmental conditions due to their unique physical and chemical parameters. The aim of the present study was to propose the establishment of new, or altered, ecological indicator values, according to Van Dam et al. classification, of species of diatoms characteristic of three post-mining aquatic ecosystems. **Methods.** In total, 36 species were identified that were characteristic of three waterbodies: a salt aquatic complex (water outflow, a drainage ditch and a pond), a mined iron ore reservoir and a mined lignite reservoir. Their ecological indicator values were calculated using OMNIDIA software, and the environmental conditions prevailing in the studied ecosystems were determined. Of the 36 characteristic species, 16 lacking at least one assigned ecological indicator value were analysed further. The analysis identified three groups of selected characteristic species which showed a correlation, or lack of such, to the tested physical and chemical parameters. **Results.** Based on this multistage study of the autecology of characteristic diatoms, comprising an analysis of environmental conditions, literature analysis and reference ecological indicator values of other species, it is proposed that 32 ecological indicator values be established or adjusted for 16 species, and that *Planothidium frequentissimum* be excluded from water quality assessments.

# **Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin**

Rafał M. Olszyński, Ewelina Szczepocka, Joanna Żelazna-Wieczorek

Department of Algology and Mycology, Faculty of Biology and Environmental Protection,  
University of Lodz, Łódź

Corresponding Author:

Rafał M. Olszyński

12/16 Banacha St, Łódź, 90-237, Poland

Email address: [rafal.olszynski@biol.uni.lodz.pl](mailto:rafal.olszynski@biol.uni.lodz.pl)

# Abstract

**Background.** The ecological state of surface waters is typically assessed by a multi-aspect approach based on a determination of its chemical and physical parameters, by hydromorphology and the use of indicator organisms such as benthic diatoms. By assigning ecological indicator values, it is possible to create diatom indices which serve as the basic tool in assessing the ecological status of surface waters. These ecological indicator values are set according to classification systems, such as the Van Dam et al. 1994 system, which classifies species of diatoms according to seven different ecological factors. However, recent studies on the autecology of diatoms have shown the need to verify and establish new ecological indicator values. To this end, aquatic ecosystems are good environments to observe the range of tolerance of benthic diatoms to environmental conditions due to their unique physical and chemical parameters. The aim of the present study was to propose the establishment of new, or altered, ecological indicator values, according to Van Dam et al. classification, of species of diatoms characteristic of three post-mining aquatic ecosystems.

**Methods.** In total, 36 species were identified that were characteristic of three waterbodies: a salt aquatic complex (water outflow, a drainage ditch and a pond), a mined iron ore reservoir and a mined lignite reservoir. Their ecological indicator values were calculated using OMNIDIA software, and the environmental conditions prevailing in the studied ecosystems were determined. Of the 36 characteristic species, 16 lacking at least one assigned ecological indicator value were analysed further. The analysis identified three groups of selected characteristic species which showed a correlation, or lack of such, to the tested physical and chemical parameters.

**Results.** Based on this multistage study of the autecology of characteristic diatoms, comprising an analysis of environmental conditions, literature analysis and reference ecological indicator values of other species, it is proposed that 32 ecological indicator values be established or adjusted for 16 species, and that *Planothidium frequentissimum* be excluded from water quality assessments.

# Introduction

Diatoms (Bacillariophyta) are one of the main biotic elements used in the biological assessment of the ecological state of surface waters (*Water Framework Directive, European Union 2000*). Due to the fact that many countries are obliged to continually engage in biomonitoring, there is a clear need to develop flawlessly functioning methods based on the standardised use of diatoms as bioindicators (*Kahlert et al. 2016; Poikane, Kelly & Cantonati 2016; Szczepocka & Żelazna-Wieczorek 2018*). Diatom indices and ecological systems based on the bioindication values of particular diatom species, derived from various environmental parameters, constitute a fundamental tool in the biological assessment of environments. Diatom indices have been commonly used to assess flowing and standing water for over 20 years (*Kelly et al. 2008; Harding & Taylor 2014; Szczepocka et al. 2014; Hutorowicz & Pasztaleniec 2014; Holms & Taylor 2015; Żelazna-Wieczorek & Nowicka-Krawczyk 2015; Kolada et al. 2016*).

Currently, many countries use the OMNIDIA program (*Lecointe et al. 1993*) as a biological assessment tool. Its latest version (version 6.0.6) allows the calculation of 18 diatom indices, and the determination of seven environmental parameters for eight ecological systems. However, the specific ecological indicator values of many of the species given in the OMNIDIA database are absent or have not been updated in response to recent research. To complete these missing values, and to verify existing ones, further studies are needed of the ecological optima and tolerance of diatom species in different types of aquatic ecosystems.

Due to their specific environmental conditions, post-mining reservoirs represent an extremely valuable source of information for the study of ecological diatom tolerance ranges. Some studies of these environments have been performed, but these have addressed diatom paleoecology and their role as indicators of past climatic or environmental change (*de Haan et al. 1993; Rakowska 1996; Thomas & John 2006; Sienkiewicz & Gqsiorowski 2016*). Until now, the autecology of diatoms in post-mining reservoirs has rarely been studied (*Van Landingham 1968; de Haan et al. 1993; Rakowska 1996; Ferreira da Silva et al. 2009; Luis et al. 2009; Luis et al. 2016; Sienkiewicz & Gqsiorowski 2016*).

The present study examines the diatom assemblages present in three post-mining reservoirs of various geological origins. Due to variations in their environmental parameters, these bodies of water serve as specific and unique habitats for the development of these algae. The diatom assemblages quickly adapt to the currently prevailing conditions, which is manifested in the presence of taxa characteristic of these specific parameters. Considering their large share of the assemblage, the index values of the assemblages constitute the most important component in the calculation of diatom indices. These species are therefore of the greatest importance for surface water biomonitoring.

The aim of the present study was to identify the species of diatoms characteristic of the three studied types of post-mine reservoirs. Following this, taxa that did not have at least one ecological indicator value specified in the OMNIDIA database, according to the environmental parameters given by Van Dam et al. (1994), were identified. New ecological indicator values were proposed based on the relationship between the occurrence of the individual species and certain selected physical and chemical parameters, or existing ones were verified.

The Van Dam et al. (1994) ecological system is one of the main systems on which the OMNIDIA programme is based. It describes the ecological indicator values of diatoms according to pH, salinity, nitrogen uptake metabolism, oxygen requirement, saprobity, trophic state and moisture aerophily. These values play a key role in calculating diatom indices, and hence need to be kept up to date to enable accurate routine biomonitoring.

## Materials & Methods

### *Study area*

The study was performed on three waterbodies created through exploration for mineral deposits or were formed by the closure of mines. All three are located in the Łódzkie and Wielkopolskie voivodeships, Central Poland.

The first complex of waterbodies - Pełczyska (PE), is situated in the village of Pełczyska, between Łódź and Łęczyca (Łódzkie voivodeship) (Fig. 1). As the local area is characterized by the presence of salt deposits, numerous wells were sunk in the eighteenth century to obtain brine. Currently, salt water flows out of one of them. This area has been studied by biologists and hydrobiologists since the 1960s (*Olaczek 1963; Pliński 1966; Pliński 1969; Pliński 1971a; Pliński 1971b; Pliński 1971c; Pliński 1973; Żelazna-Wieczorek 1996; Żelazna-Wieczorek 2002; Żelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015; Żelazna-Wieczorek & Olszyński 2016*). The waterbodies chosen for our research form the PE hydrological complex located in the vicinity of farmland; it comprises the salt water outflow, a drainage ditch and a pond, which acts as the receiver of the water.

The second complex of waterbodies - Łęczyca (LE), urban reservoir located within the city of Łęczyca (Łódzkie voivodeship) (Fig. 1). The reservoirs were created following the flooding the open-cast iron ore mine in the 1990s. This area is rich in syderite deposits, which are accompanied by other minerals. The complex consists of three connected reservoirs: two are directly connected to each other (LEP1 and LEP2), and the third (LEP3) is connected to LEP2 via a water drainage ditch (*Olszyński & Żelazna-Wieczorek 2018*). All three are located in an area with houses, garden plots and partly-wooded areas.

The third waterbody - Bogdałów reservoir (BO), created by the flooding of an opencast brown coal mine. It is located in the village of Bogdałów (Wielkopolskie voivodeship) in an area rich in lignite deposits (Fig. 1). Lignite from quaternary deposits was exploited since 1977 until 1991 to a depth of 50 meters. Due to the specific construction of the open-pit area, being characterised by the thickest layer of poorly permeable boulder clay in the region. This pit was later transformed into a storage site for quarried rocks in Koźmin. Finally, in 1993/1994, the drainage and runoff of surface waters were blocked to form a reservoir with a depth of about 12 meters surrounded by forest (*Gabryś-Godlewska et al. 2004; Gadomska et al. 2007; Orlikowski & Szwed 2009; Kasztelewicz 2011*).

## Samples

Samples of benthic diatoms from sediments and water samples were collected quarterly (once in any season) from each hydrological waterbodies. Analysis of all water samples ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^{+}$ ,  $\text{K}^{+}$ ,  $\text{Fe}^{2+/3+}$ ,  $\text{Mn}^{3+}$ ) involved flame absorption spectrophotometry SpectrAA 300 (Varian, Palo Alto, CA, USA) (detection limit is 0.05 mg/L) and UV-vis spectrophotometry S.330 (Marcel, Poland) in the case of  $\text{NH}_4^{+}$  (d.l. is 0.001 mg/L) and  $\text{PO}_4^{3-}$  (d.l. is 0.01 mg/L).  $\text{SO}_4^{2-}$  was determined by the gravimetric method (PN-C-04566-09),  $\text{Cl}^{-}$  by Mohr's method (PN-ISO 9297). The chemical analyses were performed in the Laboratory of the Department of Geology at the Faculty of Geographical Sciences, University of Łódź and the Laboratory of Computer and Analytical Techniques at the Faculty of Biology and Environmental Protection, University of



Łódź. The water temperature, pH and electric conductivity were measured *in situ* (Elmetron CP-401 and CC-401 devices). The following sampling points were established: Pełczyska (51°58'34.47"N, 19°14'21.11"E) – outflow (P.ESB.) (4 samples, both water and benthic), ditch (D.PEDB.) (1 water and 4 benthic samples) and pond (D.PEPB.) (4 samples, both water and benthic); samples were collected quarterly from July 2013 to March 2014; Łęczyca (52°3'5.30"N; 19°11'50.24"E) – reservoir 1 (D.LEP1.), reservoir 2 (D.LEP2.) and reservoir 3 (D.LEP3.), samples were collected quarterly from March 2014 to December 2015 (6 water and 8 benthic samples from each reservoirs); Bogdałów (52°2'51.29"N; 18°35'51.49"E) – reservoir (D.BOZB.), samples were collected quarterly from March 2015 to December 2016 (8 samples, both water and benthic). The water samples were subjected to chemical and physical analysis at the Laboratory of Geology at the Faculty of Geographical Sciences, University of Lodz. In total, 44 benthic samples were collected. The permanent slides were prepared according to Żelazna-Wieczorek (2011). To obtain pure diatom frustule the material was chemically treated using a H<sub>2</sub>SO<sub>4</sub> and H<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>. The cleaned diatom precipitate was mounted on permanent slides using Naphrax<sup>®</sup> synthetic resin. Qualitative and quantitative analysis of diatoms was performed using a Nikon Eclipse 50i light microscope (LM) under 1000× magnification (plan oil-immersion objective 100×/1.25): the diatoms were identified and counted for up to 500 valves in each permanent slide. Light photomicrographs were taken with an OPTA-TECH digital camera. When diatoms were difficult to identify using LM they were subjected to scanning electron microscope (SEM) analysis using a Phenom Pro X (gold layer of 8 and 20 nm, at 10 kV, low vacuum mode) at the Laboratory of Microscopy Imaging and Specialist Biological Techniques, Faculty of Biology and Environmental Protection, University of Lodz.

#### *Data processing & statistical analysis*

The average percentage (AP) for a given species was determined based on the percentage contribution (%) of the species in the samples tested for a given hydrological object (Żelazna-Wieczorek 2011). Species with AP greater than or equal to 5% for each hydrological object were identified as dominant. The incidence was determined according to the Tümping & Friedrich (1999) coefficient according to the range values: 100%–75% euconstant taxa (EC), 75%–50% constant taxa (CN), 50%–25% accessory taxa (AC) and 25%–1% accidental taxa (AD) (Tümping & Friedrich 1999). Multidimensional scaling analysis (nMDS) based on Bray-Curtis similarity coefficients was used to identify natural groupings of samples. The results are given as a 3D diagram in which the degree of similarity is represented as the distances between particular points (samples), with greater distances indicating a lower degree of similarity. The reliability of the ordering of the assemblage is represented by the stress value, which reflects how well the ordination summarizes

the observed distances among the samples. A 3-dimensional presentation, whose stress value is lower, is likely to be more satisfactory than a 2-dimensional one (Clarke & Gorley 2015).

The Shade Plot analysis, based on the Bray-Curtis similarity coefficient, was used to identify the diatom species that have the strongest influence on the similarities between the samples demonstrated in the nMDS analysis. Shade Plot analysis compares two data matrices with each other and then groups them on two levels, according to the similarity of the samples and the factors affecting their similarity, i.e. diatom species. The results are represented graphically by shading individual cells: the intensity of the shading indicates the degree of the influence of a given factor (species) on the position of its sample within a given similarity cluster. The range of the shading was determined on the basis of  $\log(x+1)$  ( $x$ —number of valves).

The SIMPER analysis was used to determine the characteristic species distinguishing the studied ecosystems. This method examines the participation of each variable in the overall similarity between groups of samples, thus indicating the species with the greatest influence on the degree of similarity, or dissimilarity, between particular samples and hydrological objects. This analysis is also based on the Bray-Curtis similarity coefficient; however, unlike the nMDS method, in which one trial is compared to all the other samples, the SIMPER analysis compares a single sample to each subsequent sample (Żelazna-Wieczorek 2011). The results indicate the species which most strongly differentiated a sampled site from the others, and to what extent. A species was regarded as being characteristic of the studied ecosystem if it was characterized by a mean dissimilarity greater than or equal to 2 according to the SIMPER analysis, and a higher mean abundance greater in one ecosystem than the other.

In total, 19 physical and chemical parameters of water were measured in the studied ecosystems. The results of the correlation analysis found 15 physical and chemical parameters indicating an environmental conditions. The parameters were subjected to principal component analysis (PCA) to determine which had the strongest effect on the selected species.

Using the information from the OMNIDIA database, the environmental conditions for each sampling point were determined according to Van Dam et al. (1994) (Table 1). Following this, the percentage share of diatom species included in each ecological indicator value class was indicated. For species found to be characteristic of the studied ecosystems, classes of ecological indicator values were assembled. Taxa which had at least one value of 0 (unknown) were selected for further analysis.

The selected ecological indicator values according to Van Dam et al. (1994) were verified, or new ones established, for the species found to be characteristic of the studied ecosystems according to three premises: previous literature reports about ecological indicator values of those species, chemical and physical conditions analysis, and the classification of the environmental conditions according to Van Dam et al. (1994).

The analyses were performed using PRIMER 7.0.13 (nMDS, Shade Plot, SIMPER), OMNIDIA 6.0.6 and STATISTICA 13 (PCA), software.

## Results & Discussion



## 209 *Chemical analysis of water samples*

210 The mean values and range of all tested parameters are given in Table 2.

211 The PE hydrological complex was characterized by elevated values of electric conductivity,  
212 reaching as high as 9230  $\mu\text{S cm}^{-1}$ . The pH changed with the direction of water outflow: a slightly  
213 acidic reaction was observed in the outflow and an alkaline one in the pond. Due to the  
214 geological profile of the region, the water flowing out of the well contained a high concentration  
215 of chloride ions, whose gradient decreased with the flow of water through the ditch to the pond.  
216 In addition, higher concentrations of the cations  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Na}^{+}$  and  $\text{K}^{+}$  were observed  
217 compared to other ecosystems, as well as the anions  $\text{HCO}_3^{-}$ ,  $\text{PO}_4^{3-}$  and  $\text{SO}_4^{2-}$ .

218 The  $\text{K}^{+}$  concentration is acknowledged parameter coming from agricultural activity, in particular  
219 animal husbandry, or municipal wastes (Macioszczyk & Dobrzyński 2002).

220 Each of the sampling points in the PE complex was characterized by different chemical  
221 parameters, resulting in differences between the habitats. The highest electrolytic conductivity  
222 was noted in the outflow, which was mainly influenced by the concentrations of  $\text{Cl}^{-}$ ,  $\text{Na}^{+}$  and  
223  $\text{HCO}_3^{-}$  ions. The maximum concentration of  $\text{HCO}_3^{-}$  ions was recorded in Pełczyska outflow in  
224 March 2014 (D.PESB.250314); in the other locations, it did not exceed 410  $\text{mg L}^{-1}$ .

225 Low concentrations of  $\text{K}^{+}$  ions were observed throughout the entire studied PE complex;  
226 however, maximum values were recorded in the locations characterised by the highest  $\text{HCO}_3^{-}$  ion  
227 content. The highest concentration of  $\text{Ca}^{+}$  ions of all ecosystems was recorded in the outflow.

228 The ditch represented an intermediate section between the PE sampling points. However, as it is  
229 susceptible to periodic drying, limited chemical data was collected from this habitat and hence it  
230 was not possible to assess its chemical and physical nature.

231 The lowest electrolytic conductivity was found in the pond, which displayed lower  
232 concentrations of  $\text{Cl}^{-}$ ,  $\text{Na}^{+}$  and, to a lesser degree,  $\text{HCO}_3^{-}$ . The pH of the water never dropped  
233 below 8, except in one case in March 2014. In the pond, the concentration of  $\text{K}^{+}$  remained  
234 relatively unchanged, which could be related to the fact that the reservoir was also a receiver of  
235 waters flowing from the surrounding arable fields. The pond was also characterized by the  
236 lowest concentration of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . In the summer periods, a significant reduction in the  
237 water table level and occasional drying of the reservoir were noted.

238 The urban reservoirs in Łęczyca (LE) were characterized by a slightly alkaline water reaction,  
239 which was similar in all reservoirs during the course of the study. No elevated concentrations of  
240  $\text{Fe}^{2+/3+}$  and  $\text{Mn}^{3+}$  ions were observed. The content of  $\text{SO}_4^{2-}$  anions was not higher than in other  
241 waterbodies studied. The concentration of  $\text{HCO}_3^{-}$  ions was lower than that observed in BO and  
242 PE. No significant differences in chemical and physical parameters were observed between the  
243 individual sampling points constituting LE.

244 The Bogdałów (BO) reservoir was characterized by an alkaline reaction. Its  $\text{K}^{+}$ ,  $\text{Cl}^{-}$  and  $\text{NH}_4^{+}$   
245 ion content was the lowest of the studied ecosystems.

## 246 *Diatom samples*

A total of 381 diatom taxa were identified in 44 benthic samples: 139 in PE, 192 in LE and 188 in BO. The dominant species in PE were *Navicula veneta*, and *Nitzschia frustulum*, in LE *Cyclostephanos dubius* and *Stephanodiscus hantzschii*, in BO *Achnantheidium minutissimum*, *Pantocsekiella ocellata* and *Mastogloia smithii*. In the examined ecosystems, the most commonly identified classes were accidental (PE-84; LE-111; BO-86), accessory (PE-25; LE-35) and euconstant taxa (BO-39) (Fig. 2).

#### *nMDS analysis*

nMDS analysis (stress level = 0.07) identified the variation between samples for each studied hydrological object (Fig. 3). The samples taken from BO constitute a separate cloud, with the samples demonstrating high similarity with each other, whereas the samples of D.LEP1, D.LEP2 and D.LEP3 constitute a distinct group, with no clear differentiation into individual reservoirs. In the case of PE, the pond group (D.PEPB) was found to be clearly distinct from the others.

#### *Shade Plot*

Shade Plot analysis identified 50 species which had the strongest influence on the degree of similarity, or non-similarity, between the samples in the studied ecosystems. Of these taxa, the three that most strongly influenced the similarity between the samples in at least two ecosystems were *Navicula veneta* (NVEN), *N. cincta* (NCCA), *N. gregaria* (NGRE), *Nitzschia frustulum* (NIFR), *N. inconspicua* (NINC), *N. palea* (NPAL), *Planothidium frequentissimum* (PLFQ), *Amphora pediculus* (APED), *Cyclotella meneghiniana* (CMEN), *Fragilaria radians* (FRAD) and *Achnantheidium minutissimum* (ADMI) (Fig. 4).

#### *SIMPER analysis*

SIMPER analysis allowed 36 species characteristic of the tested hydrological objects to be distinguished (Table 3). In addition, two species were found to be characteristic of two different ecosystems: *Achnantheidium minutissimum* for LE and BO, and *Nitzschia palea* for PE and LE.

#### *Ecological analysis based on Omnidia software*

The ecological analysis of diatom assemblages based on data obtained from the OMNIDIA program database, indicated the following:

- pH requirements: while alkaliphilic species predominate in PE (63%), a large percentage in D.PEPB are unknown (24%) or neutrophilic species (23%). The LE reservoirs were dominated by alkalibiontic (45%) and alkaliphilic (24%) organisms. In D.LEP1, 25% of species were unknown. BO was dominated by alkaliphilic (39%) and neutrophilic (29%) species, and 26 % of species were unknown (Fig 5a);
- salinity: the PE complex was characterized by the occurrence of halophilic (43%), oligohalobous (30%) and mesohalobous species (16%); the greatest proportion of the mesohalobous species were found in D.PESB (28%). The LE reservoirs were dominated by

oligohalobous (44%) and halophilic species (42%). BO was dominated by oligohalobous (43%), halophobe (24%) and unknown species (23%) (Fig 5b).

- nitrogen uptake: the most common species in the PE complex N-autotrophic tolerant (39%) followed by unknown (25%). The largest percentage of unknown species (28%) was recorded in D.PEPB and D.PEDB. In the LE reservoirs, the most common groups of species were N-autotrophic (57%) and unknown (25%). In BO, 51% species were unknown, 24% were N-autotrophic tolerant and 22% N-autotrophic sensitive (Fig 5c);

- oxygen requirements: in PE, the largest groups of species were low oxygen (30%), unknown (27%) and moderate oxygen (24%). In LE, oxybiontic species were most common (43%) followed by unknown (25%). In BO, unknown (46%) and polyoxybiontic species (42%) predominated (Fig 5d);

- sensitivity to saprobity: in PE, the largest group of taxa were  $\alpha$ -meso: polysabrobe (28%) and unknown (23%). In D.PEPB, the most abundant was  $\alpha$ -meso-polysabrobe (34%) followed by  $\beta$ -mesosaprobe (31%) and unknown (27%). LE primarily included taxa from the  $\alpha$ -mesosaprobe group (47%) and unknown (23%). In BO, unknown (34%),  $\beta$ -mesosaprobe (31%) and oligosaprobe taxa (28%) predominated (Fig 5e);

- trophic status: in PE, the largest group of diatoms were eutrophic (50%) and unknown taxa (25%), LE had the highest percentage (61%) of eutrophic species but also unknown (15%) and hypereutrophic (13%) were present. In BO, the most abundant species were unknown (42%), indifferent (19%) and meso-eutrophic (16%) (Fig 5f);

- moisture aerophily: in PE, the largest group was aquatic to aerophilic (56%), representing 66% of species in D.PESB, 61% in D.PEDB, and 42% in D.PEPB. The second largest group was unknown (23%), constituting 32% of taxa in D.PEPB. In LE, 37% of the species were aquatic (24% of taxa in D.LEP2), 54% were occasionally aerophilic and 22% were unknown. In BO, the predominant groups of species were unknown (44%) and aquatic to aerophilic (33%) (Fig 5g).

#### *Characteristic species: OMNIDIA and PCA analysis*

The analysis of species characteristic of the tested ecosystems, determined according to Van Dam et al. (1994), identified 16 taxa classified as 0 in at least one category (Table 4). The next step determined the percentage contribution of each of these species classified as class 0 for the ecological parameters defined by Van Dam et al. (1994) at each sampling point (Table 5).

The principal components analysis (PCA) was performed to find the relationships between the abiotic parameters and the characteristic species ( $n=36$ ) (Fig. 6). The Eigenvalues Plot method given eigenvalues above 1%, showed that 12 factors account for 83,2% of the total variance. The first two factors account for 31,3% of the total variance. Based on the PCA analysis for of the 16 characteristic taxa mentioned above and physical and chemical parameters, the following relationships were demonstrated:

- Group A: *Chamaepinnularia krookiformis*, *C. plinskii*, *Nitzschia liebethruthii* and *Planothidium delicatulum* demonstrate a negative correlation with pH and a positive correlation with a decrease in the concentrations of  $\text{HCO}_3^-$ ,  $\text{Ca}^{2+}$  and  $\text{Fe}^{2+/3+}$  (Fig. 6).

- Group B: *Craticula buderi*, *Planothidium frequentissimum* and *Navicula cincta* did not demonstrate any relationship with any water parameters (Fig. 6).  
- Group C: *Navicula moskalii*, *Cyclostephanos invisitatus*, *Stephanodiscus parvus*, *S. binatus*, *Diatoma moniliformis*, *Nitzschia dissipata* var. *media*, *Mastogloia smithii*, *Pantocsekiella pseudocomensis* and *Encyonopsis subminuta* demonstrated a negative correlation with a decrease in electrolyte conductivity, as well as with the concentrations of  $K^+$ ,  $Mg^{2+}$ ,  $Na^+$ ,  $SO_4^{2-}$ ,  $Cl^-$ ,  $PO_4^{3-}$  and  $Mn^{3+}$  and water pigments (Fig. 6).

*Ecological values of characteristic species*

*Characteristic species for PE*

### ***Planothidium delicatulum* (PTDE) (Figs 7. A1-A6)**

*Planothidium delicatulum* is a euconstant taxon for PE and an accidental taxon for LE. Its mean percentage share in PE was 2%, and constituted 5% in D.PESB.

*Planothidium delicatulum* does not currently have one ecological indicator value (oxygen requirements) according to Van Dam et al. (1994).

This species was more abundant in environments such as D.PESB, which was also characterized by the highest concentration of  $Cl^-$  (up to 2976 mg L<sup>-1</sup>), elevated electrolytic conductivity, and decreased  $K^+$  concentration. The pH of the water in which this species was observed did not exceed 7.

*Planothidium delicatulum* was mainly recorded in salty and brackish environments with neutral or slightly alkaline conditions (Campeau, Pienitz & Héquette 1999; Gell et al. 2005; Caballero et al. 2013; Yamamoto, Chiba & Tuji 2017; Van de Vijver et al. 2018).

Based on our findings, we suggest changing the following ecological indicator values in the Van Dam et al. (1994) classification for *P. delicatulum*:

- pH requirements: 3 (neutrophilic) (changing from 5 to 3);
- salinity: 5 (brackish-marine) (changing from 4 to 5)

### ***Chamaepinnularia krookiformis* (CHKF) (Figs 7. B1-B5) and *Chamaepinnularia plinskii* (CHPL) (Figs 7. C1-C5)**

In 2016 *Chamaepinnularia krookiformis* was divided into two separate taxa: *C. krookiformis* and *C. plinskii* (Żelazna-Wieczorek & Olszyński 2016). Both species were very often recorded together in the same ecosystem. However, the publications which identified *C. krookiformis* often do not provide appropriate photographic documentation or photos of individual specimens (Witkowski 1994; Bąk et al. 2006; Wojtal 2008; Peszek et al. 2015). Currently available documentation is insufficient to determine whether *C. krookiformis* and *C. plinskii* are both present simultaneously in a given environment or whether just one of these species exists. *Chamaepinnularia krookiformis* is an accessory taxon for PE (a constant taxon for D.PEDB), *C. plinskii* is a constant taxon for PE (a euconstant taxon for D.PEDB). The mean share of *C.*

*crookiformis* was 1.6% in all PE samples, 4% in D.PEDB; for *C. plinskii*, this amounted to 2.7% in PE, 6% in D.PEDB.

Currently, *C. crookiformis* lacks three assigned ecological indicator values. For PE, it constitutes 6% of the *unknown* group in nitrogen uptake, 6% in oxygen requirements and 6% in trophic state (respectively for D.PEDB: 16%, 16% and 15%). *C. plinskii* has no assigned ecological indicator values and represents 26% of the *unknown* group for pH requirements, 19% for salinity, 12% for nitrogen uptake, 10% for oxygen requirements, 14% for saprobity, 11% for trophic state and 10% for moisture (respectively for D.PEDB: 60%, 25%, 24%, 23%, 32%, 23% and 28%).

The conditions of the environments in which both species have been recorded indicate that they are class 3 with regard to pH range (neutrophilic). Both species were the most abundant in locations subjected to periodic drying and characterised by high concentrations of chloride ions (up to 1006 mg L<sup>-1</sup>) indicating a brackish environment (Żelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015).

On the basis of our findings and those of previous studies (Krammer & Lange-Bertalot 1986; Krammer 1992; Witkowski 1994; Bąk, Witkowski & Lange-Bertalot 2006; Wojtal 2009; Peszek et al. 2015; Żelazna-Wieczorek & Olszyński 2016), according to Van Dam et al (1994) ecological indicator system we propose:

established ecological indicator values for *Chamaepinnularia plinskii*

- pH requirements: 3 (neutrophilic);
- salinity: 4 (mesohalobous);
- trophic state: 5 (eutrophic);
- moisture aerophily: 4 (aerophilic);
- saprobity: 4 (α-meso-polysabrobe).

for *C. crookiformis*

- trophic state: 5 (eutrophic),

and the following changes for *Chamaepinnularia crookiformis*

- salinity: from 3 to 4 (mesohalobous);
- moisture aerophily: from 3 to 4 (aerophilic);
- saprobity: from 2 to 4 (β-mesosabrobe to α-meso-polysabrobe). Due to the specific conditions and locations of the studied objects, they were exposed to large fluctuations in the inflow of organic matter, mainly from runoff from arable fields and pollution caused by animal grazing. These impurities were manifested as elevated concentrations of K<sup>+</sup> ions. Therefore, our results suggest that classifying *C. crookiformis* as an oligosabrobe is inappropriate. Further tests are needed to determine the optimum occurrence of these species in areas subjected to organic matter loads.

#### *Nitzschia liebethruthii* (NLBT) (Figs 7. D1-D4)

*Nitzschia liebethruthii* is a euconstant taxon for PE. Its means percentage share was 4% in the PE samples, and 10% in the D.PEDB samples. It was most numerous in the sample from November 2013 (19%). This species has two specific ecological indicator values. The ecological indicator



value analysis for PE found *N. liebethruthii* to represent 17% of the *unknown* group in nitrogen uptake, 16% in oxygen requirements, 18% in saprobity, 17% in trophic state and 21% in moisture (respectively for D.PEDB: 37%, 37%, 43%, 36% and 49%).

*Nitzschia liebethruthii* occurred in environments subjected to periodic drying with a pH close to 7 and high concentration of chloride ions.

This species was noted in environments with increased salinity, electrolytic conductivity and high pH value (Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Lange-Bertalot et al. 2017; Földi et al. 2018).

We propose established new ecological indicator values according to Van Dam et al. (1994) assigned to *N.liebethruthii*:

- trophic state: 5 (eutrophic);

- moisture aerophily: 4 (aerophilic),

and following changes:

- pH requirements: 3 (neutrophilic) (changing from 5 to 3);

- saprobity: from 2 to 4 ( $\beta$ -mesosaprobe to  $\alpha$ -meso-polysaprobe).

#### ***Craticula buder* (CRBU) (Figs 7. E1-E5)**

*Craticula buder* is a euconstant taxon for PE and an accidental taxon for LE. Its mean percentage share was 4.3% in all samples for PE, and 12% for D.PEPB. This species has no recorded ecological indicator values. The ecological indicator value analysis for the PE found *C. buder* to constitute 19% of the *unknown* group in pH requirements, 28% in salinity, 16% in nitrogen uptake, 16% in oxygen requirements, 16% in saprobity, 19% in trophic state and 15% in moisture (respectively for D.PEPB: 53%, 73%, 46%, 46%, 46%, 54% and 41%).

Although *Craticula buder* was classified into group B (PCA), it was found to be most abundant in environments with an elevated concentration of  $\text{Cl}^-$  ions, ranging from 685 to 1090  $\text{mg L}^{-1}$ , (all samples from D.PEPB and one sample from D.PEDB in which the concentration of chloride ions was 1006  $\text{mg L}^{-1}$ ). However, relative abundance was lower in the D.PEPB sample, which was characterised by a chloride ion content of over 1500  $\text{mg L}^{-1}$ . Interestingly, the concentration of  $\text{K}^+$  ions exceeded 100  $\text{mg L}^{-1}$  at  $\text{Cl}^-$  concentrations below 1500  $\text{mg L}^{-1}$ ; therefore, it is possible that the decline of this species could be related to the concentration of  $\text{K}^+$  ions alone. Our observations indicate that the population of *C. buder* from D.PEPB favours a concentration of chloride ions from 500 to 1006  $\text{mg L}^{-1}$  which coincides with a  $\text{K}^+$  ions concentration from 50 to 70  $\text{mg L}^{-1}$ .

*Craticula buder* is widespread throughout the world and recognized as cosmopolitan (Rumrich, Lange-Bertalot & Rumrich 2000; Lange-Bertalot 2001; Bahls 2009; Soltanpour-Gargari, Lodenius & Hinz 2011; Żelazna-Wieczorek 2011; Cichoń 2016). This species was found to be dominant in environments characterised by increased electrolytic conductivity and alkaline water (Holmes & Taylor 2015). Holmes and Taylor (2015) place *C. buder* in the Bad water quality class. Their recorded values of diatom indices indicate that the environment was eutrophic.



We therefore propose the following classes of ecological indicator values according to Van Dam et al. (1994) for *Craticula buderi*:

- pH requirements: 4 (alkaliphilic);
- trophic state: 5 (eutrophic);
- salinity: 4 (mesohalobous);
- moisture aerophily: 3 (aquatic to aerophilic),

and following change:

- sabrobity: from 2 to 4 ( $\beta$ -mesosaprobe to  $\alpha$ -meso-polysaprobe).

#### ***Navicula cincta* (NCCA) (Figs 7. F1-F5)**

*Navicula cincta* is a euconstant taxon for PE, a constant taxon for LE and an accessory taxon for BO. The mean percentage share of this species for PE is 3%, of which 7% was found in D.PEPB samples.

Currently this species has been assigned two ecological indicator values. The ecological indicator values analysis for the PE found *N. cincta* to constitute 24% of the *unknown* group in pH requirements, 13% in nitrogen uptake, 12% in oxygen requirements, 13% in saprobity, and 12% in moisture (respectively for D.PEPB: 28%, 25%, 25%, 26% and 23%).

An analysis of the physical and chemical data and the variability of occurrence did not show any clear relationships between environmental parameters and the percentage share of *Navicula cincta* in the tested samples. This lack of dependence is also confirmed by the PCA analysis.

*Navicula cincta* has been recorded in various types of ecosystems, although mainly in eutrophic ones with high conductivity. It also tolerates elevated levels of organic matter. This species was also observed in habitats subjected to periodic drying (Lange-Bertalot & Genkal 1999; Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Lange-Bertalot 2001; Żelazna-Wieczorek 2011; Wojtal 2013; Lange-Bertalot, Hofmann & Werum 2017).

However, several new species from the group *N. cincta* s.l. have been described, and it can be assumed that each of these individual species in this group may be associated with narrower optimal ecological conditions (Cantonati et al. 2016).

Based on our present findings, and those of previous studies, in our opinion that it is not appropriate to classify *Navicula cincta* as an oligohalobous species with regard to *salinity*: it has been recorded in fresh (Żelazna-Wieczorek 2011; Wojtal 2013), brackish (Żelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015; Żurek et al. 2018) and salt waters (Witkowski, Lange-Bertalot & Metzeltin 2000). We propose the following ecological indicator value according to Van Dam et al. (1994) for *Navicula cincta* s.l.:

- moisture aerophily: 3 (aquatic to aerophilic).

Shade Plot analysis found that the presence of *N. cincta* s.l. can falsely indicate high similarity between samples from different environments, thus distorting the results of any environmental analysis. Therefore, with regard to the unclear taxonomic status of *Navicula cincta* s.l. and the current lack of knowledge regarding its activities, we recommend this taxon be excluded from the biological assessment of surface water quality.

#### 473 *Planothidium frequentissimum* (PLFQ) (Figs 7. G1-G5)

474 *Planothidium frequentissimum* is a euconstant taxon for PE and LE. The mean percentage of this  
 475 species for PE is 2.7%, of which 5% was found in D.PEPB samples. It was most numerous in the  
 476 D.PEDB.301113 sample (19%). The species has currently six established ecological indicator  
 477 values. The ecological indicator values analysis for PE found *P. frequentissimum* to constitute  
 478 12% of the *unknown* group in moisture aerophily (for D.PEPB 14% and D.PESB: 20%).  
 479 No relationship was observed between percentage share of *Planothidium frequentissimum* and  
 480 the changes in chemical and physical parameters in the tested samples. This lack of relationship  
 481 was confirmed by PCA analysis.

482 *Planothidium frequentissimum* is an eurytopic species that occurs globally in a variety of habitat  
 483 types, from natural springs to rivers in urban areas with high levels of pollution. Its value as an  
 484 indicator is low, as confirmed by the Shade Plot analysis, which found it to significantly affect  
 485 the degree of similarity observed between samples from different environments.  
 486 Recently *P. frequentissimum* was divided into several different species. In studied samples we  
 487 observe several species which belong to *P. frequentissimum* s. l. (Fig. 7. G1-G7) (Wetzel et al.  
 488 2019). A light microscope (LM) is still used to identify species in ecological research and  
 489 biological assessment of aquatic ecosystems. Due to the likeness of the basic morphological  
 490 features of the newly described species observed in LM, especially in the case of *P.*  
 491 *frequentissimum* s.s. and *P. straubianum*, distinguishing them will be difficult or limited, which  
 492 may lead to errors in the assessment. We therefore recommend that *P. frequentissimum* s. l. be  
 493 excluded from the biological assessment of surface water quality (Siver, et al. 2005; Levkov et al.  
 494 2007; Żelazna-Wieczorek 2011; Kulikovskiy, Lange-Bertalot & Kuznestova 2015; Szczepocka,  
 495 Nowicka-Krawczyk & Kruk 2018).

497 *Characteristic species for LE*

#### 498 *Cyclostephanos invisitatus* (CINV) (Figs 8. A1-A6)

499 *Cyclostephanos invisitatus* is a euconstant taxon for LE. Its percentage share for LE was 4.8%.  
 500 *C. invisitatus* currently has two ecological indicator values assigned. The ecological indicator  
 501 values analysis for LE found it to constitute 24% of the *unknown* group in pH requirements, 17%  
 502 in nitrogen uptake, 18% in oxygen requirements, 19% in saprobity, and 20% in moisture.

503 *Cyclostephanos invisitatus* occurs in diverse environments, however, it is most frequently  
 504 reported in aquatic ecosystems subjected to high human impact, alkaline and high conductivity  
 505 (Reavie & Smol 1998; Yang et al. 2005; Wojtal & Kwadrans, 2006; Kiss et al. 2012; Houk,  
 506 Klee & Tanaka 2014; Reavie & Kireta 2015; Olszyński & Żelazna-Wieczorek 2018).

507 We therefore propose that the following classes of ecological indicator values according to Van  
 508 Dam et al. (1994) be established for *Cyclostephanos invisitatus*:

- 509 - pH requirements: 4 (alkaliphilic);
- 510 - moisture aerophily: 1 (aquatic).

# 511 *Navicula moskalii* (NMOK) (Figs 8. B1-B6)

512 *Navicula moskalii* is an accessory taxon for LE. Its mean percentage share for LE was 1.5%. Its  
513 incidence was greatest in sample D.LEP1.250315 (26%).

514 *Navicula moskalii* has no assigned ecological indicator values. The ecological indicator value  
515 analysis for LE found it to constitute 5% of the *unknown* group in pH requirements, 7% in  
516 salinity, 5% in nitrogen uptake, 5% in oxygen requirements, 5% in saprobity, 6% in trophic state  
517 and 5% in moisture. The greatest occurrence of *N. moskalii* was observed in samples with the  
518 highest concentrations of  $\text{Ca}^{2+}$  (143.6 mg L<sup>-1</sup>),  $\text{HCO}_3^-$  (338.6 mg L<sup>-1</sup>),  $\text{SO}_4^{2-}$  (146.9 mg L<sup>-1</sup>) and  
519 with high  $\text{Mg}^{2+}$  content.

520 *Navicula moskalii* was observed in a number of ecosystems (Metzeltin & Witkowski 1996;  
521 Lange-Bertalot 2001; Żelazna-Wieczorek 2011; Noga et al. 2016; Lange-Bertalot et al. 2017),  
522 particularly in eutrophic waters with an elevated level of  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  ions. Żelazna-  
523 Wieczorek (2011) report a significant number of *N. moskalii* in springs with high levels of  
524 eutrophication, however with  $\text{Ca}^{2+}$ ,  $\text{SO}_4^{2-}$ ,  $\text{HCO}_3^-$  and  $\text{Mg}^{2+}$  concentrations lower than those in  
525 the LE samples.

526 We therefore propose that the following classes of ecological indicator values according to Van  
527 Dam et al. (1994) be established for *Navicula moskalii*:

- 528 - pH requirements: 4 (alkaliphilic);
- 529 - salinity: 2 (oligohalobous);
- 530 - trophic state: 7 (indifferent).

# 531 *Stephanodiscus binatus* (SBNT) (Figs 8. C1-C5)

532 *Stephanodiscus binatus* is a euconstant taxon for LE. Its percentage share for LE was 4.3%. *S.*  
533 *binatus* has no recorded ecological indicator values. The ecological indicator value analysis for  
534 LE found it to constitute 25% of the *unknown* group in pH requirements, 47% in salinity, 18% in  
535 nitrogen uptake, 18% in oxygen requirements, 20% in saprobity, 29% in trophic state and 21% in  
536 moisture.

537 The largest percentage share of *S. binatus* was recorded in the spring months and the lowest in  
538 autumn. Its abundance was found to be elevated in December 2014 and 2015; the same samples  
539 demonstrated the highest concentrations of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and the highest pH (above 8).

540 *Stephanodiscus binatus* has been recorded in various water ecosystems ranging from  
541 oligotrophic to eutrophic; however, all are characterised by elevated pH value (Stoermer &  
542 Håkansson 1984; Håkansson & Kling 1990; Houk, Klee & Tanaka 2014; Olszyński & Żelazna-  
543 Wieczorek 2018).

544 We therefore propose that the following classes of ecological indicator values according to Van  
545 Dam et al. (1994) be established for *Stephanodiscus binatus*:

- 546 - pH requirements: 4 (alkaliphilic);
- 547 - salinity: 2 (oligohalobous);

# 548 *Stephanodiscus parvus* (SPAV) (Figs 8. D1-D5)

*Stephanodiscus parvus* is a euconstant taxon for LE and an accidental taxon for PE. Its percentage share for LE was 2.4%. It was most abundant in the D.LEP3.260714 sample (22%). This species has three assigned ecological indicator values. The ecological indicator values analysis for LE found the taxon to constitute 9% of the *unknown* group in nitrogen uptake, 9% in oxygen requirements, 10% in saprobity, and 10% in moisture. *S. parvus* is noted mainly in eutrophic hypereutrophic ecosystems with elevated electrolytic conductivity. It is also a good indicator of waters with a strong anthropogenic impact (Reavie & Smol 1998; Reavie & Kireta, 2015; Olszyński & Żelazna-Wieczorek 2018; Reavie & Cai 2019). Based on our findings and literature data, we propose the following change in ecological indicator values according to Van Dam et al. (1994) for *Stephanodiscus parvus*:  
- pH requirements: from 5 to 4 (alkaliphilic);

*Characteristic species for BO*

#### ***Diatoma moniliformis* (DMOF) (Figs 9. A1-A6)**

*Diatoma moniliformis* is a euconstant taxon for BO. Its mean percentage share for BO was 3.9%. It currently has no assigned ecological indicator values. According to the ecological indicator values analysis for BO, this taxon constituted 10% of the *unknown* group in pH requirements, 11% in salinity, 8% in nitrogen uptake, 8% in oxygen requirements, 9% in saprobity, 9% in trophic state and 8% in moisture. *D. moniliformis* was found in 87.5 % of samples from BO. Interestingly, it constituted 28% of the share in one sample from December 2016 (D.BOZB.091216); however, its share was below 2% in the previous season, and was not higher than 1-2% in the other samples from December 2016. The chemical and physical characteristics of D.BOZB.091216 did not differ significantly from those of the other samples. This species is also found in fresh and salt water, as well as the Baltic and arctic areas with high conductivity (Potapova & Snoeijis 1997; Rumrich, Lange-Bertalot & Rumrich 2000; Levkov et al. 2007; Pniewski & Sylwestrzak 2018). One of the factors that influences the abundance of *D. moniliformis* is the water temperature. Studies indicate that temperatures above 10-15°C (Potapova & Snoeijis 1997; Pniewski & Sylwestrzak 2018) are associated with population growth. However, populations have been observed in freshwater streams and lakes in arctic areas, in which the temperature of the water is below 10°C (Antoniades Douglas & Smol 2005). Population growth was also observed at 4.7°C in sample D.BOZB.091216; therefore, low temperature may have an influence on the abundance of this species.

#### ***Encyonopsis subminuta* (ESUM) (Figs 9. B1-B6)**

*Encyonopsis subminuta* is a euconstant taxon in BO, where its mean percentage share was 4.1%. Presently, *E. subminuta* has been assigned five ecological indicator values. Ecological indicator

values analysis for BO found it to constitute 7% of the *unknown* group in *nitrogen uptake* and 8% in *moisture*.

*Encyonopsis subminuta* was found to be most abundant in sample D.BOZB.041115. The sample was also characterised by an elevated concentration of  $\text{Fe}^{2+/3+}$  ions and the lowest pH value. In subsequent samples, when the concentration of Fe ions dropped, the abundance of *E. subminuta* also decreased.

*E. subminuta* is regarded as a cosmopolitan taxon, occurring in the temperate and boreal zone. It is most abundant in oligo- to mesotrophic waters with electrolytic conductivity between 190- 250  $\mu\text{S L}^{-1}$  (Krammer 1997; Noga et al. 2014; Novais et al. 2014; Feret, Bouchez & Rimet 2017).

*E. subminuta* may be sensitive to the concentration of Fe ions; however, the increase of these ions is associated with a drop in pH. Our research confirms that the optimal pH for population size is close to 7.

#### ***Mastogloia smithii* (MSMI) (Figs 9. C1-C7)**

*Mastogloia smithii* is a euconstant taxon for BO. Its mean percentage share for BO was 6.3%. it was found in greatest numbers in D.BOZB.300615 (22%) and D.BOZB.261016 (15%). *M. smithii* has been assigned four ecological indicator values. The ecological indicator values analysis for the BO found this species to constitute 13% of the *unknown* group in nitrogen uptake, 14% in oxygen requirements and 15% in trophic state.

The environment in BO regarding *salinity* was classified according to Vam Dam et al. (1994) as oligohalobus (43% species); however, 7% of the mesohalobous species were represented by one species: *Mastogloia smithii*.

This species is recorded in fresh, brackish and salt water (Witkowski, Lange-Bertalot & Metzeltin 2000; Busse & Snoeijs 2003; Weckström & Juggins 2005; Martinzes-Goss & Evangelista 2011; Lange-Bertalot et al. 2017). Its presence in environments with varying degrees of salinity may suggest that this does not have a significant influence on population size.

Based on our present findings and literature data, we propose the following change in the ecological indicator values according to Van Dam et al. (1994) for *Mastogloia smithii*:

- salinity: from 4 to 3 (halophilic).

#### ***Nitzschia dissipata* var. *media* (NDME) (Figs 9. D1-D5)**

*Nitzschia dissipata* var. *media* is a euconstant taxon for BO. Its mean percentage share for BO was 3.8%, and the highest proportion (20%) was found in D.BOZB.041115. *N. dissipata* var. *media* has been assigned two ecological indicator values. Ecological indicator values analysis for BO found it to constitute 9% of the *unknown* group in nitrogen uptake, 10% in oxygen requirements, 11% in saprobity, 11% in trophic state and 9% in moisture.

The increase in occurrence of *N. dissipata* var. *media* is associated with an increase in the level of  $\text{Fe}^{2+/3+}$  ions, similar to *Encyonopsis subminuta*. In addition, it was found in the ecosystem, i.e. BO, with the lowest concentrations of ions indicative of the presence of organic pollutants in the environment, such as  $\text{K}^+$  and  $\text{NH}_4^+$ .



Although *Nitzschia dissipata* var. *media* is found sporadically, it is commonly found in oligo- to mesotrophic waters with a pH between 7 and 8 (Van der Vijver, Frenot & Beyens 2002; Antoniadou, Douglas & Smol 2005; Żelazna-Wieczorek 2011; Lange-Bertalot, Hofmann & Werum 2017).

Based on our findings and literature data, we propose the following ecological indicator values according to Van Dam et al. (1994) for *Nitzschia dissipata* var. *media*:

- saprobity: 2 (β-mesosaprobe)

### ***Pantocsekiella pseudocomensis* (PPCS) Figs (9. E1-E7)**

*Pantocsekiella pseudocomensis* is a euconstant taxon for BO. Its mean percentage share for BO was 4.4%. It was most abundant in D.BOZB.250315 (9%) and in D.BOZB.220616 (10%). *P. pseudocomensis* has not been assigned any ecological indicator values according to Van Dam et al. (1994). The ecological indicator values analysis for BO found it to represent 22% of the unknown group in pH requirements, 25% in salinity, 10% in nitrogen uptake, 11% in oxygen requirements, 18% in saprobity, 12% in trophic state and 13% in moisture.

The greatest amount of *P. pseudocomensis* was found in samples characterised by the highest levels of ammonium ions. Its percentage share was lowest in samples with the lowest water temperature, apart from D.BOZB.250315.

Currently, *Pantocsekiella pseudocomensis* is assigned to the *P. comensis* complex, with *P. comensis* and *P. costei*. In our opinion that assigning ecological indicator values for particular species of the *P. comensis* complex is unjustified at the current state of knowledge, and that all species within the complex should be assigned the same provisional ecological indicator values until their individual properties are better understood (Houk, Klee & Tanaka 2010; Kistenich et al. 2014; Duleba et al. 2015).

## **Conclusions**

The water ecosystems created in the post-mining areas create a complex of conditions that are not found in other natural ecosystems, and the benthic diatom species present in such environments are very often present in higher numbers than in other habitats. The specific hydro-geological conditions prevailing in the post-production reservoirs provide a unique opportunity to observe interspecies differences and intra-species variability, allowing for the verification or isolation of new taxa and a greater insight into their autecology (Żelazna-Wieczorek & Olszyński 2016; Olszyński & Żelazna-Wieczorek 2018).

The identification of species characteristic of the studied ecosystems may foster further growth of ecological research and increase the reliability of surface water quality assessment, as such knowledge is needed to verify their ecological indicator values, and hence calculate diatoms indices with greater accuracy.

Ecological indicator values as set out by Van Dam et al. (1994) are utilized in many ecological works describing the ecological conditions of the studied ecosystems. These ecological indicator values form the basis for calculating diatoms indices describing the ecological state of surface



waters. It is therefore necessary to constantly update and establish new ecological indicator values for particular diatom species. Many authors who describe new species, or encounter existing species in new ecosystems, regularly propose updates for individual ecological indicator values. However, these findings, may not be introduced and updated in the OMNIDIA program for a number of years. The OMNIDIA system is used by state institutions in many countries around the world to assess surface water quality (Campeau, Pienitz & Héquette 1999; Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Gell et al. 2005; Potapova & Ponader 2008; Wojtal & Sobczyk 2012; Caballero et al. 2013; Żelazna-Wieczorek & Olszyński 2015; Yamamoto, Chiba & Tuji 2017; Lange-Bertalot, Hofmann & Werum 2017; Földi et al. 2018; Van de Vijver, Wetzel & Ector 2018). Clearly, if these assessments are based on incomplete or outdated data, assessments of aquatic environments may be fraught with error. The present study used three principles to identify proposed changes in the classification of ecological indicator values for characteristic species according to the Van Dam et al. (1994) system, or to establish new values which were previously absent: the analysis of environmental conditions prevailing in the studied ecosystems, the analysis of relevant literature data, and references to the ecological indicator values of other species (Table 6). This mode of research can serve as a model for updating databases used to assess surface water quality.

## References

- Antoniades D., Douglas M.S.V. & Smol J.P. 2005. Benthic diatom autecology and inference model development from the Canadian High Arctic Archipelago. *Journal of Phycology* 41: 30-45 DOI: 10.1111/j.1529-8817.2005.04049.x.
- Bahls L.L. 2009. A checklist of diatoms from inland waters of the northwestern United States. *Proceedings of The Academy of Natural Sciences of Philadelphia* 158: 1-35 DOI: 10.1635/053.158.0101.
- Bąk M., Witkowski A. & Lange-Bertalot H. 2006. Diatom flora diversity in the strongly eutrophicated and  $\beta$ -mesosaprobic waters of the Szczecin Lagoon, NW Poland, southern Baltic Sea. – In: Ognjanova-Rumenova N. & Manoylov K., eds. *Advances in Phycological Studies, Festschrift in Honour of Professor Dobrina Teminskova-Topalova*. Sofia – Moscow: Pensoft Publishers & University Publishing House, 293–317.
- Busse S. & Snoeijs P. 2003. Gradient responses of diatom communities in the Bothnian Sea (northern Baltic Sea), with emphasis on responses to water movement. *Phycologia* 42(5): 451-464 DOI: 10.2216/i0031-8884-42-5-451.1.
- Caballero M., Rodríguez A., Vilaclara G., Ortega B., Roy P. & Lozano S. 2013. Hydrochemistry, ostracods and diatoms in a deep, tropical, crater lake in Western Mexico. *Journal of Limnology* 72(3): 512-523 DOI: 10.4081/jlimnol.2013.e42.
- Campeau S., Pienitz R. & Héquette A. 1999. Diatoms from the Beaufort Sea coast, southern Arctic Ocean (Canada). In: Lange-Bertalot H. & Kociolek P., eds. *Bibliotheca Diatomologica Vol. 42*. Berlin, Stuttgart: J. CRAMER 244 p.

- Cantonati M., Angeli N. Spitale D. & Lange-Bertalot H. 2016. A new *Navicula* (Bacillariophyta) species from low-elevation carbonate springs affected by anthropogenic disturbance. *Fottea, Olomouc* 16(2): 255-265 DOI: 10.5507/fot.2016.013.
- Cichoń S. 2016. *Craticula buderi* (Bacillariophyceae) in Poland. *Polish Botanical Journal* 61(2): 301–305 DOI: 10.1515/pbj-2016-0024.
- Clarke K.R. & Gorley, R.N. 2015. PRIMER v7: User Manual/Tutorial. Devon, United Kingdom: PRIMER-E Ltd 296 pp.
- De Haan H., Van Liere L., Klapwijk S.J.P. & Van Donk E. 1993. The structure and function of fen lakes in relation to water table management in The Netherlands. *Hydrobiologia* 265: 155–177 <http://dx.doi.org/10.1007/BF00007266>.
- Duleba M., Kiss K.T., Földi A., Kovács J., Borojević K.K., Molnár L.F., Plenković-Moraj A., Pohner Z., Solak C.N., Tóth B. & Ács E. 2015. Morphological and genetic variability of assemblages of *Cyclotella ocellata* Pantocsek/*C. comensis* Grunow complex (Bacillariophyta, Thalassiosirales), *Diatom Research* 30(4): 283-306 DOI: 10.1080/0269249X.2015.1101402.
- European Union. 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. Official Journal of the European Communities L327, 1–73.
- Feret L., Bouchez A. & Rimet F. 2017. Benthic diatom communities in high altitude lakes: a large scale study in the French Alps. *Annales de Limnologie - International Journal of Limnology* 53: 411-423 DOI: 10.1051/limn/2017025.
- Ferreira da Silva, E., Almeida, A.F.P., Nunes, M.L., Luís, A.T., Borg, F., Hedlund, M., Marques de Sá, C., Patinha, C. & Teixeira, P. 2009. Heavy metal pollution downstream the abandoned Coval da Mó mine (Portugal) and associated effects on epilithic diatom communities. *Science of the Total Environment* 407: 5620–5636 doi:10.1016/j.scitotenv.2009.06.047.
- Földi A., Ács É., Grigorszky I., Ector L., Wetzel C.E., Várbíró G., Kiss K.T., Dobosy P., Trábert Z., Borsodi A.K. & Duleba M. 2018. Unexpected consequences of bombing. Community level response of epiphytic diatoms to environmental stress in a saline bomb crater pond area. *PLoS ONE* 13(10): e0205343 <https://doi.org/10.1371/journal.pone.0205343>.
- Gabrys-Godlewska A., Gruszecki J., Kochanowska J., Kozula R., Pasieczna A. & Tomassi-Morawiec H. 2004. Arkusz Turek (550). In: Sikorska-Maykowska M., Zdanowski A. & Gabrys-Godlewska A., eds. *Objaśnienia do mapy geośrodowiskowej Polski 1:50 000*. Warszawa: Państwowy Instytut Geologiczny. Opracowanie Zamówione Przez Ministra Środowiska. p 30.
- Gadomska W., Kaźmierczak J., Merdzińska M., Zielińska B., Wiśniewska-Zientara G., Gajda M., Opas Z., Cieślak J., Kończak D., Kujawa S., Lukaszewicz J., Zielińska-Adasiak S., Ratajczyk G., Michalska R. & Sobczak D. 2007. Raport o stanie gminy Brudzew lata 2002-2006. Urząd Gminy Brudzew. Brudzew. p. 15.

- 741 Gell P.A., Bulpin S., Wallbrink P., Hancock G. & Bickford S. 2005. Tareena Billabong – a  
742 palaeolimnological history of an ever-changing wetland, Chowilla Floodplain, lower  
743 Murray-Darling Basin, Australia. *Marine and Freshwater Research* 56: 441-456 DOI:  
744 10.1071/MF04107.
- 745 Harding W.R. & Taylor J.C., 2014. Diatoms as indicators of historical water quality: A  
746 comparison of samples taken in the Wemmershoek catchment (Western Province, South  
747 Africa) in 1960 and 2008. *Water S.A.* 40 (4): 601–606.
- 748 Håkansson H. & Kling H. 1990. The current status of some very small freshwater diatoms of the  
749 genera *Stephanodiscus* and *Cyclostephanos*. *Diatom Research* 5: 273-287  
750 DOI:10.1080/0269249X.1990.9705119.
- 751 Houk, V., Klee R. & Tanaka H. 2010. Atlas of freshwater centric diatoms with a brief key and  
752 descriptions. Part III. Stephanodiscaceae A. *Fottea (Supplement)* 10:1–498.
- 753 Houk V., Klee R. & Tanaka H. 2014. Atlas of freshwater centric diatoms with a brief key and  
754 descriptions. Part IV. Stephanodiscaceae B. *Fottea (Supplement)* 14:1–532.
- 755 Holmes M. & Taylor J.C. 2015 Diatoms as water quality indicators in the upper reaches of the  
756 Great Fish River, Eastern Cape, South Africa. *African Journal of Aquatic Science* 1-17  
757 DOI: 10.2989/16085914.2015.1086722.
- 758 Hutorowicz A. & Pasztalenic A. 2014. Phytoplankton metric of ecological status assessment for  
759 polish lakes and its performance along nutrient gradient. *Polish Journal of Ecology*  
760 62(3): 525-540 DOI:10.3161/104.062.0312.
- 761 Kahlert M., Ács E., Almeida S.F.P., Blanco S., Dreßler M., Ector L., Karjalainen S.M., Liess A.,  
762 Mertens A., van der Wal J., Vilaste S. & Werner P. 2016. Quality assurance of diatom  
763 counts in Europe: towards harmonized datasets. *Hydrobiologia* 772: 1–14 DOI:  
764 10.1007/s10750-016-2651-8.
- 765 Kasztelewicz, Z. 2011. Assessment of Reclamation Works in Polish Open-pit Lignite Mining.  
766 *Studia KPZK* 142: 440–452.
- 767 Kelly M.G., Juggins S., Guthrie R., Pritchard S., Jamieson J., Rippey B., Hirst H. & Yallop M.  
768 2008. Assessment of ecological status in U.K. rivers using diatoms. *Freshwater Biology*  
769 53 (2): 403–422 DOI: 10.1111/j.1365-2427.2007.01903.x.
- 770 Kiss K.T., Klee R., Ector L. & Ács É. 2012. Centric diatoms of large rivers and tributaries in  
771 Hungary: morphology and biogeographic distribution. *Acta Botanica Croatica* 71: 311–  
772 363 DOI: 10.2478/v10184-011-0067-0
- 773 Kistenich S., Dreßler M., Zimmermann J., Hübener T., Bastrop R. & Jahn R. 2014. An  
774 investigation into the morphology and genetics of *Cyclotella comensis* and closely  
775 related taxa. *Diatom Research* 29(4): 423-440 DOI: 10.1080/0269249X.2014.922125.
- 776 Kolada A., Pasztaleniec A., Bielczyńska A. & Soszka H. 2016. Phytoplankton, macrophytes and  
777 benthic diatoms in lake classification: consistent, congruent, redundant? Lessons learnt  
778 from WFD-compliant monitoring in Poland. *Journal of Limnology* 59: 44–52 DOI:  
779 10.1016/j.limno.2016.05.003.

- 780 Krammer K. 1992. Pinnularia. Eine Monographie der europäischen Taxa In: *Bibliotheca*  
781 *Diatomologica* 26. Berlin–Stuttgart: J. Cramer. 353 pp.
- 782 Krammer K. 1997. Die cymbelloiden Diatomeen. Eine Monographie der weltweit bekannten  
783 Taxa. Teil 2. *Encyonema* part., *Encyonopsis* and *Cymbellopsis*. In: Lange-Bertalot H. &  
784 Kociolek P., eds. *Bibliotheca Diatomologica Vol. 37*. Berlin, Stuttgart: J. CRAMER. 96  
785 pp.
- 786 Krammer K. & Lange-Bertalot H. 1986. Bacillariophyceae. 1. Teil: Naviculaceae In: Ettl H.H,  
787 Gerloff J., Heynig H. & Mollenhauer D., eds. *Süßwasserflora von Mitteleuropa, Band*  
788 *2/1*. Jena: Gustav Fisher Verlag. 876 pp.
- 789 Krammer K. & Lange-Bertalot H. 1997. Bacillariophyceae. 1. Teil: Bacillariaceae,  
790 Epithemiaceae, Surirellaceae. In: Ettl H.H, Gerloff J., Heynig H. & Mollenhauer D.,  
791 eds. *Süßwasserflora von Mitteleuropa, Band 2/2*. Jena Stuttgart Lübeck Ulm: Gustav  
792 Fisher Verlag. 611 pp.
- 793 Kulikovskiy M.S., Lange-Bertalot H. & Kuznetsova I.V. 2015. Lake Baikal: Hotspot of Endemic  
794 Diatoms II. In: Lange-Bertalot H., ed. *Iconographia Diatomologica Vol. 26. Taxonomy-*  
795 *Biogeography-Diversity*. Germany: Koeltz Scientific Books. 656 pp.
- 796 Lange-Bertalot H. 2001. *Navicula* sensu stricto. 10 Genera Separated from *Navicula* sensu lato.  
797 *Frustulia*. In: Lange-Bertalot H., ed. *Diatom of Europe Vol. 2*. Ruggell:vA.R.G. Gantner  
798 Verlag K.G. 526 pp.
- 799 Lange-Bertalot H. & Genkal S.I. 1999. Diatoms from Siberia I. Islands in the Arctic Ocean  
800 (Yugorsky-Shar Strait). In: Lange-Bertalot H., ed. *Iconographia Diatomologica Vol. 6.*  
801 *Phytogeography-Diversity-Taxonomy*. Vaduz: A.R.G. Gantner Verlag K.G. 292 pp.
- 802 Lange-Bertalot H., Hofmann G., Werum M. & Cantonati M. 2017. Freshwater Benthic Diatoms  
803 of Central Europe: Over 800 Common Species Used in Ecological Assessment. Koeltz  
804 Botanical Books. 942 pp.
- 805 Lecointe C., Coste M. & Prygiel J. 1993. “Omnidia:” a software for taxonomy, calculation of  
806 diatom indices and inventories management. *Hydrobiologia* 269/270: 509-513.
- 807 Levkov Z., Krstic S., Metzeltin D. & Nakov T. 2007. Diatoms of Lakes Prespa and Ohrid. About  
808 500 taxa from ancient lake system. In: Lange-Bertalot H., ed. *Inconografia*  
809 *Datomologica Vol. 16*. Ruggell: A.R.G. Gantner Verlag K.G. 613 pp.
- 810 Luís A.T., Durães, N., Fernandes Pinheiro de Almeida S. & Ferreira da Silva E. 2016.  
811 Integrating geochemical (surface waters, stream sediments) and biological (diatoms)  
812 approaches to assess AMD environmental impact in a pyritic mining area: Aljustrel  
813 (Alentejo, Portugal). *Journal of Environmental Sciences* 42: 215-226  
814 <http://dx.doi.org/10.1016/j.jes.2015.07.008>.
- 815 Luís A.T., Teixeira P., Almeida S.F.P., Ector L., & Matos J.X. & Ferreira da Silva E.A. 2009.  
816 Impact of Acid Mine Drainage (AMD) on Water Quality, Stream Sediments and  
817 Periphytic Diatom Communities in the Surrounding Streams of Aljustrel Mining Area  
818 (Portugal). *Water Air Soil Pollution* 200:147-167 DOI: 10.1007/s11270-008-9900-z.



- Macioszczyk A. & Dobrzyński D. 2002. Hydrogeochemia. Wydawnictwo Naukowe PWN. Warszawa. 448 pp.
- Martinez-Goss M.R. & Evangelista L.T. 2011. A Contribution to the Taxonomy of *Mastogloia* (Class Bacillariophyceae) in the Philippines. *Philippine Journal of Science* 140(1): 7-12.
- Metzeltin D. & Witkowski A. 1996. Diatomeen der Bären-Insel. Süßwasser- und marine Arten. In: Lange-Bertalot H., ed. *Iconographia Diatomologica. Annotated Diatom Micrographs. Vol 4. Taxonomy*. Königstein/Germany: Koeltz Scientific Books 20-21.
- Noga T., Stanek-Tarkowska J., Pajączek A., Kochman N. & Peszek Ł. 2014. Ecological assessment of the San River water quality on the area of the San Valley Landscape Park. *Journal of Ecological Engineering* 15(4): 12-22 DOI: 10.12911/22998993.1125453.
- Noga T., Stanek-Tarkowska J., Rybek M., Kochaman-Kędziora N., Peszek Ł. & Pajączek A. 2016. Diversity of diatoms in the natural, mid-forest Terebowiec Stream – Bieszczady National Park. *Journal of Ecological Engineering* 17(4): 232-247 DOI: 10.12911/22998993/64506.
- Novais M.H., Morais M.M., Rosado J., Dias L.S, Hoffmann L. & Ector L. 2014. Diatoms of temporary and permanent watercourses in Southern Europe (Portugal). *River Research and Applications* 30: 1216-1232 DOI: 10.1002/rra.2818.
- Olaczek R. 1963. Zbiorowiska roślinne torfowisk niskich okolic Łęczycy. Unpublished doctoral dissertation, University of Lodz, Poland.
- Olszyński R.M. & Żelazna-Wieczorek J. 2018. *Aulacoseira pseudomuzzanensis* sp. nov. and other centric diatoms from post iron ore mining reservoirs in Poland. *Diatom Research* 33(2): 155-185 DOI: 10.1080/0269249X.2018.1509886.
- Orlikowski D. & Szwed L. 2009. Wodny kierunek rekultywacji w KWB „Adamów” SA – inwestycją w przyszłość regionu. *Górnictwo i Geoinżynieria* 33(2): 351-361.
- Peszek Ł., Noga T., Stanek-Tarkowska J., Pajączek A., Kochman-Kędziora N., Pieniążek M. 2015. The effect of anthropogenic change in the structure of diatoms and water quality of the Żołynianka and Jagielnia streams. *Journal of Ecological Engineering* 16(2): 33-51 <https://doi.org/10.12911/22998993/1856>.
- Pliński M. 1966. Glony słonego źródła w Pełczyskach pod Ozorkowem. Unpublished master dissertation, University of Lodz, Poland.
- Pliński M. 1969. Okrzemki słonego źródła w Pełczyskach pod Ozorkowem. *Zesz. Nauk. UŁ* 2(31): 123-136.
- Pliński M. 1971a. Rzadkie i nowe dla flory Polski gatunki glonów z terenów solnisk w okolicach Łęczycy. *Zesz. Nauk. UŁ* 2(41): 171-184.
- Pliński M. 1971b. Gatunki z rodzaju *Enteromorpha* (Link.) Agardh z terenu solnisk podłęczyckich. *Zesz. Nauk. UŁ* 2(41): 159-169.
- Pliński M. 1971c. System halobow w świetle współczesnych poglądów. *Wiad. Ekol.* 17(1): 18-29.
- Pliński M. 1973. Glony solnisk podłęczyckich. The algae of salt marches near Łęczycza, Central Poland. *Mon. Botanic.* 39.

- 859 Pniewski F. & Sylwestrzak Z. 2018. Influence of short periods of increased water temperature on  
860 species composition and photosynthetic activity in the Baltic periphyton communities.  
861 *Biologia* 73(11): 1067-1072 <https://doi.org/10.2478/s11756-018-0122-6>.
- 862 Poikane S., Kelly M. & Cantonati M. 2016. Benthic algal assessment of ecological status in  
863 European lakes and rivers: Challenges and opportunities. *Science of the Total*  
864 *Environment* 568: 603–613 DOI:10.1016/j.scitotenv.2016.02.027.
- 865 Potapova M. & Snoeijs P. 1997. The natural life cycle in wild populations of *Diatom*  
866 *moniliformis* (Bacillariophyceae) and its disruption in an aberrant environment. *Journal*  
867 *of Phycology* 33: 924-937 <https://doi.org/10.1111/j.0022-3646.1997.00924.x>.
- 868 Potapova M.G. & Ponader K.C. 2008. New species and combinations in the diatom genus  
869 *Sellaphora* (Sellaphoraceae) from Southeastern United States. *Harvard Papers in*  
870 *Botany* 13(1): 171–181 DOI: 10.3100/1043-4534(2008)13[171:NSACIT]2.0.CO;2.
- 871 Rakowska B. 1996. The benthic diatom community of a reservoir after the exploration of brow  
872 coal in Konin (central Poland). *Algological Studies* 82: 103–116  
873 <http://dx.doi.org/0342-1120/0116-103>.
- 874 Reavie E.D. & Cai M. 2019. Consideration of species-specific diatom indicators of  
875 anthropogenic stress in the Great Lakes. *PLoS ONE* 14(5): e0210927 DOI:  
876 <https://doi.org/10.1371/journal.pone.0210927>
- 877 Reavie E.D. & Kireta A.R. 2015. Centric, Araphid and Eunotioid Diatoms of the Coastal  
878 Laurentian Great Lakes. In: Lange-Bertalot H. & Kociolek J.P., eds. *Bibliotheca*  
879 *Diatomologica Vol. 62*. Stuttgart: J. Cramer Gebr. Borntraeger Verlagsbuchhandlung.  
880 184 pp.
- 881 Reavie E.D. & Smol J.P. 1998. Freshwater diatoms from the St. Lawrence river. In: Lange-  
882 Bertalot H. & Kociolek J.P., eds. *Bibliotheca Diatomologica Vol 41*. Berlin, Stuttgart: J.  
883 Cramer Gebr. Borntraeger Verlagsbuchhandlung, 184 pp.
- 884 Rumrich U., Lange-Bertalot H. & Rumrich, M. 2000. Diatomeen der Anden. Von Venezuela bis  
885 Patagonien/Tierra del Fuego. Und zwei weitere Beiträge. In: Lange-Bertalot H., ed.  
886 *Iconographia Diatomologica Vol. 9*. Ruggell: A.R.G. Gantner Verlag K.G. 673 pp.
- 887 Sienkiewicz E. & Gąsiorowski M. 2016. The evolution of a mining lake – From acidity to  
888 natural neutralization. *Science of the Total Environment* 557–558: 343–354  
889 <http://dx.doi.org/10.1016/j.scitotenv.2016.03.088>.
- 890 Siver P.A., Hamilton P.B., Stachura-Suchoples K. & Kociolek J.P. 2005. The Freshwater Flora  
891 of Cape Code, Massachusetts, U.S.A. In: Lange-Bertalot H., eds. *Iconographia*  
892 *Diatomologica Vol. 14. Diatoms of North America*. Ruggell: A.R.G. Gantner Verlag  
893 K.G. 463 pp.
- 894 Soltanpour-Gargari A., Lodenius M. & Hinz F. 2011. Epilithic diatoms (Bacillariophyceae) from  
895 streams in Ramsar, Iran. *Acta Botanica Croatica*. 70(2): 167-190 DOI: 10.2478/v10184-  
896 010-0006-5.



- 897 Stoermer E.F. & Håkansson H. 1984. *Stephanodiscus parvus*: validation of an enigmatic and
- 898 widely misconstrued taxon. *Nova Hedwigia* 39: 497–511
- 899 <http://www.jstor.org/stable/3226406>.
- 900 Szczepocka E., Nowicka-Krawczyk P. & Kruk A. 2018. Deceptive ecological status of urban
- 901 streams and rivers—evidence from diatom indices. *Ecosphere* 9(7):e02310 DOI:
- 902 10.1002/ecs2.2310
- 903 Szczepocka E., Szulc B., Szulc K., Rakowska B. & Żelazna-Wieczorek J. 2014. Diatom indices
- 904 in the biological assessment of the water quality based on the example of a small
- 905 lowland river. *Oceanological and Hydrobiological Studies* 43(3): 265–273 DOI:
- 906 10.2478/s13545-014-0141-z
- 907 Szczepocka E. & Żelazna-Wieczorek J. 2018. Diatom biomonitoring – scientific foundations,
- 908 commonly discussed issues and frequently committed errors. *Oceanological and*
- 909 *Hydrobiological Studies* 47(3): 313-325 DOI: 10.1515/ohs-2018-0030
- 910 Thomas E.J. & John J. 2006. Diatoms and macroinvertebrates as biomonitors of mine-like in
- 911 Collie, Western Australia. *Journal of the Royal Society of Western Australia* 89:
- 912 109–117.
- 913 Tümpling W. & Friedrich G. 1999. Methiden der Biologischen Wasseruntersuchung. Jena.
- 914 Stuttgart. Lübeck. Ulm. G. Fischer Verlag 133-153.
- 915 Weckström K. & Juggins S. 2005. Coastal diatom–environment relationships from the Gulf of
- 916 Finland, Baltic Sea. *Journal of Phycology* 42: 21-35 DOI: 10.1111/j.1529-
- 917 8817.2006.00166.x
- 918 Witkowski A. 1994. Recent and fossil diatom flora of the Gulf of Gdańsk, Southern Baltic Sea.
- 919 In: *Bibliotheca Diatomologica Vol. 28*. Cramer, Berlin, Stuttgart. 313 pp
- 920 Witkowski A., Lange-Bertalot H. & Metzeltin D. 2000. Diatom flora of marine coasts I. In:
- 921 Lange-Bertalot H., ed. *Iconographia Diatomologica. Vol. 7. Diversity-Taxonomy-*
- 922 *Identification*. Ruggell, Liechtenstein: A.R.G. Gantner Verlag K.G. 419 pp.
- 923 Wetzel C.E., Van de Vijver B., Blanco S. & Ector L. 2019. On some common and new cavum–
- 924 bearing Planorbulina (Bacillariophyta) species from freshwater. *Fottea* 19(1): 50–89
- 925 DOI: 10.5507/fot.2018.016
- 926 Wojtal A.Z. 2009. The diatoms of Kobylanka stream near Kraków (Wyżyna Krakowsko–
- 927 Częstochowska upland, S Poland). *Polish Botanical Journal* 54: 129–330.
- 928 Wojtal A.Z. 2013. Species composition and distribution of diatom assemblages in spring water
- 929 from various geological formation in Southern Poland. In: Lange-Bertalot H. &
- 930 Kociolek P., eds. *Bibliotheca Diatomologica Vol. 59*. Stuttgart: J. CRAMER in der
- 931 Gebr. Borbtraeger Verlagsbuchhandlung. 436 p.
- 932 Wojtal A.Z. & Kwadrans J. 2006. Diatoms of the Wyżyna Krakowsko-Częstochowska Upland
- 933 (S. Poland) – Coscinodiscophyceae (Thalassiositophycidae). *Polish Botanical Journal*
- 934 51: 177–207.

- Wojtal A.Z. & Sobczyk Ł. 2012. The influence of substrates and physicochemical factors on the composition of diatom assemblages in karst springs and their applicability in water-quality assessment. *Hydrobiologia* 695: 97–108 DOI 10.1007/s10750-012-1203-0.
- Van Dam H., Mertens A. & Sinkeldam, J. 1994. A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology* 28: 117–133.
- Van de Vijver B., Frenot Y. & Beyens L. 2002. Freshwater diatoms from Ile de la Possession. In: Lange-Bertalot H. & Kociolek P., eds. *Bibliotheca Diatomologica Vol. 46*. Stuttgart: J. CRAMER in der Gebr. Borbtraeger Verlagsbuchhandlung 71 p.
- Van de Vijver B., Wetzel C.E. & Ector L. 2018. Analysis of the type material of *Planothidium delicatulum* (Bacillariophyta) with the description of two new *Planothidium* species from the sub-Antarctic Region. *Fottea* 18(2): 200-211 DOI: 10.5507/fot.2018.006.
- Van Landingham S.L. 1968. Investigation of a diatom population from mine Tailings in Nye Country, Nevada, U.S.A. *Journal of Phycology* 4: 306–310.
- Yang X-D., Dong X-H., Gao G., Pan H-X. & Wu J-L. 2005. Relationship Between Surface Sediment Diatoms and Summer Water Quality in Shallow Lakes of the Middle and Lower Reaches of the Yangtze River. *Journal of Integrative Plant Biology* 4792: 153-164 <https://doi.org/10.1111/j.1744-7909.2005.00035.x>.
- Yamamoto M., Chiba T. & Tuji A. 2017. Salinity responses of benthic diatoms inhabiting tidal flats. *Diatom Research* 32: 243-250 DOI: 10.1080/0269249X.2017.1366951
- Żelazna-Wieczorek J. 1996. Studium taksonomiczne rodzaju *Vaucheria* De Candolle oraz uwagi o występowaniu zidentyfikowanych gatunków tego rodzaju. Unpublished doctoral dissertation, University of Lodz, Poland.
- Żelazna-Wieczorek J. 2002. *Vaucheria* species in selected regions in Poland. *Acta Societatis Botanicorum Poloniae* 71(2): 129-139.
- Żelazna-Wieczorek, J. 2011. Diatom flora in springs of Łódź Hills (Central Poland). In: Witkowski A., ed. *Diatom monographs. Vol. 13*. Ruggell, Liechtenstein: A.R.G. Gantner Verlag K.G. 419 pp.
- Żelazna-Wieczorek J. & Nowicka-Krawczyk P. 2015. The cascade construction of artificial ponds as a tool for urban stream restoration – The use of benthic diatoms to assess the effects of restoration practices. *Science of the Total Environment* 538: 591–599 DOI:10.1016/j.scitotenv.2015.08.110.
- Żelazna-Wieczorek J. & Olszyński R.M. 2016. Taxonomic revision of *Chamaepinnularia krookiformis* Lange-Bertalot et Krammer with a description of *Chamaepinnularia plinskii* sp. nov. *Fottea, Olomouc* 16(1): 112-121 DOI: 10.5507/fot.2016.001.
- Żelazna-Wieczorek J., Olszyński R.M. & Nowicka-Krawczyk P. 2015. Half a century of research on diatoms in athalassic habitats in central Poland. *Oceanological and Hydrobiological Studies* 441: 51–67 DOI: 10.1515/ohs-2015-0006.
- Żurek R., Diakiv V., Szarek-Gwiazda E., Kosiba J. & Wojtal A.Z. 2018. Unique Pit Lake Created in an Opencast Potassium Salt Mine (Dombrowska Pit Lake in Kalush,

975 Ukraine). *Mine Water and the Environment* 37(3): 456-469  
976 <https://doi.org/10.1007/s10230-018-0527-z>

# Figure 1


Fig. 1. Location of sampling points in the Łódzkie and Wielkopolskie voivodships. Poland. A. Pełczyska (PE). B. Łęczyca (LE). C. Bogdałów (BO).



## Figure 2

Fig. 2. Percentage share of diatoms in individual classes of prevalence according to the Tümping & Friedrich factor (1999).


PE – Pełczyska complex, LE – Łeczyca reservoir, BO – Bogdałów reservoir. EC – euconstant taxa, CN – constant taxa, AC – accessory taxa AD – accidental taxa. The number of species is shown in brackets.





# Figure 3


Fig. 3. nMDS 3D analysis. The diagram shows three distinct clouds of samples which coincide with the three hydrological objects.



# Figure 4

Fig. 4. Shade Plot analysis. The diagram shows the strength of the factor (taxon) affecting the similarity between the samples.


Upper dendrogram - samples divided according to hydrological object. Left dendrogram - 50 taxa of diatoms which have the strongest influence on the similarity between the samples.




## Figure 5(on next page)


Fig. 5. Average percentages of diatom species in individual classes of ecological values according to Van Dam et al. (1994) based on OMNIDIA 6.0.6 software.


A. pH requirements. B. Salinity. C. Nitrogen uptake. D. Oxygen requirements. E. Saprobity. F. Trophic state. G. Moisture aerophily.























# Figure 6

Fig. 6. Principal Component Analysis. The diagram presents three groups of species A, B and C, whose occurrence can be correlated with selected physical and chemical factors.




# Figure 7

Figs 7. LM microphotographs of characteristic species for Pełczyska.

A1-A6. *Planothidium delicatulum*. B1-B5. *Chamaepinnularia krookiformis*. C1-C5.

*Chamaepinnularia plinskii*. D1-D4. *Nitzschia liebethruthii*. E1-E5. *Craticula buderi*. F1-F5.


*Navicula cincta*. G1-G5. *Planothidium frequentissimum*. Scale bar = 10  $\mu\text{m}$ .



# Figure 8

Figs 8. LM microphotographs of characteristic species for Łeczyca. A1-A6. *Cyclostephanos invisitatus*. B1-B6. *Navicula moskalii*. C1-C5. *Stephanodiscus binatus*. D1-D5. *S. parvus*. Scale bar = 10  $\mu\text{m}$ .






# Figure 9

Figs 9. LM microphotographs of characteristic species for Bogdałów.

A1-A6. *Diatoma moniliformis*. B1-B6. *Encyonopsis subminuta*. C1-C7. *Mastogloia smithii* (C5-C7. Same specimen, different focal plane). D1-D5. *Nitzschia dissipata* var. *media*. E1-E7. *Pantocsekiella pseudocomensis*.



# **Table 1** (on next page)

Table 1. Classes of ecological indicators values by Van Dam et al. (1994).

		<b>Class (<i>Van Dam et al. 1994</i>)</b>							
		0	1	2	3	4	5	6	7
<b>Ecological parameters</b>	pH requirements	unknown	acidobiontic	acidophilic	neutrophilic	alkaliphilic	alkalibiontic	indifferent	
	Salinity	unknown	halophobe	oligohalobous	halophilic	mesohalobous	brackish-marine	marine-brackish	marine
	Nitrogen utake	unknown	sensitive N-autotrophic	tolerant N-autotrophic	facultative N-autotrophic	obligatory N-autotrophic			
	O <sub>2</sub> requirements	unknown	polyoxybiontic	oxybiontic	moderate	low O <sub>2</sub>	very low O <sub>2</sub>		
	Saprobity	unknown	oligosaprobe	β-mesosaprobe	a-mesosaprobe	a-meso -> polysaprobe	polysaprobe		
	Trophy state	unknown	oligotrophic	oligo-mesotrophic	mesotrophic	meso-eutrophic	eutrophic	hypereutrophic	indifferent
	Moisture aerophily	unknown	aquatic	occasionally aerophilic	aquatic to aerophilic	aerophilic	terrestrial		



## Table 2 (on next page)

Table 2. Physical and chemical parameters in the examined sampling sites. The minimum, maximum and mean values.

	Pelczyska (PE)									Leczyca (LE)									Bogdałów (BO)		
	outflow (D.PESB)			ditch (D.PEDB)			pond (D.PEPB)			L1 (D.LEP1)			L2 (D.LEP2)			L3 (D.LEP3)			reservoir (D.BOZB)		
	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.
pH	6.6	7.7	7.0	6.4	6.4	6.4	7.8	9.6	8.7	7.9	8.6	8.2	7.5	8.6	8.2	7.5	8.8	8.3	7.9	8.3	8.1
Conductivity [μS cm <sup>-1</sup> ]	4450	9230	6699	5170	5170	5170	2645	5150	3646	657	865	743	558	836	728	472	778	680	505	734	623
T [°C]	6.8	13.9	9.1	7.8	7.8	7.8	1.1	17.3	7.1	1.6	20.6	11.1	0.6	21.1	11.1	2.2	15.2	9.2	4.7	22.7	11.4
HCO <sub>3</sub> <sup>-</sup> [mg L <sup>-1</sup> ]	345	744	475	610	610	610	284	451	352	211	339	260	168	275	210	183	290	241	174	369	270
CO <sub>2</sub> [HCO <sub>3</sub> <sup>-</sup> ] [mg L <sup>-1</sup> ]	124	268	171	220	220	220	102	163	126	76	122	94	61	99	75	66	105	87	63	133	96
Cl <sup>-</sup> [mg L <sup>-1</sup> ]	1585	2976	2426	1006	1006	1006	685	1524	1053	57	93	78	60	93	76	67	106	78	41	52	45
N <sub>NH4</sub> [mg L <sup>-1</sup> ]	0.02	1.63	0.45	0.15	0.15	0.15	0.00	0.36	0.11	0.02	0.71	0.22	0.03	0.53	0.26	0.00	1.16	0.32	0.00	0.07	0.02
NH <sub>4</sub> <sup>+</sup> [mg L <sup>-1</sup> ]	0.03	2.09	0.71	0.19	0.19	0.19	0.00	0.46	0.14	0.03	0.55	0.22	0.03	0.54	0.28	0.00	0.90	0.31	0.00	0.09	0.03
PO <sub>4</sub> <sup>3-</sup> [mg L <sup>-1</sup> ]	0.60	12.46	3.73	8.57	8.57	8.57	0.85	9.10	3.99	0.22	0.53	0.36	0.05	0.57	0.41	0.09	0.57	0.34	0.27	0.45	0.34
P <sub>PO4</sub> [mg L <sup>-1</sup> ]	0.20	4.11	1.23	2.83	2.83	2.83	0.28	3.00	1.32	0.07	0.17	0.12	0.02	0.19	0.13	0.03	0.19	0.11	0.09	0.15	0.12
SO <sub>4</sub> <sup>2-</sup> [mg L <sup>-1</sup> ]	176	198	188	165	165	165	151	197	176	71	147	107	67	122	101	63	91	75	110	147	123
S <sub>SO4</sub> [mg L <sup>-1</sup> ]	58.8	66.2	62.9	55.1	55.1	55.1	50.3	65.7	58.7	23.5	49.0	35.8	22.3	40.8	33.8	21.0	30.3	25.1	37.9	49.2	41.5
COLOR [mgPt dm <sup>-3</sup> ]	25	160	81	140	140	140	50	120	78	10	60	27	9	60	27	12	60	31	4	10	6
Mn <sup>3+</sup> [mg L <sup>-1</sup> ]	0.17	0.51	0.36	0.14	0.14	0.14	0.03	0.25	0.13	0.00	0.05	0.02	0.01	0.03	0.02	0.01	0.07	0.03	0.00	0.01	0.00
Fe <sup>2+/3+</sup> [mg L <sup>-1</sup> ]	0.25	0.41	0.32	0.25	0.25	0.25	0.05	0.24	0.11	0.03	0.27	0.09	0.00	0.13	0.04	0.01	0.07	0.03	0.01	0.09	0.02
Mg <sup>2+</sup> [mg L <sup>-1</sup> ]	39.5	48.7	45.0	35.9	35.9	35.9	25.6	37.0	32.1	11.6	19.1	15.2	10.3	18.3	15.1	9.5	15.3	12.5	9.6	15.7	11.7
Ca <sup>2+</sup> [mg L <sup>-1</sup> ]	171.7	216.2	195.2	165.0	165.0	165.0	75.8	139.8	121.1	58.0	143.6	89.0	59.5	117.6	79.1	52.6	86.4	65.5	77.7	117.6	86.3
Na <sup>+</sup> [mg L <sup>-1</sup> ]	500.7	1537.4	1227.1	453.3	453.3	453.3	277.3	681.8	455.9	15.6	40.4	30.9	20.3	42.4	28.3	22.0	42.4	33.0	24.0	43.1	32.6
K <sup>+</sup> [mg L <sup>-1</sup> ]	8.5	124.8	42.4	109.8	109.8	109.8	58.6	68.8	63.5	4.5	9.5	7.2	5.1	9.3	7.3	5.6	10.7	8.2	0.1	3.6	1.5

### Table 3 (on next page)

Table 3. Characteristic species according to SIMPER analysis for studied waterbodies. EC - euconstant taxa, CN - constant taxa, CN - accessory taxa (*Tümping & Friedrich 1999*).

Pelczyska pond D.PEPB	Pelczyska dich D.PEDB	Pelczyska outflow D.PESB	Łęczyska reservoirs LE	Bogdałów reservoir BO
<i>Cocconeis placentula</i> (EC)	<i>Chamaepinnularia krookiformis</i> (CN)	<i>Fragilaria famelica</i> (EC)	<i>Achnantheidium minutissimum</i> (EC)	<i>Achnantheidium minutissimum</i> (EC)
<i>Craticula buderi</i> (EC)	<i>Chamaepinnularia plinskii</i> (EC)	<i>Navicula veneta</i> (EC)	<i>Amphora pediculus</i> (EC)	<i>Diatoma moniliformis</i> (EC)
<i>Fragilaria sopotensis</i> (EC)	<i>Craticula halophila</i> (EC)	<i>Nitzschia frustulum</i> (EC)	<i>Cyclostephanos dubius</i> (EC)	<i>Encyonopsis subminuta</i> (EC)
<i>Gomphonema parvulum</i> (EC)	<i>Navicula veneta</i> (EC)	<i>Nitzschia inconspicua</i> (EC)	<i>Cyclostephanos invisitatus</i> (EC)	<i>Mastogloia smithii</i> (EC)
<i>Hippodonta hungarica</i> (EC)	<i>Nitzschia frustulum</i> (EC)	<i>Planothidium delicatulum</i> (EC)	<i>Cyclotella meneghiniana</i> (EC)	<i>Nitzschia dissipata</i> var. <i>media</i> (EC)
<i>Navicula cincta</i> (EC)	<i>Nitzschia inconspicua</i> (EC)	<i>Tabularia fasciculata</i> (EC)	<i>Navicula gregaria</i> (EC)	<i>Pantocsekiella ocellata</i> (EC)
<i>Nitzschia palea</i> (EC)	<i>Nitzschia liebethruthii</i> (EC)		<i>Navicula moskalii</i> (AC)	<i>Pantocsekiella pseudocomensis</i> (EC)
<i>Planothidium frequentissimum</i> (EC)	<i>Nitzschia perminuta</i> (EC)		<i>Nitzschia palea</i> (EC)	
	<i>Nitzschia tubicola</i> (EC)		<i>Stephanodiscus binatus</i> (EC)	
			<i>Stephanodiscus hantzschii</i> (EC)	
			<i>Stephanodiscus parvus</i> (EC)	

# **Table 4**(on next page)

Table 4. Characteristics species with classification of ecological indicators values by Van Dam et al. (1994).

Species	Code	Moisture acrophity	Nitrogen uptake	pH requirements	Oxygen requirements	Salinity	Saprobity	Trophic state
<i>Achnantheidium minutissimum</i>	ADMI	3	2	3	1	2	2	7
<i>Amphora pediculus</i>	APED	3	2	4	2	2	2	5
<i>Chamaepinnularia krookiformis</i>	CHKF	3	0	3	0	3	1	0
<i>Chamaepinnularia plinskii</i>	CHPL	0	0	0	0	0	0	0
<i>Cocconeis placentula</i>	CPLA	2	2	4	3	2	2	5
<i>Craticula buderi</i>	CRBU	0	0	0	0	0	0	0
<i>Craticula halophila</i>	CHAL	2	2	4	2	4	3	5
<i>Cyclostephanos dubius</i>	CDUB	1	2	5	2	3	3	5
<i>Cyclostephanos invisitatus</i>	CINV	0	0	0	0	2	0	5
<i>Cyclotella meneghiniana</i>	CMEN	2	3	4	5	3	4	5
<i>Diatoma moniliformis</i>	DMOF	0	0	0	0	0	0	0
<i>Encyonopsis subminuta</i>	ESUM	0	0	3	1	1	1	1
<i>Fragilaria famelica</i>	FFAM	3	1	4	1	2	1	3
<i>Fragilaria sopotensis</i>	FSOP	1	2	4	1	2	2	4
<i>Gomphonema parvulum</i>	GPAR	3	3	3	4	2	4	5
<i>Hippodonta hungarica</i>	HHUN	3	2	4	3	2	2	4
<i>Mastogloia smithii</i>	MSMI	3	0	4	0	4	2	0
<i>Navicula cincta</i>	NCCA	0	0	0	0	2	0	7
<i>Navicula gregaria</i>	NGRE	3	2	4	4	3	3	5
<i>Navicula moskalii</i>	NMOK	0	0	0	0	0	0	0
<i>Navicula veneta</i>	NVEN	3	2	4	4	3	4	5
<i>Nitzschia dissipata</i> var. <i>media</i>	NDME	0	0	4	0	2	0	0
<i>Nitzschia frustulum</i>	NIFR	3	4	4	3	3	2	5
<i>Nitzschia inconspicua</i>	NINC	3	3	4	3	3	3	5
<i>Nitzschia liebethruthii</i>	NLBT	0	0	5	0	4	0	0
<i>Nitzschia palea</i>	NPAL	3	4	3	4	2	5	6
<i>Nitzschia perminuta</i>	NIPM	3	1	4	1	2	1	2
<i>Nitzschia tubicola</i>	NTUB	2	3	4	4	3	5	6
<i>Pantocsekiella ocellata</i>	POCL	1	1	4	1	1	1	4
<i>Pantocsekiella pseudocomensis</i>	PPCS	0	0	0	0	0	0	0
<i>Planothidium delicatulum</i>	PTDE	3	1	5	0	4	5	3
<i>Planothidium frequentissimum</i>	PLFQ	0	2	4	3	2	4	7
<i>Stephanodiscus binatus</i>	SBNT	0	0	0	0	0	0	0
<i>Stephanodiscus hantzschii</i>	SHAN	2	3	5	4	2	4	6
<i>Stephanodiscus parvus</i>	SPAV	0	0	5	0	2	0	6
<i>Tabularia fasciculata</i>	TFAS	3	2	4	3	4	3	5



## Table 5 (on next page)

Table 5. Percentage of characteristic species in the unknown (total) class according to Van Dam et al. (1994) at each sampling point.

Percentage of individual characteristic species constituting at least 10% of the 'unknown' class at each sampling point.

	pH	Salinity	Nitrogen uptake	Oxygen requirements	Saprobity	Trophic state	Moisture aerophily
	total 33%	total 44%	total 34%	total 48%	total 30%	total 22%	total 43%
<b>D.PESB.</b>	CHPL (19%) NCCA (11%)	CHPL (33%) CHKF (10%)	CHPL (11%) NCCA (10%)	PTDE (25%)	CHPL (11%)	CHPL (10%)	PLFQ (20%)
	total 92%	total 25%	total 82%	total 82%	total 80%	total 74%	total 85%
<b>D.PEDB.</b>	CHPL (60%) NCCA (32%)	CHPL (25%)	NLBT (37%) CHPL (25%) CHKF (16%)	NLBT (37%) CHPL (16%) CHKF (16%)	NLBT (43%) CHPL (32%)	NLBT (365) CHPL (23%) CHKF (15%)	NLBT (49%) CHPL (285)
	total 81%	total 73%	total 78%	total 78%	total 79%	total 60%	total 84%
<b>D.PEPB.</b>	CRBU (53%) NCCA (27%)	CRBU (73%)	CRBU (46%) NCCA (25%)	CRBU (46%) NCCA (25%)	CRBU (46%) NCCA (26%)	CRBU (54%)	CRBU (41%) NCCA (23%) PLFQ (14%)
	total 76%	total 79%	total 67%	total 68%	total 71%	total 54%	total 78%
<b>D.LEP1.</b>	CINV (36%) SBNT (23%) NMOK (16%)	SBNT (56%) NMOK (21%)	CINV (29%) SBNT (19%) NMOK (14%)	CINV (29%) SBNT (19%) NMOK (14%)	CINV (31%) SBNT (20%) NMOK (14%)	SBNT (36%) NMOK (17%)	CINV (32%) SBNT (22%) NMOK (14%)
	total 39%	total 44%	total 35%	total 35%	total 28%	total 23%	total 44%
<b>D.LEP2.</b>	SBNT (27%) CINV (11%)	SBNT (44%)	SBNT (18%)	SBNT (18%)	SBNT (20%)	SBNT (23%)	SBNT (22%) SPAV (11%) CINV (11%)
	total 57%	total 41%	total 50%	total 53%	total 60%	total 29%	total 57%
<b>D.LEP3.</b>	SBNT (25%) CINV (24%)	SBNT (41%)	SBNT (17%) CINV (15%) SPAV (13%)	SBNT (17%) CINV (16%) SPAV (14%)	SBNT (20%) CINV (18%) SPAV (15%)	SBNT (29%)	SBNT (20%) CINV (17%) SPAV (15%)
	total 33%	total 36%	total 47%	total 44%	total 38%	total 47%	total 38%
<b>D.BOZB.</b>	PPCS (22%) DMOF (10%)	PPCS (25%) DMOF (11%)	MSMI (13%) PPCS (10%)	MSMI (14%) PPCS (11%) NDME (10%)	PPCS (18%) NDME (11%)	MSMI (15%) PPCS (12%) NDME (11%)	PPCS (13%)

## Table 6 (on next page)

Table 6. Selected 16 characteristic species with the new or altered (in red) ecological indicator values according to Van Dam et al. (1994).

Species	Code	Moisture acrophity	Nitrogen uptake	pH requirements	Oxygen requirements	Salinity	Saprobity	Trophic state
<i>Chamaepinnularia krookiformis</i>	CHKF	3 4	0	3	0	3 4	1 4	0 5
<i>Chamaepinnularia plinskii</i>	CHPL	0 4	0	0 3	0	0 4	0 4	0 5
<i>Craticula buderi</i>	CRBU	0 3	0	0 4	0	0 4	0 4	0 5
<i>Cyclostephanos invisitatus</i>	CINV	0 1	0	0 4	0	2	0	5
<i>Diatoma moniliformis</i>	DMOF	0	0	0	0	0	0	0
<i>Encyonopsis subminuta</i>	ESUM	0	0	3	1	1	1	1
<i>Mastogloia smithii</i>	MSMI	3	0	4	0	4 3	2	0
<i>Navicula cincta</i>	NCCA	0 3	0	0	0	2	0	7
<i>Navicula moskalii</i>	NMOK	0	0	0 4	0	0 2	0	0 7
<i>Nitzschia dissipata</i> var. <i>media</i>	NDME	0	0	4	0	2	0 2	0
<i>Nitzschia liebethruthii</i>	NLBT	0 4	0	5 3	0	4	0 4	0 5
<i>Pantocsekiella pseudocomensis</i>	PPCS	0	0	0	0	0	0	0
<i>Planothidium delicatulum</i>	PTDE	3	1	5 3	0	4 5	5	3
<i>Planothidium frequentissimum</i>	PLFQ	0	2	4	3	2	4	7
<i>Stephanodiscus binatus</i>	SBNT	0	0	0 4	0	0 2	0	0
<i>Stephanodiscus parvus</i>	SPAV	0	0	5 4	0	2	0	6