

A peer-reviewed version of this preprint was published in PeerJ on 5 December 2019.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.8117) (peerj.com/articles/8117), which is the preferred citable publication unless you specifically need to cite this preprint.

Olszyński RM, Szczepocka E, Żelazna-Wieczorek J. 2019. Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin. PeerJ 7:e8117 <https://doi.org/10.7717/peerj.8117>

Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin

Rafał M. Olszyński^{Corresp., Equal first author, 1}, Ewelina Szczepocka¹, Joanna Żelazna-Wieczorek^{Corresp. Equal first author, 1}

¹ Department of Algology and Mycology, Faculty of Biology and Environmental Protection, University of Lodz, Łódź, Poland

Corresponding Authors: Rafał M. Olszyński, Joanna Żelazna-Wieczorek
Email address: rafal.olszynski@biol.uni.lodz.pl, joanna.zelazna@biol.uni.lodz.pl

Background. The ecological state of surface waters is typically assessed by a multi-aspect approach based on a determination of its chemical and physical parameters, by hydromorphology and the use of indicator organisms such as benthic diatoms. By assigning ecological indicator values, it is possible to create diatom indices which serve as the basic tool in assessing the ecological status of surface waters. These ecological indicator values are set according to classification systems, such as the Van Dam et al. 1994 system, which classifies species of diatoms according to seven different ecological factors. However, recent studies on the autecology of diatoms have shown the need to verify and establish new ecological indicator values. To this end, aquatic ecosystems are good environments to observe the range of tolerance of benthic diatoms to environmental conditions due to their unique physical and chemical parameters. The aim of the present study was to propose the establishment of new, or altered, ecological indicator values, according to Van Dam et al. classification, of species of diatoms characteristic of three post-mining aquatic ecosystems. **Methods.** In total, 36 species were identified that were characteristic of three waterbodies: a salt aquatic complex (water outflow, a drainage ditch and a pond), a mined iron ore reservoir and a mined lignite reservoir. Their ecological indicator values were calculated using OMNIDIA software, and the environmental conditions prevailing in the studied ecosystems were determined. Of the 36 characteristic species, 16 lacking at least one assigned ecological indicator value were analysed further. The analysis identified three groups of selected characteristic species which showed a correlation, or lack of such, to the tested physical and chemical parameters. **Results.** Based on this multistage study of the autecology of characteristic diatoms, comprising an analysis of environmental conditions, literature analysis and reference ecological indicator values of other species, it is proposed that 32 ecological indicator values be established or adjusted for 16 species, and that *Planothidium frequentissimum* be excluded from water quality assessments.

Critical multi-stranded approach for determining the ecological values of diatoms in unique aquatic ecosystems of anthropogenic origin

Rafał M. Olszyński, Ewelina Szczepocka, Joanna Żelazna-Wieczorek

Department of Algology and Mycology, Faculty of Biology and Environmental Protection,
University of Lodz, Łódź, Poland

Corresponding Authors:

Rafał M. Olszyński

12/16 Banacha St, Łódź, 90-237, Poland

Email address: rafal.olszynski@biol.uni.lodz.pl

Joanna Żelazna-Wieczorek

12/16 Banacha St, Łódź, 90-237, Poland

Email address: joanna.zelazna@biol.uni.lodz.pl

Abstract

Background. The ecological state of surface waters is typically assessed by a multi-aspect approach based on a determination of its chemical and physical parameters, by hydromorphology and the use of indicator organisms such as benthic diatoms. By assigning ecological indicator values, it is possible to create diatom indices which serve as the basic tool in assessing the ecological status of surface waters. These ecological indicator values are set according to classification systems, such as the Van Dam et al. 1994 system, which classifies species of diatoms according to seven different ecological factors. However, recent studies on the autecology of diatoms have shown the need to verify and establish new ecological indicator values. To this end, aquatic ecosystems are good environments to observe the range of tolerance of benthic diatoms to environmental conditions due to their unique physical and chemical parameters. The aim of the present study was to propose the establishment of new, or altered, ecological indicator values, according to Van Dam et al. classification, of species of diatoms characteristic of three post-mining aquatic ecosystems.

Methods. In total, 36 species were identified that were characteristic of three waterbodies: a salt aquatic complex (water outflow, a drainage ditch and a pond), a mined iron ore reservoir and a mined lignite reservoir. Their ecological indicator values were calculated using OMNIDIA software, and the environmental conditions prevailing in the studied ecosystems were determined. Of the 36 characteristic species, 16 lacking at least one assigned ecological indicator value were analysed further. The analysis identified three groups of selected characteristic species which showed a correlation, or lack of such, to the tested physical and chemical parameters.

Results. Based on this multistage study of the autecology of characteristic diatoms, comprising an analysis of environmental conditions, literature analysis and reference ecological indicator values of other species, it is proposed that 32 ecological indicator values be established or adjusted for 16 species, and that *Planothidium frequentissimum* be excluded from water quality assessments.

Introduction

Diatoms (Bacillariophyta) are one of the main biotic elements used in the biological assessment of the ecological state of surface waters (*Water Framework Directive, European Union 2000*). Due to the fact that many countries are obliged to continually engage in biomonitoring, there is a clear need to develop flawlessly functioning methods based on the standardised use of diatoms as bioindicators (*Kahlert et al. 2016; Poikane, Kelly & Cantonati 2016; Szczepocka & Żelazna-Wieczorek 2018*). Diatom indices and ecological systems based on the bioindication values of particular diatom species, derived from various environmental parameters, constitute a fundamental tool in the biological assessment of environments. Diatom indices have been commonly used to assess flowing and standing water for over 20 years (*Kelly et al. 2008; Harding & Taylor 2014; Szczepocka et al. 2014; Hutorowicz & Pasztaleniec 2014; Holms & Taylor 2015; Żelazna-Wieczorek & Nowicka-Krawczyk 2015; Kolada et al. 2016*).

Currently, many countries use the OMNIDIA program (*Lecointe et al. 1993*) as a biological assessment tool. Its latest version (version 6.0.6) allows the calculation of 18 diatom indices, and the determination of seven environmental parameters for eight ecological systems. However, the specific ecological indicator values of many of the species given in the OMNIDIA database are absent or have not been updated in response to recent research. To complete these missing values, and to verify existing ones, further studies are needed of the ecological optima and tolerance of diatom species in different types of aquatic ecosystems.

Due to their specific environmental conditions, post-mining reservoirs represent an extremely valuable source of information for the study of ecological diatom tolerance ranges. Some studies of these environments have been performed, but these have addressed diatom paleoecology and their role as indicators of past climatic or environmental change (*de Haan et al. 1993; Rakowska 1996; Thomas & John 2006; Sienkiewicz & Gqsiorowski 2016*). Until now, the autecology of diatoms in post-mining reservoirs has rarely been studied (*Van Landingham 1968; de Haan et al. 1993; Rakowska 1996; Ferreira da Silva et al. 2009; Luis et al. 2009; Luis et al. 2016; Sienkiewicz & Gqsiorowski 2016*).

The present study examines the diatom assemblages present in three post-mining reservoirs of various geological origins. Due to variations in their environmental parameters, these bodies of water serve as specific and unique habitats for the development of these algae. The diatom assemblages quickly adapt to the currently prevailing conditions, which is manifested in the presence of taxa characteristic of these specific parameters. Considering their large share of the assemblage, the index values of the assemblages constitute the most important component in the calculation of diatom indices. These species are therefore of the greatest importance for surface water biomonitoring.

The aim of the present study was to identify the species of diatoms characteristic of the three studied types of post-mine reservoirs. Following this, taxa that did not have at least one ecological indicator value specified in the OMNIDIA database, according to the environmental parameters given by Van Dam et al. (1994), were identified. New ecological indicator values were proposed based on the relationship between the occurrence of the individual species and certain selected physical and chemical parameters, or existing ones were verified.

The Van Dam et al. (1994) ecological system is one of the main systems on which the OMNIDIA programme is based. It describes the ecological indicator values of diatoms according to pH, salinity, nitrogen uptake metabolism, oxygen requirement, saprobity, trophic state and moisture aerophily. These values play a key role in calculating diatom indices, and hence need to be kept up to date to enable accurate routine biomonitoring.

Materials & Methods

Study area

The study was performed on three waterbodies created through exploration for mineral deposits or were formed by the closure of mines. All three are located in the Łódzkie and Wielkopolskie voivodeships, Central Poland.

The first complex of waterbodies - Pełczyska (PE), is situated in the village of Pełczyska, between Łódź and Łęczyca (Łódzkie voivodeship) (Fig. 1). As the local area is characterized by the presence of salt deposits, numerous wells were sunk in the eighteenth century to obtain brine. Currently, salt water flows out of one of them. This area has been studied by biologists and hydrobiologists since the 1960s (*Olaczek 1963; Pliński 1966; Pliński 1969; Pliński 1971a; Pliński 1971b; Pliński 1971c; Pliński 1973; Żelazna-Wieczorek 1996; Żelazna-Wieczorek 2002; Żelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015; Żelazna-Wieczorek & Olszyński 2016*). The waterbodies chosen for our research form the PE hydrological complex located in the vicinity of farmland; it comprises the salt water outflow, a drainage ditch and a pond, which acts as the receiver of the water.

The second complex of waterbodies - Łęczyca (LE), urban reservoir located within the city of Łęczyca (Łódzkie voivodeship) (Fig. 1). The reservoirs were created following the flooding the open-cast iron ore mine in the 1990s. This area is rich in syderite deposits, which are accompanied by other minerals. The complex consists of three connected reservoirs: two are directly connected to each other (LEP1 and LEP2), and the third (LEP3) is connected to LEP2 via a water drainage ditch (*Olszyński & Żelazna-Wieczorek 2018*). All three are located in an area with houses, garden plots and partly-wooded areas.

The third waterbody - Bogdałów reservoir (BO), created by the flooding of an opencast brown coal mine. It is located in the village of Bogdałów (Wielkopolskie voivodeship) in an area rich in lignite deposits (Fig. 1). Lignite from quaternary deposits was exploited since 1977 until 1991 to a depth of 50 meters. Due to the specific construction of the open-pit area, being characterised by the thickest layer of poorly permeable boulder clay in the region. This pit was later transformed into a storage site for quarried rocks in Koźmin. Finally, in 1993/1994, the drainage and runoff of surface waters were blocked to form a reservoir with a depth of about 12 meters surrounded by forest (*Gabryś-Godlewska et al. 2004; Gadomska et al. 2007; Orlikowski & Szwed 2009; Kasztelewicz 2011*).

Samples

Samples of benthic diatoms from sediments and water samples were collected quarterly (once in any season) from each hydrological waterbodies. Analysis of all water samples (Ca^{2+} , Mg^{2+} , Na^{+} , K^{+} , $\text{Fe}^{2+/3+}$, Mn^{3+}) involved flame absorption spectrophotometry SpectrAA 300 (Varian, Palo Alto, CA, USA) (detection limit is 0.05 mg/L) and UV-vis spectrophotometry S.330 (Marcel, Poland) in the case of NH_4^{+} (d.l. is 0.001 mg/L) and PO_4^{3-} (d.l. is 0.01 mg/L). SO_4^{2-} was determined by the gravimetric method (PN-C-04566-09), Cl^{-} by Mohr's method (PN-ISO 9297). The chemical analyses were performed in the Laboratory of the Department of Geology at the Faculty of Geographical Sciences, University of Łódź and the Laboratory of Computer and Analytical Techniques at the Faculty of Biology and Environmental Protection, University of

Łódź.. The water temperature, pH and electric conductivity were measured *in situ* (Elmetron CP-401 and CC-401 devices). The following sampling points were established: Pełczyska (51°58'34.47"N, 19°14'21.11"E) – outflow (P.ESB.) (4 samples, both water and benthic), ditch (D.PEDB.) (1 water and 4 benthic samples) and pond (D.PEPB.) (4 samples, both water and benthic); samples were collected quarterly from July 2013 to March 2014; Łęczyca (52°3'5.30"N; 19°11'50.24"E) – reservoir 1 (D.LEP1.), reservoir 2 (D.LEP2.) and reservoir 3 (D.LEP3.), samples were collected quarterly from March 2014 to December 2015 (6 water and 8 benthic samples from each reservoirs); Bogdałów (52°2'51.29"N; 18°35'51.49"E) – reservoir (D.BOZB.), samples were collected quarterly from March 2015 to December 2016 (8 samples, both water and benthic). The water samples were subjected to chemical and physical analysis at the Laboratory of Geology at the Faculty of Geographical Sciences, University of Lodz. In total, 44 benthic samples were collected. The permanent slides were prepared according to Żelazna-Wieczorek (2011). To obtain pure diatom frustule the material was chemically treated using a H₂SO₄ and H₂Cr₂O₇. The cleaned diatom precipitate was mounted on permanent slides using Naphrax[®] synthetic resin. Qualitative and quantitative analysis of diatoms was performed using a Nikon Eclipse 50i light microscope (LM) under 1000× magnification (plan oil-immersion objective 100×/1.25): the diatoms were identified and counted for up to 500 valves in each permanent slide. Light photomicrographs were taken with an OPTA-TECH digital camera. When diatoms were difficult to identify using LM they were subjected to scanning electron microscope (SEM) analysis using a Phenom Pro X (gold layer of 8 and 20 nm, at 10 kV, low vacuum mode) at the Laboratory of Microscopy Imaging and Specialist Biological Techniques, Faculty of Biology and Environmental Protection, University of Lodz.

Data processing & statistical analysis

The average percentage (AP) for a given species was determined based on the percentage contribution (%) of the species in the samples tested for a given hydrological object (Żelazna-Wieczorek 2011). Species with AP greater than or equal to 5% for each hydrological object were identified as dominant. The incidence was determined according to the Tümping & Friedrich (1999) coefficient according to the range values: 100%–75% euconstant taxa (EC), 75%–50% constant taxa (CN), 50%–25% accessory taxa (AC) and 25%–1% accidental taxa (AD) (Tümping & Friedrich 1999). Multidimensional scaling analysis (nMDS) based on Bray-Curtis similarity coefficients was used to identify natural groupings of samples. The results are given as a 3D diagram in which the degree of similarity is represented as the distances between particular points (samples), with greater distances indicating a lower degree of similarity. The reliability of the ordering of the assemblage is represented by the stress value, which reflects how well the ordination summarizes

the observed distances among the samples. A 3-dimensional presentation, whose stress value is lower, is likely to be more satisfactory than a 2-dimensional one (Clarke & Gorley 2015).

The Shade Plot analysis, based on the Bray-Curtis similarity coefficient, was used to identify the diatom species that have the strongest influence on the similarities between the samples demonstrated in the nMDS analysis. Shade Plot analysis compares two data matrices with each other and then groups them on two levels, according to the similarity of the samples and the factors affecting their similarity, i.e. diatom species. The results are represented graphically by shading individual cells: the intensity of the shading indicates the degree of the influence of a given factor (species) on the position of its sample within a given similarity cluster. The range of the shading was determined on the basis of $\log(x+1)$ (x —number of valves).

The SIMPER analysis was used to determine the characteristic species distinguishing the studied ecosystems. This method examines the participation of each variable in the overall similarity between groups of samples, thus indicating the species with the greatest influence on the degree of similarity, or dissimilarity, between particular samples and hydrological objects. This analysis is also based on the Bray-Curtis similarity coefficient; however, unlike the nMDS method, in which one trial is compared to all the other samples, the SIMPER analysis compares a single sample to each subsequent sample (Żelazna-Wieczorek 2011). The results indicate the species which most strongly differentiated a sampled site from the others, and to what extent. A species was regarded as being characteristic of the studied ecosystem if it was characterized by a mean dissimilarity greater than or equal to 2 according to the SIMPER analysis, and a higher mean abundance greater in one ecosystem than the other.

In total, 19 physical and chemical parameters of water were measured in the studied ecosystems. The results of the correlation analysis found 15 physical and chemical parameters indicating an environmental conditions. The parameters were subjected to principal component analysis (PCA) to determine which had the strongest effect on the selected species.

Using the information from the OMNIDIA database, the environmental conditions for each sampling point were determined according to Van Dam et al. (1994) (Table S1). Following this, the percentage share of diatom species included in each ecological indicator value class was indicated. For species found to be characteristic of the studied ecosystems, classes of ecological indicator values were assembled. Taxa which had at least one value of 0 (unknown) were selected for further analysis.

The selected ecological indicator values according to Van Dam et al. (1994) were verified, or new ones established, for the species found to be characteristic of the studied ecosystems according to three premises: previous literature reports about ecological indicator values of those species, chemical and physical conditions analysis, and the classification of the environmental conditions according to Van Dam et al. (1994).

The analyses were performed using PRIMER 7.0.13 (nMDS, Shade Plot, SIMPER), OMNIDIA 6.0.6 and STATISTICA 13 (PCA), software.

Results

213 *Chemical analysis of water samples*

214 The mean values and range of all tested parameters are given in Table 2.

215 The PE hydrological complex was characterized by elevated values of electric conductivity,
216 reaching as high as $9230 \mu\text{S cm}^{-1}$. The pH changed with the direction of water outflow: a slightly
217 acidic reaction was observed in the outflow and an alkaline one in the pond. Due to the
218 geological profile of the region, the water flowing out of the well contained a high concentration
219 of chloride ions, whose gradient decreased with the flow of water through the ditch to the pond.
220 In addition, higher concentrations of the cations Mg^{2+} , Ca^{2+} , Na^{+} and K^{+} were observed
221 compared to other ecosystems, as well as the anions HCO_3^{-} , PO_4^{3-} and SO_4^{2-} .

222 The K^{+} concentration is acknowledged parameter coming from agricultural activity, in particular
223 animal husbandry, or municipal wastes (Macioszczyk & Dobrzyński 2002).

224 Each of the sampling points in the PE complex was characterized by different chemical
225 parameters, resulting in differences between the habitats. The highest electrolytic conductivity
226 was noted in the outflow, which was mainly influenced by the concentrations of Cl^{-} , Na^{+} and
227 HCO_3^{-} ions. The maximum concentration of HCO_3^{-} ions was recorded in Pełczyska outflow in
228 March 2014 (D.PESB.250314); in the other locations, it did not exceed 410 mg L^{-1} .

229 Low concentrations of K^{+} ions were observed throughout the entire studied PE complex;
230 however, maximum values were recorded in the locations characterised by the highest HCO_3^{-} ion
231 content. The highest concentration of Ca^{+} ions of all ecosystems was recorded in the outflow.
232 The ditch represented an intermediate section between the PE sampling points. However, as it is
233 susceptible to periodic drying, limited chemical data was collected from this habitat and hence it
234 was not possible to assess its chemical and physical nature.

235 The lowest electrolytic conductivity was found in the pond, which displayed lower
236 concentrations of Cl^{-} , Na^{+} and, to a lesser degree, HCO_3^{-} . The pH of the water never dropped
237 below 8, except in one case in March 2014. In the pond, the concentration of K^{+} remained
238 relatively unchanged, which could be related to the fact that the reservoir was also a receiver of
239 waters flowing from the surrounding arable fields. The pond was also characterized by the
240 lowest concentration of Ca^{2+} and Mg^{2+} . In the summer periods, a significant reduction in the
241 water table level and occasional drying of the reservoir were noted.

242 The urban reservoirs in Łęczyca (LE) were characterized by a slightly alkaline water reaction,
243 which was similar in all reservoirs during the course of the study. No elevated concentrations of
244 $\text{Fe}^{2+/3+}$ and Mn^{3+} ions were observed. The content of SO_4^{2-} anions was not higher than in other
245 waterbodies studied. The concentration of HCO_3^{-} ions was lower than that observed in BO and
246 PE. No significant differences in chemical and physical parameters were observed between the
247 individual sampling points constituting LE.

248 The Bogdałów (BO) reservoir was characterized by an alkaline reaction. Its K^{+} , Cl^{-} and NH_4^{+}
249 ion content was the lowest of the studied ecosystems.

250 *Diatom samples*

A total of 381 diatom taxa were identified in 44 benthic samples: 139 in PE, 192 in LE and 188 in BO. The dominant species in PE were *Navicula veneta*, and *Nitzschia frustulum*, in LE *Cyclostephanos dubius* and *Stephanodiscus hantzschii*, in BO *Achnantheidium minutissimum*, *Pantocsekiella ocellata* and *Mastogloia smithii*. In the examined ecosystems, the most commonly identified classes were accidental (PE-84; LE-111; BO-86), accessory (PE-25; LE-35) and euconstant taxa (BO-39) (Fig. S2).

nMDS analysis

nMDS analysis (stress level = 0.07) identified the variation between samples for each studied hydrological object (Fig. 2). The samples taken from BO constitute a separate cloud, with the samples demonstrating high similarity with each other, whereas the samples of D.LEP1, D.LEP2 and D.LEP3 constitute a distinct group, with no clear differentiation into individual reservoirs. In the case of PE, the pond group (D.PEPB) was found to be clearly distinct from the others.

Shade Plot

Shade Plot analysis identified 50 species which had the strongest influence on the degree of similarity, or non-similarity, between the samples in the studied ecosystems. Of these taxa, the three that most strongly influenced the similarity between the samples in at least two ecosystems were *Navicula veneta* (NVEN), *N. cincta* (NCCA), *N. gregaria* (NGRE), *Nitzschia frustulum* (NIFR), *N. inconspicua* (NINC), *N. palea* (NPAL), *Planothidium frequentissimum* (PLFQ), *Amphora pediculus* (APED), *Cyclotella meneghiniana* (CMEN), *Fragilaria radians* (FRAD) and *Achnantheidium minutissimum* (ADMI) (Fig. 3).

SIMPER analysis

SIMPER analysis allowed 36 species characteristic of the tested hydrological objects to be distinguished (Table 2). In addition, two species were found to be characteristic of two different ecosystems: *Achnantheidium minutissimum* for LE and BO, and *Nitzschia palea* for PE and LE.

Ecological analysis based on Omnidia software

The ecological analysis of diatom assemblages based on data obtained from the OMNIDIA program database, indicated the following:

- pH requirements: while alkaliphilic species predominate in PE (63%), a large percentage in D.PEPB are unknown (24%) or neutrophilic species (23%). The LE reservoirs were dominated by alkalibiontic (45%) and alkaliphilic (24%) organisms. In D.LEP1, 25% of species were unknown. BO was dominated by alkaliphilic (39%) and neutrophilic (29%) species, and 26 % of species were unknown (Fig. S3);
- salinity: the PE complex was characterized by the occurrence of halophilic (43%), oligohalobous (30%) and mesohalobous species (16%); the greatest proportion of the mesohalobous species were found in D.PESB (28%). The LE reservoirs were dominated by

oligohalobous (44%) and halophilic species (42%). BO was dominated by oligohalobous (43%), halophobe (24%) and unknown species (23%) (Fig. S3).

- nitrogen uptake: the most common species in the PE complex N-autotrophic tolerant (39%) followed by unknown (25%). The largest percentage of unknown species (28%) was recorded in D.PEPB and D.PEDB. In the LE reservoirs, the most common groups of species were N-autotrophic (57%) and unknown (25%). In BO, 51% species were unknown, 24% were N-autotrophic tolerant and 22% N-autotrophic sensitive (Fig. S3);

- oxygen requirements: in PE, the largest groups of species were low oxygen (30%), unknown (27%) and moderate oxygen (24%). In LE, oxybiontic species were most common (43%) followed by unknown (25%). In BO, unknown (46%) and polyoxybiontic species (42%) predominated (Fig. S3);

- sensitivity to saprobity: in PE, the largest group of taxa were α -meso: polysabrobe (28%) and unknown (23%). In D.PEPB, the most abundant was α -meso-polysabrobe (34%) followed by β -mesosaprobe (31%) and unknown (27%). LE primarily included taxa from the α -mesosaprobe group (47%) and unknown (23%). In BO, unknown (34%), β -mesosaprobe (31%) and oligosaprobe taxa (28%) predominated (Fig. S3);

- trophic status: in PE, the largest group of diatoms were eutrophic (50%) and unknown taxa (25%), LE had the highest percentage (61%) of eutrophic species but also unknown (15%) and hypereutrophic (13%) were present. In BO, the most abundant species were unknown (42%), indifferent (19%) and meso-eutrophic (16%) (Fig. S3);

- moisture aerophily: in PE, the largest group was aquatic to aerophilic (56%), representing 66% of species in D.PESB, 61% in D.PEDB, and 42% in D.PEPB. The second largest group was unknown (23%), constituting 32% of taxa in D.PEPB. In LE, 37% of the species were aquatic (24% of taxa in D.LEP2), 54% were occasionally aerophilic and 22% were unknown. In BO, the predominant groups of species were unknown (44%) and aquatic to aerophilic (33%) (Fig. S3).

Characteristic species: OMNIDIA and PCA analysis

The analysis of species characteristic of the tested ecosystems, determined according to Van Dam et al. (1994), identified 16 taxa classified as 0 in at least one category (Table 3). The next step determined the percentage contribution of each of these species classified as class 0 for the ecological parameters defined by Van Dam et al. (1994) at each sampling point (Table S4).

The principal components analysis (PCA) was performed to find the relationships between the abiotic parameters and the characteristic species ($n=36$) (Fig. 4). The Eigenvalues Plot method given eigenvalues above 1%, showed that 12 factors account for 83,2% of the total variance. The first two factors account for 31,3% of the total variance. Based on the PCA analysis for of the 16 characteristic taxa mentioned above and physical and chemical parameters, the following relationships were demonstrated:

- Group A: *Chamaepinnularia krookiformis*, *C. plinskii*, *Nitzschia liebethruthii* and *Planothidium delicatulum* demonstrate a negative correlation with pH and a positive correlation with a decrease in the concentrations of HCO_3^- , Ca^{2+} and $\text{Fe}^{2+/3+}$ (Fig. 4).

- Group B: *Craticula buderi*, *Planothidium frequentissimum* and *Navicula cincta* did not demonstrate any relationship with any water parameters (Fig. 4).
- Group C: *Navicula moskalii*, *Cyclostephanos invisitatus*, *Stephanodiscus parvus*, *S. binatus*, *Diatoma moniliformis*, *Nitzschia dissipata* var. *media*, *Mastogloia smithii*, *Pantocsekiella pseudocomensis* and *Encyonopsis subminuta* demonstrated a negative correlation with a decrease in electrolyte conductivity, as well as with the concentrations of K^+ , Mg^{2+} , Na^+ , SO_4^{2-} , Cl^- , PO_4^{3-} and Mn^{3+} and water pigments (Fig. 4).

Discussion

Verification and establishing new ecological indicator values is a key step in standard biomonitoring procedure (Szczepocka & Żelazna-Wieczorek 2018). To specify ecological indicator values or establish new ones we have determined characteristic species. For these species we performed analysis of environmental condition of the ecosystem where they were noted, previous published data and co-occurring species.

Planothidium delicatulum (PTDE) (Figs 5. A-E)

Planothidium delicatulum is a euconstant taxon for PE and an accidental taxon for LE. Its mean percentage share in PE was 2%, and constituted 5% in D.PESB.

Planothidium delicatulum does not currently have one ecological indicator value (oxygen requirements) according to Van Dam et al. (1994).

This species was more abundant in environments such as D.PESB, which was also characterized by the highest concentration of Cl^- (up to 2976 mg L⁻¹), elevated electrolytic conductivity, and decreased K^+ concentration. The pH of the water in which this species was observed did not exceed 7.

Planothidium delicatulum was mainly recorded in salty and brackish environments with neutral or slightly alkaline conditions (Campeau, Pienitz & Héquette 1999; Gell et al. 2005; Caballero et al. 2013; Yamamoto, Chiba & Tuji 2017; Van de Vijver et al. 2018).

Based on our findings, we suggest changing the following ecological indicator values in the Van Dam et al. (1994) classification for *P. delicatulum*:

- pH requirements: 3 (neutrophilic) (changing from 5 to 3);
- salinity: 5 (brackish-marine) (changing from 4 to 5)

Chamaepinnularia krookiformis (CHKF) (Figs 5. F-I) and *Chamaepinnularia plinskii* (CHPL) (Figs 5. J-M)

In 2016 *Chamaepinnularia krookiformis* was divided into two separate taxa: *C. krookiformis* and *C. plinskii* (Żelazna-Wieczorek & Olszyński 2016). Both species were very often recorded together in the same ecosystem. However, the publications which identified *C. krookiformis* often do not provide appropriate photographic documentation or photos of individual specimens (Witkowski 1994; Bąk et al. 2006; Wojtal 2008; Peszek et al. 2015). Currently available

documentation is insufficient to determine whether *C. krookiformis* and *C. plinskii* are both present simultaneously in a given environment or whether just one of these species exists. *Chamaepinnularia krookiformis* is an accessory taxon for PE (a constant taxon for D.PEDB), *C. plinskii* is a constant taxon for PE (a euconstant taxon for D.PEDB). The mean share of *C. krookiformis* was 1.6% in all PE samples, 4% in D.PEDB; for *C. plinskii*, this amounted to 2.7% in PE, 6% in D.PEDB.

Currently, *C. krookiformis* lacks three assigned ecological indicator values. For PE, it constitutes 6% of the *unknown* group in nitrogen uptake, 6% in oxygen requirements and 6% in trophic state (respectively for D.PEDB: 16%, 16% and 15%). *C. plinskii* has no assigned ecological indicator values and represents 26% of the *unknown* group for pH requirements, 19% for salinity, 12% for nitrogen uptake, 10% for oxygen requirements, 14% for saprobity, 11% for trophic state and 10% for moisture (respectively for D.PEDB: 60%, 25%, 24%, 23%, 32%, 23% and 28%).

The conditions of the environments in which both species have been recorded indicate that they are class 3 with regard to pH range (neutrophilic). Both species were the most abundant in locations subjected to periodic drying and characterised by high concentrations of chloride ions (up to 1006 mg L⁻¹) indicating a brackish environment (Żelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015).

On the basis of our findings and those of previous studies (Krammer & Lange-Bertalot 1986; Krammer 1992; Witkowski 1994; Bąk, Witkowski & Lange-Bertalot 2006; Wojtal 2009; Peszek et al. 2015; Żelazna-Wieczorek & Olszyński 2016), according to Van Dam et al (1994) ecological indicator system we propose:

established ecological indicator values for *Chamaepinnularia plinskii*

- pH requirements: 3 (neutrophilic);
- salinity: 4 (mesohalobous);
- trophic state: 5 (eutrophic);
- moisture aerophily: 4 (aerophilic);
- saprobity: 4 (α-meso-polysabrobe).

for *C. krookiformis*

- trophic state: 5 (eutrophic),

and the following changes for *Chamaepinnularia krookiformis*

- salinity: from 3 to 4 (mesohalobous);
- moisture aerophily: from 3 to 4 (aerophilic);
- saprobity: from 2 to 4 (β-mesosaprobe to α-meso-polysabrobe). Due to the specific conditions and locations of the studied objects, they were exposed to large fluctuations in the inflow of organic matter, mainly from runoff from arable fields and pollution caused by animal grazing. These impurities were manifested as elevated concentrations of K⁺ ions. Therefore, our results suggest that classifying *C. krookiformis* as an oligosaprobe is inappropriate. Further tests are needed to determine the optimum occurrence of these species in areas subjected to organic matter loads.

***Nitzschia liebethruthii* (NLBT) (Figs 5. N-Q)**

Nitzschia liebethruthii is a euconstant taxon for PE. Its means percentage share was 4% in the PE samples, and 10% in the D.PEDB samples. It was most numerous in the sample from November 2013 (19%). This species has two specific ecological indicator values. The ecological indicator value analysis for PE found *N. liebethruthii* to represent 17% of the *unknown* group in nitrogen uptake, 16% in oxygen requirements, 18% in saprobity, 17% in trophic state and 21% in moisture (respectively for D.PEDB: 37%, 37%, 43%, 36% and 49%).

Nitzschia liebethruthii occurred in environments subjected to periodic drying with a pH close to 7 and high concentration of chloride ions.

This species was noted in environments with increased salinity, electrolytic conductivity and high pH value (Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Lange-Bertalot et al. 2017; Földi et al. 2018).

We propose established new ecological indicator values according to Van Dam et al. (1994) assigned to *N. liebethruthii*:

- trophic state: 5 (eutrophic);

- moisture aerophily: 4 (aerophilic),

and following changes:

- pH requirements: 3 (neutrophilic) (changing from 5 to 3);

- saprobity: from 2 to 4 (β -mesosaprobe to α -meso-polysaprobe).

***Craticula buder* (CRBU) (Figs 5. R-U)**

Craticula buder is a euconstant taxon for PE and an accidental taxon for LE. Its mean percentage share was 4.3% in all samples for PE, and 12% for D.PEPB. This species has no recorded ecological indicator values. The ecological indicator value analysis for the PE found *C. buder* to constitute 19% of the *unknown* group in pH requirements, 28% in salinity, 16% in nitrogen uptake, 16% in oxygen requirements, 16% in saprobity, 19% in trophic state and 15% in moisture (respectively for D.PEPB: 53%, 73%, 46%, 46%, 46%, 54% and 41%).

Although *Craticula buder* was classified into group B (PCA), it was found to be most abundant in environments with an elevated concentration of Cl^- ions, ranging from 685 to 1090 mg L^{-1} , (all samples from D.PEPB and one sample from D.PEDB in which the concentration of chloride ions was 1006 mg L^{-1}). However, relative abundance was lower in the D.PEPB sample, which was characterised by a chloride ion content of over 1500 mg L^{-1} . Interestingly, the concentration of K^+ ions exceeded 100 mg L^{-1} at Cl^- concentrations below 1500 mg L^{-1} ; therefore, it is possible that the decline of this species could be related to the concentration of K^+ ions alone. Our observations indicate that the population of *C. buder* from D.PEPB favours a concentration of chloride ions from 500 to 1006 mg L^{-1} which coincides with a K^+ ions concentration from 50 to 70 mg L^{-1} .

Craticula buder is widespread throughout the world and recognized as cosmopolitan (Rumrich, Lange-Bertalot & Rumrich 2000; Lange-Bertalot 2001; Bahls 2009; Soltanpour-Gargari, Lodenius & Hinz 2011; Żelazna-Wieczorek 2011; Cichoń 2016). This species was found to be dominant in environments characterised by increased electrolytic conductivity and alkaline water

(Holmes & Taylor 2015). Holmes and Taylor (2015) place *C. buder* in the Bad water quality class. Their recorded values of diatom indices indicate that the environment was eutrophic. We therefore propose the following classes of ecological indicator values according to Van Dam et al. (1994) for *Craticula buder*:

- pH requirements: 4 (alkaliphilic);
 - trophic state: 5 (eutrophic);
 - salinity: 4 (mesohalobous);
 - moisture aerophily: 3 (aquatic to aerophilic),
- and following change:
- sabrobity: from 2 to 4 (β -mesosaprobe to α -meso-polysaprobe).

***Navicula cincta* (NCCA) (Figs 5. V-Z)**

Navicula cincta is a euconstant taxon for PE, a constant taxon for LE and an accessory taxon for BO. The mean percentage share of this species for PE is 3%, of which 7% was found in D.PEPB samples.

Currently this species has been assigned two ecological indicator values. The ecological indicator values analysis for the PE found *N. cincta* to constitute 24% of the *unknown* group in pH requirements, 13% in nitrogen uptake, 12% in oxygen requirements, 13% in saprobity, and 12% in moisture (respectively for D.PEPB: 28%, 25%, 25%, 26% and 23%).

An analysis of the physical and chemical data and the variability of occurrence did not show any clear relationships between environmental parameters and the percentage share of *Navicula cincta* in the tested samples. This lack of dependence is also confirmed by the PCA analysis.

Navicula cincta has been recorded in various types of ecosystems, although mainly in eutrophic ones with high conductivity. It also tolerates elevated levels of organic matter. This species was also observed in habitats subjected to periodic drying (Lange-Bertalot & Genkal 1999; Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Lange-Bertalot 2001; Źelazna-Wieczorek 2011; Wojtal 2013; Lange-Bertalot, Hofmann & Werum 2017).

However, several new species from the group *N. cincta* s.l. have been described, and it can be assumed that each of these individual species in this group may be associated with narrower optimal ecological conditions (Cantonati et al. 2016).

Based on our present findings, and those of previous studies, in our opinion that it is not appropriate to classify *Navicula cincta* as an oligohalobous species with regard to *salinity*: it has been recorded in fresh (Źelazna-Wieczorek 2011; Wojtal 2013), brackish (Źelazna-Wieczorek, Olszyński & Nowicka-Krawczyk 2015; Źurek et al. 2018) and salt waters (Witkowski, Lange-Bertalot & Metzeltin 2000). We propose the following ecological indicator value according to Van Dam et al. (1994) for *Navicula cincta* s.l.:

- moisture aerophily: 3 (aquatic to aerophilic).

Shade Plot analysis found that the presence of *N. cincta* s.l. can falsely indicate high similarity between samples from different environments, thus distorting the results of any environmental analysis. Therefore, with regard to the unclear taxonomic status of *Navicula cincta* s.l. and the

current lack of knowledge regarding its activities, we recommend this taxon be excluded from the biological assessment of surface water quality.

***Planothidium frequentissimum* (PLFQ) (Figs 5. AA-AF)**

Planothidium frequentissimum is a euconstant taxon for PE and LE. The mean percentage of this species for PE is 2.7%, of which 5% was found in D.PEPB samples. It was most numerous in the D.PEDB.301113 sample (19%). The species has currently six established ecological indicator values. The ecological indicator values analysis for PE found *P. frequentissimum* to constitute 12% of the *unknown* group in moisture aerophily (for D.PEPB 14% and D.PESB: 20%).

No relationship was observed between percentage share of *Planothidium frequentissimum* and the changes in chemical and physical parameters in the tested samples. This lack of relationship was confirmed by PCA analysis.

Planothidium frequentissimum is an eurytopic species that occurs globally in a variety of habitat types, from natural springs to rivers in urban areas with high levels of pollution. Its value as an indicator is low, as confirmed by the Shade Plot analysis, which found it to significantly affect the degree of similarity observed between samples from different environments.

Recently *P. frequentissimum* was divided into several different species. In studied samples we observe several species which belong to *P. frequentissimum* s. l. (Fig. 5. AA-AF) (Wetzel et al. 2019). A light microscope (LM) is still used to identify species in ecological research and biological assessment of aquatic ecosystems. Due to the likeness of the basic morphological features of the newly described species observed in LM, especially in the case of *P. frequentissimum* s.s. and *P. straubianum*, distinguishing them will be difficult or limited, which may lead to errors in the assessment. We therefore recommend that *P. frequentissimum* s. l. be excluded from the biological assessment of surface water quality (Siver, et al. 2005; Levkov et al. 2007; Żelazna-Wieczorek 2011; Kulikovskiy, Lange-Bertalot & Kuznestova 2015; Szczepocka, Nowicka-Krawczyk & Kruk 2018).

***Cyclostephanos invisitatus* (CINV) (Figs 5. AG-AJ)**

Cyclostephanos invisitatus is a euconstant taxon for LE. Its percentage share for LE was 4.8%. *C. invisitatus* currently has two ecological indicator values assigned. The ecological indicator values analysis for LE found it to constitute 24% of the *unknown* group in pH requirements, 17% in nitrogen uptake, 18% in oxygen requirements, 19% in saprobity, and 20% in moisture.

Cyclostephanos invisitatus occurs in diverse environments, however, it is most frequently reported in aquatic ecosystems subjected to high human impact, alkaline and high conductivity (Reavie & Smol 1998; Yang et al. 2005; Wojtal & Kwadrans, 2006; Kiss et al. 2012; Houk, Klee & Tanaka 2014; Reavie & Kireta 2015; Olszyński & Żelazna-Wieczorek 2018).

We therefore propose that the following classes of ecological indicator values according to Van Dam et al. (1994) be established for *Cyclostephanos invisitatus*:

- pH requirements: 4 (alkaliphilic);
- moisture aerophily: 1 (aquatic).

518 *Navicula moskalii* (NMOK) (Figs 5. AK-AN)

519 *Navicula moskalii* is an accessory taxon for LE. Its mean percentage share for LE was 1.5%. Its
520 incidence was greatest in sample D.LEP1.250315 (26%).

521 *Navicula moskalii* has no assigned ecological indicator values. The ecological indicator value
522 analysis for LE found it to constitute 5% of the *unknown* group in pH requirements, 7% in
523 salinity, 5% in nitrogen uptake, 5% in oxygen requirements, 5% in saprobity, 6% in trophic state
524 and 5% in moisture. The greatest occurrence of *N. moskalii* was observed in samples with the
525 highest concentrations of Ca^{2+} (143.6 mg L⁻¹), HCO_3^- (338.6 mg L⁻¹), SO_4^{2-} (146.9 mg L⁻¹) and
526 with high Mg^{2+} content.

527 *Navicula moskalii* was observed in a number of ecosystems (Metzeltin & Witkowski 1996;
528 Lange-Bertalot 2001; Żelazna-Wieczorek 2011; Noga et al. 2016; Lange-Bertalot et al. 2017),
529 particularly in eutrophic waters with an elevated level of Ca^{2+} and HCO_3^- ions. Żelazna-
530 Wieczorek (2011) report a significant number of *N. moskalii* in springs with high levels of
531 eutrophication, however with Ca^{2+} , SO_4^{2-} , HCO_3^- and Mg^{2+} concentrations lower than those in
532 the LE samples.

533 We therefore propose that the following classes of ecological indicator values according to Van
534 Dam et al. (1994) be established for *Navicula moskalii*:

- 535 - pH requirements: 4 (alkaliphilic);
- 536 - salinity: 2 (oligohalobous);
- 537 - trophic state: 7 (indifferent).

538 *Stephanodiscus binatus* (SBNT) (Figs 5. AO-AR)

539 *Stephanodiscus binatus* is a euconstant taxon for LE. Its percentage share for LE was 4.3%. *S.*
540 *binatus* has no recorded ecological indicator values. The ecological indicator value analysis for
541 LE found it to constitute 25% of the *unknown* group in pH requirements, 47% in salinity, 18% in
542 nitrogen uptake, 18% in oxygen requirements, 20% in saprobity, 29% in trophic state and 21% in
543 moisture.

544 The largest percentage share of *S. binatus* was recorded in the spring months and the lowest in
545 autumn. Its abundance was found to be elevated in December 2014 and 2015; the same samples
546 demonstrated the highest concentrations of Ca^{2+} , Mg^{2+} and the highest pH (above 8).

547 *Stephanodiscus binatus* has been recorded in various water ecosystems ranging from
548 oligotrophic to eutrophic; however, all are characterised by elevated pH value (Stoermer &
549 Håkansson 1984; Håkansson & Kling 1990; Houk, Klee & Tanaka 2014; Olszyński & Żelazna-
550 Wieczorek 2018).

551 We therefore propose that the following classes of ecological indicator values according to Van
552 Dam et al. (1994) be established for *Stephanodiscus binatus*:

- 553 - pH requirements: 4 (alkaliphilic);
- 554 - salinity: 2 (oligohalobous);

555 *Stephanodiscus parvus* (SPAV) (Figs 5. AS-AV)

Stephanodiscus parvus is a euconstant taxon for LE and an accidental taxon for PE. Its percentage share for LE was 2.4%. It was most abundant in the D.LEP3.260714 sample (22%). This species has three assigned ecological indicator values. The ecological indicator values analysis for LE found the taxon to constitute 9% of the *unknown* group in nitrogen uptake, 9% in oxygen requirements, 10% in saprobity, and 10% in moisture. *S. parvus* is noted mainly in eutrophic hypereutrophic ecosystems with elevated electrolytic conductivity. It is also a good indicator of waters with a strong anthropogenic impact (Reavie & Smol 1998; Reavie & Kireta, 2015; Olszyński & Żelazna-Wieczorek 2018; Reavie & Cai 2019). Based on our findings and literature data, we propose the following change in ecological indicator values according to Van Dam et al. (1994) for *Stephanodiscus parvus*:
- pH requirements: from 5 to 4 (alkaliphilic);

***Diatoma moniliformis* (DMOF) (Figs 5. AW-BA)**

Diatoma moniliformis is a euconstant taxon for BO. Its mean percentage share for BO was 3.9%. It currently has no assigned ecological indicator values. According to the ecological indicator values analysis for BO, this taxon constituted 10% of the *unknown* group in pH requirements, 11% in salinity, 8% in nitrogen uptake, 8% in oxygen requirements, 9% in saprobity, 9% in trophic state and 8% in moisture. *D. moniliformis* was found in 87.5 % of samples from BO. Interestingly, it constituted 28% of the share in one sample from December 2016 (D.BOZB.091216); however, its share was below 2% in the previous season, and was not higher than 1-2% in the other samples from December 2016. The chemical and physical characteristics of D.BOZB.091216 did not differ significantly from those of the other samples. This species is also found in fresh and salt water, as well as the Baltic and arctic areas with high conductivity (Potapova & Snoeijis 1997; Rumrich, Lange-Bertalot & Rumrich 2000; Levkov et al. 2007; Pniewski & Sylwestrzak 2018). One of the factors that influences the abundance of *D. moniliformis* is the water temperature. Studies indicate that temperatures above 10-15°C (Potapova & Snoeijis 1997; Pniewski & Sylwestrzak 2018) are associated with population growth. However, populations have been observed in freshwater streams and lakes in arctic areas, in which the temperature of the water is below 10°C (Antoniades Douglas & Smol 2005). Population growth was also observed at 4.7°C in sample D.BOZB.091216; therefore, low temperature may have an influence on the abundance of this species.

***Encyonopsis subminuta* (ESUM) (Figs 5. BB-BG)**

Encyonopsis subminuta is a euconstant taxon in BO, where its mean percentage share was 4.1%. Presently, *E. subminuta* has been assigned five ecological indicator values. Ecological indicator values analysis for BO found it to constitute 7% of the *unknown* group in nitrogen uptake and 8% in moisture.

Encyonopsis subminuta was found to be most abundant in sample D.BOZB.041115. The sample was also characterised by an elevated concentration of $\text{Fe}^{2+/3+}$ ions and the lowest pH value. In subsequent samples, when the concentration of Fe ions dropped, the abundance of *E. subminuta* also decreased.

E. subminuta is regarded as a cosmopolitan taxon, occurring in the temperate and boreal zone. It is most abundant in oligo- to mesotrophic waters with electrolytic conductivity between 190- 250 $\mu\text{S L}^{-1}$ (Krammer 1997; Noga et al. 2014; Novais et al. 2014; Feret, Bouchez & Rimet 2017). *E. subminuta* may be sensitive to the concentration of Fe ions; however, the increase of these ions is associated with a drop in pH. Our research confirms that the optimal pH for population size is close to 7.

***Mastogloia smithii* (MSMI) (Figs 5. BH-BM)**

Mastogloia smithii is a euconstant taxon for BO. Its mean percentage share for BO was 6.3%. It was found in greatest numbers in D.BOZB.300615 (22%) and D.BOZB.261016 (15%). *M. smithii* has been assigned four ecological indicator values. The ecological indicator values analysis for the BO found this species to constitute 13% of the *unknown* group in nitrogen uptake, 14% in oxygen requirements and 15% in trophic state.

The environment in BO regarding *salinity* was classified according to Vam Dam et al. (1994) as oligohalobus (43% species); however, 7% of the mesohalobous species were represented by one species: *Mastogloia smithii*.

This species is recorded in fresh, brackish and salt water (Witkowski, Lange-Bertalot & Metzeltin 2000; Busse & Snoeijis 2003; Weckström & Juggins 2005; Martinzes-Goss & Evangelista 2011; Lange-Bertalot et al. 2017). Its presence in environments with varying degrees of salinity may suggest that this does not have a significant influence on population size.

Based on our present findings and literature data, we propose the following change in the ecological indicator values according to Van Dam et al. (1994) for *Mastogloia smithii*:

- salinity: from 4 to 3 (halophilic).

***Nitzschia dissipata* var. *media* (NDME) (Figs 5. BN-BQ)**

Nitzschia dissipata var. *media* is a euconstant taxon for BO. Its mean percentage share for BO was 3.8%, and the highest proportion (20%) was found in D.BOZB.041115. *N. dissipata* var. *media* has been assigned two ecological indicator values. Ecological indicator values analysis for BO found it to constitute 9% of the *unknown* group in nitrogen uptake, 10% in oxygen requirements, 11% in saprobity, 11% in trophic state and 9% in moisture.

The increase in occurrence of *N. dissipata* var. *media* is associated with an increase in the level of $\text{Fe}^{2+/3+}$ ions, similar to *Encyonopsis subminuta*. In addition, it was found in the ecosystem, i.e. BO, with the lowest concentrations of ions indicative of the presence of organic pollutants in the environment, such as K^+ and NH_4^+ .

Although *Nitzschia dissipata* var. *media* is found sporadically, it is commonly found in oligo- to mesotrophic waters with a pH between 7 and 8 (Van der Vijver, Frenot & Beyens 2002;

Antoniades, Douglas & Smol 2005; Żelazna-Wieczorek 2011; Lange-Bertalot, Hofmann & Werum 2017).

Based on our findings and literature data, we propose the following ecological indicator values according to Van Dam et al. (1994) for *Nitzschia dissipata* var. *media*:
- saprobity: 2 (β -mesosaprobe)

***Pantocsekiella pseudocomensis* (PPCS) (Figs 5. BR-BX)**

Pantocsekiella pseudocomensis is a euconstant taxon for BO. Its mean percentage share for BO was 4.4%. It was most abundant in D.BOZB.250315 (9%) and in D.BOZB.220616 (10%). *P. pseudocomensis* has not been assigned any ecological indicator values according to Van Dam et al. (1994). The ecological indicator values analysis for BO found it to represent 22% of the unknown group in pH requirements, 25% in salinity, 10% in nitrogen uptake, 11% in oxygen requirements, 18% in saprobity, 12% in trophic state and 13% in moisture. The greatest amount of *P. pseudocomensis* was found in samples characterised by the highest levels of ammonium ions. Its percentage share was lowest in samples with the lowest water temperature, apart from D.BOZB.250315.

Currently, *Pantocsekiella pseudocomensis* is assigned to the *P. comensis* complex, with *P. comensis* and *P. costei*. In our opinion that assigning ecological indicator values for particular species of the *P. comensis* complex is unjustified at the current state of knowledge, and that all species within the complex should be assigned the same provisional ecological indicator values until their individual properties are better understood (Houk, Klee & Tanaka 2010; Kistenich et al. 2014; Duleba et al. 2015).

Conclusions

The water ecosystems created in the post-mining areas create a complex of conditions that are not found in other natural ecosystems, and the benthic diatom species present in such environments are very often present in higher numbers than in other habitats. The specific hydro-geological conditions prevailing in the post-production reservoirs provide a unique opportunity to observe interspecies differences and intra-species variability, allowing for the verification or isolation of new taxa and a greater insight into their autecology (Żelazna-Wieczorek & Olszyński 2016; Olszyński & Żelazna-Wieczorek 2018).

The identification of species characteristic of the studied ecosystems may foster further growth of ecological research and increase the reliability of surface water quality assessment, as such knowledge is needed to verify their ecological indicator values, and hence calculate diatoms indices with greater accuracy.

Ecological indicator values as set out by Van Dam et al. (1994) are utilized in many ecological works describing the ecological conditions of the studied ecosystems. These ecological indicator values form the basis for calculating diatoms indices describing the ecological state of surface waters. It is therefore necessary to constantly update and establish new ecological indicator values for particular diatom species.

Many authors who describe new species, or encounter existing species in new ecosystems, regularly propose updates for individual ecological indicator values. However, these findings, may not be introduced and updated in the OMNIDIA program for a number of years. The OMNIDIA system is used by state institutions in many countries around the world to assess surface water quality (Campeau, Pienitz & Héquette 1999; Rumrich, Lange-Bertalot & Rumrich 2000; Witkowski, Lange-Bertalot & Metzeltin 2000; Gell et al. 2005; Potapova & Ponader 2008; Wojtal & Sobczyk 2012; Caballero et al. 2013; Żelazna-Wieczorek & Olszyński 2015; Yamamoto, Chiba & Tuji 2017; Lange-Bertalot, Hofmann & Werum 2017; Földi et al. 2018; Van de Vijver, Wetzel & Ector 2018). Clearly, if these assessments are based on incomplete or outdated data, assessments of aquatic environments may be fraught with error. The present study used three principles to identify proposed changes in the classification of ecological indicator values for characteristic species according to the Van Dam et al. (1994) system, or to establish new values which were previously absent: the analysis of environmental conditions prevailing in the studied ecosystems, the analysis of relevant literature data, and references to the ecological indicator values of other species (Table 4). This mode of research can serve as a model for updating databases used to assess surface water quality.

References

- Antoniades D., Douglas M.S.V. & Smol J.P. 2005. Benthic diatom autecology and inference model development from the Canadian High Arctic Archipelago. *Journal of Phycology* 41: 30-45 DOI: 10.1111/j.1529-8817.2005.04049.x.
- Bahls L.L. 2009. A checklist of diatoms from inland waters of the northwestern United States. *Proceedings of The Academy of Natural Sciences of Philadelphia* 158: 1-35 DOI: 10.1635/053.158.0101.
- Bąk M., Witkowski A. & Lange-Bertalot H. 2006. Diatom flora diversity in the strongly eutrophicated and β -mesosaprobic waters of the Szczecin Lagoon, NW Poland, southern Baltic Sea. – In: Ognjanova-Rumenova N. & Manoylov K., eds. *Advances in Phycological Studies, Festschrift in Honour of Professor Dobrina Teminskova-Topalova*. Sofia – Moscow: Pensoft Publishers & University Publishing House, 293–317.
- Busse S. & Snoeijs P. 2003. Gradient responses of diatom communities in the Bothnian Sea (northern Baltic Sea), with emphasis on responses to water movement. *Phycologia* 42(5): 451-464 DOI: 10.2216/i0031-8884-42-5-451.1.
- Caballero M., Rodríguez A., Vilaclara G., Ortega B., Roy P. & Lozano S. 2013. Hydrochemistry, ostracods and diatoms in a deep, tropical, crater lake in Western Mexico. *Journal of Limnology* 72(3): 512-523 DOI: 10.4081/jlimnol.2013.e42.
- Campeau S., Pienitz R. & Héquette A. 1999. Diatoms from the Beaufort Sea coast, southern Arctic Ocean (Canada). In: Lange-Bertalot H. & Kociolek P., eds. *Bibliotheca Diatomologica Vol. 42*. Berlin, Stuttgart: J. CRAMER 244 p.

- Cantonati M., Angeli N. Spitale D. & Lange-Bertalot H. 2016. A new *Navicula* (Bacillariophyta) species from low-elevation carbonate springs affected by anthropogenic disturbance. *Fottea, Olomouc* 16(2): 255-265 DOI: 10.5507/fot.2016.013.
- Cichoń S. 2016. *Craticula buderi* (Bacillariophyceae) in Poland. *Polish Botanical Journal* 61(2): 301–305 DOI: 10.1515/pbj-2016-0024.
- Clarke K.R. & Gorley, R.N. 2015. PRIMER v7: User Manual/Tutorial. Devon, United Kingdom: PRIMER-E Ltd 296 pp.
- De Haan H., Van Liere L., Klapwijk S.J.P. & Van Donk E. 1993. The structure and function of fen lakes in relation to water table management in The Netherlands. *Hydrobiologia* 265: 155–177 <http://dx.doi.org/10.1007/BF00007266>.
- Duleba M., Kiss K.T., Földi A., Kovács J., Borojević K.K., Molnár L.F., Plenković-Moraj A., Pohner Z., Solak C.N., Tóth B. & Ács E. 2015. Morphological and genetic variability of assemblages of *Cyclotella ocellata* Pantocsek/*C. comensis* Grunow complex (Bacillariophyta, Thalassiosirales), *Diatom Research* 30(4): 283-306 DOI: 10.1080/0269249X.2015.1101402.
- European Union. 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. Official Journal of the European Communities L327, 1–73.
- Feret L., Bouchez A. & Rimet F. 2017. Benthic diatom communities in high altitude lakes: a large scale study in the French Alps. *Annales de Limnologie - International Journal of Limnology* 53: 411-423 DOI: 10.1051/limn/2017025.
- Ferreira da Silva, E., Almeida, A.F.P., Nunes, M.L., Luís, A.T., Borg, F., Hedlund, M., Marques de Sá, C., Patinha, C. & Teixeira, P. 2009. Heavy metal pollution downstream the abandoned Coval da Mó mine (Portugal) and associated effects on epilithic diatom communities. *Science of the Total Environment* 407: 5620–5636 doi:10.1016/j.scitotenv.2009.06.047.
- Földi A., Ács É., Grigorszky I., Ector L., Wetzel C.E., Várbíró G., Kiss K.T., Dobosy P., Trábert Z., Borsodi A.K. & Duleba M. 2018. Unexpected consequences of bombing. Community level response of epiphytic diatoms to environmental stress in a saline bomb crater pond area. *PLoS ONE* 13(10): e0205343 <https://doi.org/10.1371/journal.pone.0205343>.
- Gabrys-Godlewska A., Gruszecki J., Kochanowska J., Kozula R., Pasieczna A. & Tomassi-Morawiec H. 2004. Arkusz Turek (550). In: Sikorska-Maykowska M., Zdanowski A. & Gabrys-Godlewska A., eds. *Objaśnienia do mapy geośrodowiskowej Polski 1:50 000*. Warszawa: Państwowy Instytut Geologiczny. Opracowanie Zamówione Przez Ministra Środowiska. p 30.
- Gadomska W., Kaźmierczak J., Merdzińska M., Zielińska B., Wiśniewska-Zientara G., Gajda M., Opas Z., Cieślak J., Kończak D., Kujawa S., Lukaszewicz J., Zielińska-Adasiak S., Ratajczyk G., Michalska R. & Sobczak D. 2007. Raport o stanie gminy Brudzew lata 2002-2006. Urząd Gminy Brudzew. Brudzew. p. 15.

- 747 Gell P.A., Bulpin S., Wallbrink P., Hancock G. & Bickford S. 2005. Tareena Billabong – a
748 palaeolimnological history of an ever-changing wetland, Chowilla Floodplain, lower
749 Murray-Darling Basin, Australia. *Marine and Freshwater Research* 56: 441-456 DOI:
750 10.1071/MF04107.
- 751 Harding W.R. & Taylor J.C., 2014. Diatoms as indicators of historical water quality: A
752 comparison of samples taken in the Wemmershoek catchment (Western Province, South
753 Africa) in 1960 and 2008. *Water S.A.* 40 (4): 601–606.
- 754 Håkansson H. & Kling H. 1990. The current status of some very small freshwater diatoms of the
755 genera *Stephanodiscus* and *Cyclostephanos*. *Diatom Research* 5: 273-287
756 DOI:10.1080/0269249X.1990.9705119.
- 757 Houk, V., Klee R. & Tanaka H. 2010. Atlas of freshwater centric diatoms with a brief key and
758 descriptions. Part III. Stephanodiscaceae A. *Fottea (Supplement)* 10:1–498.
- 759 Houk V., Klee R. & Tanaka H. 2014. Atlas of freshwater centric diatoms with a brief key and
760 descriptions. Part IV. Stephanodiscaceae B. *Fottea (Supplement)* 14:1–532.
- 761 Holmes M. & Taylor J.C. 2015 Diatoms as water quality indicators in the upper reaches of the
762 Great Fish River, Eastern Cape, South Africa. *African Journal of Aquatic Science* 1-17
763 DOI: 10.2989/16085914.2015.1086722.
- 764 Hutorowicz A. & Pasztalenic A. 2014. Phytoplankton metric of ecological status assessment for
765 polish lakes and its performance along nutrient gradient. *Polish Journal of Ecology*
766 62(3): 525-540 DOI:10.3161/104.062.0312.
- 767 Kahlert M., Ács E., Almeida S.F.P., Blanco S., Dreßler M., Ector L., Karjalainen S.M., Liess A.,
768 Mertens A., van der Wal J., Vilaste S. & Werner P. 2016. Quality assurance of diatom
769 counts in Europe: towards harmonized datasets. *Hydrobiologia* 772: 1–14 DOI:
770 10.1007/s10750-016-2651-8.
- 771 Kasztelewicz, Z. 2011. Assessment of Reclamation Works in Polish Open-pit Lignite Mining.
772 *Studia KPZK* 142: 440–452.
- 773 Kelly M.G., Juggins S., Guthrie R., Pritchard S., Jamieson J., Rippey B., Hirst H. & Yallop M.
774 2008. Assessment of ecological status in U.K. rivers using diatoms. *Freshwater Biology*
775 53 (2): 403–422 DOI: 10.1111/j.1365-2427.2007.01903.x.
- 776 Kiss K.T., Klee R., Ector L. & Ács É. 2012. Centric diatoms of large rivers and tributaries in
777 Hungary: morphology and biogeographic distribution. *Acta Botanica Croatica* 71: 311–
778 363 DOI: 10.2478/v10184-011-0067-0
- 779 Kistenich S., Dreßler M., Zimmermann J., Hübener T., Bastrop R. & Jahn R. 2014. An
780 investigation into the morphology and genetics of *Cyclotella comensis* and closely
781 related taxa. *Diatom Research* 29(4): 423-440 DOI: 10.1080/0269249X.2014.922125.
- 782 Kolada A., Pasztaleniec A., Bielczyńska A. & Soszka H. 2016. Phytoplankton, macrophytes and
783 benthic diatoms in lake classification: consistent, congruent, redundant? Lessons learnt
784 from WFD-compliant monitoring in Poland. *Journal of Limnology* 59: 44–52 DOI:
785 10.1016/j.limno.2016.05.003.

- 786 Krammer K. 1992. Pinnularia. Eine Monographie der europäischen Taxa In: *Bibliotheca*
787 *Diatomologica* 26. Berlin–Stuttgart: J. Cramer. 353 pp.
- 788 Krammer K. 1997. Die cymbelloiden Diatomeen. Eine Monographie der weltweit bekannten
789 Taxa. Teil 2. *Encyonema* part., *Encyonopsis* and *Cymbellopsis*. In: Lange-Bertalot H. &
790 Kociolek P., eds. *Bibliotheca Diatomologica Vol. 37*. Berlin, Stuttgart: J. CRAMER. 96
791 pp.
- 792 Krammer K. & Lange-Bertalot H. 1986. Bacillariophyceae. 1. Teil: Naviculaceae In: Ettl H.H,
793 Gerloff J., Heynig H. & Mollenhauer D., eds. *Süßwasserflora von Mitteleuropa, Band*
794 *2/1*. Jena: Gustav Fisher Verlag. 876 pp.
- 795 Krammer K. & Lange-Bertalot H. 1997. Bacillariophyceae. 1. Teil: Bacillariaceae,
796 Epithemiaceae, Surirellaceae. In: Ettl H.H, Gerloff J., Heynig H. & Mollenhauer D.,
797 eds. *Süßwasserflora von Mitteleuropa, Band 2/2*. Jena Stuttgart Lübeck Ulm: Gustav
798 Fisher Verlag. 611 pp.
- 799 Kulikovskiy M.S., Lange-Bertalot H. & Kuznetsova I.V. 2015. Lake Baikal: Hotspot of Endemic
800 Diatoms II. In: Lange-Bertalot H., ed. *Iconographia Diatomologica Vol. 26. Taxonomy-*
801 *Biogeography-Diversity*. Germany: Koeltz Scientific Books. 656 pp.
- 802 Lange-Bertalot H. 2001. *Navicula* sensu stricto. 10 Genera Separated from *Navicula* sensu lato.
803 *Frustulia*. In: Lange-Bertalot H., ed. *Diatom of Europe Vol. 2*. Ruggell:vA.R.G. Gantner
804 Verlag K.G. 526 pp.
- 805 Lange-Bertalot H. & Genkal S.I. 1999. Diatoms from Siberia I. Islands in the Arctic Ocean
806 (Yugorsky-Shar Strait). In: Lange-Bertalot H., ed. *Iconographia Diatomologica Vol. 6.*
807 *Phytogeography-Diversity-Taxonomy*. Vaduz: A.R.G. Gantner Verlag K.G. 292 pp.
- 808 Lange-Bertalot H., Hofmann G., Werum M. & Cantonati M. 2017. Freshwater Benthic Diatoms
809 of Central Europe: Over 800 Common Species Used in Ecological Assessment. Koeltz
810 Botanical Books. 942 pp.
- 811 Lecointe C., Coste M. & Prygiel J. 1993. “Omnidia:” a software for taxonomy, calculation of
812 diatom indices and inventories management. *Hydrobiologia* 269/270: 509-513.
- 813 Levkov Z., Krstic S., Metzeltin D. & Nakov T. 2007. Diatoms of Lakes Prespa and Ohrid. About
814 500 taxa from ancient lake system. In: Lange-Bertalot H., ed. *Inconografia*
815 *Datomologica Vol. 16*. Ruggell: A.R.G. Gantner Verlag K.G. 613 pp.
- 816 Luís A.T., Durães, N., Fernandes Pinheiro de Almeida S. & Ferreira da Silva E. 2016.
817 Integrating geochemical (surface waters, stream sediments) and biological (diatoms)
818 approaches to assess AMD environmental impact in a pyritic mining area: Aljustrel
819 (Alentejo, Portugal). *Journal of Environmental Sciences* 42: 215-226
820 <http://dx.doi.org/10.1016/j.jes.2015.07.008>.
- 821 Luís A.T., Teixeira P., Almeida S.F.P., Ector L., & Matos J.X. & Ferreira da Silva E.A. 2009.
822 Impact of Acid Mine Drainage (AMD) on Water Quality, Stream Sediments and
823 Periphytic Diatom Communities in the Surrounding Streams of Aljustrel Mining Area
824 (Portugal). *Water Air Soil Pollution* 200:147-167 DOI: 10.1007/s11270-008-9900-z.

- Macioszczyk A. & Dobrzyński D. 2002. Hydrogeochemia. Wydawnictwo Naukowe PWN. Warszawa. 448 pp.
- Martinez-Goss M.R. & Evangelista L.T. 2011. A Contribution to the Taxonomy of *Mastogloia* (Class Bacillariophyceae) in the Philippines. *Philippine Journal of Science* 140(1): 7-12.
- Metzeltin D. & Witkowski A. 1996. Diatomeen der Bären-Insel. Süßwasser- und marine Arten. In: Lange-Bertalot H., ed. *Iconographia Diatomologica. Annotated Diatom Micrographs. Vol 4. Taxonomy*. Königstein/Germany: Koeltz Scientific Books 20-21.
- Noga T., Stanek-Tarkowska J., Pajączek A., Kochman N. & Peszek Ł. 2014. Ecological assessment of the San River water quality on the area of the San Valley Landscape Park. *Journal of Ecological Engineering* 15(4): 12-22 DOI: 10.12911/22998993.1125453.
- Noga T., Stanek-Tarkowska J., Rybek M., Kochaman-Kędziora N., Peszek Ł. & Pajączek A. 2016. Diversity of diatoms in the natural, mid-forest Terebowiec Stream – Bieszczady National Park. *Journal of Ecological Engineering* 17(4): 232-247 DOI: 10.12911/22998993/64506.
- Novais M.H., Morais M.M., Rosado J., Dias L.S, Hoffmann L. & Ector L. 2014. Diatoms of temporary and permanent watercourses in Southern Europe (Portugal). *River Research and Applications* 30: 1216-1232 DOI: 10.1002/rra.2818.
- Olaczek R. 1963. Zbiorowiska roślinne torfowisk niskich okolic Łęczycy. Unpublished doctoral dissertation, University of Lodz, Poland.
- Olszyński R.M. & Żelazna-Wieczorek J. 2018. *Aulacoseira pseudomuzzanensis* sp. nov. and other centric diatoms from post iron ore mining reservoirs in Poland. *Diatom Research* 33(2): 155-185 DOI: 10.1080/0269249X.2018.1509886.
- Orlikowski D. & Szwed L. 2009. Wodny kierunek rekultywacji w KWB „Adamów” SA – inwestycją w przyszłość regionu. *Górnictwo i Geoinżynieria* 33(2): 351-361.
- Peszek Ł., Noga T., Stanek-Tarkowska J., Pajączek A., Kochman-Kędziora N., Pieniążek M. 2015. The effect of anthropogenic change in the structure of diatoms and water quality of the Żołynianka and Jagielnia streams. *Journal of Ecological Engineering* 16(2): 33-51 <https://doi.org/10.12911/22998993/1856>.
- Pliński M. 1966. Glony słonego źródła w Pełczyskach pod Ozorkowem. Unpublished master dissertation, University of Lodz, Poland.
- Pliński M. 1969. Okrzemki słonego źródła w Pełczyskach pod Ozorkowem. *Zesz. Nauk. UŁ* 2(31): 123-136.
- Pliński M. 1971a. Rzadkie i nowe dla flory Polski gatunki glonów z terenów solnisk w okolicach Łęczycy. *Zesz. Nauk. UŁ* 2(41): 171-184.
- Pliński M. 1971b. Gatunki z rodzaju *Enteromorpha* (Link.) Agardh z terenu solnisk podłęczyckich. *Zesz. Nauk. UŁ* 2(41): 159-169.
- Pliński M. 1971c. System halobow w świetle współczesnych poglądów. *Wiad. Ekol.* 17(1): 18-29.
- Pliński M. 1973. Glony solnisk podłęczyckich. The algae of salt marches near Łęczycza, Central Poland. *Mon. Botanic.* 39.

- 865 Pniewski F. & Sylwestrzak Z. 2018. Influence of short periods of increased water temperature on
866 species composition and photosynthetic activity in the Baltic periphyton communities.
867 *Biologia* 73(11): 1067-1072 <https://doi.org/10.2478/s11756-018-0122-6>.
- 868 Poikane S., Kelly M. & Cantonati M. 2016. Benthic algal assessment of ecological status in
869 European lakes and rivers: Challenges and opportunities. *Science of the Total*
870 *Environment* 568: 603–613 DOI:10.1016/j.scitotenv.2016.02.027.
- 871 Potapova M. & Snoeijs P. 1997. The natural life cycle in wild populations of *Diatom*
872 *moniliformis* (Bacillariophyceae) and its disruption in an aberrant environment. *Journal*
873 *of Phycology* 33: 924-937 <https://doi.org/10.1111/j.0022-3646.1997.00924.x>.
- 874 Potapova M.G. & Ponader K.C. 2008. New species and combinations in the diatom genus
875 *Sellaphora* (Sellaphoraceae) from Southeastern United States. *Harvard Papers in*
876 *Botany* 13(1): 171–181 DOI: 10.3100/1043-4534(2008)13[171:NSACIT]2.0.CO;2.
- 877 Rakowska B. 1996. The benthic diatom community of a reservoir after the exploration of brow
878 coal in Konin (central Poland). *Algological Studies* 82: 103–116
879 <http://dx.doi.org/0342-1120/0116-103>.
- 880 Reavie E.D. & Cai M. 2019. Consideration of species-specific diatom indicators of
881 anthropogenic stress in the Great Lakes. *PLoS ONE* 14(5): e0210927 DOI:
882 <https://doi.org/10.1371/journal.pone.0210927>
- 883 Reavie E.D. & Kireta A.R. 2015. Centric, Araphid and Eunotioid Diatoms of the Coastal
884 Laurentian Great Lakes. In: Lange-Bertalot H. & Kociolek J.P., eds. *Bibliotheca*
885 *Diatomologica Vol. 62*. Stuttgart: J. Cramer Gebr. Borntraeger Verlagsbuchhandlung.
886 184 pp.
- 887 Reavie E.D. & Smol J.P. 1998. Freshwater diatoms from the St. Lawrence river. In: Lange-
888 Bertalot H. & Kociolek J.P., eds. *Bibliotheca Diatomologica Vol 41*. Berlin, Stuttgart: J.
889 Cramer Gebr. Borntraeger Verlagsbuchhandlung, 184 pp.
- 890 Rumrich U., Lange-Bertalot H. & Rumrich, M. 2000. Diatomeen der Anden. Von Venezuela bis
891 Patagonien/Tierra del Fuego. Und zwei weitere Beiträge. In: Lange-Bertalot H., ed.
892 *Iconographia Diatomologica Vol. 9*. Ruggell: A.R.G. Gantner Verlag K.G. 673 pp.
- 893 Sienkiewicz E. & Gąsiorowski M. 2016. The evolution of a mining lake – From acidity to
894 natural neutralization. *Science of the Total Environment* 557–558: 343–354
895 <http://dx.doi.org/10.1016/j.scitotenv.2016.03.088>.
- 896 Siver P.A., Hamilton P.B., Stachura-Suchoples K. & Kociolek J.P. 2005. The Freshwater Flora
897 of Cape Code, Massachusetts, U.S.A. In: Lange-Bertalot H., eds. *Iconographia*
898 *Diatomologica Vol. 14. Diatoms of North America*. Ruggell: A.R.G. Gantner Verlag
899 K.G. 463 pp.
- 900 Soltanpour-Gargari A., Lodenius M. & Hinz F. 2011. Epilithic diatoms (Bacillariophyceae) from
901 streams in Ramsar, Iran. *Acta Botanica Croatica*. 70(2): 167-190 DOI: 10.2478/v10184-
902 010-0006-5.

- 903 Stoermer E.F. & Håkansson H. 1984. *Stephanodiscus parvus*: validation of an enigmatic and
- 904 widely misconstrued taxon. *Nova Hedwigia* 39: 497–511
- 905 <http://www.jstor.org/stable/3226406>.
- 906 Szczepocka E., Nowicka-Krawczyk P. & Kruk A. 2018. Deceptive ecological status of urban
- 907 streams and rivers—evidence from diatom indices. *Ecosphere* 9(7):e02310 DOI:
- 908 10.1002/ecs2.2310
- 909 Szczepocka E., Szulc B., Szulc K., Rakowska B. & Żelazna-Wieczorek J. 2014. Diatom indices
- 910 in the biological assessment of the water quality based on the example of a small
- 911 lowland river. *Oceanological and Hydrobiological Studies* 43(3): 265–273 DOI:
- 912 10.2478/s13545-014-0141-z
- 913 Szczepocka E. & Żelazna-Wieczorek J. 2018. Diatom biomonitoring – scientific foundations,
- 914 commonly discussed issues and frequently committed errors. *Oceanological and*
- 915 *Hydrobiological Studies* 47(3): 313-325 DOI: 10.1515/ohs-2018-0030
- 916 Thomas E.J. & John J. 2006. Diatoms and macroinvertebrates as biomonitors of mine-like in
- 917 Collie, Western Australia. *Journal of the Royal Society of Western Australia* 89:
- 918 109–117.
- 919 Tümpling W. & Friedrich G. 1999. Methiden der Biologischen Wasseruntersuchung. Jena.
- 920 Stuttgart. Lübeck. Ulm. G. Fischer Verlag 133-153.
- 921 Weckström K. & Juggins S. 2005. Coastal diatom–environment relationships from the Gulf of
- 922 Finland, Baltic Sea. *Journal of Phycology* 42: 21-35 DOI: 10.1111/j.1529-
- 923 8817.2006.00166.x
- 924 Witkowski A. 1994. Recent and fossil diatom flora of the Gulf of Gdańsk, Southern Baltic Sea.
- 925 In: *Bibliotheca Diatomologica Vol. 28*. Cramer, Berlin, Stuttgart. 313 pp
- 926 Witkowski A., Lange-Bertalot H. & Metzeltin D. 2000. Diatom flora of marine coasts I. In:
- 927 Lange-Bertalot H., ed. *Iconographia Diatomologica. Vol. 7. Diversity-Taxonomy-*
- 928 *Identification*. Ruggell, Liechtenstein: A.R.G. Gantner Verlag K.G. 419 pp.
- 929 Wetzel C.E., Van de Vijver B., Blanco S. & Ector L. 2019. On some common and new cavum–
- 930 bearing Planorbulina (Bacillariophyta) species from freshwater. *Fottea* 19(1): 50–89
- 931 DOI: 10.5507/fot.2018.016
- 932 Wojtal A.Z. 2009. The diatoms of Kobylanka stream near Kraków (Wyżyna Krakowsko–
- 933 Częstochowska upland, S Poland). *Polish Botanical Journal* 54: 129–330.
- 934 Wojtal A.Z. 2013. Species composition and distribution of diatom assemblages in spring water
- 935 from various geological formation in Southern Poland. In: Lange-Bertalot H. &
- 936 Kociolek P., eds. *Bibliotheca Diatomologica Vol. 59*. Stuttgart: J. CRAMER in der
- 937 Gebr. Borbtraeger Verlagsbuchhandlung. 436 p.
- 938 Wojtal A.Z. & Kwadrans J. 2006. Diatoms of the Wyżyna Krakowsko-Częstochowska Upland
- 939 (S. Poland) – Coscinodiscophyceae (Thalassiositophycidae). *Polish Botanical Journal*
- 940 51: 177–207.

- 941 Wojtal A.Z. & Sobczyk Ł. 2012. The influence of substrates and physicochemical factors on the
942 composition of diatom assemblages in karst springs and their applicability in water-
943 quality assessment. *Hydrobiologia* 695: 97–108 DOI 10.1007/s10750-012-1203-0.
- 944 Van Dam H., Mertens A. & Sinkeldam, J. 1994. A coded checklist and ecological indicator
945 values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic*
946 *Ecology* 28: 117–133.
- 947 Van de Vijver B., Frenot Y. & Beyens L. 2002. Freshwater diatoms from Ile de la Possession. In:
948 Lange-Bertalot H. & Kociolek P., eds. *Bibliotheca Diatomologica Vol. 46*. Stuttgart: J.
949 CRAMER in der Gebr. Borchtraeger Verlagsbuchhandlung 71 p.
- 950 Van de Vijver B., Wetzel C.E. & Ector L. 2018. Analysis of the type material of *Planothidium*
951 *delicatulum* (Bacillariophyta) with the description of two new *Planothidium* species
952 from the sub-Antarctic Region. *Fottea* 18(2): 200-211 DOI: 10.5507/fot.2018.006.
- 953 Van Landingham S.L. 1968. Investigation of a diatom population from mine Tailings in Nye
954 Country, Nevada, U.S.A. *Journal of Phycology* 4: 306–310.
- 955 Yang X-D., Dong X-H., Gao G., Pan H-X. & Wu J-L. 2005. Relationship Between Surface
956 Sediment Diatoms and Summer Water Quality in Shallow Lakes of the Middle and
957 Lower Reaches of the Yangtze River. *Journal of Integrative Plant Biology* 4792: 153-
958 164 <https://doi.org/10.1111/j.1744-7909.2005.00035.x>.
- 959 Yamamoto M., Chiba T. & Tuji A. 2017. Salinity responses of benthic diatoms inhabiting tidal
960 flats. *Diatom Research* 32: 243-250 DOI: 10.1080/0269249X.2017.1366951
- 961 Żelazna-Wieczorek J. 1996. Studium taksonomiczne rodzaju *Vaucheria* De Candolle oraz uwagi
962 o występowaniu zidentyfikowanych gatunków tego rodzaju. Unpublished doctoral
963 dissertation, University of Lodz, Poland.
- 964 Żelazna-Wieczorek J. 2002. *Vaucheria* species in selected regions in Poland. *Acta Societatis*
965 *Botanicorum Poloniae* 71(2): 129-139.
- 966 Żelazna-Wieczorek, J. 2011. Diatom flora in springs of Łódź Hills (Central Poland). In:
967 Witkowski A., ed. *Diatom monographs. Vol. 13*. Ruggell, Liechtenstein: A.R.G.
968 Gantner Verlag K.G. 419 pp.
- 969 Żelazna-Wieczorek J. & Nowicka-Krawczyk P. 2015. The cascade construction of artificial
970 ponds as a tool for urban stream restoration – The use of benthic diatoms to assess the
971 effects of restoration practices. *Science of the Total Environment* 538: 591–599
972 DOI:10.1016/j.scitotenv.2015.08.110.
- 973 Żelazna-Wieczorek J. & Olszyński R.M. 2016. Taxonomic revision of *Chamaepinnularia*
974 *krookiformis* Lange-Bertalot et Krammer with a description of *Chamaepinnularia*
975 *plinskii* sp. nov. *Fottea, Olomouc* 16(1): 112-121 DOI: 10.5507/fot.2016.001.
- 976 Żelazna-Wieczorek J., Olszyński R.M. & Nowicka-Krawczyk P. 2015. Half a century of
977 research on diatoms in athalassic habitats in central Poland. *Oceanological and*
978 *Hydrobiological Studies* 441: 51–67 DOI: 10.1515/ohs-2015-0006.
- 979 Żurek R., Diakiv V., Szarek-Gwiazda E., Kosiba J. & Wojtal A.Z. 2018. Unique Pit Lake
980 Created in an Opencast Potassium Salt Mine (Dombrowska Pit Lake in Kalush,

981 Ukraine). *Mine Water and the Environment* 37(3): 456-469
982 <https://doi.org/10.1007/s10230-018-0527-z>

Figure 1

Location of sampling points in the Łódzkie and Wielkopolskie voivodships. Poland.

A. Pełczyńska (PE). B. Łęczyca (LE). C. Bogdałów (BO).



Figure 2

nMDS 3D analysis. The diagram shows three distinct clouds of samples which coincide with the three hydrological objects.




Figure 3

Shade Plot analysis. The diagram shows the strength of the factor (taxon) affecting the similarity between the samples.

Upper dendrogram - samples divided according to hydrological object. Left dendrogram - 50 taxa of diatoms which have the strongest influence on the similarity between the samples.




Figure 4

Principal Component Analysis.

The diagram presents three groups of species A, B and C, whose occurrence can be correlated with selected physical and chemical factors.




Figure 5

LM microphotographs of characteristic diatom species.

A-E. *Planothidium delicatulum*. F-I. *Chamaepinnularia krookiformis*. J-M. *Chamaepinnularia plinskii*. N-Q. *Nitzschia liebethruthii*. R-U. *Craticula buderi*. V-Z. *Navicula cincta*. AA-AF. *Planothidium frequentissimum*. AG-AJ. *Cyclostephanos invisitatus*. AK-AN. *Navicula moskalii*. AO-AR. *Stephanodiscus binatus*. AS-AV. *S. parvus*. AW-BA. *Diatoma moniliformis*. BB-BG. *Encyonopsis subminuta*. BH-BM. *Mastogloia smithii* (BK-BM. Same specimen, different focal plane). BN-BQ. *Nitzschia dissipata* var. *media*. BR-BX. *Pantocsekiella pseudocomensis*. Scale bar = 10µm.




Table 1 (on next page)

Physical and chemical parameters in the examined sampling sites. The minimum, maximum and mean values.

	Pelczyska (PE)									Leczyca (LE)									Bogdałów (BO)		
	outflow (D.PESB)			ditch (D.PEDB)			pond (D.PEPB)			L1 (D.LEP1)			L2 (D.LEP2)			L3 (D.LEP3)			reservoir (D.BOZB)		
	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.	min	max	avar.
pH	6.6	7.7	7.0	6.4	6.4	6.4	7.8	9.6	8.7	7.9	8.6	8.2	7.5	8.6	8.2	7.5	8.8	8.3	7.9	8.3	8.1
Conductivity [μS cm ⁻¹]	4450	9230	6699	5170	5170	5170	2645	5150	3646	657	865	743	558	836	728	472	778	680	505	734	623
T [°C]	6.8	13.9	9.1	7.8	7.8	7.8	1.1	17.3	7.1	1.6	20.6	11.1	0.6	21.1	11.1	2.2	15.2	9.2	4.7	22.7	11.4
HCO ₃ ⁻ [mg L ⁻¹]	345	744	475	610	610	610	284	451	352	211	339	260	168	275	210	183	290	241	174	369	270
CO ₂ [HCO ₃ ⁻] [mg L ⁻¹]	124	268	171	220	220	220	102	163	126	76	122	94	61	99	75	66	105	87	63	133	96
Cl ⁻ [mg L ⁻¹]	1585	2976	2426	1006	1006	1006	685	1524	1053	57	93	78	60	93	76	67	106	78	41	52	45
N _{NH4} [mg L ⁻¹]	0.02	1.63	0.45	0.15	0.15	0.15	0.00	0.36	0.11	0.02	0.71	0.22	0.03	0.53	0.26	0.00	1.16	0.32	0.00	0.07	0.02
NH ₄ ⁺ [mg L ⁻¹]	0.03	2.09	0.71	0.19	0.19	0.19	0.00	0.46	0.14	0.03	0.55	0.22	0.03	0.54	0.28	0.00	0.90	0.31	0.00	0.09	0.03
PO ₄ ³⁻ [mg L ⁻¹]	0.60	12.46	3.73	8.57	8.57	8.57	0.85	9.10	3.99	0.22	0.53	0.36	0.05	0.57	0.41	0.09	0.57	0.34	0.27	0.45	0.34
P _{PO4} [mg L ⁻¹]	0.20	4.11	1.23	2.83	2.83	2.83	0.28	3.00	1.32	0.07	0.17	0.12	0.02	0.19	0.13	0.03	0.19	0.11	0.09	0.15	0.12
SO ₄ ²⁻ [mg L ⁻¹]	176	198	188	165	165	165	151	197	176	71	147	107	67	122	101	63	91	75	110	147	123
S _{SO4} [mg L ⁻¹]	58.8	66.2	62.9	55.1	55.1	55.1	50.3	65.7	58.7	23.5	49.0	35.8	22.3	40.8	33.8	21.0	30.3	25.1	37.9	49.2	41.5
COLOR [mgPt dm ⁻³]	25	160	81	140	140	140	50	120	78	10	60	27	9	60	27	12	60	31	4	10	6
Mn ³⁺ [mg L ⁻¹]	0.17	0.51	0.36	0.14	0.14	0.14	0.03	0.25	0.13	0.00	0.05	0.02	0.01	0.03	0.02	0.01	0.07	0.03	0.00	0.01	0.00
Fe ^{2+/3+} [mg L ⁻¹]	0.25	0.41	0.32	0.25	0.25	0.25	0.05	0.24	0.11	0.03	0.27	0.09	0.00	0.13	0.04	0.01	0.07	0.03	0.01	0.09	0.02
Mg ²⁺ [mg L ⁻¹]	39.5	48.7	45.0	35.9	35.9	35.9	25.6	37.0	32.1	11.6	19.1	15.2	10.3	18.3	15.1	9.5	15.3	12.5	9.6	15.7	11.7
Ca ²⁺ [mg L ⁻¹]	171.7	216.2	195.2	165.0	165.0	165.0	75.8	139.8	121.1	58.0	143.6	89.0	59.5	117.6	79.1	52.6	86.4	65.5	77.7	117.6	86.3
Na ⁺ [mg L ⁻¹]	500.7	1537.4	1227.1	453.3	453.3	453.3	277.3	681.8	455.9	15.6	40.4	30.9	20.3	42.4	28.3	22.0	42.4	33.0	24.0	43.1	32.6
K ⁺ [mg L ⁻¹]	8.5	124.8	42.4	109.8	109.8	109.8	58.6	68.8	63.5	4.5	9.5	7.2	5.1	9.3	7.3	5.6	10.7	8.2	0.1	3.6	1.5

Table 2 (on next page)

Characteristic species according to SIMPER analysis for studied waterbodies.

EC - euconstant taxa, CN - constant taxa, CN - accessory taxa (Tümping & Friedrich 1999).

Pelczyska dich D.PEDB	Łęczyca reservoirs LE	Bogdałów reservoir BO
<i>Chamaepinnularia krookiformis</i> (AC)	<i>Achnantheidium minutissimum</i> (EC)	<i>Achnantheidium minutissimum</i> (EC)
<i>Chamaepinnularia plinskii</i> (CN)	<i>Amphora pediculus</i> (EC)	<i>Diatoma moniliformis</i> (EC)
<i>Cocconeis placentula</i> (CN)	<i>Cyclostephanos dubius</i> (EC)	<i>Encyonopsis subminuta</i> (EC)
<i>Craticula buderii</i> (EC)	<i>Cyclostephanos invisitatus</i> (EC)	<i>Mastogloia smithii</i> (EC)
<i>Craticula halophila</i> (EC)	<i>Cyclotella meneghiniana</i> (EC)	<i>Nitzschia dissipata</i> var. <i>media</i> (EC)
<i>Fragilaria famelica</i> (EC)	<i>Navicula gregaria</i> (EC)	<i>Pantocsekiella ocellata</i> (EC)
<i>Fragilaria sopotensis</i> (CN)	<i>Navicula moskalii</i> (AC)	<i>Pantocsekiella pseudocomensis</i> (EC)
<i>Gomphonema parvulum</i> (EC)	<i>Nitzschia palea</i> (EC)	
<i>Hippodonta hungarica</i> (CN)	<i>Stephanodiscus binatus</i> (EC)	
<i>Navicula cincta</i> (EC)	<i>Stephanodiscus hantzschii</i> (EC)	
<i>Navicula veneta</i> (EC)	<i>Stephanodiscus parvus</i> (EC)	
<i>Nitzschia frustulum</i> (EC)		
<i>Nitzschia inconspicua</i> (EC)		
<i>Nitzschia liebetruthii</i> (EC)		
<i>Nitzschia palea</i> (CN)		
<i>Nitzschia perminuta</i> (EC)		
<i>Nitzschia tubicola</i> (CN)		
<i>Planothidium delicatulum</i> (EC)		
<i>Planothidium frequentissimum</i> (EC)		
<i>Tabularia fasciculata</i> (EC)		

Table 3(on next page)

Characteristics species with classification of ecological indicators values by Van Dam et al. (1994).

Species	Code	Moisture acrophity	Nitrogen uptake	pH requirements	Oxygen requirements	Salinity	Saprobity	Trophic state
<i>Achnantheidium minutissimum</i>	ADMI	3	2	3	1	2	2	7
<i>Amphora pediculus</i>	APED	3	2	4	2	2	2	5
<i>Chamaepinnularia krookiformis</i>	CHKF	3	0	3	0	3	1	0
<i>Chamaepinnularia plinskii</i>	CHPL	0	0	0	0	0	0	0
<i>Cocconeis placentula</i>	CPLA	2	2	4	3	2	2	5
<i>Craticula buderi</i>	CRBU	0	0	0	0	0	0	0
<i>Craticula halophila</i>	CHAL	2	2	4	2	4	3	5
<i>Cyclostephanos dubius</i>	CDUB	1	2	5	2	3	3	5
<i>Cyclostephanos invisitatus</i>	CINV	0	0	0	0	2	0	5
<i>Cyclotella meneghiniana</i>	CMEN	2	3	4	5	3	4	5
<i>Diatoma moniliformis</i>	DMOF	0	0	0	0	0	0	0
<i>Encyonopsis subminuta</i>	ESUM	0	0	3	1	1	1	1
<i>Fragilaria famelica</i>	FFAM	3	1	4	1	2	1	3
<i>Fragilaria sopotensis</i>	FSOP	1	2	4	1	2	2	4
<i>Gomphonema parvulum</i>	GPAR	3	3	3	4	2	4	5
<i>Hippodonta hungarica</i>	HHUN	3	2	4	3	2	2	4
<i>Mastogloia smithii</i>	MSMI	3	0	4	0	4	2	0
<i>Navicula cincta</i>	NCCA	0	0	0	0	2	0	7
<i>Navicula gregaria</i>	NGRE	3	2	4	4	3	3	5
<i>Navicula moskalii</i>	NMOK	0	0	0	0	0	0	0
<i>Navicula veneta</i>	NVEN	3	2	4	4	3	4	5
<i>Nitzschia dissipata</i> var. <i>media</i>	NDME	0	0	4	0	2	0	0
<i>Nitzschia frustulum</i>	NIFR	3	4	4	3	3	2	5
<i>Nitzschia inconspicua</i>	NINC	3	3	4	3	3	3	5
<i>Nitzschia liebethruthii</i>	NLBT	0	0	5	0	4	0	0
<i>Nitzschia palea</i>	NPAL	3	4	3	4	2	5	6
<i>Nitzschia perminuta</i>	NIPM	3	1	4	1	2	1	2
<i>Nitzschia tubicola</i>	NTUB	2	3	4	4	3	5	6
<i>Pantocsekiella ocellata</i>	POCL	1	1	4	1	1	1	4
<i>Pantocsekiella pseudocomensis</i>	PPCS	0	0	0	0	0	0	0
<i>Planothidium delicatulum</i>	PTDE	3	1	5	0	4	5	3
<i>Planothidium frequentissimum</i>	PLFQ	0	2	4	3	2	4	7
<i>Stephanodiscus binatus</i>	SBNT	0	0	0	0	0	0	0
<i>Stephanodiscus hantzschii</i>	SHAN	2	3	5	4	2	4	6
<i>Stephanodiscus parvus</i>	SPAV	0	0	5	0	2	0	6
<i>Tabularia fasciculata</i>	TFAS	3	2	4	3	4	3	5

Table 4(on next page)

Selected 16 characteristic species with the new or altered (in red) ecological indicator values according to Van Dam et al. (1994).

Species	Code	Moisture acrophity	Nitrogen uptake	pH requirements	Oxygen requirements	Salinity	Saprobity	Trophic state
<i>Chamaepinnularia krookiformis</i>	CHKF	3 4	0	3	0	3 4	1 4	0 5
<i>Chamaepinnularia plinskii</i>	CHPL	0 4	0	0 3	0	0 4	0 4	0 5
<i>Craticula buderi</i>	CRBU	0 3	0	0 4	0	0 4	0 4	0 5
<i>Cyclostephanos invisitatus</i>	CINV	0 1	0	0 4	0	2	0	5
<i>Diatoma moniliformis</i>	DMOF	0	0	0	0	0	0	0
<i>Encyonopsis subminuta</i>	ESUM	0	0	3	1	1	1	1
<i>Mastogloia smithii</i>	MSMI	3	0	4	0	4 3	2	0
<i>Navicula cincta</i>	NCCA	0 3	0	0	0	2	0	7
<i>Navicula moskalii</i>	NMOK	0	0	0 4	0	0 2	0	0 7
<i>Nitzschia dissipata</i> var. <i>media</i>	NDME	0	0	4	0	2	0 2	0
<i>Nitzschia liebethruthii</i>	NLBT	0 4	0	5 3	0	4	0 4	0 5
<i>Pantocsekiella pseudocomensis</i>	PPCS	0	0	0	0	0	0	0
<i>Planothidium delicatulum</i>	PTDE	3	1	5 3	0	4 5	5	3
<i>Planothidium frequentissimum</i>	PLFQ	0	2	4	3	2	4	7
<i>Stephanodiscus binatus</i>	SBNT	0	0	0 4	0	0 2	0	0
<i>Stephanodiscus parvus</i>	SPAV	0	0	5 4	0	2	0	6