

A peer-reviewed version of this preprint was published in PeerJ on 25 October 2019.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.7608) (peerj.com/articles/7608), which is the preferred citable publication unless you specifically need to cite this preprint.

Šťovíček A, Cohen-Chalamish S, Gillor O. 2019. The effect of reverse transcription enzymes and conditions on high throughput amplicon sequencing of the 16S rRNA. PeerJ 7:e7608
<https://doi.org/10.7717/peerj.7608>

The effect of reverse transcription enzymes and conditions on high throughput amplicon sequencing of the 16S rRNA

Adam Šťovíček¹, Smadar Cohen-Chalamish², and Osnat Gillor¹

¹Zuckerberg Institute for Water Research, Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Israel

²The Mina and Everard Goodman Faculty of Life Sciences and Advanced Materials and Nanotechnology Institute, Bar-Ilan University, Ramat-Gan, Israel

Corresponding author:
Osnat Gillor¹

Email address: gilloro@bgu.ac.il

ABSTRACT

It is assumed that the sequencing of ribosomes better reflects the active microbial community than the sequencing of the ribosomal RNA encoding genes. Yet, many studies exploring microbial communities in various environments, ranging from the human gut to deep oceans, questioned the validity of this paradigm due to the discrepancies between the DNA and RNA based communities. Here we focus on an often neglected key step in the analysis, the reverse transcription (RT) reaction. Previous studies showed that RT may introduce biases when expressed genes and ribosomal rRNA are quantified, yet its effect on microbial diversity and community composition was never tested. High throughput sequencing of ribosomal RNA is a valuable tool to understand microbial communities as it better describes the active population than DNA analysis. However, the necessary step of RT may introduce biases that have so far been poorly described. In this manuscript, we compare three RT enzymes, commonly used in soil microbiology, in two temperature modes to determine a potential source of bias due to non-standardized RT conditions. In our comparisons, we have observed up to 6 fold differences in bacterial class abundance. A temperature induced bias can be partially explained by G-C content of the affected bacterial groups, thus pointing towards a need for higher reaction temperatures. However, another source of bias was due to enzyme processivity differences. This bias is potentially hard to overcome and thus mitigating it might require the use of one enzyme for the sake of cross-study comparison.

INTRODUCTION

Massively parallel amplicon sequencing revolutionized our view of microbial world: by sequencing a taxonomic tag such as 16S rRNA encoding gene, it allows taxonomic description of the microbial communities (Quammen, 2018). However, the existing approaches introduce caveats: the DNA amplicon sequencing may capture 'relic DNA', which is a recalcitrant genetic material from dead cells or naked DNA (Carini et al., 2016) in addition, amplicon sequencing carries technical biases due to sample preparation, DNA extraction methods (Pan et al., 2010), amplification reaction (Pfeiffer et al., 2014) and analysis (Pollock et al., 2018). Moreover, DNA-based microbiome information can describe the total community, but it cannot report which members are metabolically active (Blazewicz et al., 2013). In contrast to DNA based tools, analysis of ribosomes can describe the metabolically active members of a given community. The combination of data generated from rRNA encoding genes and ribosomes led to a wide range of ecological insights, including the response to climatic changes (Angel et al., 2013), pH and water availability (Romanowicz et al., 2016), and biogeochemical processes (Freedman et al., 2015).

Ribosomal analysis studies are based on an assumption that ribosomes are more abundant in active cells compared to dormant ones (Blazewicz et al., 2013; Lennon and Jones, 2011). However, this assumption may not always be correct. Dormant bacteria may be misclassified as active, when ribosomes are present in cells and spores that are inactive (Segev et al., 2013; Blagodatskaya and Kuzyakov, 2013). In contrast,

active bacteria with low metabolic turnover and low ribosomal count could be labeled as dormant when sequencing depth is insufficient (Steven et al., 2017; Joergensen and Wichern, 2018). In spite of various biases that introduce discrepancies in the community structure (Forney et al., 2004), ribosomal analysis can capture the biological variability highlighting large differences between samples. However, if more subtle differences are of interest, technical biases could confound biological interpretations (Lever et al., 2015; McCarthy et al., 2015). This is due to specific challenges introduced RNA based analysis (Bustin and Nolan, 2004, 2017). Therefore, to confidently compare results across ribosome-based amplicon sequencing studies, we must determine which component of the analysis: RNA extraction, processing or data analysis may influence the outcome and introduce biases.

Prior studies focused on biases in the steps of RNA extraction, amplification and sequencing, but disregard any biases that may occur during RT (Creer et al., 2016). At the crucial step of RT, most researches simply 'follow the manufacture instructions' (Table 1). However, RT kits typically detail a wide range of temperatures, primer, template and reaction options, which may lead to different results. The reverse transcriptase (RT) enzyme requires sequence priming to initiate a reaction. Primers could be poly-A complementary, random or sequence specific. Poly-A priming is limited to eukaryotic mRNA which makes it unsuitable for use with ribosomal taxonomic tags. Opinions vary about the usefulness of random and sequence-specific priming for the analysis of microbiomes: Random priming may produce higher yield of cDNA and improve the detection limit (Zhang and Byrne, 2015; Ståhlberg et al., 2004a), but may decrease the reproducibility and introduce bias (Bustin and Nolan, 2004; Hansen et al., 2010). Sequence specific primes require fine tuning of the reaction conditions and higher template concentration than random priming (Ståhlberg et al., 2004b). Moreover, the results of the RT reaction is determined not only by the type of the RT enzyme used, but also by the reaction conditions (Curry et al., 2002; Ståhlberg et al., 2004a,b; Bustin and Nolan, 2004; Sieber et al., 2010).

Ideally RT efficiency is near 100 %, but in practice it varies dramatically: 90 % efficiency was reported for SuperScript III (mutated MMLV RT) (Ståhlberg et al., 2004a), 20 % for Murine Leukemia Virus (M-MLV) RT (Curry et al., 2002), and as low as 2 % for Avian Myeloblastosis Virus (AMV) RT (Ståhlberg et al., 2004a). Moreover almost two order of magnitude difference were reported between mutated and wild-type AMV RT (Ståhlberg et al., 2004a).

To the best of our knowledge no study has yet compared the RT reaction conditions for environmental microbiome profile. We hypothesize that during RT reactions, varying RT enzyme types and temperature conditions will yield different results in microbial diversity and community composition. We further predict that variations in communities will be G-C dependent. To test our prediction, we present a comparative study of commonly used RT enzymes in the field of environmental microbiology as well as a comparison of two different reaction temperatures.

Manufacturer	RT Enzyme	RT origin	Temperature [°C] Suggested Used	RNA type	Primer type	Ref
Promega	MMLV	MMLV	37–42	NA	rRNA	926R
				NA	rRNA & mRNA	Random hexamers
	ImProm-II	AMV	37–55	42	rRNA & mRNA	Random hexamers
				NA	rRNA	Random hexamers
Qiagen	QuantiTect	Quantiscript	42–50	NA	rRNA	Unique RT Primer Mix
				37	rRNA	Random hexamers
	Omniscript	Quantiscript	37	NA	mRNA	Random hexamers
				NA	mRNA	invA-R
Takara	PrimeScript II	AMV	42–50	NA	mRNA	Random hexamers
				NA	rRNA	Random hexamers
Roche	Roche reverse transcription kit	AMV	42–60	42 & 50	rRNA	Random hexamers
				42 & 50	rRNA	Random hexamers
Thermo Fisher	MMLV	MMLV	37–42	45	rRNA	900R
				NA	rRNA	Random hexamers
	SuperScript-II	MMLV	42–55	NA	rRNA	1492R
				NA	mRNA	Random hexamers
	SuperScript-III	MMLV	42–55	NA	rRNA	Random hexamers
				NA	rRNA	27F & LR3
						Romanowicz et al. (2016)

Table 1. Literature overview of RT conditions applied in soil microbiological studies.

MATERIALS AND METHODS

Study site and sample collection scheme

Soil samples were collected at the central Negev Desert highlands, Israel (Zin Plateau, 30°86'N, 34°80'E) at an established ecological research site. The mean annual precipitation at the sampling site is 90 mm and the mean annual temperature is 30 °C (LTER data). Samples were collected under the canopy of perennial shrub *Hammada scoparia* in October 2015 at the end of the dry season as previously described (Baubin et al., 2019). Briefly, sampling was conducted in seven random blocks, each providing two technical replicates resulting in 14 samples. Samples were collected from the top 5 cm of the soil, following the removal of crust and debris. The soil samples were processed within 24 h of collection. Samples were homogenized using 2 mm sieve and the duplicates from each block were composited. This resulted in seven final replicates.

RNA preparation

Total RNA was extracted from each of the seven samples using a phenol-chlorophorm extraction previously described by Angel (2012). The reaction buffer pH was adjusted to 5. The total RNA was subsequently purified with the MagListo™ Total RNA Extraction Kit (Bioneer, Daejeon, Republic of Korea). Contaminant DNA was removed using a DNase I from the MasterPure RNA Purification Kit (Epicenter, Madison, WI, USA) with two successive treatments of 30 min according to manufacturer's instructions. The reaction mixture was purified using the MagListo™ Total RNA Extraction Kit (Bioneer). The absence of contaminant DNA was verified using total bacterial primers 341F (5' CCTACGGGAG-GCAGCAG 3') and 515R (5' TTACCGCGGCTGCTGGCAC 3') (Klindworth et al., 2013) and DreamTaq DNA polymerase (Thermo Scientific, Waltham, MA, USA) under the following conditions: 95 °C for 5 min, followed by 26 cycles of 95 °C for 15 s, 60 °C for 30 s and 72 °C for 30 s for extension followed by 72 °C for 5 min for final extension. If amplification was detected the sample was discarded, re-extracted, purified and tested. Only DNA-free samples were used in this study.

Reverse transcription reaction conditions

RT kits used in this experiment were chosen to represent the most commonly used enzymes in the field (Figure 1). All seven samples were reverse transcribed by the same kit to reduce batch effects. Each RT kit used in this study originated from a different source: (I.) ImProm-II Reverse Transcription System enzyme (Promega, Madison, WI, USA) originates from AMV RT, (II.) SuperScript IV Reverse Transcriptase Kit enzyme (ThermoFisher Scientific, Waltham, MA, USA) originates from MMLV RT and (III.) TGIRT originates from the mobile group II introns reverse transcriptase (Mohr et al., 2013) TGIRT™-III Enzyme (InGex, St. Louis, MO, USA). Each reaction consisted of 50 ng of total RNA template, measured by Quanti-iT™ RNA Assay Kit (ThermoFisher), and random hexamer primers (0.5 µg/reaction). Template and primer mix were heated to 70 °C (ImProm-II) or 65 °C (SuperScript IV). Each reaction was subsequently cooled to 4 °C for 5 min and incubated at 42 °C (ImProm-II), 55 °C (ImProm-II and Superscript) or 57 °C (TGIRT) for 60 min (ImProm-II), 120 min (TGIRT) or 10 min (SuperScript IV). All reactions were terminated and DNA was removed by alkaline lysis using 2 µl of 1 M NaOH, incubating for 12 min at 70 °C. After which the reaction was neutralized using 4 µl of 0.5 M acetic acid (Table 2).

Illumina sequence preparation

The V3 and V4 regions of the resulting cDNA were amplified using 341F (5' CCTACGGGAGGCAGCAG 3') and 806R (5' GGTCTGGACTACHVGGGTWTCTAAT 3') (Klindworth et al., 2013) primers. Each reaction was performed in triplicate and consisted of 1 mM bovine serum albumin (Takara, Kusatsu, Japan), 2.5 µl of 10x standard buffer, 5 µM primers, 0.8 mM dNTPs, 0.4 µl DreamTaq DNA polymerase (Thermo Scientific, Waltham, MA, USA), and 4 µl of template cDNA. The reaction mixtures were subsequently amplified using the following PCR conditions: 95 °C for 30 s, 27 cycles of 95 °C for 15 s, 50 °C for 30 s, 68 °C for 30 s and 68 °C for 5 min. Resulting amplicon presence was verified using 1.5 % agarose gel electrophoresis. Resulting technical triplicates were combined, and the sequencing libraries were constructed using the TruSeq® DNA Sample Preparation Kit (Illumina, San Diego, CA, USA) following the manufacturer's recommendations. The amplicon libraries were sequenced (250x2 base pairs, pair-end) on the Illumina MiSeq System platform at the Research Resources Centre at the University of Illinois.

Manufacturer	RT kit	Primers	Thermo cycling		Reaction mix	
			Temperature [°C]	Time [min]	Reactant	Amount
Promega	Im-Prom II	Random Hexamers (500 ng/reaction)	70	5	DTT	10 µM
			4	5	Tris-HCl	50 mM
			25	5	KCl	75 mM
			42	60	MgCl ₂	2.5 mM
			70	15	dNTP	0.5 mM
Promega	Im-Prom II	Random Hexamers (500 ng/reaction)			RNase inhibitor	0.5 µl/20 µl
			70	5	DTT	10 µM
			4	5	Tris-HCl	50 mM
			25	5	KCl	75 mM
			55	60	MgCl ₂	2.5 mM
ThermoFisher	SuperScriptIV	Random Hexamers (2.5 µM)	70	15	dNTP	0.5 mM
					RNase inhibitor	0.5 µl/20 µl
			65	5	DTT	5 µM
			0	1	Tris-HCl	50 mM
			23	10	KCl	50 mM
TGIRT	TGIRT-III	Random Hexamers (500 ng/reaction)	55	10	MgCl ₂	4 mM
			80	10	dNTP	0.5 mM
					RNase inhibitor	0.5 µl/20 µl
			65	5	DTT	5 µM
			0	1	Tris-HCl	10 mM

Table 2. A summary of conditions applied to the different reaction conditions. .

Sequence analysis

Resulting paired end sequences were merged using the CASPER program (Kwon et al., 2014), and the resulting merged reads were clustered using the UPARSE pipeline according to the recommended settings (Edgar, 2013). The resulting OTU representative sequences were taxonomically assigned with the SINA incremental aligner using a lowest common ancestor algorithm (Pruesse et al., 2007) and the SILVA database version 132 (Quast et al., 2013). All sequences retrieved in this study were uploaded to European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) submission number PRJEB32237.

Class enrichment plot data preparation

To explore whether G-C content contributed to differences in relative abundances of different taxa, we ran the following analysis: for each reaction condition we calculated pairwise comparisons at the class level: we normalized the proportional enrichment in each respective reaction conditions following Equation 1 (Figure 2 and Supplementary Figure S2), where the A and B represent a class at the different conditions.

$$Class_{normalized} = \frac{A - B}{A + B} \quad (1)$$

The error bars represent a standard deviation, which have been calculated as a standard deviation of each category and normalized according to the Equation 2. The δA and δB represent the standard deviation of class A and B. Details of deriving this equations are specified in Supplementary Equation S1.

$$\delta Class_{normalized} = \frac{2}{(A + B)^2} \sqrt{B^2 \delta A^2 + A^2 \delta B^2} \quad (2)$$

Statistical analysis

All data analysis was performed in R v3.4.3 (R Core Team, 2018). The dataset was sub-sampled (rarified) to an even depth of 9000 sequences per sample using python numpy package v1.15.4 (Van Der Walt et al., 2011). The subsampling removed five samples from the dataset. Additionally, three more samples were removed as outliers (<https://gitlab.com/stovicek/rt/article/blob/master/data/preparation.ipynb>). In order to equalize the number of replicates, two random samples were removed from the TGIRT dataset. This reduced the number of replicates to four samples per experimental category. The sample diversity was analyzed using the vegan package v2.5-2 (Oksanen et al., 2018). The data was visualized using the R package ggplot2 package v2.2.1 (Wickham, 2016) and python package matplotlib (Hunter, 2007).

RESULTS

Sample preparation and diversity analysis

After removing obvious outliers, four samples from each condition were analyzed. The summary of the analysis and all the code used to produce each figure is included in the Supplementary File S1 and S2. The changes in bacterial diversity among the tested conditions were expressed using species count, Pielou's evenness index (Pielou, 1967) and Shannon diversity index (Shannon and Weaver, 1949). Despite observed trends in the diversity indices, no statistically significant differences were detected (Supplementary Figure S1, Supplementary diversity_statistics.ipynb).

Relative abundance plot

A relative abundance of major taxonomic classes is depicted in the Figure 1. Each column is an average of four biological replicates. OTUs (operational taxonomic units) that were not taxonomically assigned at this level are summarized as "Unclassified" ($\approx 2\%$). Various patterns (detailed below) were detected among the experimental conditions: some patterns could be attributed to differences in reaction temperature (which ranged from 42°C to 55°C and 57°C). Other patterns could be linked to enzyme type. RT reactions with Superscript IV and TGIRT RTs yielded no significant differences in class relative abundances. However, transcription with ImProm-II RT at a similar temperature ($\approx 55^\circ\text{C}$) yielded different abundances: specifically, the abundances of Alphaproteobacteria, Bacteroidia, Deltaproteobacteria, Oxyphotobacteria, Rubrobacteria and Verrucomicrobiae decreased. However, Chloflexia, Gammaproteobacteria and Thermoleophilia abundances increased when their ribosomes were transcribed with ImProm-II RT (Figure 1, Supplementary Table S1). When transcription occurred at lower temperature (42°C), relative abundances of Bacilli, Deltaproteobacteria and Oxyphotobacteria were enriched, while Actinobacteria, Chloroflexia and Acidimicrobia were depleted under the same conditions (Figure 1 and Supplementary Table S1).

Class enrichment plot

The Figure 2 also depicts the weighted average of the G-C content in each class. The proportional comparison is interpreted as follows: A value of zero in the proportional comparison represents the taxonomic class count that is exactly equal between the two compared groups (Figure 2). A value of 1 or -1 is assigned when a given taxonomic class is only present in one category and absent in another, respectively. In general, there was a tendency towards lower G-C content lower temperature of ImProm-II (Figure 2,a, and Supplementary Figure S2, a and b). No statistically significant differences were detected between the profiles resulting from RT of SuprScript-IV or TGIRT (Supplementary Figure S2, d). The taxa Gemmatimonadetes, Fibrobacteria and Thermoanaerobaculia were relatively insensitive to the RT conditions. However the majority of the classes were enriched in some conditions. (I.) the rate of RT was only sensitive to temperature for the classes Alphaproteobacteria, Gemmatimonadetes, Fibrobacteria, Thermoanaerobaculia, TK10, and Blastocatellia (Figure 2, a). These groups tend to have extreme GC content (both high and low). (II.) Gammaproteobacteria, Planctomycetacia and Phycisphaerae are relatively insensitive to the reaction temperature (Figure 2, a), but their abundances vary with different RT enzymes (Figure 2, b, Supplementary figure S2).

Class enrichment statistics

We calculated a linear regression, where the response variable was the relative proportion of each class between two tested categories, and the explanatory variable was the G-C content (the assumptions tests and plots can be found in the Supplementary file S2). The linear regression assumptions were tested: in case of two condition pairs (ImProm-II at 42°C & SuperScript-IV as well as ImProm-II at 55°C & SuperScript-IV), the assumption conditions were not met (Supplementary file S2). Since we cannot confidently discard the null hypothesis in these cases, we do not consider these two tests significant. Therefore, we are considering only the ImProm-II 42°C & ImProm-II 55°C as well as ImProm-II 42°C & TGIRT as a significant outcome. Differences in the remaining cases cannot be explained by the weighted G-C content alone (Table 3).

DISCUSSION

As high throughput sequencing has become increasingly accessible in recent years, researchers urgently call for method standardization to allow for accurate cross-study comparisons (Pan et al., 2010; Blago-




Figure 1. Relative abundance of main classes across each tested condition. Only top 15 % of the most abundant classes are represented and the rest is summarized in the 'Low abundance' category. The x axis show different enzymes and conditions. The y axis shows an average relative abundance. Each category is an average of 4 samples.

209 datskaya and Kuzyakov, 2013). With this motivation, researchers developed new platforms that offer
 210 protocols and standardized methods for data acquisition from DNA resources, such as the Earth Micro-
 211 biome Project (<http://www.earthmicrobiome.org/>) that standardizes DNA amplicon sequencing. Despite
 212 the success with standardizing methods and protocols for DNA-based analysis, to this date, there is
 213 RNA-based methods have not been standardized, despite the discrepancies repeatedly reported between
 214 the RNA and DNA based analyses (Blazewicz et al., 2013; Carini et al., 2016; Dlott et al., 2015) and
 215 the plethora of methods used in these studies (Table 1). The analysis of rRNA adds specific biases to
 216 high throughput sequencing analysis, such as reduced template stability compared to DNA, RT priming
 217 bias, and linearity of RT reaction (Bustin and Nolan, 2004). These biases need to be either minimized or
 218 standardized.

219 In this study, we focused on one crucial step in the RNA analysis that was previously overlooked:
 220 the transcription of RNA to cDNA (Table 1). Several biases connected to the RT reaction have been
 221 described for RT-qPCR, such as quantification of expressed genes (Bustin and Nolan, 2004, 2017; Zhang
 222 and Byrne, 2015) in RNA-Seq, i.e., primer related bias of expressed transcripts (Hansen et al., 2010). Yet,
 223 the role of RT in diversity patterns was not yet investigated in the context of high throughput sequencing
 224 of ribosomes. Here, we focus on the role of enzyme and reaction temperature in shaping the diversity and
 225 composition of ribosome-based communities. We have compared four RT enzymes commonly used in soil




Figure 2. A proportional comparison of most abundant classes between the ImProm-II at 42 °C and 55 °C (a) and the ImProm-II at 55 °C and SuperScriptIV at 55 °C (b). Enrichment is expressed such that a class that is equally proportional in both conditions, has a value of 0. If the class shows in one condition but is absent from another, its value would be equal to 1 or -1 respectively. Furthermore, a weighted average of the GC content of each class is expressed as the bar color. Each value is an average of 4 biological replicates.

Condition 1	Condition 2	Adjusted R2	t value	p value	Significance	Note
ImProm-II 42 °C	ImProm-II 55 °C	0.429	5.366	4.89E-06	***	
ImProm-II 42 °C	SuperScript IV	0.1373	2.692	0.0127	*	•
ImProm-II 42 °C	TGIRT	0.2032	3.231	0.00264	**	
ImProm-II 55 °C	SuperScript IV	0.0841	-2.097	0.0431	*	•
ImProm-II 55 °C	TGIRT	0.03624	-1.546	0.131		
SuperScript IV	TGIRT	0.03634	1.548	0.130		

Table 3. The linear regression statistics.

We used a GC content as an explanatory variable of a the class enrichment. The rows marked with a • do not fulfill all test assumptions (see supplementary diversity_statistics.ipynb).

microbiology (Table 1) at two distinct temperature modes (42 °C and 55 °C–57 °C). Then we analyzed temperature and RT enzyme-related effects on the resulting community profiles (Table 2).

Under different reaction conditions, we detected differences in the relative abundance of bacterial classes portrayed by different reaction conditions (Figure 1). Some observed differences can be attributed to the combined effect of reaction temperature and average template G-C content (Figure 2 and Supplementary Figure S2). As expected, this effect is clearest when we applied the same enzyme (ImProm-II) at two reaction temperatures (42 °C and 55 °C), then the G-C content had the highest prediction power ($t = 5.366$, $p = 4.8 \times 10^{-5}$, Table 3). Likewise, in every comparison of RT enzymes at low and high temperature, G-C content affected the relative abundance of taxonomic classes with statistical significance. Although the RT reactions are commonly performed at 42 °C (Table 1), our results indicate that this reaction temperature is too low to allow successful RT of some soil community taxa, in particular species with higher G-C content.

When transcription was performed with different enzymes under similar reaction temperatures, relative abundances of taxa differed notably (Figure 1 and 2). Although the RTs of SuperScript-IV and TGIRT originate from different organisms, they yielded similar taxa abundances. Reactions with ImProm-II yielded different profiles. These differences cannot be explained by the G-C content (Table 3), but could be attributed to ribosome properties and the efficiency of RT. The ribosomes extracted from the soil environment were diverse and probably differ in their secondary and tertiary structures (Yilmaz et al., 2006) post-transcriptional modifications (Schwartz and Motorin, 2017). Thus RTs kinetics would differ. The discrepancies reported in this study raise further questions: how would one decide which enzymes or temperatures best reflect the active community composition? It has been suggested that the total

community could be used as a reference to accurately deduce the diversity. Furthermore, this study was performed on desert soil samples and the effects in other environments remain to be determined.

We previously demonstrated that the total and active communities in the Negev soil used in this study differ in both abundance profiles and community composition during the dry season, while during the wet season, no differences were detected (Baubin et al., 2019). During the dry season, a "phantom taxa" (Klein et al., 2016), *Deinococcus-Thermus*, comprised $\approx 30\%$ of the total soil community (Baubin et al., 2019) but was undetected in the active community of the dry season by any of the methods used here. These results suggest that the DNA-based total community may differ from the RNA-based community and thus cannot be used as a reliable reference for diversity. Furthermore, our results underline a need to standardize and specify the RT conditions that allow cross-study comparisons. The scale of the effect of RT conditions on the RNA-based community might vary with a studied biome. Depending on obvious factors such as GC content (discussed above), as well as poorly studied factors such as ribosomal post-transcriptional modifications. Therefore, we recommend verifying each case separately before attempting a cross-study comparison.

CONCLUSION

We have tested commonly utilized RT enzymes at assorted temperatures and observed marked differences in the output community structure. These differences were attributed to RT type and reaction conditions. We suggest that RT reaction conditions may dictate the diversity of a given community and therefore the exact conditions should be detailed in full [i.e., the common notation, "according to manufacturer instructions" does not provide sufficient information (Table 1)]. Furthermore, RT should be performed at a sufficiently high temperature to minimize the G-C bias, preferably at 55°C . Lastly, we suggest that the same RT enzyme should be used across comparable studies, since we detected discrepancies between RT enzymes performing at equivalent conditions (Figure 2, b). Here, we highlight, for the first time, the need for standardisation and careful consideration of RT reaction conditions in studies describing ribosome-based diversity and community composition.

ACKNOWLEDGEMENT

The authors would like to give thanks to the help and advice of Prof. Shula Michaeli from the Mina an Everard Faculty of Life Sciences. We thank Dr. Lusine Ghazaryan from the Zuckerberg Institute for Water Research for her help with the sequencing preparation, and Rachel Lichtenberg for her invaluable advice on English stylistics and grammar.

REFERENCES

- Angel, R. (2012). Total Nucleic Acid Extraction from Soil. *Protocol Exchange*.
- Angel, R. and Conrad, R. (2013). Elucidating the microbial resuscitation cascade in biological soil crusts following a simulated rain event. *Environmental Microbiology*, 15:2799–2815.
- Angel, R., Pasternak, Z., Soares, M. I. M., Conrad, R., and Gillor, O. (2013). Active and total prokaryotic communities in dryland soils. *FEMS Microbiology Ecology*, 86(1):130–138.
- Baldrian, P., Kolář, M., Štursová, M., Kopecký, J., Valášková, V., Větrovský, T., Žifčáková, L., Šnajdr, J., Rídl, J., Vlček, Č., and Voříšková, J. (2012). Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *ISME Journal*, 6(2):248–258.
- Barnard, R. L., Osborne, C. a., and Firestone, M. K. (2015). Changing precipitation pattern alters soil microbial community response to wet-up under a Mediterranean-type climate. *The ISME Journal*, 9(4):946–957.
- Baubin, C., Farrell, A. M., Šťovíček, A., Ghazaryan, L., Giladi, I., and Gillor, O. (2019). Seasonal and spatial variability in total and active bacterial communities from desert soil. *Pedobiologia*, 74(February):7–14.
- Blagodatskaya, E. and Kuzyakov, Y. (2013). Active microorganisms in soil: Critical review of estimation criteria and approaches. *Soil Biology and Biochemistry*, 67:192–211.
- Blazewicz, S. J., Barnard, R. L., Daly, R. a., and Firestone, M. K. (2013). Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses. *The ISME journal*, 7(11):2061–2068.

- 297 Bustin, S. A. and Nolan, T. (2004). Pitfalls of quantitative real- time reverse-transcription polymerase
298 chain reaction. *Journal of Biomolecular Techniques*, 15(3):155–166.
- 299 Bustin, S. A. and Nolan, T. (2017). Talking the talk, but not walking the walk: RT-qPCR as a paradigm
300 for the lack of reproducibility in molecular research. *European Journal of Clinical Investigation*,
301 47(10):756–774.
- 302 Carini, P., Marsden, P. J., Leff, J. W., Morgan, E. E., Strickland, M. S., and Fierer, N. (2016). Relic DNA
303 is abundant in soil and obscures estimates of soil microbial diversity. *Nature Microbiology*, 2(16242).
- 304 Carson, J. K., Gonzalez-Quinones, V., Murphy, D. V., Hinz, C., Shaw, J. A., and Gleeson, D. B. (2010).
305 Low Pore Connectivity Increases Bacterial Diversity in Soil. *Applied and Environmental Microbiology*,
306 76(12):3936–3942.
- 307 Che, R., Jinling, Q., Tahmasbian, I., Wang, F., Zhou, S., Xu, Z., Cui, X., and Che, R. (2018). Litter
308 amendment rather than phosphorus can dramatically change inorganic nitrogen pools in a degraded
309 grassland soil by affecting nitrogen-cycling microbes. *Soil biology & biochemistry*, v. 120:pp. 145–
310 152–2018 v.120.
- 311 Creer, S., Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W. K., Potter, C., and Bik, H. M.
312 (2016). The ecologist's field guide to sequence-based identification of biodiversity. *Methods in Ecology
313 and Evolution*, 7(9):1008–1018.
- 314 Curry, J., McHale, C., and Smith, M. T. (2002). Low efficiency of the moloney murine leukemia virus
315 reverse transcriptase during reverse transcription of rare t(8;21) fusion gene transcripts. *BioTechniques*,
316 32(4):768–775.
- 317 Degelmann, D. M., Kolb, S., Dumont, M., Murrell, J. C., and Drake, H. L. (2009). Enterobacteriaceae
318 facilitate the anaerobic degradation of glucose by a forest soil. *FEMS Microbiology Ecology*, 68(3):312–
319 319.
- 320 Dlott, G., Maul, J. E., Buyer, J., and Yarwood, S. (2015). Microbial rRNA: RDNA gene ratios may be
321 unexpectedly low due to extracellular DNA preservation in soils. *Journal of Microbiological Methods*,
322 115:112–120.
- 323 Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature
324 methods*, 10(10):996–8.
- 325 Forney, L. J., Zhou, X., and Brown, C. J. (2004). Molecular microbial ecology: Land of the one-eyed
326 king. *Current Opinion in Microbiology*, 7(3):210–220.
- 327 Freedman, Z. B., Romanowicz, K. J., Upchurch, R. A., and Zak, D. R. (2015). Differential responses of
328 total and active soil microbial communities to long-term experimental N deposition. *Soil Biology and
329 Biochemistry*, 90:275–282.
- 330 García, R., Bælum, J., Fredslund, L., Santorum, P., and Jacobsen, C. S. (2010). Influence of temperature
331 and predation on survival of salmonella enterica serovar typhimurium and expression of invA in soil
332 and manure-amended soil. *Applied and Environmental Microbiology*, 76(15):5025–5031.
- 333 Hansen, K. D., Brenner, S. E., and Dudoit, S. (2010). Biases in Illumina transcriptome sequencing caused
334 by random hexamer priming. *Nucleic Acids Research*, 38(12):1–7.
- 335 Huang, K., Chen, C., Zhang, J., Tang, Z., Shen, Q., Rosen, B. P., and Zhao, F.-J. (2016). Efficient Arsenic
336 Methylation and Volatilization Mediated by a Novel Bacterium from an Arsenic-Contaminated Paddy
337 Soil. *Environ. Sci. Technol.*, 50(12):6389–6396.
- 338 Hunter, J. D. (2007). Matplotlib: A 2d graphics environment. *Computing In Science & Engineering*,
339 9(3):90–95.
- 340 Joergensen, R. G. and Wichern, F. (2018). Alive and kicking: Why dormant soil microorganisms matter.
341 *Soil Biology and Biochemistry*, 116(October 2017):419–430.
- 342 Jurburg, S. D., Nunes, I., Brejnrod, A., Jacquiod, S., Priemé, A., Sørensen, S. J., Van Elsas, J. D.,
343 and Salles, J. F. (2017). Legacy effects on the recovery of soil bacterial communities from extreme
344 temperature perturbation. *Frontiers in Microbiology*, 8(SEP):1–13.
- 345 Ke, X., Lu, W., and Conrad, R. (2015). High Oxygen Concentration Increases the Abundance and Activity
346 of Bacterial Rather than Archaeal Nitrifiers in Rice Field Soil. *Microbial Ecology*, 70(4):961–970.
- 347 Klein, A. M., Bohannan, B. J., Jaffe, D. A., Levin, D. A., and Green, J. L. (2016). Molecular evidence for
348 metabolically active bacteria in the atmosphere. *Frontiers in Microbiology*, 7(MAY):1–11.
- 349 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glöckner, F. O. (2013). Eval-
350 uation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-
351 based diversity studies. *Nucleic acids research*, 41:e1.

- 352 Kwon, S., Lee, B., and Yoon, S. (2014). CASPER: Context-aware scheme for paired-end reads from
353 high-throughput amplicon sequencing. *BMC Bioinformatics*, 15(9):S10.
- 354 Lennon, J. T. and Jones, S. E. (2011). Microbial seed banks: the ecological and evolutionary implications
355 of dormancy. *Nature reviews. Microbiology*, 9(2):119–30.
- 356 Lever, M. A., Torti, A., Eickenbusch, P., Michaud, A. B., Šantl-Temkiv, T., and Jørgensen, B. B. (2015).
357 A modular method for the extraction of DNA and RNA, and the separation of DNA pools from diverse
358 environmental sample types. *Frontiers in Microbiology*, 6(MAY).
- 359 Lillis, L., Doyle, E., and Clipson, N. (2009). Comparison of DNA- and RNA-based bacterial community
360 structures in soil exposed to 2,4-dichlorophenol. *Journal of Applied Microbiology*, 107(6):1883–1893.
- 361 McCarthy, A., Chiang, E., Schmidt, M. L., and Deneff, V. J. (2015). RNA preservation agents and nucleic
362 acid extraction method bias perceived bacterial community composition. *PLoS ONE*, 10(3):1–14.
- 363 Mohr, S., Ghanem, E., Smigh, W., Sheeter, D., Qin, Y., King, O., Polioudakis, D., Iyer, V. R., Hunicke-
364 Smigh, S., Swamy, S., Kuersten, S., and Lambowitz, A. M. (2013). Thermostable group II intron reverse
365 transcriptase fusion proteins and their use in cDNA synthesis and next-generation RNA sequencing.
366 *Rna*, 19(7):958–970.
- 367 Nacke, H., Fischer, C., Thürmer, A., Meinicke, P., and Daniel, R. (2014). Land Use Type Significantly
368 Affects Microbial Gene Transcription in Soil. *Microbial Ecology*, 67(4):919–930.
- 369 Nunes, I., Jurburg, S., Jacquiod, S., Brejnrod, A., Falcão Salles, J., Priemé, A., and Sørensen, S. J. (2018).
370 Soil bacteria show different tolerance ranges to an unprecedented disturbance. *Biology and Fertility of*
371 *Soils*, 54(2):189–202.
- 372 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara,
373 R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H. (2018). *vegan*:
374 *Community Ecology Package*.
- 375 Pan, Y., Bodrossy, L., Frenzel, P., Hestnes, A. G., Krause, S., Lüke, C., Meima-Franke, M., Siljanen,
376 H., Svenning, M. M., and Bodelier, P. L. (2010). Impacts of inter- and intralaboratory variations
377 on the reproducibility of microbial community analyses. *Applied and Environmental Microbiology*,
378 76(22):7451–7458.
- 379 Paulin, M. M., Nicolaisen, M. H., Jacobsen, C. S., Gimsing, A. L., Sørensen, J., and Bælum, J. (2013).
380 Improving Griffith's protocol for co-extraction of microbial DNA and RNA in adsorptive soils. *Soil*
381 *Biology and Biochemistry*, 63:37–49.
- 382 Pfeiffer, S., Pastar, M., Mitter, B., Lippert, K., Hackl, E., Lojan, P., Oswald, A., and Sessitsch, A.
383 (2014). Improved group-specific primers based on the full SILVA 16S rRNA gene reference database.
384 *Environmental Microbiology*, 16(8):2389–2407.
- 385 Placella, S. A., Brodie, E. L., and Firestone, M. K. (2012). Rainfall-induced carbon dioxide pulses
386 result from sequential resuscitation of phylogenetically clustered microbial groups. *Proceedings of the*
387 *National Academy of Sciences of the United States of America*, 109(27):10931–10936.
- 388 Pollock, J., Laura, G., Trong, W., and Mick, W. (2018). The Madness of Microbiome : Attempting To
389 Find Consensus. *Applied and Environmental Microbiology*, 84(7):1–12.
- 390 Pratscher, J., Dumont, M. G., and Conrad, R. (2011). Ammonia oxidation coupled to CO₂ fixation
391 by archaea and bacteria in an agricultural soil. *Proceedings of the National Academy of Sciences*,
392 108(10):4170–4175.
- 393 Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., and Glöckner, F. O. (2007).
394 SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence
395 data compatible with ARB. *Nucleic acids research*, 35(21):7188–96.
- 396 Quammen, D. (2018). *The Tangled Tree: A Radical New History of Life*. Simon and Schuster.
- 397 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.
398 (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based
399 tools. *Nucleic acids research*, 41:D590–D596.
- 400 R Core Team (2018). *R: A Language and Environment for Statistical Computing*. R Foundation for
401 Statistical Computing, Vienna, Austria.
- 402 Romanowicz, K. J., Freedman, Z. B., Upchurch, R. A., Argiroff, W. A., and Zak, D. R. (2016). Active
403 microorganisms in forest soils differ from the total community yet are shaped by the same environmental
404 factors: The influence of pH and soil moisture. *FEMS Microbiology Ecology*, 92(10):1–9.
- 405 Schwartz, S. and Motorin, Y. (2017). Next-generation sequencing technologies for detection of modified
406 nucleotides in RNAs. *RNA Biology*, 14(9):1124–1137.

- 407 Segev, E., Rosenberg, A., Mamou, G., Sinai, L., and Ben-Yehuda, S. (2013). Molecular kinetics of
408 reviving bacterial spores. *Journal of Bacteriology*, 195(9):1875–1882.
- 409 Shannon, C. E. and Weaver, W. (1949). *The mathematical theory of information*, volume 97.
- 410 Sieber, M. W., Recknagel, P., Glaser, F., Witte, O. W., Bauer, M., Claus, R. A., and Frahm, C. (2010).
411 Substantial performance discrepancies among commercially available kits for reverse transcription quan-
412 titative polymerase chain reaction: A systematic comparative investigator-driven approach. *Analytical*
413 *Biochemistry*, 401(2):303–311.
- 414 Ståhlberg, A., Håkansson, J., Xian, X., Semb, H., and Kubista, M. (2004a). Properties of the Reverse
415 Transcription Reaction in mRNA Quantification. *Clinical Chemistry*, 50(3):509–515.
- 416 Ståhlberg, A., Kubista, M., and Michael, P. (2004b). Comparison of Reverse Transcriptases in Gene
417 Expres- sion Analysis. *Clinical Chemistry*, 50(9):1675–1678.
- 418 Steven, B., Hesse, C., Soghigian, J., Gallegos-Graves, L. V., and Dunbar, J. (2017). Simulated rRNA/DNA
419 Ratios Show Potential To Misclassify Active Populations as Dormant. *Applied and Environmental*
420 *Microbiology*, 83(11):1–11.
- 421 Van Der Walt, S., Colbert, S. C., and Varoquaux, G. (2011). The NumPy array: A structure for efficient
422 numerical computation. *Computing in Science and Engineering*, 13(2):22–30.
- 423 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
- 424 Yilmaz, S. L., Okten, H. E., and Noguera, D. R. (2006). Making All Parts of the 16S rRNA of. *Applied*
425 *and environmental microbiology*, 72(1):733–744.
- 426 Zhang, J. and Byrne, C. D. (2015). Differential priming of RNA templates during cDNA synthesis
427 markedly affects both accuracy and reproducibility of quantitative competitive reverse-transcriptase
428 PCR. *Biochemical Journal*, 337(2):231–241.