

A peer-reviewed version of this preprint was published in PeerJ on 10 August 2017.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.3612) (peerj.com/articles/3612), which is the preferred citable publication unless you specifically need to cite this preprint.

Beckham JL, Atkinson S. 2017. An updated understanding of Texas bumble bee (Hymenoptera: Apidae) species presence and potential distributions in Texas, USA. PeerJ 5:e3612
<https://doi.org/10.7717/peerj.3612>

An updated understanding of Texas bumble bee (Hymenoptera: Apidae) species presence and potential distributions in Texas, USA

Jessica L Beckham ^{Corresp., 1}, Samuel Atkinson ²

¹ Advanced Environmental Research Institute, University of North Texas, Denton, Texas, United States

² Advanced Environmental Research Institute and Department of Biological Sciences, University of North Texas, Denton, Texas, United States

Corresponding Author: Jessica L Beckham

Email address: jessica.beckham@unt.edu

Texas is the second largest state in the United States of America, and the largest state in the contiguous USA at nearly 700,000 sq. km. Several Texas bumble bee species have shown evidence of declines in portions of their continental ranges, and conservation initiatives targeting these species will be most effective if species distributions are well established. To date, statewide bumble bee distributions for Texas have been inferred primarily from specimen records housed in natural history collections. To improve upon these maps, and help inform conservation decisions, this research aimed to (1) update existing Texas bumble bee presence databases to include recent (2007-2016) data from citizen science repositories and targeted field studies, (2) model statewide species distributions of the most common bumble bee species in Texas using MaxEnt, and (3) identify conservation target areas for the state that are most likely to contain habitat suitable for multiple declining species. The resulting Texas bumble bee database is comprised of 3,580 records, to include previously compiled museum records dating from 1897, recent field survey data, and vetted records from citizen science repositories. These data yielded an updated state species list that includes 11 species, as well as species distribution models (SDMs) for the most common Texas bumble bee species, including two that have shown evidence of range-wide declines: *B. fraternus* (Smith, 1854) and *B. pensylvanicus* (DeGeer, 1773). Based on analyses of these models, we have identified conservation priority areas within the Texas Cross Timbers, Texas Blackland Prairies, and East Central Texas Plains ecoregions where suitable habitat for both *B. fraternus* and *B. pensylvanicus* are highly likely to co-occur.

An updated understanding of Texas bumble bee (Hymenoptera: Apidae) species presence and potential distributions in Texas, USA

Jessica L. Beckham¹, Sam F. Atkinson²

¹ Advanced Environmental Research Institute, University of North Texas, Denton, TX, USA

² Advanced Environmental Research Institute and Department of Biological Sciences, University of North Texas, Denton, TX, USA

Corresponding Author:

Jessica L. Beckham

Email address: jessica.beckham@unt.edu

15 **Abstract:**

16 Texas is the second largest state in the United States of America, and the largest state in the
 17 contiguous USA at nearly 700,000 sq. km. Several Texas bumble bee species have shown
 18 evidence of declines in portions of their continental ranges, and conservation initiatives targeting
 19 these species will be most effective if species distributions are well established. To date,
 20 statewide bumble bee distributions for Texas have been inferred primarily from specimen
 21 records housed in natural history collections. To improve upon these maps, and help inform
 22 conservation decisions, this research aimed to (1) update existing Texas bumble bee presence
 23 databases to include recent (2007-2016) data from citizen science repositories and targeted field
 24 studies, (2) model statewide species distributions of the most common bumble bee species in
 25 Texas using MaxEnt, and (3) identify conservation target areas for the state that are most likely
 26 to contain habitat suitable for multiple declining species. The resulting Texas bumble bee
 27 database is comprised of 3,580 records, to include previously compiled museum records dating
 28 from 1897, recent field survey data, and vetted records from citizen science repositories. These
 29 data yielded an updated state species list that includes 11 species, as well as species distribution
 30 models (SDMs) for the most common Texas bumble bee species, including two that have shown
 31 evidence of range-wide declines: *B. fraternus* (Smith, 1854) and *B. pensylvanicus* (DeGeer,
 32 1773). Based on analyses of these models, we have identified conservation priority areas within
 33 the Texas Cross Timbers, Texas Blackland Prairies, and East Central Texas Plains ecoregions
 34 where suitable habitat for both *B. fraternus* and *B. pensylvanicus* are highly likely to co-occur.

Introduction:

Bumble bees (*Bombus* spp.) are an important group of insect pollinators that provide sustaining pollination services for both agricultural systems and biological communities (Corbet et al., 1991; Klein et al., 2007; Potts et al., 2010; Garibaldi et al., 2013). Unfortunately, bumble bees face many modern challenges, including those posed by pesticide use (Gill et al., 2012; Whitehorn et al., 2012; Rundlöf et al., 2015), climate change (Kerr et al., 2015), disease (Szabo et al., 2012), and habitat loss (Kearns et al., 1998; Gixti et al., 2009; Winfree et al., 2009). Consequently, bumble bee losses have been noted across Europe (Carvell, 2002; Sarospataki et al., 2005; Goulson et al., 2006; Fitzpatrick et al., 2007; Kosior et al., 2007; Goulson, 2010), Asia (Yang, 1999; Matsumura et al., 2004; Inoue et al., 2008; Xie et al., 2008)), and North America (Thorp, 2005; Thorp & Shepherd, 2005; Colla & Packer, 2008; Gixti et al., 2009; Colla & Ratti, 2010; Cameron et al., 2011; Colla et al., 2012). Furthermore, evidence presented by Bartomeus et al. (2013) suggests that bumble bees are among the most vulnerable of North American bee taxa.

Because bumble bees are the primary pollinators of several agricultural crops, declines in bumble bee populations may threaten the permanence of foods such as blueberries, tomatoes, and peppers (Hatfield et al., 2012; Shipp et al., 1994; Whittington & Winston, 2004). Additionally, research has indicated that the loss of bumble bees from wild ecosystems could result in a pronounced decline in overall plant diversity (Memmott et al., 2004), limited seed production by native plants, and less effective pollination by other pollinators (Brosi & Briggs, 2013).

Historically, eight species of bumble bees have been documented in Texas (Franklin, 1913; Warriner, 2012): *B. auricomus* (Robertson, 1903), *B. bimaculatus* Cresson, 1863, *B.*

fervidus (Fabricius, 1798), *B. fraternus* (Smith, 1854), *B. griseocollis* (DeGeer, 1773), *B. impatiens* Cresson, 1863, *B. pensylvanicus* (DeGeer, 1773), and *B. variabilis* (Cresson, 1872). Additionally, *B. pensylvanicus sonorus*, sometimes classified as the distinct species *B. sonorus* Say, 1837 (Franklin, 1913; Stephen, 1957; Thorp et al., 1983; Warriner, 2012), has been documented in Texas. For the purposes of this research we have treated this taxon as a western morphological variant of *B. pensylvanicus* following Milliron (1973), Labougle (1990), Poole (1996), and Williams et al. (2014), and refer to it hereafter as *B. pensylvanicus sonorus*. However, it should be noted that the taxonomic status of this taxon is still debatable, and genetic studies are warranted to address this question.

Though bumble bee declines can be difficult to assess owing to a lack of long-term monitoring data (Berenbaum et al., 2007), some studies have suggested regional and national declines for several species whose ranges extend into Texas, including *B. auricomus* (Colla et al., 2012), *B. fervidus* (Colla & Packer, 2008; Colla et al., 2011; Colla et al., 2012), *B. fraternus* (Colla & Packer, 2008; Grixti et al., 2009; Hatfield et al., 2014), *B. pensylvanicus* (Berenbaum et al., 2007; Colla & Packer, 2008; Grixti et al., 2009; Cameron et al., 2011; Colla et al., 2011; Colla et al., 2012), and *B. variabilis* (Grixti et al., 2009; Colla et al., 2011; Colla et al., 2012). To our knowledge, the only recent attempt to assess the current status of Texas bumble bee populations took place in a 24-county region of northeast Texas, where five species had historically been found (Beckham et al., 2016). In that study, the results of targeted field surveys (2010-2014) were compared to natural history specimen records. While the researchers did not reconfirm presence of the regionally rare *B. bimaculatus* or *B. variabilis* in northeast Texas, they showed that the current relative abundances of the potentially declining *B. fraternus* and *B. pensylvanicus* were equivalent to historic levels in the study region, as was that of the nationally

stable *B. griseocollis*. Nevertheless, compelling evidence for bumble bee declines across North America has raised awareness of the need for conservation measures. In Texas this has led to three taxa (*B. pensylvanicus*, *B. pensylvanicus sonorus*, and *B. variabilis*) being designated as species of greatest conservation need in the Texas Conservation Action Plan (Texas Parks & Wildlife Department, 2012).

An effective species conservation program requires delineation of the distribution of the species of interest (Eken et al., 2004), as well as an understanding of their habitat. Prior to the current study, efforts to establish Texas bumble bee distributions included the compilation and mapping of Texas specimen data from natural history collections by Warriner (2012) and targeted field studies in the northeastern portion of the state (Beckham et al., 2016). Williams et al. (2014) also modeled range-wide species distributions for North American bumble bees, to include Texas species. While informative and foundational, these studies left some question as to the fine-scale distributions of Texas bumble bee species, as well as gaps in data for portions of the state.

Unlike many insect taxa, bumble bees can often be identified to species in the field and in photographs, and so data collected by citizen scientists can provide invaluable information for bumble bee research. These data have been used to monitor populations and nesting densities (Osborne et al., 2008; Lye et al., 2011), and have served as presence data for species distribution models (SDMs) (Kadoya et al., 2009). In recent efforts to spread awareness and fill gaps in presence data, two citizen scientist projects were established specifically for Texas volunteers to submit their bumble bee sightings, in the form of photographs with locality data: the Texas Bumblebees Facebook page, founded in 2011, and the iNaturalist Bees and Wasps of Texas project, founded in 2014. These repositories now contain hundreds of sightings recorded by

numerous volunteers from across the state that can be used as additional presence data for SDMs, complementing those provided by natural history collection records and field surveys.

SDMs estimate the geographic distribution of a species using field observations and associated environmental predictor variables. Over the last few decades, SDMs have become increasingly useful tools for conservation planning (Guisan & Thuiller, 2005). The Maximum Entropy (MaxEnt) algorithm is attractive because its implementation is straightforward and makes use of presence-only data, along with environmental variables associated with presence localities, to produce robust distribution models (Phillips et al., 2004; Phillips et al., 2006).

In recent years MaxEnt has successfully been used for a variety of applications related to bumble bees. To aid in conservation planning, Koch & Strange (2009) used historic specimen data to construct range maps for four North American species, and Penado et al. (2016) predicted the potential ranges of rare species in the Iberian Peninsula, also revealing the possibility of undiscovered populations in mountainous areas. Others have used MaxEnt to evaluate species trends, including Cameron et al. (2011), who assessed the rangewide persistence of eight North American species, and Dellicour et al. (2017), who inferred past and present distributions of European bumble bees to investigate range shifts. And, to aid in invasive species management decisions, Kadoya et al. (2009) implemented citizen science presence data in MaxEnt to predict the potential for invasion by *B. terrestris* in Japan. These studies underscore the utility of models produced by MaxEnt for bumble bee species policy and planning purposes.

In an effort to update and increase the understanding of statewide bumble bee species presence in Texas, and to inform the planning and implementation of state conservation actions, this study's aims were to:

1) Compile statewide bumble bee presence records from novel data sources (citizen science repositories and recent field studies) and natural history collections to produce an updated database of known bumble bee presence in Texas.

2) Create SDMs using MaxEnt to illustrate the potential ranges of common bumble bee species in Texas.

3) Identify conservation target areas that are likely to contain multiple declining bumble bee species.

Materials & Methods:

Species Presence Data—

Bumble bee presence data were gathered from natural history museum collections, citizen science repositories, and recent field collection data. These were compiled into one database for mapping and species distribution modeling purposes.

Previously compiled natural history collection records (Warriner, 2012; L. Richardson, unpublished dataset) included data from the following institutions: Elm Fork Natural Heritage Museum, Texas A&M University Insect Collection, Texas Memorial Museum, Illinois Natural History Survey Insect Collection, Cornell University Insect Collection, Florida State University Collection of Arthropods, Mississippi State University Entomological Museum, K.C. Emerson Entomology Museum, Purdue Entomological Research Collection, Smithsonian Natural History Museum, University of Arkansas Arthropod Museum, University of Georgia Collection of Arthropods, University of Michigan Museum of Zoology, University of Minnesota Insect Collection, American Museum of Natural History, Canadian National Collection, Essig Museum of Entomology, Los Angeles County Museum, Ohio State University, Bohart Museum of

Entomology, University of Colorado Museum of Natural History, California State Collection of
Arthropods, University of California Riverside Entomology Research Museum, Lyman
Entomological Collection, Bee Biology and Systematics Laboratory, USGS Patuxent Wildlife
Research Center, and the Yale Peabody Museum. When possible, records lacking geographic
coordinates were georeferenced using Google Earth; record localities given as county names
were georeferenced to coordinates corresponding to the county seat.

Two citizen scientist repositories, iNaturalist and the Texas Bumblebees Facebook page
(hereafter referred to as “Texas Bumblebees”), were also mined for presence records. Only
records that included photographs that could be identified to species, as well as location data,
were incorporated into the modeling database. The iNaturalist platform allows contributors to
include locality coordinates with their submissions; when available, these coordinates were
included in the presence database. Records lacking coordinates, as well as all submissions from
Texas Bumblebees, were georeferenced based on locality information provided by the observer
using Google Earth. Several citizen scientist records were also directly submitted as photographs
to J. Beckham. Species determinations followed Williams et al. (2008) and Colla et al. (2011).
Additionally, field collection data were gathered from recent research projects performed by J.
Beckham and M. Warriner in 2010-2016 (Beckham et al., 2016; Beckham, unpublished dataset).

Correcting for Sampling Bias—

A fundamental assumption in MaxEnt is that the presence data upon which models are
built have been acquired from random sampling across the study area. However, museum and
citizen science data are not typically collected in a systematic fashion, resulting in spatially
biased data. Running MaxEnt without correcting for this sampling bias results in overfitting of
models to reflect survey effort rather than actual species distributions. When sample size is

sufficient, spatial filtering is recommended to minimize errors of omission and commission in MaxEnt resulting from using spatially biased presence data (Kramer-Schadt et al., 2013). Additionally, because MaxEnt uses presence-only data to produce SDMs, background points are selected to contrast against the presence locations. By default, MaxEnt assigns every pixel the same probability of being selected as a background point. Selecting background points so that they carry the same bias as (unfiltered) presence data aids in the production of more accurate distribution models (Phillips et al., 2009; Merow et al., 2013). In order to achieve this end, a bias grid with cell values weighted to reflect non-uniform sampling effort (following Elith et al. 2010) can be implemented in MaxEnt for biased apriori background point selection.

The presence data included in the present study were biased towards heavily populated areas, especially the Dallas-Fort Worth metroplex, San Antonio, Houston, and Austin. To account for this bias, presence data were spatially filtered and a bias grid was produced to modify the random selection of background points using tools from the SDMtoolbox (Brown, 2014) implemented in ArcMap 10.2.2 (ESRI, Redlands, CA). Presence data were spatially filtered using the SDMtoolbox *rarefy occurrence data for SDMs* tool that reduces spatial autocorrelation by removing duplicate occurrence points and reduces occurrence records to a single point within a specified area; for our data we used a 15-km² resolution. The bias grid was created using the SDMtoolbox *Gaussian kernel density of sampling localities* tool loaded with presence points from all species; a sampling bias distance of 30 km was chosen. The resulting spatially filtered presence data and bias file were utilized in MaxEnt.

Preparation of Environmental Variables—

Environmental layers of 19 derived bioclimatic variables were downloaded from the WorldClim global climate database (Hijmans et al., 2005, available at www.worldclim.org) at a

resolution of 10 minutes ($\sim 344 \text{ km}^2$) as ESRI grids. This resolution was chosen because it allowed us to include lower resolution data that had been georeferenced to county. All rasters were clipped to match the boundaries of the state of Texas using the *extract by mask* tool in ArcMap 10.2.2; the cartographic boundary file for the mask was downloaded from <https://www.census.gov/geo/maps-data/data/tiger-cart-boundary.html> and a shapefile of Texas was produced and projected to WGS 1984. All clipped environmental rasters were converted to ASCII files in ArcMap for use in MaxEnt.

It is recommended that correlation be minimized between predictor variables implemented in MaxEnt, as would be done for a traditional statistical model (Merow et al., 2013). A Pearson's correlation analysis was performed on all 19 layers using the *correlation and summary stats* tool available for ArcMap in the SDMtoolbox, and pairs of environmental layers with correlation coefficients >0.80 were identified (Table S1). Variables were ranked prior to analyses according to perceived ecological importance, with derived variables (BIO02, BIO03, BIO04, BIO07, and BIO15) being given lowest ranking. Variables pertaining to the warmest and wettest months and quarters were ranked above similar variables involving coldest and driest months and quarters given that bumble bees are very generally more active in warmer, wetter months. When variable importance could not be intuited based on the aforementioned criteria, ranking was based upon the arbitrary WorldClim code. The lower-ranking predictor of each correlated pair was removed, resulting in a total of ten environmental predictors being maintained for species distribution analyses (Table 1).

[TABLE 1 approximately HERE]

216 *Production of Species Distribution Models (SDMs)—*

217 Input data for species distribution modeling in MaxEnt consisted of the aforementioned
 218 spatially filtered presence data (occurrence records) and the 10 environmental layers (predictor
 219 variables). The Gaussian kernel density bias file was also loaded to alter selection of background
 220 points to reflect sampling bias. MaxEnt was run for 100 replicates for each species with default
 221 settings, except as follows: jackknifing was selected to measure variable importance, 30% test
 222 percentage was implemented, and 5000 max iterations were allowed. The average logistic output
 223 over 100 replicates was calculated in MaxEnt for each species, and resulting SDM rasters were
 224 visualized in ArcMap 10.2.2. MaxEnt's logistic output assigns each pixel in the study area a
 225 value between 0 and 1, and is the closest approximation of the probability of species presence
 226 (Elith et al., 2011). The default MaxEnt species prevalence value (an estimate of the probability
 227 of species presence at “typical” presence locations for the target species) was used to transform
 228 the raw MaxEnt output into the logistic output for each SDM. The reasoning behind using
 229 MaxEnt's default value of 0.5, which can be interpreted as there being a 50% chance of the
 230 target species being present in a suitable location, is that the actual species prevalence values are
 231 rarely known and hard to approximate. Thus, calculated probability values from MaxEnt's
 232 logistic output are interpreted not as the probability of species occurrence, but instead as the
 233 probability of suitable habitat being present for the target species. For detailed mathematical
 234 explanations of MaxEnt's logistic output see Elith et al. (2011) and Phillips et al. (2006).

235 *SDM Raster Processing and Analyses—*

236 For visualization and analyses purposes, SDM rasters were reclassified using the
 237 *Reclassify* tool in ArcMap's Spatial Analyst toolbox from the continuous scale of 0 – 1
 238 probability calculated in MaxEnt to a categorical 0 – 2 scale, with 0 being low probability of

suitable habitat being present (< 0.25), 1 being medium probability ($0.25 - 0.49$), and 2 being high probability (> 0.5). These rasters were then displayed with the underlying EPA Level III ecoregions (spatial layer downloaded from <https://www.epa.gov/eco-research/ecoregion-download-files-state-region-6>; see Figure 1). Identifying the ecoregions associated with each bumble bee species will help inform decisions regarding the planning and implementation of conservation plans. Furthermore, because funds and manpower dedicated to conservation can be limited, we identified areas where suitable habitat was likely to be present for two of the declining species found in Texas (*B. fraternus* and *B. pensylvanicus*). In order to do this, the *Combine* tool in ArcMap's Spatial Analyst toolbox was used to overlay the reclassified *B. fraternus* and *B. pensylvanicus* rasters, and cells with high (>0.5) probabilities of suitable habitat for both were identified.

[FIGURE 1 approximately HERE]

Model Validation—

The accuracy of each SDM was evaluated using values of the area under the Receiver Operating Characteristic (ROC) curve, or AUC value, plotted and calculated in MaxEnt based on the training and test data. These values convey the predictive performance of the model as compared to a selection of random background points, and can be interpreted as the probability that any presence site is ranked higher in terms of habitat suitability for the target species than a random background site (Merow et al., 2013). An AUC value of 0.5 reflects a model that is no better than random, while an AUC of 1 would be considered a “perfect” model. We trained each model on 70% of the presence data points, and then tested it with the remaining 30% of the data, resulting in both training and test AUC values for each target species' SDM.

We then tested the null hypothesis that the average test AUC values generated for each species were significantly different from those predicted by a null model, as described in Raes & ter Steege (2007). We generated a null distribution for each species by first randomly drawing 999 sets of random collection localities from the study area; the number of points in each set was equal to the corresponding number of spatially filtered presence points used for the target species' SDM. A model was then produced in MaxEnt for each set of points, with all settings identical to those used to produce the actual SDMs. The test AUC was calculated for each null SDM, and a null distribution of the 999 test AUCs for the random replicates was produced. The actual model's average test AUC was then compared to the upper 95% confidence limit of the null distribution to test the null hypothesis (alpha level of 0.05). Random draws were performed in R (version 3.3.3, <https://cran.r-project.org/index.html>, 2017) using the *RandomPoints* function contained in the *dismo* package (Hijmans et al., 2017), and confidence limits were calculated in SAS (version 9.4, SAS Institute, 2013).

Results:

Species Presence—

A total of 3,580 Texas records spanning 1897-2016 were compiled from museum collections, citizen science repositories and recent fieldwork (Table 2). These data included 747 previously unpublished citizen science records from the past ten years that were recorded in iNaturalist and Texas Bumblebees (Figure 2). Eleven species were identified from these records, including one specimen of *B. appositus* Cresson, 1878, five specimens of *B. morrisoni* Cresson, 1878, and two specimens of *B. nevadensis* Cresson, 1874, which had not previously been documented in the list of Texas bumble bee species compiled by Warriner (2012). The updated species list for Texas, along with numbers of specimens records, is as follows: *B. appositus*

(n=1), *B. auricomus* (n=13), *B. bimaculatus* (n=5), *B. fervidus* (n=4), *B. fraternus* (n= 239), *B. griseocollis* (n=108), *B. impatiens* (n=173), *B. morrisoni* (n=5), *B. nevadensis* (n=2), *B. pensylvanicus* (n=3,010), and *B. variabilis* (n=20). Notably, a subset of 314 *B. pensylvanicus* records were also logged as presenting the *sonorus* morphology.

[TABLE 2 approximately HERE]

[FIGURE 2 approximately HERE]

Species Distribution Models—

Of the twelve species represented by the data, four species comprised 99% of the records, and so we only modeled the distributions for these most abundant species: *B. fraternus*, *B. griseocollis*, *B. impatiens*, and *B. pensylvanicus*. Sample sizes for the other species known from Texas were deemed insufficient for spatial filtering and modeling purposes. The logistic outputs of SDMs produced in MaxEnt, as well as the reclassified visualizations of these models, for *B. fraternus*, *B. griseocollis*, *B. impatiens*, and *B. pensylvanicus* are shown in Figures 3-6, respectively.

[FIGURES 3-6 approximately HERE]

The contribution of each bioclimatic variable to each SDM varied across the four species, as did the most important variables for each model. Annual precipitation (BIO12) was the top

contributor to SDMs for *B. griseocollis* (71.6%), *B. impatiens* (73.8%), and *B. pensylvanicus* (20.3%). Isothermality (BIO03), calculated as $[(\text{mean diurnal temperature range} / \text{mean annual temperature}) * 100]$ was the top contributor to the SDM for *B. fraternus* (51% contribution), and was the second and fifth most important variable for *B. griseocollis* (8.5%) and *B. impatiens* (2.1%) SDMs, respectively. The mean temperature of the driest quarter (BIO09) was also a top-five contributor for all models. The top five bioclimatic variables contributing to each SDM, as well as their percent contributions, are presented in Table 3.

[TABLE 3 approximately HERE]

To evaluate the validity of each SDM, we calculated both training and test AUC values on 70% and 30% of the presence data, respectively. As expected, test AUC values were lower than training AUCs, but all values were above 0.65. We then tested the hypothesis that the average test AUC for each SDM was significantly different from that of a random null distribution ($\alpha=0.05$). All average test AUC values from actual SDMs were significantly higher than the values calculated from the null distributions ($p<0.01$), and so we concluded that all models were statistically significant. The average training and test AUC values for each SDM, as well as the upper 95% confidence intervals of the null distributions, are contained in Table 3.

According to our results, *B. pensylvanicus* is both the most common and the most widespread bumble bee species in Texas, representing 84% of the specimens in this study. While it is most likely to be found in the central and eastern portions of the state, there have been

specimens recorded in every ecoregion. *B. fraternus* (~7% of specimens) is most likely to be found in the northeast portion of the state, but its range may extend to far north and far west Texas. By contrast, the ranges of both *B. impatiens* (5% of specimens) and *B. griseocollis* (3% of specimens) are currently limited primarily to the eastern portions of Texas.

Priority areas for actions targeting the conservation of declining species were identified by overlaying the reclassified maps of *B. fraternus* and *B. pensylvanicus* in ArcMap to ascertain whether there were areas of the state that carried high probabilities of suitable habitat being present for both species (Figure 7). This analysis revealed a region of northeast Texas that covers portions of the Cross Timbers, Texas Blackland Prairies, and East Central Texas Plains ecoregions where habitat is likely to be highly suitable for both species.

[FIGURE 7 approximately HERE]

Discussion:

The species presence records documented in this study included three new, albeit historically very rare, species from the list of Texas bumble bees previously published by Warriner (2012): *B. appositus* (n=1), *B. morrisoni* (n=5), and *B. nevadensis* (n=2). These records were retrieved from the database of museum records managed by Leif Richardson (Williams et al., 2014; L. Richardson, unpublished dataset), and records were validated with museums of origin. As such, we have revised the species list of Texas bumble bees to include 11 species.

In comparison to the Texas portion of the range map presented in Williams et al. (2014), our model for *B. fraternus* was somewhat contracted, lacking the high likelihood of presence

predicted by Williams et al. across the Western Gulf Coastal Plain, and throughout the Southwestern Tablelands and High Plains. By contrast, we identified novel presence records for both *B. impatiens* and *B. griseocollis* in the East Central Texas Plains and South Central Plains, resulting in a shift of the western edges of these species' Texas ranges from those predicted by Williams et al. (2014). Because *B. griseocollis* populations are considered stable across its range (Colla & Packer, 2008; Colla et al., 2010), and *B. impatiens* populations are considered stable (Cameron et al., 2011) or possibly expanding (Colla & Packer, 2008), our results suggest that Texas populations should be monitored for further evidence of range expansion for these species. The model of the range of *B. pensylvanicus* presented by Williams et al. (2014) is similar to our model, which is virtually statewide. Had we included presence records from the neighboring states of Louisiana, Oklahoma, and New Mexico, our models, especially the model for *B. fraternus*, might have been extended further into border ecoregions. Future work should include further improvement of SDMs for these species that include records from neighboring states.

One drawback to presence-only species distribution modeling is that models can be biased towards areas that contain more presence records (Kramer-Schadt et al., 2013). In the case of our data, presence records from areas with larger human populations (i.e., the Dallas-Fort Worth metroplex, Austin, San Antonio, and Houston) were more common than less populated areas of the state. In order to address this problem, presence records were spatially filtered and a Gaussian bias file was implemented in MaxEnt. However, these techniques did not completely resolve the issue of sampling bias, and individual species distributions were still biased towards highly populated areas. Conversely, SDMs for areas that were not well sampled, such as the Southwestern Tablelands and High Plains ecoregions located in north central Texas, predicted low or moderate likelihood of any bumble bee species presence. Syfert et al. (2013) similarly

showed that, save an intentional sampling strategy across the study area, SDMs will be somewhat biased towards clustered data. Thus, we consider our SDMs to be conservative, and suggest cautious interpretation of species presence in marginally sampled ecoregions. Targeted field surveys in these areas are needed to achieve both a more uniform statewide bumble bee sampling effort and less clustered data for SDMs.

Of the ten predictor variables included in the SDMs, we interpret our results to mean that annual precipitation, isothermality, and the mean temperature of the driest quarter are the most important bioclimatic factors for predicting bumble bee habitat suitability in Texas. Precipitation was the top variable contributing to the SDMS for three of the four species studied (*B. griseocollis*, *B. impatiens*, and *B. pensylvanicus*), and isothermality was the top contributor to the *B. fraternus* model, as well as a top-five contributor to two other SDMs. The mean temperature of the driest quarter was one of the top five environmental predictors for all SDMs. As global climate change threatens to disrupt temperature and precipitation patterns, the current models may change drastically. Furthermore, though these environmental factors are undoubtedly important for bumble bee habitat, it should be noted that other factors for which data were not available may also influence the suitability of habitat, such as pesticide use, available nesting and foraging sites, and habitat connectivity.

While multiple factors have been implicated in bumble bee declines, habitat loss, often associated with urban sprawl and agricultural intensification, is considered to be a leading cause (Kearns et al., 1998; Winfree et al., 2009). Bumble bee conservation initiatives designed to curb these declines should include actions that establish, enhance, and maintain habitat. The declining *B. fraternus* and *B. pensylvanicus* persist in northeast Texas, where a substantial area of native rangeland remains (Beckham et al., 2016); across the state, native rangelands comprise

approximately 63% of the state's nonfederal rural land (U.S. Department of Agriculture, 2013), and encouraging the conservation of these lands amidst rapid population growth and urbanization may prove beneficial to bumble bees. Additionally, practices such as targeted agri-environment schemes that increase floral biodiversity have been shown to increase bumble bee diversity and abundance in agricultural systems (Pywell et al. 2006; Carvell et al. 2007; Carvell et al. 2011) and, within urban environments, green spaces such as parks and community gardens can provide valuable habitats for bumble bees in otherwise unfavorable landscapes (McFrederick & LeBuhn, 2006; Ahrne et al., 2009; Goulson et al., 2010; Beckham, 2016). In Texas, where over 95% of land is privately owned (Texas Land Trends, 2014), the conservation of bumble bee species will require a multifaceted approach, including public education and outreach, as well as working with municipalities and private landowners to create and maintain pollinator habitat.

The results of our multi-species analyses suggest starting points for conservation programs when funding and manpower is limited. The target area includes portions of Texas' Cross Timbers, Blackland Prairies, and East Central Plains, where suitable habitat for both *B. fraternus* and *B. pensylvanicus* is highly likely to co-occur. Notably, this target area fully contains the Dallas-Fort Worth metroplex, a heavily urbanized area that saw a decrease in working lands acreage of almost 316,000 acres and whose human population grew by about 2.3 million people from 1997-2012 (Texas Land Trends, 2015). However, the conservation target area also contains surrounding rural areas across the region, and so our results have underscored the need for conservation strategies that will address both urban and rural environments.

Conclusions:

Our study has updated the knowledge of presence and distribution of the four most common bumble bee species in Texas, to include *B. griseocollis*, *B. impatiens*, and the declining

B. fraternus and *B. pensylvanicus*. By combining novel data from vetted citizen science records reported over the last ten years (i.e., 2007-2016) and data from recent field surveys (2010-2016), with previously compiled museum specimen records, the most complete database of Texas bumble bee records available has been established. These records have resulted in a revised species list for the state that includes three species previously not known to Texas. Additionally, while continental range maps have previously been modeled for bumble bee species found in Texas, we have produced fine-scale SDMs using MaxEnt and identified priority areas for bumble bee conservation efforts that will be practical for state-level conservation planning.

Acknowledgments:

Thank you to Leif Richardson and Michael Warriner, who provided historical bumble bee records from the state of Texas, and to the many individuals and institutions who contributed to those databases. We also thank the following individuals for providing insight into individual historic records: John Ascher and Hadel Go regarding records from the American Museum of Natural History, Karen Wright regarding records from the Texas A&M University Insect Collection, and Doug Yanega regarding records from the Essig Museum of Entomology at Berkeley. Additionally, thank you to the many citizen scientist volunteers who contributed hundreds of bumble bee records to iNaturalist, Texas Bumblebees, and directly to this project.

References:

- Ahrne K, Bengtsson J, & Elmqvist T. 2009. Bumble Bees (*Bombus* spp.) along a gradient of increasing urbanization. *PLoS ONE* 4: e5574.
- Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R. 2013. Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences* 110: 4656–4660.
- Beckham J. 2016. The Influence of Urban Green Spaces on Declining Bumble Bees (Hymenoptera: Apidae). D. Phil. Dissertation, University of North Texas.
- Beckham J. 2016. Texas *Bombus* records from 2016 field surveys. Unpublished raw data.
- Beckham J, Warriner MD, Atkinson SF, Kennedy JH. 2016. The persistence of bumble bees (Hymenoptera: Apidae) in northeastern Texas. *Proceedings of the Washington Entomological Society* 118: 481–497.
- Berenbaum M, Bernhardt P, Buchmann S, Calderone NW, Goldstein P, Inouye DW, ... Thompson FC. 2007. Status of Pollinators in North America. Washington, DC: The National Academies Press.
- Brosi BJ, Briggs HM. 2013. Single pollinator species losses reduce floral fidelity and plant reproductive function. *Proceedings of the National Academy of Sciences* 110: 13044–13048.
- Brown JL. 2014. SDMtoolbox: a python-based GIS toolkit for landscape genetic, biogeographic and species distribution model analyses. *Methods in Ecology and Evolution* 5: 694–700.
- Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, Griswold TL. 2011. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences* 108: 662–667.
- Carvell C. 2002. Habitat use and conservation of bumblebees (*Bombus* spp.) under different grassland management regimes. *Biological Conservation* 103: 33–49.
- Carvell C, Osborne JL, Bourke AFG, Freeman SN, Pywell RF, Heard MS. 2011. Bumble bee species' responses to a targeted conservation measure depend on landscape context and habitat quality. *Ecological Applications* 21: 1760–1771.
- Carvell C, Meek WR, Pywell RF, Goulson D, Nowakowski M. 2007. Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *Journal of Applied Ecology* 44: 29–40.

- 479 Colla S, Packer L. 2008. Evidence for decline in eastern North American bumblebees
480 (Hymenoptera: Apidae), with special focus on *Bombus affinis* Cresson. *Biodiversity and*
481 *Conservation* 17: 1379–1391.
- 482
- 483 Colla S, Gadallah F, Richardson L, Wagner D, Gall L. 2012. Assessing declines of North
484 American bumble bees (*Bombus* spp.) using museum specimens. *Biodiversity and Conservation*
485 21: 3585–3595.
- 486
- 487 Colla S, Ratti CM. 2010. Evidence for the decline of the western bumble bee (*Bombus*
488 *occidentalis* Greene) in British Columbia. *The Pan-Pacific Entomologist* 86: 32–34.
- 489
- 490 Colla S, Richardson L, Williams P. 2011. Bumble bees of the eastern United States. Washington,
491 D.C.: U.S. Department of Agriculture Pollinator Partnership.
- 492
- 493 Corbet SA, Williams IH, Osborne JL. 1991. Bees and the pollination of crops and wild flowers
494 in the European Community. *Bee World* 72: 47–59.
- 495
- 496 Dellicour S, Kastally C, Varela S, Michez D, Rasmont P, Mardulyn P, Lecocq T. 2017.
497 Ecological niche modelling and coalescent simulations to explore the recent geographical range
498 history of five widespread bumblebee species in Europe. *Journal of Biogeography* 44: 39–50.
- 499
- 500 Eken G, Bennun L, Brooks TM, Darwall W, Fishpool LD, Foster M, ... Salaman P. 2004. Key
501 biodiversity areas as site conservation targets. *BioScience* 54: 1110–1118.
- 502
- 503 Elith J, Kearney M, Phillips SJ. 2010. The art of modelling range-shifting species. *Methods in*
504 *Ecology and Evolution* 1: 330–342.
- 505
- 506 Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ. 2011. A statistical
507 explanation of MaxEnt for ecologists. *Diversity and Distributions* 17: 43–57.
- 508
- 509 ESRI. 2013. ArcMap 10.2.2. Environmental Systems Research Institute, Inc., Redlands,
510 California.
- 511
- 512 Fitzpatrick U, Murray MG, Paxton RJ, Breen J, Cotton D, Santorum V, Brown MJF. 2007.
513 Rarity and decline in bumblebees – a test of causes and correlates in the Irish fauna. *Biological*
514 *Conservation* 136: 185–194.
- 515
- 516 Franklin HJ. 1913. The Bombidae of the New World. Part I. *Transactions of the American*
517 *Entomological Society* 38: 177–486.
- 518
- 519 Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R,
520 Cunningham SA ... Bartomeus I. 2013. Wild pollinators enhance fruit set of crops regardless of
521 honey bee abundance. *Science* 339: 1608–1611.
- 522
- 523 Gill RJ, Ramos-Rodriguez O, Raine NE. 2012. Combined pesticide exposure
524 severely affects individual- and colony-level traits in bees. *Nature* 491: 105–109.

- Goulson D. 2010. *Bumblebees: Behaviour, Ecology, and Conservation* [2nd ed.]. Oxford: Oxford University Press.
- Goulson D, Hanley ME, Darvill B, Ellis JS. 2006. Biotope associations and the decline of bumblebees (*Bombus* spp.). *Journal of Insect Conservation* 10: 95–103.
- Goulson D, Lepais O, O'Connor S, Osborne JL, Sanderson RA, Cussans J, ... Darvill B. 2010. Effects of land use at a landscape scale on bumblebee nest density and survival. *Journal of Applied Ecology* 47: 1207–1215.
- Grixti JC, Wong LT, Cameron SA, Favret C. 2009. Decline of bumble bees (*Bombus*) in the North American Midwest. *Biological Conservation* 142: 75–84.
- Guisan A, Thuiller W. 2005. Predicting species distribution: offering more than simple habitat models. *Ecology Letters* 8: 993–1009.
- Hatfield R, Jepsen S, Mader E, Black SH, Shepherd M. 2012. Conserving Bumble Bees: Guidelines for Creating and Managing Habitat for America's Declining Pollinators. 32 pp. Portland, OR: The Xerces Society for Invertebrate Conservation.
- Hatfield R, Jepsen S, Thorp R, Richardson L, Colla S. 2014. *Bombus fraternus*. The IUCN Red List of Threatened Species 2014: e.T44937623A69001851. <http://dx.doi.org/10.2305/IUCN.UK.2014-3.RLTS.T44937623A69001851.en>
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965–1978.
- Hijmans RJ, Phillips S, Leathwick J, Elith J. 2017. dismo: Species Distribution Modeling. R package version 1.1-4. <https://CRAN.R-project.org/package=dismo>
- Inoue MN, Yokoyama J, Washitani I. 2008. Displacement of Japanese native bumblebees by the recently introduced *Bombus terrestris* (L.) (Hymenoptera: Apidae). *Journal of Insect Conservation* 12: 135–146.
- Kadoya T, Ishii HS, Kikuchi R, Suda SI, Washitani I. 2009. Using monitoring data gathered by volunteers to predict the potential distribution of the invasive alien bumblebee *Bombus terrestris*. *Biological Conservation* 142: 1011–1017.
- Kearns CA, Inouye DW, Waser NM. 1998. Endangered mutualisms: the conservation of plant-pollinator interactions. *Annual Review of Ecology and Systematics* 29: 83–112.
- Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, ... Wagner DL. 2015. Climate change impacts on bumblebees converge across continents. *Science* 349: 177–180.
- Klein A-M, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen,

- 571 C, Tscharrntke T. 2007. Importance of pollinators in changing landscapes for world crops.
572 *Proceedings of the Royal Society B: Biological Sciences* 274: 303–313.
- 573
- 574 Koch JB, Strange JP. 2009 Constructing a species database and historic range maps for North
575 American Bumblebees (*Bombus sensu stricto* Latreille) to inform conservation decisions.
576 *Uludag Bee Journal* 9: 97–108.
- 577
- 578 Kosior A, Celary W, Olejniczak P, Fijal J, Król W, Solarz W, Plonka P. 2007. The decline of the
579 bumble bees and cuckoo bees (Hymenoptera: Apidae: Bombini) of western and central Europe.
580 *Oryx* 41: 79–88.
- 581
- 582 Kramer-Schadt S, Niedballa J, Pilgrim JD, Schröder B, Lindenborn J, Reinfelder V, ...
583 Cheyne SM. 2013. The importance of correcting for sampling bias in MaxEnt species
584 distribution models. *Diversity and Distributions* 19: 1366–1379.
- 585
- 586 Labougle, JM. 1990. *Bombus* of Mexico and Central America (Hymenoptera, Apidae).
587 *University of Kansas Science Bulletin* 54:35–73.
- 588
- 589 Lye GC, Osborne JL, Park KJ, Goulson D. 2012. Using citizen science to monitor *Bombus*
590 populations in the UK: nesting ecology and relative abundance in the urban
591 environment. *Journal of Insect Conservation* 16: 697–707.
- 592
- 593 Matsumura C, Yokoyama J, Washitani I. 2004. Invasion status and potential impacts of an
594 invasive alien bumblebee, *Bombus terrestris* L. (Hymenoptera: Apidae) naturalized in southern
595 Hokkaido, Japan. *Global Environmental Research* 8: 51–66.
- 596
- 597 McFrederick QS, LeBuhn G. 2006. Are urban parks refuges for bumble bees *Bombus* spp.
598 (Hymenoptera: Apidae)? *Biological Conservation* 129: 372–382.
- 599
- 600 Memmott J, Waser NM, Price MV. 2004. Tolerance of pollination networks to species
601 extinctions. *Proceedings of the Royal Society B-Biological Sciences* 271: 2605–2611.
- 602
- 603 Merow C, Smith MJ, Silander Jr. JA. 2013. A practical guide to MaxEnt for modeling species'
604 distributions: what it does, and why inputs and settings matter. *Ecography* 36: 1058–1069.
- 605
- 606 Milliron HE. 1973. A monograph of the Western Hemisphere bumblebees (Hymenoptera:
607 Apidae; Bombinae). II. The genus *Megabombus* subgenus *Megabombus*. *Memoirs of the*
608 *Entomological Society of Canada* 89:81–237.
- 609
- 610 Osborne JL, Martin AP, Shortall CR, Todd AD, Goulson D, Knight ME, ... Sanderson RA. 2008.
611 Quantifying and comparing bumblebee nest densities in gardens and countryside
612 habitats. *Journal of Applied Ecology* 45: 784–792.
- 613
- 614 Penado A, Rebelo H, Goulson D. 2016. Spatial distribution modelling reveals climatically
615 suitable areas for bumblebees in undersampled parts of the Iberian Peninsula. *Insect*
616 *Conservation and Diversity* 9: 391–401.

- Phillips SJ, Anderson RP, Schapire RE. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190: 231–259.
- Phillips SJ, Dudík M, Schapire RE. 2004. A maximum entropy approach to species distribution modeling. In *Proceedings of the Twenty-First International Conference on Machine Learning*, pp. 655–662.
- Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S. 2009. Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecological Applications* 19: 181–197.
- Poole, RW. 1996. Nomina insecta nearctica: a check list of the insects of North America. Volume 2. Hymenoptera, Mecoptera, Megaloptera, Neuroptera, Raphidioptera, Trichoptera. Entomological Information Services, Rockville, Maryland.
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010. Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* 25: 345–353.
- Pywell RF, Warman EA, Hulmes L, Hulmes S, Nuttall P, Sparks TH, ... Sherwood A. 2006. Effectiveness of new agri-environment schemes in providing foraging resources for bumblebees in intensively farmed landscapes. *Biological Conservation* 129: 192–206.
- Raes N, ter Steege H. 2007. A null-model for significance testing of presence-only species distribution models. *Ecography* 30: 727–736.
- Richardson L. 2017. Texas *Bombus* records from natural history collections. Unpublished raw data.
- Rundlöf M, Andersson GK, Bommarco R, Fries I, Hederström V, Herbertsson L, ... Smith HG. 2015. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* 521: 77–80.
- Sarospataki M, Novak J, Viktoria M. 2005. Assessing the threatened status of bumble bee species (Hymenoptera: Apidae) in Hungary, Central Europe. *Biodiversity and Conservation* 14: 2437–2446.
- Shipp JL, Whitfield GH, Papadopoulos AP. 1994. Effectiveness of the bumble bee, *Bombus impatiens* Cresson (Hymenoptera, Apidae), as a pollinator of greenhouse sweet pepper. *Scientia Horticulturae* 57: 29–39.
- Stephen WP. 1957. Bumblebees of western America (Hymenoptera: Apoidea). *Oregon College Agricultural Experiment Station Technical Bulletin* 40:1–163.
- Syfert MM, Smith MJ, Coomes DA. 2013. The Effects of Sampling Bias and Model Complexity on the Predictive Performance of MaxEnt Species Distribution Models. *PLoS ONE* 8: 10.1371.

- Szabo ND, Colla SR, Wagner DL, Gall LF, Kerr JT. 2012. Do pathogen spillover, pesticide use, or habitat loss explain recent North American bumblebee declines? *Conservation Letters* 5: 232–239.
- Texas Land Trends. 2014. Status update and trends of Texas rural working lands. Texas A&M Institute of Renewable Natural Resources, College Station, Texas. 1 (1): 1 - 13.
- Texas Land Trends Web Data Explorer. 2015. Texas A&M Institute of Renewable Natural Resources, College Station, Texas. URL: <http://txlandtrends.org/data>.
- Texas Parks and Wildlife Department. 2012. *Texas Conservation Action Plan 2012 – 2016: Statewide/Multi-region Handbook*. Editor, Wendy Connally, Texas Conservation Action Plan Coordinator. Austin, Texas.
- Thorp, R. 2005. *Bombus franklini* Frison 1921 Franklin’s Bumble Bee (Hymenoptera: Apidae: Apinae: Bombinae). In: Shepherd, MD, Vaughan DM, Black SH (Eds.), Red List of Pollinator Insects of North America. CD-ROM Version 1 (May 2005). The Xerces Society for Invertebrate Conservation, Portland, OR.
- Thorp RW, Horning DS, Dunning LL. 1983. Bumble bees and cuckoo bumble bees of California (Hymenoptera: Apidae). *Bulletin of the California Insect Survey* 23:1–79.
- Thorp RW, Shepherd MD. 2005. Profile: subgenus *Bombus*. In: Shepherd, MD, Vaughan DM, Black SH (Eds.), Red List of Pollinator Insects of North America, CD-ROM Version 1 (May 2005). The Xerces Society for Invertebrate Conservation, Portland, OR.
- U.S. Department of Agriculture. 2013. Summary Report: 2010 National Resources Inventory. Natural Resources Conservation Service, Washington, DC, and Center for Survey Statistics and Methodology, Iowa State University, Ames, Iowa.
- Warriner MD. 2012. Bumble bees (Hymenoptera: Apidae) of Texas: Historical distributions. *The Southwestern Naturalist* 57: 442–445.
- Whitehorn P, O’Connor S, Wackers FL, Goulson D. 2012. Neonicotinoid pesticide reduces colony growth and queen production. *Science* 336: 351–352.
- Whittington R, Winston ML. 2004. Comparison and examination of *Bombus occidentalis* and *Bombus impatiens* (Hymenoptera: Apidae) in tomato greenhouses. *Journal of Economic Entomology* 97: 1384–1389.
- Williams P, Cameron SA, Hines HM, Cederberg B, Rasmont P. 2008. A simplified subgeneric classification of the bumblebees (genus *Bombus*). *Apidologie* 39:46–74.
- Williams P, Thorp RW, Richardson L, Colla SR. 2014. Bumble Bees of North America: An Identification Guide. Princeton, New Jersey: Princeton University Press.

709

710 Winfree R, Aguilar R, Vazquez DP, LeBuhn G, Aizen MA. 2009. A meta-analysis of bees'
711 responses to anthropogenic disturbance. *Ecology* 90: 2068–2076.

712

713 Xie Z, Williams PH, Tang Y. 2008. The effect of grazing on bumblebees in the high rangelands
714 of the eastern Tibetan Plateau of Sichuan. *Journal of Insect Conservation* 12: 695–703.

715

716 Yang DR. 1999. The status of species diversity and conservation strategy of bumble bees, a
717 pollination insect in Lancang River Basin of Yunnan, China. *Chinese Biodiversity* 7: 170–174.

718

Table 1 (on next page)

WorldClim bioclimatic environmental predictor variables used for MaxEnt analyses.

Variables are presented in the order of perceived environmental importance that was used to eliminate correlated variables, with derived variables (BIO02, BIO03, BIO04, BIO07, and BIO15) being given lowest ranking. Variables pertaining to the warmest and wettest months and quarters were ranked above similar variables involving coldest and driest months and quarters given that bumble bees are generally more active in warmer months, wetter months. When variable importance could not be intuited based on the aforementioned criteria, ranking was based upon the arbitrary WorldClim code. Variables included in further analyses are highlighted in bold. When applicable, the correlate upon which removal of a variable was based, as well as the Pearson's correlation coefficient (r), is indicated. A Pearson's correlation coefficient of 0.8 or higher was used as the cutoff for variable removal. The full results of the Pearson's correlation analysis are available in Supplement Table 1.

Rank for Pearson's correlation analysis	WorldClim Code	Environmental Predictor Variable	Basis for removal from analysis (correlate and Pearson's correlation coefficient, if applicable)
1	BIO01	Annual Mean Temperature	
2	BIO12	Annual Precipitation	
3	BIO10	Mean Temperature of Warmest Quarter	
4	BIO08	Mean Temperature of Wettest Quarter	
5	BIO09	Mean Temperature of Driest Quarter	
6	BIO11	Mean Temperature of Coldest Quarter	
7	BIO5	Max Temperature of Warmest Month	BIO10; 0.92216
8	BIO6	Minimum Temperature of Coldest Month	BIO11; 0.97781
9	BIO13	Precipitation of Wettest Month	BIO12; 0.88484
10	BIO14	Precipitation of Driest Month	BIO12; 0.9619
11	BIO18	Precipitation of Warmest Quarter	
12	BIO19	Precipitation of Coldest Quarter	BIO12; 0.9529
13	BIO16	Precipitation of Wettest Quarter	BIO12; 0.9028
14	BIO17	Precipitation of Driest Quarter	BIO12; 0.96877
15	BIO02	Mean Diurnal Range (Mean of monthly (max temp - min temp))	
16	BIO03	Isothermality (BIO2/BIO7) (*100)	
17	BIO04	Temperature Seasonality (Standard Deviation*100)	
18	BIO07	Temperature Annual Range (BIO5-BIO6)	BIO11; -0.85041
19	BIO15	Precipitation Seasonality (Coefficient of Variation)	BIO12; -0.81672

Table 2 (on next page)

Sources of bumble bee presence data.

1

Data Source (Time Period)	Number of Records
Museum Collections (1897-2012)	2,105
iNaturalist (2007-2016)	377
Texas Bumble Bees Facebook (2007-2016)	370
Field Data (2010-2016)	728
<i>Total</i>	3,580

2

Table 3 (on next page)

Analysis of species distribution model validity and variable contributions.

The total number of occurrence records (N) and the number of spatially filtered occurrence records used for SDM production (N') are shown for each species, along with average training and test AUCs for each species distribution model. The upper 95% confidence interval of the null distribution's test AUC for each species is also shown for model validation purposes; note that $p < 0.01$ for all species, meaning that the modeled distributions are significantly different from the simulated null distributions. Lastly, top variables contributing to the SDM of each bumble bee species are included, along with percent contribution of each.

Species	N	N'	Avg. Training AUC $\pm$ S.D. (number of training points in each replicate)	Avg. Test AUC $\pm$ S.D. (number of test points in each replicate)	Upper 95% C.I. of Null Distribution's Test AUC (p-value of comparison of actual SDM vs. null)	Top five variables contributing to model (percent contribution)
<i>B. fraternus</i>	239	69	0.826 $\pm$ 0.027 (49)	0.745 $\pm$ 0.06 (20)	0.6102 (p<0.01)	bio03 (51%); bio09 (11%); bio08 (10.2%); bio18 (9.2%); bio01 (4.4%)
<i>B. griseocollis</i>	108	39	0.903 $\pm$ 0.02 (28)	0.866 $\pm$ 0.04 (11)	0.6572 (p<0.01)	bio12 (71.6%); bio03 (8.5%); bio08 (5.6%); bio09 (4.2%); bio04 (2.8%)
<i>B. impatiens</i>	173	24	0.931 $\pm$ 0.015 (17)	0.9 $\pm$ 0.05 (7)	0.6958 (p<0.01)	bio12 (73.8%); bio09 (8.6%); bio08 (7.6%); bio01 (4.1%); bio03 (2.1%)
<i>B. pensylvanicus</i>	3,010	322	0.747 $\pm$ 0.012 (226)	0.667 $\pm$ 0.03 (96)	0.5217 (p<0.01)	bio12 (20.3%); bio09 (15.2%); bio11 (12.9%); bio02 (12.5%); bio04 (11.7%)

1

2

Figure 1

EPA Level III ecoregions in the state of Texas (shapefile downloaded from <https://www.epa.gov/eco-research/ecoregion-download-files-state-region-6>).




Figure 2

Novel presence data (n=747) compiled from the citizen science repositories iNaturalist and Texas Bumblebees Facebook page.

Included species: *B. auricomus* (n=11), *B. fraternus* (n=48), *B. griseocollis* (n=60), *B. impatiens* (n=29), *B. pensylvanicus* (n=557), and *B. pensylvanicus sonorus* (n=42). EPA Level III ecoregions are shown for reference (a=Arizona/New Mexico Mountains; b=Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains).




Figure 3

Modeling results for *B. fraternus*.

A) Mean logistic output from MaxEnt, averaged over 100 replicates, showing probability of suitable habitat being present for *B. fraternus* (average training AUC over 100 runs = 0.826 ± 0.027 and average test AUC 0.745 ± 0.06). Unfiltered species presence data are represented as black dots. B) Reclassified SDM for *B. fraternus* in Texas shown with EPA Level III ecoregions outlined (a=Arizona/New Mexico Mountains; b= Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains). Suitable habitat for *B. fraternus* is likely across the northern half of Texas, with highest modeled probabilities found in the Cross Timbers, Texas Blackland Prairies, and East Central Texas Plains ecoregions.





Figure 4

Modeling results for *B. griseocollis*.

A) Mean logistic output from MaxEnt, averaged over 100 replicates, showing probability of suitable habitat being present for *B. griseocollis* (average training AUC over 100 runs = 0.903 ± 0.02 and average test AUC 0.866 ± 0.04). Unfiltered species presence data points are represented as black dots. B) Reclassified SDM for *B. griseocollis* in Texas shown with EPA Level III ecoregions outlined (a=Arizona/New Mexico Mountains; b= Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains). The eastern half of the state is most likely to contain suitable habitat for *B. griseocollis*, with the highest probabilities in the northern portions of the Texas Blackland Prairies, East Central Texas Plains, and South Central Plains ecoregions.





Figure 5

Modeling results for *B. impatiens*.

A) Mean logistic output from MaxEnt, averaged over 100 replicates, showing probability of suitable habitat being present for *B. impatiens* (average training AUC over 100 runs = 0.931 ± 0.015 and average test AUC 0.9 ± 0.05). Unfiltered species presence data points are represented as black dots. B) Reclassified SDM for *B. impatiens* in Texas shown with EPA Level III ecoregions outlined (a=Arizona/New Mexico Mountains; b= Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains). The highest probabilities of suitable habitat modeled for *B. impatiens* in Texas are in the eastern portion of the state, especially in the South Central Plains ecoregion.



Figure 6

Modeling results for *B. pensylvanicus*.

A) Mean logistic output from MaxEnt, averaged over 100 replicates, showing probability of suitable habitat being present for *B. pensylvanicus* (average training AUC over 100 runs = 0.747 ± 0.012 and average test AUC 0.667 ± 0.03). Unfiltered species presence data are represented as black dots. B) Reclassified SDM for *B. pensylvanicus* in Texas shown with EPA Level III ecoregions outlined (a=Arizona/New Mexico Mountains; b= Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains). Suitable habitat for this species is likely to be observed in most parts of the state.





Figure 7

Target areas for Texas bumble bee conservation.

Areas where there is a high (>0.5) probability of suitable habitat being present for the declining *B. fraternus* and *B. pensylvanicus* species are highlighted, and EPA Level III ecoregions are outlined (a=Arizona/New Mexico Mountains; b= Chihuahuan Deserts; c=High Plains; d=Southwestern Tablelands; e=Central Great Plains; f=Cross Timbers; g=Edwards Plateau; h=Southern Texas Plains; i=Texas Blackland Prairies; j=East Central Texas Plains; k=Western Gulf Coastal Plain; l=South Central Texas Plains). Conservation target areas were calculated using the *combine* tool in ArcMap's Spatial Analyst Toolbox; reclassified distributions of *B. fraternus* and *B. pensylvanicus* were overlaid in implementation of this tool to achieve results. A region of northeast Texas including portions of the Cross Timbers, Texas Blackland Prairies, and East Central Texas Plains ecoregions were identified as target areas for conservation actions.

