

A peer-reviewed version of this preprint was published in PeerJ on 16 August 2017.

[View the peer-reviewed version](https://peerj.com/articles/3674) (peerj.com/articles/3674), which is the preferred citable publication unless you specifically need to cite this preprint.

Ettinger CL, Williams SL, Abbott JM, Stachowicz JJ, Eisen JA. 2017. Microbiome succession during ammonification in eelgrass bed sediments. PeerJ 5:e3674 <https://doi.org/10.7717/peerj.3674>

Microbiome succession during ammonification in eelgrass bed sediments

Cassandra L Ettinger¹, Susan L Williams^{2,3}, Jessica M Abbott², John J Stachowicz², Jonathan A Eisen^{Corresp. 1, 2, 4}

¹ Genome Center, University of California, Davis, Davis, CA, United States

² Department of Evolution and Ecology, University of California, Davis, Davis, CA, United States

³ Bodega Marine Lab, University of California, Davis, Bodega Bay, CA, United States

⁴ Department of Medical Microbiology and Immunology, University of California, Davis, Davis, CA, United States

Corresponding Author: Jonathan A Eisen

Email address: jaeisen@ucdavis.edu

Background. Eelgrass (*Zostera marina*) is a marine angiosperm and foundation species that plays an important ecological role in primary production, food web support, and elemental cycling in coastal ecosystems. As with other plants, the microbial communities living in, on, and near eelgrass are thought to be intimately connected to the ecology and biology of eelgrass. Here we characterized the microbial communities in eelgrass sediments throughout an experiment to quantify the rate of ammonification, the first step in early remineralization of organic matter, or diagenesis, from plots at a field site in Bodega Bay, CA.

Methods. Sediment was collected from 72 plots from a 15 month long field experiment in which eelgrass genotypic richness and relatedness were manipulated. In the laboratory, we placed sediment samples (n= 4 per plot) under a N₂ atmosphere, incubated them at *in situ* temperatures (15 °C) and sampled them initially and after 4, 7, 13, and 19 days to determine the ammonification rate. Comparative microbiome analysis using high throughput sequencing of 16S rRNA genes was performed on sediment samples taken initially and at 7, 13 and 19 days to characterize the relative abundances of microbial taxa and how they changed throughout early diagenesis.

Results. Within-sample diversity of the sediment microbial communities across all plots decreased after the initial timepoint using both richness based (observed number of OTUs, Chao1) and richness and evenness based diversity metrics (Shannon, Inverse Simpson). Additionally, microbial community composition changed across the different timepoints. Many of the observed changes in relative abundance of taxonomic groups between timepoints appeared driven by sulfur cycling with observed decreases in sulfur reducers (*Desulfobacterales*) and corresponding increases in sulfide oxidizers (*Alteromonadales* and *Thiotrichales*). None of these changes in composition or richness were associated with ammonification rates.

Discussion. Overall, our results showed that the microbiome of sediment from different plots followed similar successional patterns, which we surmise to be due to changes related to sulfur metabolism. These large changes likely overwhelmed any potential changes in sediment microbiome related to ammonification rate. We found no relationship between eelgrass presence or genetic composition and the microbiome. This was likely due to our sampling of bulk sediments to measure ammonification rates rather than sampling microbes in sediment directly in contact with the plants and suggests that eelgrass influence on the sediment microbiome may be limited in spatial extent. More in-depth functional studies associated with eelgrass microbiome will be required in order to fully understand the implications of

these microbial communities in broader host-plant and ecosystem functions (e.g. elemental cycling and eelgrass-microbe interactions).

1 **Microbiome succession during ammonification in eelgrass bed sediments**

2
3
4 Cassandra L. Ettinger¹, Susan L. Williams^{2,3}, Jessica M. Abbott², John J. Stachowicz², Jonathan
5 A. Eisen^{1,2,4}

6
7
8 ¹. Genome Center, University of California, Davis, CA. 95616

9 ². Department of Evolution and Ecology, University of California, Davis, CA. 95616

10 ³. Bodega Marine Lab, University of California Davis, Bodega Bay, CA. 94923

11 ⁴. Department of Medical Microbiology and Immunology, University of California, Davis, CA.
12 95616

13
14
15 **Corresponding Author:**

16 Jonathan Eisen^{1,2,4}

17 Email address: jaeisen@ucdavis.edu

40 **Abstract:**

41 **Background.** Eelgrass (*Zostera marina*) is a marine angiosperm and foundation species that
42 plays an important ecological role in primary production, food web support, and elemental
43 cycling in coastal ecosystems. As with other plants, the microbial communities living in, on, and
44 near eelgrass are thought to be intimately connected to the ecology and biology of eelgrass. Here
45 we characterized the microbial communities in eelgrass sediments throughout an experiment to
46 quantify the rate of ammonification, the first step in early remineralization of organic matter, or
47 diagenesis, from plots at a field site in Bodega Bay, CA.

48 **Methods.** Sediment was collected from 72 plots from a 15 month long field experiment in which
49 eelgrass genotypic richness and relatedness were manipulated. In the laboratory, we placed
50 sediment samples (n= 4 per plot) under a N₂ atmosphere, incubated them at *in situ* temperatures
51 (15 °C) and sampled them initially and after 4, 7, 13, and 19 days to determine the
52 ammonification rate. Comparative microbiome analysis using high throughput sequencing of
53 16S rRNA genes was performed on sediment samples taken initially and at 7, 13 and 19 days to
54 characterize the relative abundances of microbial taxa and how they changed throughout early
55 diagenesis.

56 **Results.** Within-sample diversity of the sediment microbial communities across all plots
57 decreased after the initial timepoint using both richness based (observed number of OTUs,
58 Chao1) and richness and evenness based diversity metrics (Shannon, Inverse Simpson).
59 Additionally, microbial community composition changed across the different timepoints. Many
60 of the observed changes in relative abundance of taxonomic groups between timepoints appeared
61 driven by sulfur cycling with observed decreases in sulfur reducers (*Desulfobacterales*) and
62 corresponding increases in sulfide oxidizers (*Alteromonadales* and *Thiotrichales*). None of these
63 changes in composition or richness were associated with ammonification rates.

64 **Discussion.** Overall, our results showed that the microbiome of sediment from different plots
65 followed similar successional patterns, which we surmise to be due to changes related to sulfur
66 metabolism. These large changes likely overwhelmed any potential changes in sediment
67 microbiome related to ammonification rate. We found no relationship between eelgrass presence
68 or genetic composition and the microbiome. This was likely due to our sampling of bulk
69 sediments to measure ammonification rates rather than sampling microbes in sediment directly in
70 contact with the plants and suggests that eelgrass influence on the sediment microbiome may be
71 limited in spatial extent. More in-depth functional studies associated with eelgrass microbiome
72 will be required in order to fully understand the implications of these microbial communities in
73 broader host-plant and ecosystem functions (e.g. elemental cycling and eelgrass-microbe
74 interactions).

75

76

77

78

79

80 Introduction

81 Eelgrass (*Zostera marina* L.) is a widely-distributed marine angiosperm that supports
82 ecologically and economically valuable functions (Williams & Heck, 2001), including high rates
83 of primary production, higher trophic levels, and elemental cycling (Hemminga & Duarte, 2000).
84 Much of the high primary production of eelgrass and its associated algal community ends up as
85 detritus (Cebrian & Lartigue, 2004), which fuels high rates of ammonification, the first step in
86 the early diagenesis of organic matter (Bernier, 1980), in the sediments of eelgrass beds.
87 Although the role of microbes in the decomposition of organic matter and remineralization in
88 marine sediments is broadly appreciated (Arndt et al., 2013), the extent to which microbial
89 community composition and process rates are influenced by the characteristics of eelgrass beds
90 is unclear.

91
92 The microorganisms associated with eelgrass have been found to be distinct for different eelgrass
93 parts (e.g. roots, leaves, rhizomes) and appear to vary within and between host plants
94 (Fahimipour et al., 2017; Bengtsson et al., 2017; Ettinger et al., 2017; Holland-Moritz et al.,
95 2017). Many of the dominant taxa found in association with eelgrass beds are predicted to be
96 involved in nitrogen and sulfur cycling (Capone, 1982; Welsh, 2000; Nielsen et al., 2001; Lovell,
97 2002; Sun et al., 2015; Cúcio et al., 2016; Ettinger et al., 2017; Holland-Moritz et al., 2017). The
98 microbial communities in eelgrass bed sediment are significantly different from that of
99 surrounding unvegetated sediment (Cúcio et al., 2016) and even from eelgrass roots collected
100 within the same bed (Fahimipour et al., 2017, Ettinger et al., 2017). Furthermore, even within
101 grass beds, sediment community composition differences are correlated with eelgrass density
102 (Ettinger et al., 2017), suggesting the potential for eelgrass influence of microbial processes.

103
104 Seagrass density, biomass, growth and resilience are all known to be influenced by the genetic
105 composition and diversity of eelgrass assemblages (Hughes & Stachowicz, 2004; Reusch et al.,
106 2005; Hughes & Stachowicz, 2011; Stachowicz et al., 2013). At the conclusion of a larger
107 experiment testing the effects of eelgrass genotypic richness and relatedness on eelgrass biomass
108 accumulation and other ecosystem functions (Abbott, 2015, Abbott et al., in review), we sampled
109 the microbial communities in eelgrass sediments in plots that varied in genetic diversity. We
110 characterized the relative abundances of microbial taxa and how they changed as early
111 diagenesis proceeded during a laboratory experiment that quantified the rate of ammonification
112 as a function of plant genotypic diversity and abundance.

114 Methods

115 *Ammonification Experiment*

116 The rate of ammonification was determined in sediments collected from plots of a field
117 experiment lasting 15 months in which eelgrass genotypic richness and relatedness were
118 manipulated and various ecosystem functions were measured (Abbott, 2015, Abbott et al., in
119 review). The experiment initially crossed two levels of genotypic richness levels (2, 6) with three

120 levels of genetic relatedness (more, less, and as closely related as expected by chance (Frasier,
121 2008; Stachowicz et al., 2013)) with 6 replicates per richness x relatedness combination for a
122 total of 72 plots. Plots were 40.4 cm long x 32.7 cm wide x 15.2 cm deep. Genotypic
123 composition changed in the treatments as a result of mortality of some planted genotypes early in
124 the experiment and some plots lost all eelgrass by the end of the experiment; this mortality was
125 independent of treatment. Because samples for ammonification were taken at the end of the
126 experiment, we used final genotypic composition to calculate realized diversity and relatedness
127 for each plot for use in analysis.

128

129 In October 2014, prior to the harvest of eelgrass from the experiment, we collected ~500 cm³ of
130 sediment from the top 10 cm of the sediment surface in each plot to determine the rate of
131 ammonification (see Williams et al., in revision for more details). In the laboratory, we placed
132 sediment samples in a N₂- filled glove box, removed macroscopic pieces of eelgrass and animals
133 using forceps, and then filled opaque glass centrifuge tubes with sediments (n = 4 per plot).
134 Tubes were incubated at *in situ* temperatures (15 °C) and sampled for porewater and adsorbed
135 ammonium and sediment porosity initially and after 4, 7, 13, and 19 days of incubation.
136 Ammonium production rates were calculated by linear regression of $\mu\text{mol NH}_4\text{-N}_{\text{porewater} +}$
137 $\text{adsorbed}/\text{L}$ sediment versus incubation time (days) (Mackin & Aller, 1984; Dennison, Aller &
138 Alberte, 1987; Williams, 1990). We also removed belowground and aboveground eelgrass
139 biomass from each plot, cleaned it of sediments and epiphytes, and dried it to constant mass (see
140 Abbott 2015, Abbott et al. in review for more details).

141

142 *Molecular Analysis*

143 Sediment was collected at each timepoint during the ammonification experiment for microbial
144 analysis. DNA was extracted from the sediment taken initially and at 7, 13 and 19 days (herein
145 referred to as timepoints 1, 2, 3 and 4 respectively) using the PowerSoil DNA Isolation kit (MO
146 BIO Laboratories, Inc., Carlsbad, CA, USA) according to the manufacturer's protocol. The V4
147 region of the 16S rRNA gene was amplified using the "universal" 515F and 806R primers
148 (Caporaso et al., 2012) with a modified barcode system as in Fahimipour et al (2017). A detailed
149 amplification protocol can be found here

150 (https://seagrassmicrobiome.files.wordpress.com/2015/01/16s_library_pcr_protocol_pnas.pdf).

151 Molecular libraries were sent to the UC Davis Genome Center Core Facilities for sequencing on
152 an Illumina MiSeq (Illumina, Inc. San Diego, CA, USA) to generate 250 bp paired-end data.

153

154 *Sequence Processing*

155 A custom in-house script was used to demultiplex, quality check and merge paired-end reads

156 (https://github.com/gjospin/scripts/blob/master/Demul_trim_prep.pl). Sequences were then

157 analyzed using the Quantitative Insights Into Microbial Ecology (QIIME) v. 1.9.0 workflow

158 (Caporaso et al., 2010). For a detailed walkthrough of the following analysis using QIIME see

159 the IPython notebook

160 (<http://nbviewer.jupyter.org/gist/casett/a42c64ca4b74b1d414f59eb5362e63a3>). A total of
161 10,958,285 reads obtained from the sequencing run passed quality filtering (Q20), of which
162 7,856,501 paired-end reads merged successfully (71.69%). Chimeras were identified using
163 USEARCH v. 6.1 and filtered out. Sequences were then *denovo* clustered into operational
164 taxonomic units (OTUs) at 97 percent similarity using UCLUST (Edgar, 2010) and taxonomy
165 was assigned using the the GreenGenes database (v.13_8) (DeSantis et al., 2006). Using the
166 `filter_taxa_from_otu_table.py` and `filter_otus_from_otu_table.py` QIIME scripts, chloroplast
167 DNA, mitochondrial DNA, singletons and reads classified as "Unassigned" at the domain level
168 were filtered out of the dataset before downstream analysis.

169

170 *Data Analysis and Visualization*

171 Data manipulation, visualization and statistical analyses were performed in R (R Core Team,
172 2016) using the `ggplot2` (Wickham & Hadley, 2009), `vegan` (Dixon, 2003), `phyloseq` (McMurdie
173 & Holmes, 2013), `coin` (Hothorn et al., 2008) and `FSA` packages (Ogle, 2016). For statistical
174 comparisons and visualization, the dataset was subsampled without replacement to an even depth
175 of 5000 sequences. As a result eight samples were removed from downstream analysis due to
176 low sequence counts (SampleID: I4T4, C5T4, K3T3, J4T3, J2T3, D5T3, G5T4 and K2T3).

177

178 A variety of metrics, including observed OTUs, Chao1 (Chao, 1984), Shannon (Shannon &
179 Weaver, 1949) and Inverse Simpson (Simpson, 1949) indices, were used to calculate the within-
180 sample (alpha) diversity for the dataset. Kruskal-Wallis tests with 9999 permutations were used
181 to test for significant differences in alpha diversity between different sample categories including
182 timepoint, plot location, block, spot, eelgrass diversity, eelgrass richness, eelgrass evenness,
183 eelgrass relatedness, eelgrass status in the plot and treatment. For categories where the Kruskal-
184 Wallis test resulted in a rejected null hypothesis ($p < 0.05$), Bonferroni corrected post-hoc Dunn
185 tests were performed to identify which groups showed stochastic dominance.

186

187 To assess between-sample (beta) diversity, the Unifrac (weighted and unweighted) (Lozupone et
188 al., 2007; Hamady, Lozupone & Knight, 2010) and Bray-Curtis (Bray & Curtis, 1957)
189 dissimilarities were calculated. These diversity metrics were then compared using permutational
190 manovas (PERMANOVAs) to test for significant differences between sample categories (see
191 above) with 9999 permutations using the Bonferroni correction (Anderson, 2001). Mantel tests
192 were used to test for correlations between Bray-Curtis dissimilarities calculated for the microbial
193 data and euclidean distances calculated for continuous variables such as aboveground eelgrass
194 biomass (g/plot), belowground eelgrass biomass (g/plot), total eelgrass biomass (g/plot), plot
195 decomposition rate, detritus standing stock (g/plot), ammonification rate ($\mu\text{mol NH}_4\text{-N/L}$
196 sediment/d) and eelgrass plot final genotypic diversity and relatedness (assessed previously in
197 Abbott 2015, Abbott et al. in review - Shannon Diversity, Rao's Q, average relatedness,
198 genotypic evenness). These tests were performed in R with `vegan` using 9999 permutations.

199

200 To compare microbial community composition among timepoints, we collapsed OTUs into
201 taxonomic orders using the `tax_glom` function in `phyloseq` and then removed groups with a mean
202 abundance of less than two percent. Rare groups were removed to avoid false positives from low
203 abundance taxa and to focus analysis on abundant groups that may influence sediment
204 biogeochemistry. The average relative abundance of taxonomic orders was compared between
205 timepoints using Bonferroni corrected Kruskal-Wallis tests in R. For taxonomic groups where
206 the Kruskal-Wallis test resulted in a rejected null hypothesis, Bonferroni corrected post-hoc
207 Dunn tests were performed to identify which timepoint comparisons for each taxonomic order
208 showed stochastic dominance. To determine the nature of the relationship between
209 ammonification rate and specific taxonomic groups whose mean relative abundances differed
210 significantly between timepoints, we built linear models in R. We focused specifically on the
211 three taxonomic groups with the largest variance in relative abundance and the models were built
212 using the timepoint where the largest variance was observed.

213

214 **Results**

215 *Alpha Diversity Metrics: Within sample diversity decreases after initial timepoint*

216 Alpha diversity was significantly different between timepoints (K-W test; $p < 0.001$, Figure 1,
217 Table S1) for all metrics and post-hoc Dunn tests identified that the alpha diversity for timepoint
218 1 was consistently greater than that for other timepoints ($p < 0.001$, Table S2). The decrease in
219 diversity from timepoint 1 to the subsequent timepoints is expected as obligate aerobes are not
220 likely to survive after initial inoculation in sealed tubes. Plot location, eelgrass relatedness and
221 eelgrass richness did not affect any estimate of alpha-diversity across timepoints or within single
222 timepoints (K-W test, $p > 0.05$, Table S1).

223

224 *Beta Diversity Metrics: Microbial community composition changes over time*

225 Microbial community composition differed between timepoints for all three dissimilarity
226 metrics, Bray-Curtis, unweighted and weighted Unifrac (PERMANOVA, $p < 0.001$, Figure 2,
227 Table S3). Subsequent pair-wise PERMANOVA test results found that all pair-wise timepoint
228 comparisons differ significantly in composition ($p < 0.001$, Table S4). PERMANOVA test
229 results for other sample categories were not significantly different ($p > 0.05$, Table S3).
230 Surprisingly, we did not detect any associations of the initial microbiome (timepoint 1) with plot
231 level features such as eelgrass genotypic richness or eelgrass presence/absence (Figure 3, Table
232 S5).

233

234 *Microbial composition effects on ammonification rate*

235 Ammonification rates ranged from 12 to 640 $\mu\text{mol NH}_4\text{-N/L sediment/d}$, values typical for
236 eelgrass (Iizumi, Hattori & McRoy, 1982; Dennison, Aller & Alberte, 1987, Williams et al., in
237 revision). Using the full dataset, we tested for correlations between Bray-Curtis dissimilarities
238 and euclidean distances of several measured variables including ammonification rate and
239 eelgrass final genotypic diversity and relatedness. None of these measured variables were

240 correlated with microbial dissimilarities (Mantel test, $p > 0.05$, Table S6). We then focused our
241 analyses on testing for correlations between these measures and the dissimilarities of only the
242 initial or final timepoints, but still found no correlations (Mantel test, $p > 0.05$, Table S7).

243

244 *Taxonomic composition*

245 The orders *Pirellulales*, *Chromatiales*, *Desulfobacterales*, *Bacteroidales*, *Alteromonadales*,
246 *Campylobacterales* and *Thiotrichales* had mean relative abundances that were significantly
247 different across all timepoints (K-W test, $p < 0.001$, Table S8). Since we were interested in the
248 significance of the directional changes in the observed succession pattern, we focused our
249 investigation on the sequential timepoint comparisons during post-hoc analysis.

250

251 We saw a clear succession in eelgrass sediment microbiota during the experiment, which was
252 characterized by several significant differences (Figure 4, Table S9, Table S10). The strongest
253 among these involved several main observations:

254 1) An initial increase in the mean relative abundance of *Campylobacterales*, mainly members of
255 the family *Helicobacteraceae*, between timepoints 1 and 2 (4.8 to 12.57%), followed by a
256 decrease in relative abundance (12.57 to 9.36%) from timepoint 2 to 3.

257 2) An increase in relative abundance from 3.12 to 6.21% in *Alteromonadales* between timepoints
258 2 - 4.

259 3) A doubling of the average relative abundance of *Thiotrichales*, specifically the genus
260 *Thiomicrospira*, from 9.36 to 18.53% between timepoint 3 and 4.

261

262 In our linear model analysis, we did not detect a significant relationship between ammonification
263 rate and the relative abundance of *Thiotrichales* (timepoint 4, F-statistic = 0.323, adjusted r-
264 squared = -0.01, $p = 0.517$), *Alteromonadales* (timepoint 4, F-statistic = 0.167, adjusted r-squared
265 = -0.012, $p = 0.684$) or *Campylobacterales* (timepoint 2, F-statistic = 1.962, adjusted r-squared =
266 0.013, $p = 0.166$).

267

268 **Discussion**

269 We did not detect any association of the microbiome with plot level features such as eelgrass
270 genotypic richness or eelgrass presence/absence (Figure 3). This result originally seemed
271 surprising given previous work indicating a correlation between eelgrass presence and sediment
272 microbiota (Cúcio et al., 2016; Ettinger et al., 2017). However, it is important to note that
273 microbiome samples came from homogenized bulk sediment collected from whole plots rather
274 than sediment specifically in close association with eelgrass roots. This suggests that associations
275 between microbiota and eelgrass are localized to plant surfaces or immediately adjacent
276 sediments do not extend far from the plant itself. Indeed, Fahimipour et al. found that the root
277 microbiome differed substantially from that found in sediments taken from within the eelgrass
278 bed, but not specifically associated with roots. Alternatively, it is possible that the
279 homogenization and transport from field to the lab fundamentally altered the microbiome,

280 causing the differences with previous studies. One alternative possibility is that these plots do not
281 differ because eelgrass has a lasting effect on the sediment microbiome and the plots without
282 eelgrass, since they previously, although briefly, had eelgrass, have just not yet returned to a
283 non-eelgrass microbiome state.

284

285 To conduct the ammonification experiment, the sediment was moved from its natural setting, in
286 which a micro-oxic zone exists around eelgrass roots (Jensen et al., 2005), into an anaerobic,
287 enclosed system. Seagrass sediments are highly anaerobic below the very top layers and thus,
288 organic matter diagenesis is predominantly an anaerobic process (Harrison, 1989; Marbà et al.,
289 2006). This procedure enabled us not only to quantify ammonification rates but also to study
290 successional shifts in communities under these conditions during which we observed reductions
291 in alpha diversity and changes in taxonomic composition. Overall, the different samples,
292 regardless of the ammonification rate, followed similar successional patterns, which we infer to
293 be due largely to a response to sulfur metabolism, based on the likely functional role of the
294 taxonomic groups that exhibited the greatest change in relative abundance across timepoints. The
295 relative decrease in *Desulfobacterales* (an order for which most of the characterized species are
296 known to be sulfate reducers), concomitant with an increase in *Alteromonadales* and
297 *Thiotrichales* (both groups dominated by sulfide oxidizers), supports this hypothesis, suggesting
298 that sulfate reduction and sulfide oxidation were coupled during the experiment. We note that
299 each “replicate” sample does not follow the exact sample succession pattern. This can be seen
300 especially in timepoint 4 samples which are widely scattered on the PCoA plot (Figure 2). The
301 variation between these samples appears to be due in large part to differences in relative
302 abundance of specific likely sulfide oxidizers (e.g. *Thiotrichales*). We also note that by
303 conducting this process in an anaerobic setting and only focusing on 16S rRNA gene sequence
304 analysis, we are unable to detect the role of microbial eukaryotes (e.g. fungi, ciliates, amoeba)
305 during and throughout early diagenesis in seagrass bed sediments. This may be of little
306 consequence as, in contrast to in terrestrial systems where microbial eukaryotes are known to
307 participate in ammonification, these groups are historically thought to contribute little to the
308 primarily anaerobic process of organic matter diagenesis in seagrass sediments (Newell, 1981 for
309 *Z. marina*; Blum et al., 1988 for tropical seagrass leaf litter, Harrison, 1989), although they have
310 been observed in seagrass detritus (Harrison & Mann, 1975; Harrison, 1989).

311

312 We did not detect any major correlations between the microbiome and ammonification rate.
313 There are multiple explanations for this including that ammonium production can occur as a
314 byproduct of a variety of microbial processes and metabolic pathways (Herbert, 1999; Zehr &
315 Kudela, 2011). General microbial activity has been previously linked with rates of seagrass
316 decomposition (Blum & Mills, 1991), so perhaps what we observe here is a broader community
317 process that cannot be linked to any one taxonomic group. A more likely explanation is that the
318 effects of ammonium production may be present in our dataset, but are masked here by stronger
319 processes (e.g. sulfur metabolism). In marine sediments, sulfate reduction can be attributed as

320 responsible for a large part of organic carbon oxidation and the dominant anaerobic process as it
321 is more thermodynamically favorable than methanogenesis (Berner, 1980; Capone & Kiene,
322 1988; Marbà et al., 2006). Thus, the overall succession pattern that we are seeing is likely an
323 accurate representation of what occurs during early remineralization of organic matter in anoxic
324 seagrass sediments even if we cannot link it to the ammonification rate here.

325

326 **Conclusions**

327 Seagrass beds are known as hotspots of primary production, organic matter degradation, and
328 elemental cycling and previous work has suggested that sulfur metabolism can play an important
329 ecological role in these beds. In this study, we wanted to identify if successional patterns in
330 microbial communities during early diagenesis were correlated with the rate of ammonification.
331 We found no such correlation, instead, observing a successional pattern consistent with sulfur
332 cycling. Future work should endeavor to use metagenomic techniques to investigate the
333 abundance of genes associated with sulfur metabolism to confirm this observation. Additionally,
334 all though no correlation was found between ammonification rate and 16S rRNA gene sequence
335 data, metagenomics might identify functional genes that are enriched in samples with a higher
336 rate of ammonification. Seagrass beds have important ecosystem functions, but our knowledge of
337 the microbial communities inhabiting these beds and their functions is still fragmentary. This
338 work contributes to the growing body of knowledge on the eelgrass microbiome, providing some
339 contextual functional framework for the sediment associated generally within these beds and
340 highlighting a growing need for functional studies in this and other host-microbe-environment
341 systems.

342

343 **Acknowledgments**

344 Illumina sequencing was performed at the DNA Technologies Core facility in the UC Davis
345 Genome Center in Davis, California. We thank Qingyi “John” Zhang for his help with the DNA
346 extractions and Illumina library preparation.

347

348 **References**

- 349 Abbott JM. 2015. The effects of genotypic richness, genetic relatedness, and trait variation in a
350 seagrass community. PhD thesis. University of California, Davis.
- 351 Abbott, JM; Grosberg RK, Williams SL, and Stachowicz JJ. In review. Multiple dimensions of
352 intraspecific diversity affect biomass of eelgrass and its associated community. Submitted to
353 *Ecology*.
- 354 Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance. *Austral*
355 *ecology* 26:32–46.

- 356 Arndt S., Jørgensen BB., LaRowe DE., Middelburg JJ., Pancost RD., Regnier P. 2013.
357 Quantifying the degradation of organic matter in marine sediments: A review and synthesis.
358 *Earth-Science Reviews* 123:53–86.
- 359 Bengtsson MM., Buhler A., Brauer A., Dahlke S., Schubert H., Blindow I. 2017. Eelgrass leaf
360 surface microbiomes are locally variable and highly correlated with epibiotic eukaryotes.
361 DOI: 10.1101/111559.
- 362 Berner RA. 1980. *Early Diagenesis: A Theoretical Approach*. Princeton University Press.
- 363 Blum LK., Mills AL. 1991. Microbial growth and activity during the initial stages of seagrass
364 decomposition. *Marine ecology progress series* 70:73–82.
- 365 Blum LK., Mills AL., Zieman JC., Zieman RT. 1988. Abundance of bacteria and fungi in
366 seagrass and mangrove detritus. *Marine ecology progress series* 42:73–78.
- 367 Bray JR., Curtis JT. 1957. An Ordination of the Upland Forest Communities of Southern
368 Wisconsin. *Ecological monographs* 27:325–349.
- 369 Capone DG. 1982. Nitrogen Fixation (Acetylene Reduction) by Rhizosphere Sediments of the
370 Eelgrass *Zostera marina*. *Marine ecology progress series* 10:67–75.
- 371 Capone DG., Kiene RP. 1988. Comparison of microbial dynamics in marine and freshwater
372 sediments: Contrasts in anaerobic carbon catabolism. *Limnology and oceanography*
373 33:725–749.
- 374 Caporaso JG., Kuczynski J., Stombaugh J., Bittinger K., Bushman FD., Costello EK., Fierer N.,
375 Peña AG., Goodrich JK., Gordon JL., Huttley GA., Kelley ST., Knights D., Koenig JE., Ley
376 RE., Lozupone CA., McDonald D., Muegge BD., Pirrung M., Reeder J., Sevinsky JR.,
377 Turnbaugh PJ., Walters WA., Widmann J., Yatsunenko T., Zaneveld J., Knight R. 2010.
378 QIIME allows analysis of high-throughput community sequencing data. *Nature methods*

- 379 7:335–336.
- 380 Caporaso JG., Lauber CL., Walters WA., Berg-Lyons D., Huntley J., Fierer N., Owens SM.,
381 Betley J., Fraser L., Bauer M., Gormley N., Gilbert JA., Smith G., Knight R. 2012. Ultra-
382 high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.
383 *The ISME journal* 6:1621–1624.
- 384 Cebrian J., Lartigue J. 2004. Patterns of herbivory and decomposition in aquatic and terrestrial
385 ecosystems. *Ecological monographs* 74:237–259.
- 386 Chao A. 1984. Non-parametric estimation of the number of classes in a population. *Scandinavian*
387 *journal of statistics, theory and applications* 11:265–270.
- 388 Cúcio C., Engelen AH., Costa R., Muyzer G. 2016. Rhizosphere Microbiomes of European +
389 Seagrasses Are Selected by the Plant, But Are Not Species Specific. *Frontiers in*
390 *microbiology* 7:440.
- 391 Dennison WC., Aller RC., Alberte RS. 1987. Sediment ammonium availability and eelgrass
392 (*Zostera marina*) growth. *Marine biology* 94:469–477.
- 393 DeSantis TZ., Hugenholtz P., Larsen N., Rojas M., Brodie EL., Keller K., Huber T., Dalevi D.,
394 Hu P., Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and
395 workbench compatible with ARB. *Applied and environmental microbiology* 72:5069–5072.
- 396 Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of vegetation*
397 *science: official organ of the International Association for Vegetation Science* 14:927–930.
- 398 Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
399 26:2460–2461.
- 400 Ettinger CL., Voerman SE., Lang JM., Stachowicz JJ., Eisen JA. 2017. Microbial communities
401 in sediment from *Zostera marina* patches, but not the *Z. marina* leaf or root microbiomes,

- 402 vary in relation to distance from patch edge. *PeerJ*.
- 403 Fahimipour AK., Kardish MR., Eisen JA., Lang JM., Green JL., Stachowicz JJ. 2017. Global-
404 scale structure of the eelgrass microbiome. *Applied and Environmental Microbiology*. DOI:
405 10.1128/AEM.03391-16.
- 406 Frasier TR. 2008. STORM: software for testing hypotheses of relatedness and mating patterns.
407 *Molecular ecology resources* 8:1263–1266.
- 408 Hamady M., Lozupone C., Knight R. 2010. Fast UniFrac: facilitating high-throughput
409 phylogenetic analyses of microbial communities including analysis of pyrosequencing and
410 PhyloChip data. *The ISME journal* 4:17–27.
- 411 Harrison PG. 1989. Detrital processing in seagrass systems: A review of factors affecting decay
412 rates, remineralization and detritivory. *Aquatic botany* 35:263–288.
- 413 Harrison PG., Mann KH. 1975. Detritus formation from eelgrass (*Zostera marina* L.): The
414 relative effects of fragmentation, leaching, and decay. *Limnology and oceanography*
415 20:924–934.
- 416 Hemminga MA., Duarte CM. 2000. *Seagrass Ecology*. Cambridge University Press.
- 417 Herbert RA. 1999. Nitrogen cycling in coastal marine ecosystems. *FEMS microbiology reviews*
418 23:563–590.
- 419 Holland-Moritz HE., Lang JM., Stachowicz JJ., Eisen JA. In prep. Exploring the Biogeography
420 of Microbial Communities on the Surface of Seagrasses.
- 421 Hothorn T., Hornik K., van de Wiel MA., Zeileis A. 2008. Implementing a Class of Permutation
422 Tests: The coin Package. *Journal of statistical software* 28. DOI: 10.18637/jss.v028.i08.
- 423 Hughes AR., Stachowicz JJ. 2004. Genetic diversity enhances the resistance of a seagrass
424 ecosystem to disturbance. *Proceedings of the National Academy of Sciences of the United*

- 425 *States of America* 101:8998–9002.
- 426 Hughes AR., Stachowicz JJ. 2011. Seagrass genotypic diversity increases disturbance response
427 via complementarity and dominance. *The Journal of ecology* 99:445–453.
- 428 Iizumi H., Hattori A., McRoy CP. 1982. Ammonium regeneration and assimilation in eelgrass
429 (*Zostera marina*) beds. *Marine biology* 66:59–65.
- 430 Jensen SI., Kühl M., Glud RN., Jørgensen LB., Priemé A. 2005. Oxic microzones and radial
431 oxygen loss from roots of *Zostera marina*. *Marine ecology progress series* 293:49–58.
- 432 Lovell CR. 2002. Plant-Microbe Interactions in the Marine Environment. In: Bitton G eds.
433 *Encyclopedia of environmental microbiology*, Wiley, New York, NY. 5: 2539–2554
- 434 Lozupone CA., Hamady M., Kelley ST., Knight R. 2007. Quantitative and Qualitative Diversity
435 Measures Lead to Different Insights into Factors That Structure Microbial Communities.
436 *Applied and environmental microbiology* 73:1576–1585.
- 437 Mackin JE., Aller RC. 1984. Ammonium adsorption in marine sediments. *Limnology and*
438 *oceanography* 29:250–257.
- 439 Marbà N., Holmer M., Gacia E., Barrón C. 2006. Seagrass Beds and Coastal Biogeochemistry.
440 In: *Seagrasses: Biology, Ecology and Conservation*. 135–157.
- 441 McMurdie PJ., Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis
442 and Graphics of Microbiome Census Data. *PloS one* 8:e61217.
- 443 Newell SY. 1981. Fungi and Bacteria in or on Leaves of Eelgrass (*Zostera marina* L.) from
444 Chesapeake Bay. *Applied and environmental microbiology* 41:1219–1224.
- 445 Nielsen LB., Finster K., Welsh DT., Donnelly A., Herbert RA., de Wit R., Lomstein BA. 2001.
446 Sulphate reduction and nitrogen fixation rates associated with roots, rhizomes and
447 sediments from *Zostera noltii* and *Spartina maritima* meadows. *Environmental*

- 448 *microbiology* 3:63–71.
- 449 Ogle DH. 2016. *FSA: Fisheries Stock Analysis*. R package version 0.8.12.
- 450 R Core Team. 2016. *R: A language and environment for statistical computing*. R Foundation for
451 Statistical Computing, Vienna, Austria.
- 452 Reusch TBH., Ehlers A., Hämmerli A., Worm B. 2005. Ecosystem recovery after climatic
453 extremes enhanced by genotypic diversity. *Proceedings of the National Academy of
454 Sciences of the United States of America* 102:2826–2831.
- 455 Shannon CE., Weaver W. 1949. *The Mathematical Theory of Communication*. Urbana:
456 University of Illinois Press, 29.
- 457 Simpson EH. 1949. Measurement of Diversity. *Nature* 163:688–688.
- 458 Stachowicz JJ., Kamel SJ., Hughes AR., Grosberg RK. 2013. Genetic Relatedness Influences
459 Plant Biomass Accumulation in Eelgrass (*Zostera marina*). *The American naturalist*
460 181:715–724.
- 461 Sun F., Zhang X., Zhang Q., Liu F., Zhang J., Gong J. 2015. Seagrass (*Zostera marina*)
462 Colonization Promotes the Accumulation of Diazotrophic Bacteria and Alters the Relative
463 Abundances of Specific Bacterial Lineages Involved in Benthic Carbon and Sulfur Cycling.
464 *Applied and environmental microbiology* 81:6901–6914.
- 465 Welsh DT. 2000. Nitrogen fixation in seagrass meadows: Regulation, plant-bacteria interactions
466 and significance to primary productivity. *Ecology letters* 3:58–71.
- 467 Wickham H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
- 468 Williams SL. 1990. Experimental Studies of Caribbean Seagrass Bed Development. *Ecological
469 monographs* 60:449–469.
- 470 Williams SL., Heck Jr. KL. 2001. Seagrass community ecology. In: Bertness, Gaines SD, Hay

- 471 ME eds. *Marine community ecology*. Sinauer Associates, 317–337.
- 472 Willams, SL; Reynolds LK, Abbott JM., Stachowicz JJ. In revision. Marine macrophyte detritus
473 and decomposition: the role of intraspecific genetic variation. *Estuaries and Coasts*.
- 474 Zehr JP., Kudela RM. 2011. Nitrogen cycle of the open ocean: from genes to ecosystems. *Annual*
475 *review of marine science* 3:197–225.
- 476

Figure 1

Alpha diversity decreases over time.

Four alpha diversity metrics (observed number of OTUs, Chao1, Shannon and Inverse Simpson diversity indices) are shown here as box plots grouped and colored by timepoint. Timepoint 1 (initial samples), 2 (7 days), 3 (13 days), and 4 (19 days).




Figure 2

Microbial community composition changes over time.

Principal Coordinates Analysis (PCoA) of Weighted Unifrac distances of microbial communities are shown here with shapes and colors representative of respective timepoint. Timepoint 1 (initial samples), 2 (7 days), 3 (13 days), and 4 (19 days).




Figure 3

Initial microbial community composition is not correlated with eelgrass presence/absence.

Principal Coordinates Analysis (PCoA) of Weighted Unifrac distances of microbial communities at the initial timepoint (timepoint 1) are shown here. Points in the ordination are colored by eelgrass status in each plot (one genotype, multiple genotypes, absent).




Figure 4

Taxonomic composition varies over time.

The average relative abundance of taxonomic orders with a mean greater than two percent are shown across timepoints with the standard error of the mean represented by error bars. Lines are grouped by phylum and colored by taxonomic order. Timepoint 1 (initial samples), 2 (7 days), 3 (13 days), and 4 (19 days).

