

**A peer-reviewed version of this preprint was published in PeerJ on 7 October 2014.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.611) (peerj.com/articles/611), which is the preferred citable publication unless you specifically need to cite this preprint.

Rumberger JM, Arch JRS, Green A. 2014. Butyrate and other short-chain fatty acids increase the rate of lipolysis in 3T3-L1 adipocytes. PeerJ 2:e611 <https://doi.org/10.7717/peerj.611>

1 **Butyrate and other Short-Chain Fatty Acids Increase the Rate of Lipolysis in 3T3-L1**  
2 **Adipocytes**

3  
4 **John M. Rumberger<sup>1,2</sup>, Jonathan R. S. Arch<sup>2</sup> and Allan Green<sup>3</sup>**

5  
6 From Bassett Healthcare, Cooperstown, NY 13326<sup>1</sup>, the Clore Laboratory, University of  
7 Buckingham, Buckingham MK18 1EG UK<sup>2</sup> and the Department of Chemistry and Biochemistry,  
8 SUNY Oneonta, Oneonta, NY 13820<sup>3</sup>

9  
10 Running Title: Short-Chain Fatty Acid Regulation of Lipolysis

11 **Address correspondence to:**

12 Allan Green, D.Phil., Department of Chemistry and Biochemistry, SUNY College at Oneonta,  
13 120 Physical Science Building, 108 Ravine Parkway, Oneonta, NY 13820.

14 Tel: (607) 436-2604

15 Email: allan.green@oneonta.edu

16 **Key words:** lipolysis, adipose tissue, butyrate, short-chain fatty acids, histone deacetylase

17 *The abbreviations used are:* FFA, free fatty acids; SCFA, short-chain fatty acids; ABA, L-2-  
18 aminobutyric acid; HDAC, histone deacetylase; TSA, trichostatin A; ERK, extracellular signal-  
19 regulated kinase; MAPK, mitogen-activated protein kinase; AMPK, AMP-activated protein  
20 kinase; TNF- $\alpha$ , Tumor necrosis factor alpha; PDH, pyruvate dehydrogenase; GPCR, G protein-  
21 coupled receptor; PKA, protein kinase A; PDK4, pyruvate dehydrogenase kinase 4; DCA,  
22 dichloroacetate; H-89, N-[2-(p-Bromocinnamylamino)ethyl]-5-isoquinolinesulfonamide  
23 dihydrochloride

24 **Abstract**

We determined the effect of butyrate and other short-chain fatty acids (SCFA) on rates of lipolysis in 3T3-L1 adipocytes. Prolonged treatment with butyrate (5 mM) increased the rate of lipolysis approximately 2-3-fold. Aminobutyric acid and acetate had little or no effect on lipolysis, however propionate stimulated lipolysis, suggesting that butyrate and propionate act through their shared activity as histone deacetylase (HDAC) inhibitors. Consistent with this, the HDAC inhibitor trichostatin A (1  $\mu$ M) also stimulated lipolysis to a similar extent as did butyrate. Western blot data suggested that neither mitogen-activated protein kinase (MAPK) activation nor perilipin down-regulation are necessary for SCFA-induced lipolysis. Stimulation of lipolysis with butyrate and trichostatin A was glucose-dependent. Changes in AMP-activated protein kinase (AMPK) phosphorylation mediated by glucose were independent of changes in rates of lipolysis. The glycolytic inhibitor iodoacetate prevented both butyrate- and Tumor necrosis factor- $\alpha$ -(TNF- $\alpha$ ) mediated increases in rates of lipolysis indicating glucose metabolism is required. However, unlike TNF- $\alpha$ -, butyrate-stimulated lipolysis was not associated with increased lactate release or inhibited by activation of pyruvate dehydrogenase (PDH) with dichloroacetate. These data demonstrate an important relationship between lipolytic activity and reported HDAC inhibitory activity of butyrate, other short-chain fatty acids and trichostatin A. Given that HDAC inhibitors are presently being evaluated for the treatment of diabetes and other disorders, more work will be essential to determine if these effects on lipolysis are due to inhibition of HDAC.

## INTRODUCTION

High rates of adipose tissue lipolysis can lead to production of excess free fatty acids (FFA). Excess FFA increase the rate of hepatic glucose output, induce skeletal muscle insulin resistance, and have other adverse effects that contribute to development of diabetes and cardiovascular disease [1-3]. Much is known about how adipose tissue lipolysis is regulated acutely or minute-to-minute but little is known about long-term regulation over time periods relevant to the progression of chronic disease. A better understanding of the mechanisms regulating rates of lipolysis over the long term may reveal new targets for therapeutic intervention [4-6].

Significant quantities of short-chain fatty acids (SCFA) are produced through fermentation of dietary fibers in the lower intestinal tract. In humans SCFA constitute approximately 10% of the caloric energy absorbed [7]. Near millimolar concentrations of butyrate are found in the hepatic portal vein, and concentrations *in vivo* may be physiologically significant for the regulation of adipocyte  $\beta$ -adrenergic receptor gene expression [7;8]. It has been reported that SCFA influence lipid metabolism,  $\beta$ -adrenergic receptor concentrations, and leptin production [8-11].

SCFA have a number of effects on cells, many of which, especially those of butyrate, are mediated through inhibition of histone deacetylases (HDACs) [12-14]. HDACs are involved in the pathogenesis of diabetes and are currently of interest as targets for the treatment of several diseases including diabetes and cancer [15-17]. In addition, SCFA have been shown to be ligands for the orphan G protein-coupled receptors (GPCR) GPR41 and GPR43 [18]. GPR41 has been reported to mediate the effects of SCFA on leptin production in adipocytes [11].

Given that both GPCR and HDACs are under active investigation as therapeutic targets for a wide spectrum of diseases, we conducted this study to determine whether SCFA affect rates of lipolysis in adipocytes.

## METHODS AND MATERIALS

*Materials:* 3T3-L1 cells (ATCC CL-173) were obtained from ATCC (Manassas, VA); glucose-containing DMEM and antibiotics were from Atlanta Biologicals (Norcross, GA); glucose-free DMEM was from Irvine Scientific (Santa Anna, CA); fetal bovine serum was from Hyclone Laboratories, Inc. (Logan, UT); insulin (Humulin<sup>®</sup>R7) was from Eli Lilly and Co. (Indianapolis, IN); BSA was from Interger Co. (Purchase, NY); I-Block was from Pierce (Rockford, IL); glutamine was from Gibco (Grand Island, NY); anti-ERK 1/2 and anti-active MAP kinase antibodies were from Promega (Madison, WI); anti-AMPK $\alpha$  and anti-phospho-AMPK $\alpha$  (T172) antibodies were from Cell Signaling Technology (Beverly, MA); secondary antibody (donkey anti-rabbit HRP conjugate) was from Santa Cruz Biotechnology (Santa Cruz, CA); and glycerol reagent for glycerol release assay was from Amresco (Solon, OH) and lactate assay reagent was from Trinity Biotech (St. Louis, MO). All other reagents were from Sigma (St. Louis, MO). Short-chain fatty acids were purchased as sodium salts and dissolved in DMEM.

*Cell Culture:* 3T3-L1 cells were cultured in 24-well plates and maintained as previously described [19] in standard medium (DMEM with high glucose, supplemented with 10% fetal bovine serum and with PSA (penicillin 100 units/ml, streptomycin 100  $\mu$ g/ml, and amphotericin 0.25  $\mu$ g/ml). Medium was changed every 2-3 days. At 2-4 days after confluence, differentiation

into adipocytes was initiated as follows: standard medium was supplemented with 5 µg/ml insulin, 0.5 µg/ml dexamethasone, and 0.5 mmol/l 3-isobutyl-1-methylxanthine for 2 days. The medium was then changed and supplemented with insulin only for 2-3 days. Thereafter, the cells were maintained in standard medium only. Cells were used 3-10 days post-differentiation. For experimental conditions without glucose, cells were incubated in DMEM without glucose, supplemented with 1% BSA, 4 mM glutamine, 44 mM NaHCO<sub>3</sub>, 20 mM HEPES and 0.01% pyruvic acid.

*Glycerol Assay:* Lipolysis was measured as the rate of glycerol release. After the various treatments, cells were washed three times with DMEM, and then incubated for another 1 hour.. Media were then collected from the cells and heated at 65°C for 8 minutes to inactivate any enzymes released from the cells. Samples (50 µl) were then assayed for glycerol using 150 µl glycerol reagent in a flat bottom 96-well plate. Absorption was measured at 500 nm on a Molecular Devices plate reader.

*Lactate Assay:* lactate concentrations were determined colorimetrically, using a kit from Trinity Bioech (St. Louis, MO), by following instructions provided by the manufacturer.

*Western Blots:* Cells were harvested in Laemmli sample buffer [20] and aspirated with a syringe five times through a 25g needle. The samples were centrifuged (16,000 g, 30 seconds) to remove fat, and then heated at 95°C for 5 minutes prior to being resolved on SDS polyacrylamide gels (10%). Proteins were transferred to nitrocellulose membranes. Membranes were blocked with 5% blotto, 1% BSA or 0.2% I-Block, and probed with polyclonal rabbit antibodies raised against perilipin (gift of Dr. Andrew Greenberg, Human Nutrition Research Center, Tufts University, Boston, MA, USA), AMPKα, phospho-AMPKα (T172), ERK1/2 or active MAP Kinase (each at dilutions of 1:3,000). After incubation with anti-rabbit IgG-HRP

(diluted 1:10,000), the blots were developed with ECL Plus and visualized with Hyperfilm ECL (Amersham Pharmacia Biotech, Piscataway, NJ).

*Statistics:* Differences between pairs of treatments were analyzed by Student's t-test. A P-value of less than 0.05 was considered statistically significant. P-values are given in the figure legends

## RESULTS

To investigate the effect of butyrate on lipolysis, 3T3-L1 adipocytes were incubated for up to 4 h with 5 mM butyrate then washed, and the rate of glycerol release was determined over the next hour (figure 1). After 1 h treatment with butyrate the rate of lipolysis was similar to that of control cells. However, there was then a time-dependent increase in the rate of lipolysis in butyrate-treated cells, with maximal stimulation occurring between 3 and 4 hours (figure 1). Maximal rates of lipolysis with butyrate were similar in time course studies for at least 48 hours (data not shown).

We next investigated the effect of a series of related SCFA (figure 2). Similar to the 4 h studies, 5 mM butyrate caused a 2-3-fold increase in the rate of lipolysis whereas 5 mM acetate or 2-aminobutyric acid (ABA) had little or no effect. We hypothesized that the known HDAC inhibitory activity of butyrate underlies its lipolytic effect, and so we evaluated the effects of 20 mM propionate, a less potent HDAC inhibitor than butyrate, and 1  $\mu$ M trichostatin A (TSA), a potent and specific small molecule HDAC inhibitor. Both propionate and TSA increased rates of lipolysis to a similar extent as did butyrate. Together these findings suggest that HDAC inhibition is involved in the lipolytic effect of these compounds.

Because we have used SCFA at concentrations at which HDAC inhibitory activity is maximal [13], the possibility remains that lower concentrations of SCFA also increase rates of lipolysis, which would imply action through an alternative mechanism. Figure 3 shows the dose-dependence of the various SCFA on rates of lipolysis. The half-maximal concentrations of butyrate and propionate were in the low millimolar range, and the relative potencies of the SCFA were butyrate>propionate>acetate. Both the absolute and the relative potencies of the SCFA indicate that inhibition of HDAC is important for increasing rates of lipolysis, and that it is unlikely they are having this effect through activation of a G protein-coupled receptor.

Butyrate has been shown to alter  $\beta$ -adrenergic receptor profiles in adipocytes [8;9]. Therefore, we investigated whether this altered  $\beta$ -receptor profile might account for the effect of butyrate on rates of lipolysis. Assuming that  $\beta$ -adrenergic receptors have some constitutive activity in the absence of agonist [21], greater concentrations of  $\beta$ -adrenergic receptors would be expected to increase rates of lipolysis by increasing cellular concentrations of cyclic AMP. The increased concentration of cyclic AMP would in turn activate PKA and hence increase rates of lipolysis. To investigate this possibility we used the PKA inhibitor H-89, which would be expected to prevent the increase in lipolysis with butyrate treatment if the mechanism involves increases in cyclic AMP concentrations. Figure 4 depicts an experiment where 3T3-L1 adipocytes were treated for 4 h with glucose, with or without 5 mM butyrate. The PKA inhibitor H-89 (50  $\mu$ M) was added for the last 30 minutes of the incubation, cells were washed, and rates of lipolysis were measured. H-89 prevented the stimulation of lipolysis by isoproterenol, the classic  $\beta$ -adrenergic receptor agonist. Similarly H-89 prevented the increased rates of lipolysis seen with butyrate suggesting that increased cyclic AMP concentrations underlie the butyrate effect on lipolysis.

Long term regulation of lipolysis by other mediators, such as TNF- $\alpha$ , is thought to involve activation of the MAP kinases ERK1&2 and the down-regulation of perilipin [22;23]. To see whether these cellular events are important for butyrate-stimulated lipolysis we treated cells with or without butyrate for 18 h and then performed Western blots on total cell extracts for these proteins (figure 5). Although TNF- $\alpha$  treatment resulted in increased lipolysis along with activation of MAP kinase (shown by phosphorylation of ERK 1/2) and down-regulation of perilipin, butyrate treatment had no such effect. These data demonstrate that neither MAP kinase activation nor perilipin down-regulation is necessary for increasing rates of lipolysis in 3T3-L1 adipocytes. This was of interest because we have previously reported that these events are also not sufficient to allow increased lipolysis in the presence of TNF- $\alpha$  [19]. In those studies we showed that the effects of TNF- $\alpha$  on rates of lipolysis were dependent on the presence of glucose although the effects on perilipin and MAPK phosphorylation were independent of glucose (see figure 5).

We have reported that the stimulatory effect of TNF- $\alpha$  on lipolysis occurs only in the presence of glucose [19]. Therefore we determined whether this glucose dependence is true also for the lipolytic effect of butyrate. Figure 6A shows that butyrate stimulates lipolysis only when glucose is present in the incubation medium. Similarly, the lipolytic effect of the HDAC inhibitor trichostatin A occurred only in the presence of glucose (figure 6B). The lipolytic effect of propionate was also glucose-dependent (data not shown). Changes in energy status due to glucose deprivation are reflected in increased AMP/ATP ratios which in turn lead to phosphorylation of AMPK. Shown in figure 7 is a western blot of total protein extracts from cells treated for 6 hours with TNF- $\alpha$  or butyrate in the presence or absence of glucose. As seen in the upper panel, phosphorylation of AMPK (T172) was decreased in cells treated with glucose

182 compared to those without, regardless of the presence of TNF- $\alpha$  or butyrate. So although  
183 AMP/ATP ratios appear to be affected by short periods of glucose deprivation, phosphorylation  
184 of AMPK is dependent only on the presence of glucose and therefore cannot explain the  
185 increased rates of lipolysis, which also require the presence of TNF- $\alpha$  or butyrate.

186 To investigate whether the glucose requirement is identical for TNF- $\alpha$  and for butyrate  
187 we investigated the requirement for glucose metabolism using the glycolytic inhibitor  
188 iodoacetate. Glyceraldehyde-3-phosphate dehydrogenase is specifically inhibited by 100  $\mu$ M  
189 iodoacetate, whereas other glycolytic enzymes are inhibited by iodoacetate only at millimolar  
190 concentrations [24].

191 Iodoacetate (100  $\mu$ M) prevented the increased rates of lipolysis with TNF- $\alpha$  and glucose  
192 (figure 8B) supporting our previous data that glucose metabolism is required for the glucose  
193 effect [19]. Iodoacetate also prevented the increased rates of lipolysis with butyrate and glucose  
194 (figure 8C). However, the glucose-dependence of TNF- $\alpha$  correlated well with the ability of  
195 glucose to be metabolized to lactate, whereas the glucose-dependence of butyrate did not.  
196 Treatment of 3T3-L1 adipocytes with TNF- $\alpha$  but not butyrate resulted in increased lactate release  
197 into the media (figure 8A). When cells were incubated in the presence of dichloroacetate (a  
198 pyruvate dehydrogenase kinase 4 inhibitor (PDK4) that promotes glucose oxidation over  
199 conversion to lactate) TNF- $\alpha$  no longer increased rates of lipolysis (figure 8B). In marked  
200 contrast, dichloroacetate had no effect on butyrate-stimulated lipolysis (figure 8C), suggesting  
201 that the glucose effects on TNF- $\alpha$ - and butyrate-stimulated lipolysis are mechanistically distinct.

## 202 DISCUSSION

203 We have demonstrated that butyrate increases the rate of glycerol release in 3T3-L1  
204 adipocytes. This effect of butyrate was slow to develop, suggesting that changes in gene

expression are involved, rather than rapid mechanisms that would affect, for example, production of a second messenger.

Butyrate and other HDAC inhibitors, such as trichostatin A, have been used for many years in the laboratory to enhance expression from viral promoters [13;25;26]. In addition, the effects of butyrate and certain other short-chain fatty acids on endogenous gene expression have been studied extensively and their effects have been attributed to their inhibition of histone deacetylase [14;27]. The histone deacetylase inhibitory activity of butyrate alters  $\beta$ -adrenergic receptor profiles in adipocytes [8;9]. Although receptor profiles were not assessed in the present study, the PKA inhibitor H-89 prevented butyrate-mediated increases in rates of lipolysis, suggesting that changes in intracellular cyclic AMP are important in the effect. The ability of H-89 to prevent butyrate-mediated increases in rates of lipolysis is in contrast to its lack of effect on serum-stimulated lipolysis in primary adipocytes [28]

While acute regulation of lipolysis (e.g., by hormones such as insulin) is well documented and factors that regulate lipolysis over longer time periods are poorly understood. However, it is known that lipolysis can be regulated chronically in vivo. For example, we have reported that expression of hormone-sensitive lipase decreases in obese subjects after weight loss [29], and short-term fasting has been reported to increase expression of this enzyme [30]. Aging and obesity are both associated with chronic alterations in adipose tissue lipolysis [31-33].

While butyrate is established as an HDAC inhibitor, we considered various other explanations for its stimulatory effect on lipolysis. One possibility was that the mechanism is similar to that of  $\text{TNF}\alpha$ , one of the few other agents known to have long-term effects on lipolysis in 3T3-L1 adipocytes. However,  $\text{TNF}\alpha$  has been reported to stimulate lipolysis through the ERK1/2 pathway in these cells, leading to down-regulation of perilipin [34;35]. We found

similar effects of  $\text{TNF}\alpha$ , but not butyrate, essentially eliminating this as a mechanism for the butyrate effect.

The small molecule HDAC inhibitor, trichostatin A mimicked the effect of butyrate on lipolysis, including the glucose-dependence of the effect. We found that the lipolytic activity of other SCFA was not directly related to carbon chain length per se, but to their relative potency as HDAC inhibitors. It is important to note that at the concentrations used, butyrate and propionate, but not acetate or ABA, exhibit significant HDAC inhibitory activity [13;14]. Propionate (three carbons instead of four) inhibits HDAC less potently than butyrate, showed a similar response with lipolysis. The effect of propionate on lipolysis was also glucose-dependent. By contrast two closely related molecules that inhibit HDAC comparatively poorly (four-carbon ABA and two-carbon acetate) did not stimulate lipolysis at concentrations where related compounds with HDAC inhibitory activity has maximal effects on lipolysis. Thus the lipolytic activity of SCFA cannot be directly attributed to carbon chain length and instead appears to be correlated with HDAC inhibitory activity.

Another possibility is that butyrate acts through a GPCR. SCFA have been shown to be ligands for the orphan G protein-coupled receptors (GPCR) GPR41 and GPR43 [18]. However, neither the absolute concentrations we have used (i.e., millimolar) nor the order of potency of the SCFA we have observed are consistent with an effect on GPR41 or GPR43. For HDAC inhibition the order of potency is butyrate>propionate>acetate [13;14], which is consistent with effects we observed on lipolysis, whereas for GPCR activation the relative order is acetate>propionate> butyrate GPR43 [18]. Second, activation of GPCR is rapid (minutes or less) whereas the effect of butyrate on lipolysis was slow to develop (hours), consistent with a

requirement for new protein synthesis that would be expected as a manifestation of the HDAC inhibitory activity.

The stimulatory effect of SCFA on lipolysis was dependent on the presence of glucose in the incubation medium. The requirement for glucose cannot be readily explained by alterations in cellular energy status. First, pyruvate was present as an alternative energy source and we previously reported that cellular ATP concentrations were similar to control after 16 hours glucose deprivation [19]. Second, phosphorylation of AMPK, a biosensor for increased intracellular AMP/ATP ratios, was dependent only on the presence of glucose, whereas increases in rates of lipolysis also required either butyrate or TNF- $\alpha$ . That AMPK activation is not causal is consistent with a recent report that concluded that activation of AMPK in adipocytes by agents that increase cyclic AMP levels is a consequence of lipolysis and not the direct result of increases in cyclic AMP levels or PKA activity [36]. Our data, previous and reported here, suggest that the primary mechanism by which TNF- $\alpha$  increases rates of lipolysis is through enhancing glucose uptake and metabolism to lactate. Indeed, both TNF- $\alpha$  and HDAC inhibitors have been shown to increase glucose uptake, but unlike TNF- $\alpha$ , the HDAC inhibitors appear to be muscle specific and do not affect glucose uptake in 3T3-L1 adipocytes [37;38]. These reports are consistent with our data showing that TNF- $\alpha$  but not butyrate increases release of lactate into the culture media. Although the glycolytic inhibitor iodoacetate prevented increased lipolysis with both TNF- $\alpha$  and butyrate, suggesting glucose metabolism is important, the differential effect with the PDK4 inhibitor dichloroacetate suggests that the mechanism of glucose action is distinct.

In conclusion, these data demonstrate that certain SCFA, as well as trichostatin A, increase the rate of lipolysis in 3T3-L1 adipocytes. Further work will be necessary to establish a

273 causal relationship between HDAC inhibition and lipolysis, and details of the mechanisms  
 274 involved. Nevertheless, because HDAC inhibitors are being actively investigated as potential  
 275 therapeutic agents for a number of diseases including diabetes [17], it will be important to  
 276 determine whether such inhibitors increase circulating free fatty acid concentrations, which  
 277 would likely worsen insulin resistance and possibly have adverse effects in diabetes

# Reference List

- [1] Bergman RN, Ader M (2000) Free fatty acids and pathogenesis of type 2 diabetes mellitus. *Trends Endocrinol Metab* 11:351-356
- [2] Ginsberg HN (2000) Insulin resistance and cardiovascular disease. *J Clin Invest* 106:453-458
- [3] Egan BM, Greene EL, Goodfriend TL (2001) Insulin resistance and cardiovascular disease. *Am J Hypertens* 14:116S-125S
- [4] Bergman RN, Mittelman SD (1998) Central role of the adipocyte in insulin resistance. *J Basic Clin Physiol Pharmacol* 9:205-221
- [5] Large V, Arner P (1998) Regulation of lipolysis in humans. Pathophysiological modulation in obesity, diabetes, and hyperlipidaemia. *Diabetes Metab* 24:409-418
- [6] Green A (2006) Cellular regulation of lipolysis. In: Runge MS, Pattererson C (eds.) *Principles of Molecular Medicine*, 2 ed. Humana Press, Inc., Totoya, NJ, pp 518-523
- [7] Bergman EN (1990) Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev* 70:567-590
- [8] Krief S, Feve B, Baude B, Zilberfarb V, Strosberg AD, Pairault J, Emorine LJ (1994) Transcriptional modulation by n-butyric acid of beta-1, beta-2 and beta-3 adrenergic receptor balance in 3T3-F442A adipocytes. *J Biol Chem* 269:6664-6670
- [9] Ding ST, Smith EO, McNeel RL, Mersmann HJ (2000) Modulation of porcine adipocyte beta-adrenergic receptors by hormones and butyrate. *J Anim Sci* 78:927-933
- [10] Metz SHM, Lopes-Cardoza M, Van den Berg G (1974) Inhibition of lipolysis in bovine adipose tissue by butyrate and  $\beta$ -hydroxybutyrate. *FEBS Lett* 47:19-22
- [11] Xiong YM, Miyamoto N, Shibata K, Valasek MA, Motoike T, Kedzierski RM, Yanagisawa M (2004) Short-chain fatty acids stimulate leptin production in adipocytes through the G protein-coupled receptor GPR41. *Proc Natl Acad Sci USA* 101:1045-1050

- [12] Waldecker M, Kautenburger T, Daumann H, Busch C, Schrenk D (2008) Inhibition of histone-deacetylase activity by short-chain fatty acids and some polyphenol metabolites formed in the colon. *J Nutr Biochem* 19:587-593
- [13] Marshall S, Duong T, Wu T, Hering MA, Yada J, Higgins S, Orbus RJ, Yan ZH, Rumberger JM (2003) Enhanced expression of uridine diphosphate-N-acetylglucosaminyl transferase (OGT) in a stable, tetracycline-inducible HeLa cell line using histone deacetylase inhibitors: kinetics of cytosolic OGT accumulation and nuclear translocation. *Anal Biochem* 319:304-313
- [14] Kruh J (1982) Effects of sodium butyrate, a new pharmacological agent, on cells in culture. *Mol Cell Biochem* 42:65-82
- [15] Das C, Kundu TK (2005) Transcriptional regulation by the acetylation of nonhistone proteins in humans -- a new target for therapeutics. *IUBMB Life* 57:137-149
- [16] Gray SG, De Meyts P (2005) Role of histone and transcription factor acetylation in diabetes pathogenesis. *Diabetes Metab Res Rev* 21:416-433
- [17] Christensen DP, Dahlhoff M, Lundh M, Rasmussen DM, Nielsen MD, Billestrup N, Grunnet LG, Mandrup-Poulsen TM (2011) Histone Deacetylase (HDAC) Inhibition as a Novel Treatment for Diabetes Mellitus. *Mol Med* 17:378-390
- [18] Brown AJ, Goldsworthy SM, Barnes AA, Eilert MM, Tcheang L, Daniels D, Muir AI, Wigglesworth MJ, Kinghorn I, Fraser NJ, Pike NB, Strum JC, Steplewski KM, Murdock PR, Holder JC, Marshall FH, Szekeres PG, Wilson S, Ignar DM, Foord SM, Wise A, Dowell SJ (2003) The orphan G protein-coupled receptors GPR41 and GPR43 are activated by propionate and other short chain carboxylic acids. *J Biol Chem* 278:11312-11319
- [19] Green A, Rumberger JM, Stuart CA, Ruhoff MS (2004) Stimulation of lipolysis by TNF- $\alpha$  in 3T3-L1 adipocytes is glucose-dependent: implications for long-term regulation of lipolysis. *Diabetes* 53:74-81
- [20] Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685
- [21] Chidiac P, Hebert TE, Valiquette M, Dennis M, Bouvier M (1994) Inverse agonist activity of beta-adrenergic antagonists. *Mol Pharmacol* 45:490-499
- [22] Souza SC, De Vargas LM, Yamamoto MT, Lien P, Franciosa MD, Moss LG, Greenberg AS (1998) Overexpression of perilipin A and B blocks the ability of tumor necrosis factor  $\alpha$  to increase lipolysis in 3T3-L1 adipocytes. *J Biol Chem* 273:24665-24669
- [23] Gronning LM, Cederberg A, Miura N, Enerbäck S, Taskén K (2002) Insulin and TNF $\alpha$  induce expression of the forkhead transcription factor gene *Foxc2* in 3T3-L1 adipocytes via PI3K and ERK 1/2-dependent pathways. *Mol Endocrinol* 16:873-883
- [24] Webb JL. Enzyme and metabolic inhibitors. New York, NY: Academic Press, Inc.; 1966.
- [25] Li SH, Burns DK, Rumberger JM, Presky DH, Wilkinson VL, Anostario MJr, Wolitzky BA, Norton CR, Familletti PC, Kim JJ (1994) The consensus repeat domains of E-selectin enhance ligand binding. *J Biol Chem* 269:4431-4437
- [26] Condreay JP, Witherspoon SM, Clay WC, Kost TA (1999) Transient and stable gene expression in mammalian cells transduced with a recombinant baculovirus vector. *Proc Natl Acad Sci USA* 96:127-132
- [27] Sealy L, Chalkley R (1978) The effect of sodium butyrate on histone modification. *Cell* 14:115-121

- [28] Rumberger JM, Peters T, Jr., Burrington C, Green A (2004) Transferrin and iron contribute to the lipolytic effect of serum in isolated adipocytes. *Diabetes* 53:2535-2541
- [29] Klein S, Luu K, Gasic S, Green A (1996) Effect of weight loss on whole body and cellular lipid metabolism in severely obese humans. *Am J Physiol Endocrinol Metab* 270:E739-E745
- [30] Sztalryd C, Kraemer FB (1994) Regulation of hormone-sensitive lipase during fasting. *Am J Physiol* 266:E179-E185
- [31] Arner P (1999) Catecholamine-induced lipolysis in obesity. *Int J Obesity Rel Metab Dis* 23 Suppl 1:10-13
- [32] Dax EM, Partilla JS, Gregerman RI (1981) Mechanism of the age-related decrease of epinephrine-stimulated lipolysis in isolated rat adipocytes: beta-adrenergic receptor binding, adenylate cyclase activity and cyclic AMP accumulation. *J Lipid Res* 22:934-943
- [33] Hickner RC, Racette SB, Binder EF, Fisher JS, Kohrt WM (1999) Suppression of whole body and regional lipolysis by insulin: Effects of obesity and exercise. *J Clin Endocrinol Metab* 84:3886-3895
- [34] Rosenstock M, Greenberg AS, Rudich A (2001) Distinct long-term regulation of glycerol and non-esterified fatty acid release by insulin and TNF- $\alpha$  in 3T3-L1 adipocytes. *Diabetologia* 44:55-62
- [35] Souza SC, Palmer HJ, Kang YH, Yamamoto MT, Muliro KV, Paulson KE, Greenberg AS (2003) TNF- $\alpha$  induction of lipolysis is mediated through activation of the extracellular signal related kinase pathway in 3T3-L1 adipocytes. *J Cell Biochem* 89:1077-1086
- [36] Gauthier MS, Miyoshi H, Souza SC, Cacicedo JM, Saha AK, Greenberg AS, Ruderman NB (2008) AMP-activated protein kinase (AMPK) is activated as a consequence of lipolysis in the adipocyte: potential mechanism and physiological relevance. *J Biol Chem* M708177200
- [37] Wang CN, O'Brien L, Brindley DN (1998) Effects of cell-permeable ceramides and tumor necrosis factor- $\alpha$  on insulin signaling and glucose uptake in 3T3-L1 adipocytes. *Diabetes* 47:24-31
- [38] Takigawa-Imamura H, Sekine T, Murata M, Takayama K, Nakazawa N, Nakagawa J (2003) Stimulation of Glucose Uptake in Muscle Cells by Prolonged Treatment with Scriptide, a Histone Deacetylase Inhibitor. *Bioscience, Biotechnology, and Biochemistry* 67:1499-1506

## FIGURE LEGENDS

*Figure 1. Time dependence of butyrate on rates of lipolysis in 3T3-L1 adipocytes.* 3T3 L1 adipocytes were incubated with nothing (●-●) or 5 mM butyrate (○-○). At the indicated times cells were washed and incubated for one hour, and then glycerol was measured as an index of the rate of lipolysis, as described in the methods section. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3). \* P<0.05; \*\* P<0.003; \*\*\* P<0.001.

*Figure 2. Effect of Short-Chain fatty acids and trichostatin A on lipolysis.* 3T3-L1 adipocytes were treated for 18 hours with no additions (Ctl); 5 mM Butyrate (But), 5 mM 2-amino-butyrate (ABA), 5 mM acetate (Ace); 20 mM propionate (Pro); or 1  $\mu$ M Trichostatin A (TSA). Rates of glycerol release were then determined as in the legend to figure 1. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3). \*\* P<0.05; \* P<0.0001 (TSA compared to control).

*Figure 3. Relative lipolytic potencies of Short-Chain fatty acids.* 3T3-L1 adipocytes were treated for 24 hrs with the indicated concentrations of Short-Chain fatty acids. After incubation, cells were washed and rates of lipolysis determined as in the legend to figure 1. Panel (A) Comparison of butyrate (○-○) with ABA (●-●). The concentrations were 0, 1, 2, 3 and 5 mM. Panel (B) Comparison of propionate (○-○) with acetate (●-●). Concentrations were 0, 1, 3, 10 and 30 mM. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3).

*Figure 4. Effect of the PKA inhibitor H-89 on rates of lipolysis.* 3T3-L1 adipocytes were treated for 4 hr with glucose plus no further additions (Ctl) or 5 mM butyrate (But). Isoproterenol (1  $\mu$ M) and H-89 (50  $\mu$ M) were added as indicated and glycerol was measured after another 30 min. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3). \* P<0.01

*Figure 5. Effect of butyrate on MAPK and perilipin.* 3T3-L1 adipocytes were treated with or without 25 mM glucose and 50 ng/ml TNF- $\alpha$  or 5 mM butyrate for 24 hr. Total protein extracts were prepared and Western blots were performed with antibodies raised against (A) perilipin, (B) phosphorylated MAP Kinase or (C) total MAP Kinase. Data depicted are representative of three or more independent experiments.

*Figure 6. Effect of glucose on the lipolytic effect of butyrate and trichostatin A* (Panel A) 3T3-L1 adipocytes were incubated in glucose free media supplemented with pyruvate for 24 hrs with no additions (Ctl), 50 ng/ml TNF- $\alpha$  (TNF), 5 mM butyrate (But), 25 mM glucose (Glc), or in combinations as indicated. Cells were washed and glycerol release measured as in the legend to figure 1. (Panel B) 3T3-L1 adipocytes were treated with 1  $\mu$ M Trichostatin A (TSA) in the presence or absence of 25 mM glucose. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3). \* P<0.0001; \*\* P<0.001.

*Figure 7. Effect of butyrate on phosphorylation of AMPK.* 3T3-L1 adipocytes were treated with or without 25 mM glucose and 50 ng/ml TNF- $\alpha$  or 5 mM butyrate for 6 hr. Total protein extracts were prepared and Western blots were performed with antibodies raised against phospho-AMPK

(T172) (upper panel)) or total AMPK (lower panel). Data depicted are representative of three or more independent experiments.

*Figure 8. Effect of iodoacetate and dichloroacetate.* (A) 3T3-L1 adipocytes were treated in 25 mM glucose for 6 hr with no additions (ctl), 50 ng/ml TNF- $\alpha$  (TNF), or 5 mM butyrate. (B) Adipocytes were treated with TNF and glucose and either 100  $\mu$ M iodoacetate (Iod) or 100  $\mu$ M dichloroacetate (DCA) and then glycerol release was measured as in the legend to figure 1. (C) Same conditions as panel B, but with butyrate. Data depicted are representative of three or more independent experiments. Data shown are means  $\pm$  SE (n=3). \* P<0.05; \*\* P<0.003

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8

