

A peer-reviewed version of this preprint was published in PeerJ on 1 December 2017.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.4120) (peerj.com/articles/4120), which is the preferred citable publication unless you specifically need to cite this preprint.

Ara T, Sogawa N. 2017. Effects of shinbuto and ninjinto on prostaglandin E₂ production in lipopolysaccharide-treated human gingival fibroblasts. PeerJ 5:e4120 <https://doi.org/10.7717/peerj.4120>

Effects of shinbuto and ninjinto on prostaglandin E₂ production in lipopolysaccharide-treated human gingival fibroblasts

Toshiaki Ara¹ and Norio Sogawa¹

¹Department of Pharmacology, Matsumoto Dental University, 1780 Gobara Hirooka, Shiojiri, Nagano 399-0781, Japan

Corresponding author:
Toshiaki Ara¹

Email address: ara.t@po.mdu.ac.jp

ABSTRACT

Previously, we revealed that several kampo medicines that are used for patients with excess and/or medium patterns [kakkonto (TJ-1), shosaikoto (TJ-9), hangeshashinto (TJ-14), and orento (TJ-120)] reduced prostaglandin (PG)E₂ levels using LPS-treated human gingival fibroblasts (HGFs). Recently, we examined other kampo medicines used for patients with the deficiency pattern [bakumondoto (TJ-29), shinbuto (TJ-30), ninjinto (TJ-32), and hochuekkito (TJ-41)] and the herbs comprising shinbuto and ninjinto using the same experimental model. Shinbuto and ninjinto concentration-dependently reduced LPS-induced PGE₂ production by HGFs, whereas hochuekkito weakly reduced and bakumondoto did not reduce PGE₂ production. Shinbuto and ninjinto did not alter cyclooxygenase (COX) activity or the expression of molecules involved in the arachidonic acid cascade. Therefore, we next examined which herbs comprising shinbuto and ninjinto reduce LPS-induced PGE₂ production. Among these herbs, shokyo (*Zingiberis Rhizoma*) and kankyo (*Zingiberis Processum Rhizoma*) strongly and concentration-dependently decreased LPS-induced PGE₂ production. However, both shokyo and kankyo increased the expression of cytosolic phospholipase (cPL)A₂ but did not affect annexin1 or COX-2 expression. These results suggest that shokyo and kankyo suppress cPLA₂ activity. We demonstrated that kampo medicines suppress inflammatory responses in patients with the deficiency pattern, and in those with excess or medium patterns. Moreover, kampo medicines that contain shokyo or kankyo are considered to be effective for the treatment of inflammatory diseases.

INTRODUCTION

Periodontal disease is an inflammatory disease of the gingiva that destroys periodontal tissues. In severe cases, alveolar bone is absorbed. In inflammatory responses and tissue degradation, prostaglandin E₂ (PGE₂), interleukin (IL)-6, and IL-8 play important roles. As PGE₂ has several functions in vasodilation, the enhancement of vascular permeability and pain, and osteoclastogenesis induction, PGE₂ participates in inflammatory responses and alveolar bone resorption in periodontal disease (Noguchi and Ishikawa, 2007).

Previously, we reported that several kampo medicines, shosaikoto (TJ-9) (Ara et al., 2008b), orento (TJ-120) (Ara et al., 2010), hangeshashinto (TJ-14) (Nakazono et al., 2010), and kakkonto (TJ-1) (Kitamura et al., 2014), suppress lipopolysaccharide (LPS)-induced PGE₂ production by human gingival fibroblasts (HGFs). Moreover, we found that shokyo, kanzo, and keihi, which are herbs contained in kakkonto, reduce PGE₂ production (Ara and Sogawa, 2016). These results suggested that these kampo medicines and herbs have anti-inflammatory effects in periodontal disease.

However, these kampo medicines are used for patients with the excess pattern or medium pattern. Kampo medicine used for those with the deficiency pattern remains to be elucidated. In the present study, we therefore examined the anti-inflammatory effects of the kampo medicines for patients with the

deficiency pattern [bakumondoto (TJ-29), shinbuto (TJ-30), ninjinto (TJ-32), and hochuekkito (TJ-41)], which are used for the treatment of inflammatory diseases. Furthermore, we examined the effects on PGE₂ production using herbs comprising the kampo medicines that reduce PGE₂ production.

MATERIALS AND METHODS

Reagents

Kampo medicines (bakumondoto, shinbuto, ninjinto, and hochuekkito) were purchased from Tsumura & Co. (Tokyo, Japan). Powders of 8 herbs (bukuryo, bushi, kankyo, kanzo, ninjin, shakuyaku, shokyo, and sojutsu) were provided by Tsumura & Co. The ingredients in shinbuto and ninjinto formulas are shown in Tables 1 and 2. Powders of kampo medicines or herbs were suspended in Dulbecco's modified Eagle's medium (D-MEM, Sigma, St. Louis, MO) containing 10% heat-inactivated fetal calf serum, 100 units/ml penicillin, and 100 mg/ml streptomycin (culture medium), and were rotated at 4°C overnight. Then, the suspensions were centrifuged and the supernatants were filtrated through a 0.45 μm-pore membrane. Lipopolysaccharide (LPS) from *Porphyromonas gingivalis* 381 was provided by Professor Nobuhiro Hanada (School of Dental Medicine, Tsurumi University, Japan). Arachidonic acid was purchased from Cayman Chemical (Ann Arbor, MI). Other reagents were purchased from Nacalai tesque (Kyoto, Japan).

Cells

HGFs were prepared as described previously (Nakazono et al., 2010). In brief, HGFs were prepared from free gingiva during the extraction of an impacted tooth with the informed consent of the subjects who consulted Matsumoto Dental University Hospital. The free gingival tissues were cut into pieces and seeded onto 24-well plates (AGC Techno Glass Co., Chiba, Japan). HGFs were maintained in culture medium at 37°C in a humidified atmosphere of 5% CO₂. For passage, HGFs were trypsinized, suspended, and plated into new cultures in a 1:3 dilution ratio. HGFs were used between the 10th to 15th passages in the assays. This study was approved by the Ethical Committee of Matsumoto Dental University (No. 0063).

Measurement of cell viability

The numbers of cells were measured using WST-8 (Cell Counting Kit-8; Dojindo, Kumamoto, Japan) according to the manufacturer's instructions. In brief, the media were removed by aspiration and the cells were treated with a 100-μl mixture of WST-8 with culture medium for 2 h at 37°C in CO₂ incubator. Optical density was measured (measured wavelength at 450 nm and reference wavelength at 655 nm) using an iMark microplate reader (Bio-Rad, Hercules, CA), and the mean background value was subtracted from each value. Data is represented as means ± S.D. (n = 4).

Measurement of prostaglandin E₂ (PGE₂), interleukin (IL)-6, and IL-8

HGFs were seeded in 96-well plates (10,000 cells/well) and incubated in serum-containing medium at 37°C overnight. Then, the cells were treated with varying concentrations of each kampo medicine (0, 0.01, 0.1, or 1 mg/ml) or each herb (0, 10, 30, or 100 μg/ml) in the absence or presence of LPS (10 ng/ml) for 24 h (200 μl per well) in triplicate or quadruplicate for each sample. After the culture supernatants were collected, viable cell numbers were measured using WST-8 as described above.

The concentrations of PGE₂, IL-6, and IL-8 in the culture supernatants were measured by enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions (PGE₂, Cayman Chemical; IL-6 and IL-8, Thermo Fisher Scientific Inc., Camarillo, MA, USA), and were adjusted by the number of viable cells. Data are represented as pg or ng per 10,000 cells (mean ± S.D.).

Measurement of cyclooxygenase (COX)-2 activity

COX-2 activity was evaluated as reported previously (Wilborn et al., 1995) with slight modification. In brief, to estimate COX-2 activity, HGFs were treated with LPS and herbs for 8 h, washed, and incubated in culture medium containing exogenous arachidonic acid (10 μM). The concentrations of PGE₂ in the supernatants were measured by ELISA. Data are represented as pg per 10,000 cells (mean ± S.D.).

Preparation of cell lysates

HGFs were cultured in 60-mm dishes and treated with combinations of LPS and herbs for the indicated times. Then, cells were washed twice with Tris-buffered saline, transferred into microcentrifuge tubes,

and centrifuged at $6,000 \times g$ for 5 min at 4°C . Supernatants were aspirated and cells were lysed on ice in lysis buffer [50 mM Tris-HCl, pH 7.4, 1% Nonidet P-40, 0.25% sodium deoxycholate, 150 mM NaCl, 1 mM ethyleneglycol bis(2-aminoethylether)tetraacetic acid (EGTA), 1 mM sodium orthovanadate, 10 mM sodium fluoride, 1/100 volume of protease inhibitor cocktail (Nacalai tesque)] for 30 min at 4°C . Samples were next centrifuged at $12,000 \times g$ for 15 min at 4°C , and supernatants were collected. The protein concentration was measured using a BCA Protein Assay Reagent kit (Pierce Chemical Co., Rockford, IL).

Western blotting

The samples (10 μg of protein) were fractionated in a polyacrylamide gel under reducing conditions and transferred onto a polyvinylidene difluoride (PVDF) membrane (Hybond-P; GE Healthcare, Uppsala, Sweden). The membranes were blocked with 5% ovalbumin for 1 h at room temperature and incubated with primary antibody for an additional 1 h. The membranes were further incubated with horseradish peroxidase-conjugated secondary antibodies for 1 h at room temperature. Protein bands were visualized with an ECL kit (GE Healthcare). Densitometric values of each band were calculated using ImageJ software.

Antibodies against COX-2 (sc-1745, 1:500 dilution), cytosolic PLA₂ (cPLA₂) (sc-438, 1:200 dilution), annexin 1 (sc-11387, 1:1,000 dilution), and actin (sc-1616, 1:1,000 dilution), which detects a broad range of actin isoforms, were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Antibodies against extracellular signal-regulated kinase (ERK; p44/42 MAP kinase antibody, 1:1,000 dilution) and phosphorylated ERK [Phospho-p44/42 MAPK (Thr202/Tyr204) (E10) monoclonal antibody, 1:2,000 dilution] were from Cell Signaling Technology (Danvers, MA). Horseradish peroxidase-conjugated anti-goat IgG (sc-2020, 1:20,000 dilution) was from Santa Cruz, and anti-rabbit IgG (1:20,000 dilution) and anti-mouse IgG (1:20,000 dilution) were from DakoCytomation (Glostrup, Denmark).

Statistical analysis

Differences between groups were evaluated by the two-tailed pairwise comparison test with a pooled variance, followed by correction with the Holm method (total 10 null hypotheses; 5 null hypotheses without kampo vs. with kampo in the absence or presence of LPS in Fig. 1, total 10 null hypotheses; 3 null hypotheses without kampo vs. with kampo in the absence of LPS, 3 null hypotheses without kampo vs. with kampo in the presence of LPS, and 4 null hypotheses without LPS vs. with LPS in Fig. 2). Differences between the control group and experimental groups were evaluated by a two-tailed Dunnett's test (Figs. 3 and 6).

All computations were performed with the statistical program R (<http://www.r-project.org/>). Dunnett's test was performed using the 'glht' function in the 'multcomp' package. Values with $P < 0.05$ were considered significantly different.

RESULTS

Effects of kampo medicines on HGFs viability

First, we examined the effects of four kampo medicines (bakumondoto, shinbuto, ninjinto, and hochuekkito) on HGFs viability. Bakumondoto did not affect the viability up to 10 mg/ml at 24 h treatment (Fig. 1). In contrast, Shinbuto, ninjinto, and hochuekkito did not affect the viability up to 2 mg/ml, but decreased at 5 mg/ml and 10 mg/ml (Fig. 1). Therefore, up to 1 mg/ml of kampo medicines was used in further experiments because we used the same concentration of kampo medicines in previous studies (Ara et al., 2008b, 2010; Nakazono et al., 2010; Kitamura et al., 2014).

Effects of kampo medicines on prostaglandin (PGE)₂, interleukin (IL)-6, and IL-8 production

We examined whether these kampo medicines affected the production of PGE₂ and inflammatory cytokines (IL-6 and IL-8) by HGFs. The concentrations of PGE₂, IL-6, and IL-8 were adjusted according to viable cell number. HGFs treated with 10 ng/ml of LPS produced large amounts of PGE₂, IL-6, and IL-8. Shinbuto and ninjinto strongly and concentration-dependently reduced LPS-induced PGE₂ production (Fig. 2). In contrast, bakumondoto and hochuekkito had no or little effect on PGE₂ production. Bakumondoto weakly, and shinbuto, ninjinto, and hochuekkito strongly increased LPS-induced IL-6 production (Fig. 2). Bakumondoto and hochuekkito weakly increased LPS-induced IL-8 production, but shinbuto and ninjinto did not affect IL-8 production (Fig. 2).

From these results, we selected two kampo medicines, shinbuto and ninjinto, which decreased PGE₂ production and used them in the following experiments.

Effects of shinbuto and ninjinto on the arachidonic acid cascade

To clarify the mechanism of how shinbuto and ninjinto reduced LPS-induced PGE₂ production more directly, we examined the effects of these two kampo medicines on the arachidonic acid cascade. First, we examined the effects of shinbuto and ninjinto on COX activity. In order to bypass PLA₂, we added exogenous arachidonic acid to HGFs treated with LPS alone or LPS plus kampo medicine (shinbuto or ninjinto). Then, we measured the PGE₂ level produced by COX. However, shinbuto and ninjinto did not affect LPS-induced PGE₂ production (Fig. 3).

Next, we examined whether shinbuto and ninjinto affected the expression of molecules in the arachidonic acid cascade. cPLA₂, which is the most upstream enzyme in the arachidonic acid cascade, releases arachidonic acid from plasma membranes. Shinbuto slightly reduced cPLA₂ expression and ninjinto slightly increased cPLA₂ expression (Fig. 4A). COX-2 was weakly expressed in the absence of LPS, and the treatment with LPS alone increased COX-2 expression. However, shokyo did not alter but kankyo slightly increased LPS-induced COX-2 expression (Fig. 4). Annexin1 (also named lipocortin1) is produced by glucocorticoids and inhibits cPLA₂ activity (Gupta et al., 1984; Wallner et al., 1986). Shinbuto and ninjinto slightly increased annexin1 expression (Fig. 4A) in a concentration-dependent manner (Fig. 4B).

Lastly, we evaluated the effects of shinbuto and ninjinto on ERK phosphorylation. cPLA₂ is directly phosphorylated and activated by phosphorylated ERK (Lin et al., 1993; Gijón et al., 1999). Therefore, we examined whether shinbuto and ninjinto suppressed LPS-induced ERK phosphorylation. LPS treatment enhanced ERK phosphorylation at 0.5 h and its phosphorylation was attenuated. However, 1 mg/ml of shinbuto or ninjinto did not affect LPS-induced ERK phosphorylation (Fig. 5).

Effects of herbs on PGE₂ production and molecular expression in the arachidonic acid cascade

We examined whether herbs which comprising shinbuto and ninjinto affected LPS-induced PGE₂, IL-6 and IL-8 production by HGFs. When HGFs cells were treated with 10 ng/ml of LPS, HGFs cells produced large amounts of PGE₂. Bukuryo increased LPS-induced PGE₂ production. Shokyo, kankyo and kanzo strongly and significantly reduced LPS-induced PGE₂ production (Fig. 6A). Moreover, shokyo and kankyo decreased PGE₂ production in a concentration-dependent manner (Fig. 6B). Other herbs had little or no effect on PGE₂ production. Bukuryo increased LPS-induced IL-6 and IL-8 production, and kankyo increased IL-8 production. Kanzo reduced IL-6 production (Fig. 6A).

We then examined whether shokyo and kankyo affected the expression of molecules in the arachidonic acid cascade. Both shokyo and kankyo increased the expression of cPLA₂ but did not affect annexin1 or COX-2 expression (Fig. 7).

DISCUSSION

In our previous studies, we reported the importance of HGFs in the study of periodontal disease (Kamemoto et al., 2009; Ara et al., 2010; Nakazono et al., 2010; Ara et al., 2012; Kitamura et al., 2014; Ara and Sogawa, 2016), because HGFs are the most prominent cells in periodontal tissue. Moreover, LPS-treated HGFs produce inflammatory chemical mediators, such as PGE₂ and inflammatory cytokines such as IL-6 and IL-8 (Sismey-Durrant and Hopps, 1991; Bartold and Haynes, 1991; Tamura et al., 1992). Moreover, HGFs continue to produce PGE₂ (Ara et al., 2008a), IL-6, and IL-8 (Ara et al., 2009) in the presence of LPS. Therefore, the large amount of chemical mediators and cytokines derived from HGFs may be contained in periodontal tissues. From these findings, we believe that examining the effects of drugs on HGFs is needed in the study of periodontal disease.

In the present study, we examined the effects of kampo medicines on LPS-induced PGE₂, IL-6, and IL-8 production by HGFs in patients with the deficiency pattern. Shinbuto and ninjinto dose-dependently reduced LPS-induced PGE₂ production (Fig. 2), similar with shosaikoto, hangeshashinto, orento, and kakkonto (Ara et al., 2008b; Nakazono et al., 2010; Ara et al., 2010; Kitamura et al., 2014). However, shinbuto and ninjinto increased LPS-induced IL-6 and IL-8 production (Fig. 2). In general, acid non-steroidal anti-inflammatory drugs (NSAIDs) exhibit anti-inflammatory effects by suppressing PGE₂

production even though they do not affect IL-6 or IL-8 production. Therefore, our results suggest that shinbuto and ninjinto have anti-inflammatory effects in periodontal disease similar with acid NSAIDs.

In the experiments at the herb level, shokyo (*Zingiberis Rhizoma*), kankyo (*Zingiberis Processum Rhizoma*), and kanzo (*Glycyrrhizae Radix*) reduced PGE₂ production (Fig. 6). Shokyo is contained in shinbuto (Table 1), and kankyo and kanzo are contained in ninjinto (Table 2). Shokyo is the powdered rhizome of ginger (*Zingiber officinale* Roscoe), whereas, kankyo is the steamed and powdered rhizome of ginger. Many reports demonstrated that ginger has anti-inflammatory effects in human (Afzal et al., 2001; Lakhan et al., 2015), animal (Thomson et al., 2002; Aimbire et al., 2007; El-Abhar et al., 2008), and *in vitro* models (Ara and Sogawa, 2016; Podlogar and Verspohl, 2012). Shokyo contains gingerols such as 6-, 8-, and 10-gingerols. With prolonged storage or heat-treatment of ginger, gingerols are converted to shogaols, which are the dehydrated form of the gingerols (Afzal et al., 2001). Therefore, kankyo contains the largest amount of shogaols.

Recently, we found that shokyo suppressed LPS-induced PGE₂ production by HGFs and that shokyo may suppress PLA₂ activity (Ara and Sogawa, 2016). In the present study, we examined the effects of kankyo in comparison with shokyo. Shokyo and kankyo increased cPLA₂ expression but did not alter annexin 1 expression (Fig. 7). Moreover, we revealed that shinbuto and ninjinto, which contain shokyo and kankyo respectively, did not alter PGE₂ production when arachidonic acid was added to bypass the upstream pathway (Fig. 3). These data suggest that shokyo and kankyo did not affect the downstream pathway of arachidonic acid, which includes COX-2 and PGE synthase. In addition, shinbuto and ninjinto did not affect ERK phosphorylation (Fig. 5). From our findings described above, we were unable to explain the mechanism of the reduction in PGE₂ production. As gingerols in ginger are reported to inhibit both calcium-independent PLA₂ (iPLA₂) and cPLA₂ activities (Nievergelt et al., 2011), shokyo and kankyo are suggested to inhibit PLA₂ as discussed in the previous study (Ara and Sogawa, 2016). Previously, we reported that cPLA₂ is the main isoform in HGFs (Ara and Sogawa, 2016) among the subtypes such as cPLA₂, iPLA₂, and secretory PLA₂ (sPLA₂) (Burke and Dennis, 2009). Therefore, shokyo and kankyo may mainly inhibit cPLA₂ activity in HGFs. We found that orento decreases LPS-induced PGE₂ production via the suppression of ERK phosphorylation (Ara et al., 2010). However, orento may also reduce LPS-induced PGE₂ production by inhibition of cPLA₂ activity because orento contains kankyo.

We demonstrated that shokyo and kankyo concentration-dependently reduced LPS-induced PGE₂ production (Fig. 6A), and that the effects of kankyo are slightly stronger than those of shokyo (Fig. 6B). In previous study, 6- and 8-gingerols were found to not inhibit cPLA₂ activity, but 10-gingerol and 6-, 8-, and 10-shogaols did (Nievergelt et al., 2011). Therefore, the difference in these effects on PGE₂ production between shokyo and kankyo may be due to the amount of shogaols in these herbs.

We demonstrated that shinbuto and ninjinto slightly increased annexin1 expression (Fig. 4). However, the involvement of annexin1 in the reduction in PGE₂ production is unlikely. Shokyo and kankyo did not alter annexin1 expression (Fig. 7). All 4 herbs other than shokyo in shinbuto did not reduce PGE₂ production, but rather, bukuryo increased PGE₂ production (Fig. 6A). Similarly, kanzo in ninjinto increased annexin1 expression in HGFs, and kanzo also inhibited COX activity (Ara and Sogawa, 2016). The 2 residual herbs other than kankyo and kanzo did not reduce PGE₂ production (Fig. 6A). Therefore, the increased annexin1 expression did not contribute to decreased PGE₂ production.

At the herb level, we were unable to clarify which herbs affect cytokine production. Bukuryo in shinbuto increased LPS-induced IL-6 and IL-8 production. Therefore, this effect of shinbuto on increased IL-6 production may be due to bukuryo. However, shinbuto did not alter IL-8 production even though it contains bukuryo. Moreover, although ninjinto increased LPS-induced IL-6 production, kanzo reduced IL-6 production, and the other three herbs, kankyo, sojutsu, and ninjin, did not alter IL-6 production. Similarly, although ninjinto did not alter IL-8 production, kankyo increased IL-8 production. Therefore, the effects of herbs on IL-6 and IL-8 production are considered to not be due to a single herb but to the combination of herbs.

Both the expression of COX-2, and the production of IL-6 and IL-8 are widely known to be regulated by NF-κB. Ginger and its components gingerol and shogaol have been reported to suppress NF-κB activation, and to reduce COX-2 expression and the production of IL-6 and IL-8. For example, ginger suppressed NF-κB activation in ovarian cancer cells (Rhode et al., 2007), and 6-gingerol suppressed NF-κB activation in mouse macrophage RAW264.7 cells (Pan et al., 2008), TPA-treated mouse skin *in vivo* (Kim et al., 2005), and in intestinal epithelial cells (Saha et al., 2016). Similarly, 6-shogaol

suppressed NF- κ B activation in mouse macrophage RAW264.7 cells (Pan et al., 2008) and microglia cells (Ha et al., 2012). 6-Gingerol and 6-shogaol suppressed COX-2 expression in mouse macrophage RAW264.7 cells (Pan et al., 2008) and primary rat astrocytes (Shim et al., 2011). 6-Gingerol reduced the production of IL-1 α , IL-1 β , IL-6, and IL-8 in intestinal epithelial cells (Saha et al., 2016). However, shinbuto and ninjinto, which contain shokyo and kankyo, respectively, increased LPS-induced IL-6 and IL-8 production by HGFs (Fig. 2) similar with kakkonto (Kitamura et al., 2014). Moreover, these two kampo medicines, shokyo, and kankyo did not suppress COX-2 expression (Figs. 4A and 7). These findings raised the possibility that shokyo and kankyo, their components, gingerols and shogaols, do not suppress the NF- κ B pathway in HGFs. The assumption is able to explain why shokyo and kankyo did not suppress COX-2 expression, which is also regulated by the NF- κ B pathway. Furthermore, 6-gingerol and 6-shogaol had no effect on LPS-induced IL-8 production in human bronchial epithelial cells (Podlogar and Verspohl, 2012). Therefore, the effects of gingerols and shogaols may be different among cell types.

CONCLUSION

We demonstrated that shinbuto and ninjinto reduced LPS-induced PGE₂ production by HGFs. Moreover, shokyo and kankyo, which are included in these kampo medicines respectively, concentration-dependently reduced LPS-induced PGE₂ production. However, shokyo and kankyo did not alter the expression of the molecules in the arachidonic acid cascade, suggesting that shokyo and kankyo inhibit cPLA₂ activity. Therefore, the kampo medicines that contain shokyo or kankyo may have the ability to reduce PGE₂ production. We found that the kampo medicines used for patients with the deficiency pattern also have anti-inflammatory effects in those with the excess pattern or medium pattern. We expect kampo medicines to be used for improving inflammatory diseases, such as periodontal disease and stomatitis, in patients with any pattern.

ETHICAL APPROVAL

This study was approved by the Ethical Committee of Matsumoto Dental University (No. 0063).

CONFLICT OF INTERESTS

The authors have no conflicts of interests to disclose.

ACKNOWLEDGMENTS

We thank Prof. Nobuo Yoshinari (Department of Periodontology, Matsumoto Dental University) for HGFs preparation. We also thank Prof. Nobuyuki Udagawa (Department of Biochemistry, Matsumoto Dental University) and Prof. Naoyuki Takahashi (Institute for Oral Science, Matsumoto Dental University) for their advice on our work.

FUNDING

The study was supported by funding from JSPS KAKENHI Grant Number JP16H05144, the Nagano Society for the Promotion of Science, and a Scientific Research Special Grant from Matsumoto Dental University.

REFERENCES

- Afzal, M., Al-Hadidi, D., Menon, M., Pesek, J., and Dhimi, M. (2001). Ginger: an ethnomedical, chemical and pharmacological review. *Drug Metabol Drug Interact*, 18(3-4):159–190.
- Aimbire, F., Penna, S., Rodrigues, M., Rodrigues, K., Lopes-Martins, R., and Sertié, J. (2007). Effect of hydroalcoholic extract of *Zingiber officinalis* rhizomes on LPS-induced rat airway hyperreactivity and lung inflammation. *Prostaglandins Leukot Essent Fatty Acids*, 77(3-4):129–138.
- Ara, T., Fujinami, Y., Imamura, Y., and Wang, P. (2008a). Lipopolysaccharide-treated human gingival fibroblasts continuously produce PGE₂. *J Hard Tissue Biol*, 17(3):121–124.
- Ara, T., Fujinami, Y., Urano, H., Hirai, K., Hattori, T., and Miyazawa, H. (2012). Protein kinase A enhances lipopolysaccharide-induced IL-6, IL-8, and PGE₂ production by human gingival fibroblasts. *J Negat Results Biomed*, 11:10.

- Ara, T., Honjo, K., Fujinami, Y., Hattori, T., Imamura, Y., and Wang, P. (2010). Preventive effects of a kampo medicine, orento on inflammatory responses in lipopolysaccharide treated human gingival fibroblasts. *Biol Pharm Bull*, 33(4):611–616.
- Ara, T., Kurata, K., Hirai, K., Uchihashi, T., Uematsu, T., Imamura, Y., Furusawa, K., Kurihara, S., and Wang, P. (2009). Human gingival fibroblasts are critical in sustaining inflammation in periodontal disease. *J Periodontal Res*, 44(1):21–27.
- Ara, T., Maeda, Y., Fujinami, Y., Imamura, Y., Hattori, T., and Wang, P. (2008b). Preventive effects of a kampo medicine, shosaikoto, on inflammatory responses in LPS-treated human gingival fibroblasts. *Biol Pharm Bull*, 31(6):1141–1144.
- Ara, T. and Sogawa, N. (2016). Studies on shokyo, kanzo, and keihi in kakkonto medicine on prostaglandin E₂ production in lipopolysaccharide-treated human gingival fibroblasts. *Int Sch Res Notices*, 2016:9351787.
- Bartold, P. and Haynes, D. (1991). Interleukin-6 production by human gingival fibroblasts. *J Periodontal Res*, 26(4):339–345.
- Burke, J. and Dennis, E. (2009). phospholipase A₂ biochemistry. *Cardiovasc Drugs Ther*, 23(1):49–59.
- El-Abhar, H., Hammad, L., and Gawad, H. (2008). Modulating effect of ginger extract on rats with ulcerative colitis. *J Ethnopharmacol*, 118(3):367–372.
- Gijón, M., Spencer, D., Kaiser, A., and Leslie, C. (1999). Role of phosphorylation sites and the C2 domain in regulation of cytosolic phospholipase A₂. *J Cell Biol*, 145(6):1219–1232.
- Gupta, C., Katsumata, M., Goldman, A., Herold, R., and Piddington, R. (1984). Glucocorticoid-induced phospholipase A₂-inhibitory proteins mediate glucocorticoid teratogenicity *in vitro*. *Proc Natl Acad Sci U S A*, 81(4):1140–1143.
- Ha, S., Moon, E., Ju, M., Kim, D., Ryu, J., Oh, M., and Kim, S. (2012). 6-shogaol, a ginger product, modulates neuroinflammation: a new approach to neuroprotection. *Neuropharmacology*, 63(2):211–223.
- Kamemoto, A., Ara, T., Hattori, T., Fujinami, Y., Imamura, Y., and PL, W. (2009). Macrolide antibiotics like azithromycin increase lipopolysaccharide-induced IL-8 production by human gingival fibroblasts. *Eur J Med Res*, 14(7):309–314.
- Kim, S., Kundu, J., Shin, Y., Park, J., Cho, M., Kim, T., and Surh, Y. (2005). [6]-gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF- κ B in phorbol ester-stimulated mouse skin. *Oncogene*, 24(15):2558–2567.
- Kitamura, H., Urano, H., and Ara, T. (2014). Preventive effects of a kampo medicine, kakkonto, on inflammatory responses via the suppression of extracellular signal-regulated kinase phosphorylation in lipopolysaccharide-treated human gingival fibroblasts. *ISRN Pharmacol*, 2014:784019.
- Lakhan, S., Ford, C., and Tepper, D. (2015). *Zingiberaceae* extracts for pain: a systematic review and meta-analysis. *Nutr J*, 14:50.
- Lin, L., Wartmann, M., Lin, A., Knopf, J., Seth, A., and Davis, R. (1993). cPLA₂ is phosphorylated and activated by MAP kinase. *Cell*, 72(2):269–278.
- Nakazono, Y., Ara, T., Fujinami, Y., Hattori, T., and Wang, P. (2010). Preventive effects of a kampo medicine, hangeshashinto on inflammatory responses in lipopolysaccharide-treated human gingival fibroblasts. *J Hard Tissue Biol*, 19(1):43–50.
- Nievergelt, A., Marazzi, J., Schoop, R., Altmann, K., and Gertsch, J. (2011). Ginger phenylpropanoids inhibit IL-1 β and prostanoid secretion and disrupt arachidonate-phospholipid remodeling by targeting phospholipases A₂. *J Immunol*, 187(8):4140–4150.
- Noguchi, K. and Ishikawa, I. (2007). The roles of cyclooxygenase-2 and prostaglandin E₂ in periodontal disease. *Periodontol 2000*, 43:85–101.
- Pan, M., Hsieh, M., Hsu, P., Ho, S., Lai, C., Wu, H., Sang, S., and Ho, C. (2008). 6-shogaol suppressed lipopolysaccharide-induced up-expression of iNOS and COX-2 in murine macrophages. *Mol Nutr Food Res*, 52(12):1467–1477.
- Podlogar, J. and Verspohl, E. (2012). Antiinflammatory effects of ginger and some of its components in human bronchial epithelial (BEAS-2B) cells. *Phytother Res*, 26(3):333–336.
- Rhode, J., Fogoros, S., Zick, S., Wahl, H., Griffith, K., Huang, J., and Liu, J. (2007). Ginger inhibits cell growth and modulates angiogenic factors in ovarian cancer cells. *BMC Complement Altern Med*, 7:44.
- Saha, P., Katarkar, A., Das, B., Bhattacharyya, A., and Chaudhuri, K. (2016). 6-gingerol inhibits *Vibrio cholerae*-induced proinflammatory cytokines in intestinal epithelial cells via modulation of NF- κ B.

- Pharm Biol*, 54(9):1606–1615.
- Shim, S., Kim, S., Choi, D., Kwon, Y., and Kwon, J. (2011). Anti-inflammatory effects of [6]-shogaol: potential roles of HDAC inhibition and HSP70 induction. *Food Chem Toxicol*, 49(11):2734–2740.
- Sismey-Durrant, H. and Hopps, R. (1991). Effect of lipopolysaccharide from *Porphyromonas gingivalis* on prostaglandin E₂ and interleukin-1 β release from rat periosteal and human gingival fibroblasts *in vitro*. *Oral Microbiol Immunol*, 6(6):378–380.
- Tamura, M., Tokuda, M., Nagaoka, S., and Takada, H. (1992). Lipopolysaccharides of *Bacteroides intermedius* (*Prevotella intermedia*) and *Bacteroides* (*Porphyromonas*) *gingivalis* induce interleukin-8 gene expression in human gingival fibroblast cultures. *Infect Immun*, 60(1):4932–4937.
- Thomson, M., Al-Qattan, K., Al-Sawan, S., Alnaqeeb, M., Khan, I., and Ali, M. (2002). The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostaglandins Leukot Essent Fatty Acids*, 67(6):475–478.
- Wallner, B., Mattaliano, R., Hession, C., Cate, R., Tizard, R., Sinclair, L., Foeller, C., Chow, E., Browning, J., Ramachandran, K., and Pepinsky, R. (1986). Cloning and expression of human lipocortin, a phospholipase A₂ inhibitor with potential anti-inflammatory activity. *Nature*, 320(6057):77–81.
- Wilborn, J., Crofford, L., Burdick, M., Kunkel, S., Strieter, R., and Peters-Golden, M. (1995). Cultured lung fibroblasts isolated from patients with idiopathic pulmonary fibrosis have a diminished capacity to synthesize prostaglandin E₂ and to express cyclooxygenase-2. *J Clin Invest*, 95(4):1861–1868.

Table 1. The ingredients in the shinbuto formula

Japanese name	Latin name	Amount (g)	Amount (g/g of product)*
bukuryo	<i>Poria Sclerotium</i>	4.0	0.089
shakuyaku	<i>Paeoniae Radix</i>	3.0	0.067
sojutsu	<i>Atractylodis Lanceae Rhizoma</i>	3.0	0.067
shokyo	<i>Zingiberis Rhizoma</i>	1.5	0.033
bushi	<i>Processi Aconiti Radix</i>	0.5	0.011
total		12.0	0.267

*7.5 g of shinbuto product contains 2.0 g of a dried extract of the mixed crude drugs.

Table 2. The ingredients in the ninjinto formula

Japanese name	Latin name	Amount (g)	Amount (g/g of product)*
kankyo	<i>Zingiberis Processum Rhizoma</i>	3.0	0.083
kanzo	<i>Glycyrrhizae Radix</i>	3.0	0.083
sojutsu	<i>Atractylodis Lanceae Rhizoma</i>	3.0	0.083
ninjin	<i>Ginseng Radix</i>	3.0	0.083
total		12.0	0.333

*7.5 g of ninjinto product contains 2.5 g of a dried extract of the mixed crude drugs.

FIGURE LEGENDS

Figure 1

Effects of kampo medicines on cytotoxicity. HGFs were treated with combinations of LPS (0 or 10 ng/ml) and kampo medicine (0, 0.5, 1, 2, 5, or 10 mg/ml) for 24 h. Then, the numbers of viable cells were measured with WST-8. Open circles, treatment without LPS; closed circles, treatment with 10 ng/ml of LPS. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (without vs. with kampo medicine). P values were calculated by pairwise comparisons and corrected with the Holm method (10 null hypotheses).

Figure 2

Effects of kampo medicines on PGE₂, IL-6, and IL-8 production. HGFs were treated with combinations of LPS (0 or 10 ng/ml) and kampo medicine (0, 0.01, 0.1, or 1 mg/ml) for 24 h. Concentrations of PGE₂, IL-6, and IL-8 were measured by ELISA, adjusted by cell number, and expressed as per 10,000 cells (mean \pm S.D., $n = 3$). Open circles, treatment without LPS; closed circles, treatment with 10 ng/ml of LPS. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (without vs. with kampo medicine). # $P < 0.05$, ## $P < 0.01$, ### $P < 0.001$ (without LPS vs. with LPS). P values were calculated by pairwise comparisons and corrected with the Holm method (10 null hypotheses).

Figure 3

Effects of kampo medicines on COX activity. HGFs were treated with LPS (10 ng/ml) and kampo medicine (1 mg/ml) for 8 h, washed, and then treated with 10 μ M arachidonic acid for 30 min. Concentrations of PGE₂ were measured by ELISA, adjusted by cell number, and expressed as per 10,000 cells (mean \pm S.D., $n = 4$). P values by Dunnett's test are indicated.

Figure 4

Effects of kampo medicines on cPLA₂, annexin 1, and COX-2 expression. HGFs were treated with a combination of LPS (0 or 10 ng/ml) and kampo medicines (0 or 1 mg/ml) for 8 h, and protein levels were examined by Western blotting. The band densities were normalized against LPS alone and actin, and indicated below each band. shi, shinbuto; ni, ninjinto.

Figure 5

Effects of kampo medicines on LPS-induced ERK phosphorylation. HGFs were untreated (0 h), treated with LPS (10 ng/ml), or treated with both LPS and kampo medicines (1 mg/ml) for 0.5, 1, or 2 h. PMA was used as a positive control. Western blotting was performed using anti-phosphorylated ERK or anti-ERK antibodies. pERK, phosphorylated ERK. The upper band indicates ERK1 (p44 MAPK) and lower band ERK2 (p42 MAPK).

Figure 6

Effects of herbs on LPS-induced PGE₂, IL-6, and IL-8 production. (A) HGFs were treated with combinations of LPS (0 or 10 ng/ml) and each herb (100 μ g/ml) for 24 h. Concentrations of PGE₂, IL-6, and IL-8 were measured by ELISA, adjusted by cell number, and expressed as per 10,000 cells (mean \pm S.D., $n = 3$). (B) HGFs were treated with combinations of LPS (10 ng/ml) and herbs (0, 1, 10, or 100 μ g/ml) for 24 h. Concentrations of PGE₂ were measured by ELISA, adjusted by cell number, and expressed as per 10,000 cells (mean \pm S.D., $n = 3$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (LPS alone vs. LPS plus herb, Dunnett's test).

Figure 7

Effects of shokyo and kankyo on cPLA₂, annexin 1, and COX-2 expression. HGFs were treated with a combination of LPS (0 or 10 ng/ml) and herbs (1 mg/ml) for 8 h, and protein levels were examined by Western blotting. The band densities were normalized against LPS alone and actin, and indicated below each band. s, shokyo; k, kankyo.




Figure 1. Effects of kampo medicines on cytotoxicity




Figure 2. Effects of kampo medicines on PGE₂, IL-6, and IL-8 production




Figure 3. Effects of kampo medicines on COX activity




Figure 4. Effects of kampo medicines on cPLA₂, annexin 1, and COX-2 expression





Figure 5. Effects of kampo medicines on LPS-induced ERK phosphorylation






Figure 7. Effects of shokyo and kankyo on cPLA₂, annexin 1, and COX-2 expression