

A peer-reviewed version of this preprint was published in PeerJ on 11 December 2018.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.6094) (peerj.com/articles/6094), which is the preferred citable publication unless you specifically need to cite this preprint.

Montero-Pau J, Gómez A, Serra M. 2018. Founder effects drive the genetic structure of passively dispersed aquatic invertebrates. PeerJ 6:e6094 <https://doi.org/10.7717/peerj.6094>

Founder effects drive the genetic structure of passively dispersed aquatic invertebrates

Javier Montero-Pau^{1,2, 3,#}, Africa Gómez² and Manuel Serra¹

¹ *Institut Cavanilles de Biodiversitat i Biologia Evolutiva, Universitat de València, A.O.22085, 46071, Valencia, Spain*

² *Department of Biological Sciences, University of Hull, Hull, HU6 7RX, UK*

³ *Institute for the Conservation and Breeding of Agricultural Biodiversity (COMAV-UPV), Universitat Politècnica de València, Camino de Vera s/n, 46022, Valencia, Spain*

#Corresponding author:

Javier Montero-Pau, Institute for the Conservation and Breeding of Agricultural Biodiversity (COMAV-UPV), Universitat Politècnica de València, Camino de Vera s/n, 46022, Valencia (Spain)

Phone: (+34) 963878847

jmonteropau@upv.es

Running title (45 characters):

Founder effects and local adaptation

Abstract

Populations of passively dispersed organisms in continental aquatic habitats typically show high levels of neutral genetic differentiation despite their high dispersal capabilities. Several evolutionary factors, including founder events, local adaptation, and life cycle features such as high population growth rates and the presence of propagule banks, have been proposed to be responsible for this paradox. Here, we have modeled the colonization process in these organisms to assess the impact of migration rate, growth rate, population size, local adaptation and life-cycle features on their population genetic structure. Our simulation results show that the strongest effect on population structure is caused by persistent founder effects, resulting from the interaction of a few population founders, high population growth rates, large population sizes and the presence of diapausing egg banks. In contrast, the role of local adaptation, genetic hitchhiking and migration is limited to small populations in these organisms. Our results indicate that local adaptation could have different impact on genetic structure in different groups of zooplankters.

Keywords: migration, local adaptation, genetic differentiation, zooplankton, Monopolization Hypothesis, Rotifera, Cladocera.

40 Introduction

41 Successful dispersal and colonization are essential for the establishment and persistence of
42 species, and an understanding of these processes is crucial in the face of changing climate ,
43 habitat destruction, and biological invasions, which are rapidly affecting the abundance and
44 distribution patterns of many species (Parmesan & Yohe, 2003; Chen et al., 2011; Blackburn,
45 Lockwood & Cassey, 2015). The evolutionary outcome of dispersal and colonization results from
46 a potentially complex interplay of neutral and selective factors, including local adaptation,
47 founder effects or bottlenecks causing genetic drift during the first stages of colonization,
48 inbreeding depression, or high levels of gene flow that could erode local adaptation
49 (Lenormand, 2002; Kliber & Eckert, 2005; Rosenblum, Hickerson & Moritz, 2007; Keller &
50 Taylor, 2008; Verhoeven et al., 2011). In addition, life-cycle features and demographic
51 characteristics may act as modulators and lead to different evolutionary outcomes (Burton,
52 Phillips & Travis, 2010). For example, species with high population growth rates after a
53 bottleneck are more likely to maintain their genetic variability (“founder-flush” model) (Carson,
54 1968; Templeton, 2008), and populations of organisms with resistant life stages (e.g. diapausing
55 eggs) are more likely to be connected by migration even at long distances (Frisch, Green &
56 Figuerola, 2007). Predicting the outcome of these factors is a major question in evolutionary
57 and conservation ecology, and requires an understanding of the effect of each factor and their
58 interactions.

59 Populations of passively dispersing aquatic invertebrates (e.g., freshwater bryozoans,
 60 rotifers, cladocerans, copepods, anostracans, notostracans) and macrophytes inhabiting lentic
 61 habitats typically present a high level of neutral genetic differentiation (Freeland, Romualdi &
 62 Okamura, 2000; Zierold, Hanfling & Gómez, 2007; Mills, Lunt & Gómez, 2007; Muñoz et al.,
 63 2008; Makino & Tanabe, 2009; Xu et al., 2009; Escudero et al., 2010), despite their high
 64 dispersal capabilities through diapausing propagules (Allen, 2007; Frisch, Green & Figuerola,
 65 2007; Vanschoenwinkel et al., 2011). This has been termed the “migration-gene flow paradox”.
 66 Regardless of their taxonomic disparity, these organisms share biological features promoting a
 67 rapid monopolization of resources in the new environment: high population growth rates, large
 68 population sizes and the production of resistant stages in their life cycle. The latter can
 69 accumulate in sediments and form dormant propagule banks (Hairston, 1996; Brendonck & De
 70 Meester, 2003) and constitute the dispersal stage. As a result, once a habitat becomes available
 71 and is colonized, the population can grow very quickly creating a numerical advantage that
 72 dilutes the genetic impact of further immigrants (Waters, Fraser & Hewitt, 2013), resulting in a
 73 persistent founder effect (Boileau, Hebert & Schwartz, 1992). This explanation was expanded
 74 by (De Meester et al., 2002) into the so called “Monopolization Hypothesis” (MH hereafter) to
 75 include local adaptation as an important force contributing to reduce effective gene flow and
 76 therefore maintaining the genetic structure of passively dispersed aquatic organisms. The MH
 77 postulates that the migration-gene flow paradox could be explained by a combination of three
 78 factors: (1) persistent founder effects, (2) selection against immigrants due to local adaptation
 79 and (3) buildup of linkage disequilibrium between neutral markers and genes under selection.

Local adaptation is an important and rapid process in many zooplanktonic organisms (Cousyn et al., 2001; Decaestecker et al., 2007; Costanzo & Taylor, 2010; Declerck et al., 2015; Tarazona, García-Roger & Carmona, 2017). The impact of local adaptation on population genetic structure is diverse as it is dependent on the impact of other evolutionary forces, not only selection (Kawecki & Ebert, 2004). For instance, it can promote genetic differentiation, 'isolation-by-adaptation' sensu (Nosil, 2007) or reinforce the existing genetic differentiation by reducing effective gene flow (Orsini et al., 2013). Irrespective of local adaptation, populations recently founded by a small number of propagules can be highly inbred and show inbreeding depression (De Meester, 1993; Tortajada, Carmona & Serra, 2009). This could give migrants a fitness advantage and favor gene flow into the population (Ebert et al., 2002; Haag et al., 2006). The accumulation of large numbers of resistant stages (i.e. propagule banks) in sediments is also a characteristic of many aquatic species inhabiting temporary habitats. These banks have an important role in community-level ecological (Chesson, 1983; Cáceres, 1997; Montero-Pau & Serra, 2011) and evolutionary processes (Brendonck & De Meester, 2003). They increase the effective population size due to postponed reproduction in the bank, and thus reduce genetic drift (Kaj, Krone & Lascoux, 2001). However, this effect may be indirect, as gene flow is also postponed in the bank (Kaj, Krone & Lascoux, 2001; Berg, 2005).

The relative importance of and the interactions between the demographic features with neutral and selective processes acting during colonization has remained poorly understood. Therefore, an explicit analysis of the effects of local adaptation, persistent founder effects, and their interplay on the differentiation of populations of aquatic organisms is due, especially

during the first stages of colonization when populations are still small and, thus, more sensitive to stochastic effects.

Here, we have modeled the colonization process of zooplanktonic organisms to clarify how migration rate, growth rate, population size, local adaptation and the existence of a propagule bank shape the population genetic structure during the first stages of colonization. Our primary interest is to gain insights into the relative importance of (1) persistent founder effects, (2) selection against immigrants as a consequence of local adaptation, and (3) random associations between neutral genes and genes under selection (linkage disequilibrium).

Materials and Methods

We developed a genetic and demographic model to analyze the effects of population growth rate, population size, presence of a diapausing egg bank and local adaptation on the population genetic structure of aquatic organisms. We assumed a geographic scenario with two habitats, with local populations connected through reciprocal migration. These local populations are founded simultaneously after a single event of migration from a source population.

The model was based on the life cycle of rotifers and cladocerans (i.e. cyclical parthenogenesis), which are major taxonomic groups in the zooplankton. Cyclical parthenogenesis combines parthenogenesis with episodic sexual reproduction and typically consists of several asexual generations followed by a sexual generation, generally associated with habitat degradation. The sexual generation produces diapausing eggs that hatch into

asexual individuals once the habitat becomes suitable again. As not all eggs hatch from one growth period to the next, they may accumulate in the sediment and form extensive diapausing egg banks (Brendonck & De Meester, 2003).

The demographic submodel is outlined in Fig. 1. Briefly, it consists in six steps:

- step 1. Hatching of diapausing eggs (resident and immigrant)
- step 2. Asexual proliferation
- step 3. Sexual reproduction and production of diapausing eggs
- step 4. Diapausing eggs survival in the sediment
- step 5. Migration of diapausing eggs
- step 6. Back to step 1

Note that this demography implies two time scales: (1) a within-planktonic growth period (often within-year; index, t), and (2) an among-sexual generations scale (often among-years; index, y).

Migration, either from the source population or between habitats, is assumed to occur via diapausing eggs, which are passively transferred between habitats, and their hatching time is assumed to be the same as for locally produced diapausing eggs. As migration rates are low relative to the size of diapausing egg bank, emigration is assumed to have negligible effects on the source bank.

Genetic submodel

All individuals are considered to have n neutral loci and n loci under selection. All loci are biallelic and no mutation is assumed. Absence of linkage is assumed among neutral loci and

among loci under selection. Contrarily, the model accounts for physical linkage between each neutral locus and a corresponding selected locus. For this case, several recombination levels, including absence of linkage, were assumed. Loci under selection act additively on growth rate. Consequently, no dominance and no epistatic effects are assumed. Local adaptation requires a genotype-environment interaction on fitness. This is modeled through $\delta_{i,j,l}$, which is the effect on the intrinsic growth rate (see below) of allele i ($i: 1, 2$) at locus j ($j: 1, \dots, n$) in locality l ($l: 1, 2$). The assumptions are (1) $\delta_{1,j,1} = \delta_{2,j,2}$, and (2) $\delta_{i,j,l} = -\delta_{j \neq i,j,l}$; so, in the case of homozygotes for a given selected locus, they will experience an increase or decrease of their growth rate by $|2\delta|$ depending on the locality. Hence, the growth rate for each genotype g in each locality l ($r_{g,l}$) can be decomposed into r (basal growth rate) and θ (deviation of each genotype), so that

$$r_{g,l} = r + \theta_{g,l}$$

where g is the genotype, l is the locality, and $\theta_{g,l}$ is the summation of the fitness components (δ) in locality l of the alleles carried by a genotype g in the n loci under selection. Thus, in any given locality, the growth rate during the asexual reproduction will vary between the limits $r \pm 2n\delta$.

Sexual reproduction is assumed to be panmictic and, for simplicity, is considered to be synchronic and at the end of the growing season ($t = \tau$). As linkage disequilibrium can occur due to selection and genetic drift, gametic frequencies are computed. Gametes are then drawn to produce the diapausing eggs.

Genetic distance between populations was estimated based on neutral loci as

$$F_{ST} = \frac{\bar{H}_T - \bar{H}_S}{\bar{H}_T}$$

where $\bar{H}_T$ is the average expected heterozygosity for the two populations considered as a single one for the neutral loci, and $\bar{H}_S$ the average of the mean expected heterozygosity within each populations for the neutral loci (Hedrick, 2011). Allelic frequencies for each locus were computed using the total number of alleles. Similarly, a genetic distance for loci under selection (F_{STQ}) was computed (Le Corre & Kremer, 2012). F_{ST} and F_{STQ} values were obtained just after hatching of diapausing eggs.

Population growth

The asexual phase spans from time $t = 0$ to τ , which is the moment when sexual reproduction takes place. During the asexual phase the population grows deterministically according to a logistic growth model:

$$\frac{dN_{i,g}}{dt} = N_{i,g} r_{i,g} \left[1 - \frac{\sum_g N_{i,g}}{K} \right],$$

where $N_{i,g}$ is the density of the genotype g in the locality i , $r_{i,g}$ is its intrinsic population growth rate during the asexual phase, and K the carrying capacity. Note that K is genotype-independent. At the onset of each asexual growth season ($t = 0$), $N_{i,g}$ is the sum of the hatched diapausing eggs, a fraction of them having been locally produced $H_{i,g}$, and the rest being immigrants $M_{i,g}$.

At $t = \tau$ of the sexual generation y , the number of diapausing eggs produced $P_{i,g}(y)$ is stochastically computed from $N_{i,g}(\tau, y)$ assuming a sexual proportion m (fraction of the females

that becomes sexual), a sex ratio sr and an effective fecundity e (number of diapausing eggs produced per sexual female).

Mortality of diapausing eggs in the sediment with egg bank was assumed to be age-independent (annual survival rate γ). Empirical information supporting this assumption for field populations is not available. However, our model can account for fast senescence when it assumes the absence of egg bank. When a new planktonic growing season starts ($t = 0$) a fraction λ of the diapausing eggs in the sediment hatches.

Source population and local population founding

The two populations are founded at time $y = t = 0$ by F diapausing eggs randomly drawn from a single source population. The source population is assumed to be in Hardy-Weinberg equilibrium and of infinite size, so that extraction of migrants does not change genotype frequencies. All loci are considered neutral in the source population, so no preadaptation to any of the populations exists.

Model implementation

The impact of carrying capacity (K), growth rate (r), migration (M), selection pressure (δ) and recombination rate on F_{ST} were analyzed by exploring a range of realistic values for zooplanktonic organisms. K was varied from $2 \cdot 10^2$ to $2 \cdot 10^7$ individuals, which is equivalent to densities from 0.001 to 100 individuals/L in a small pond of 200 m² and 1 m depth, in good agreement with reported average densities of cladocerans and rotifers (Carmona, Gómez & Serra, 1995; Ortells, Gómez & Serra, 2003; Tavernini, 2008). r was explored from 0.05 to 1 day⁻¹.

Cladocerans show maximum r of 0.2-0.6 days⁻¹ and rotifers 0.2-1.5 days⁻¹ (Allan, 1976). The number of population founders (F) was set to 1 diapausing egg across most simulations. That is, foundation is considered a rare event. Note that as the model assumes cyclical parthenogenesis, a single diapausing egg is enough for population foundations. The effect of numbers of founders (F) was also explored (1, 2, 5, 50 diapausing eggs). Other parameter values used in the simulations are shown in Table 1.

Simulations considered two scenarios regarding diapausing egg banks: (1) an annual, age-independent, diapausing egg survival rate on the sediment ($\gamma = 0.763$) (i.e., existence of a diapausing egg bank); and (2) $\gamma = 0.763$ for eggs of age = 1 and a $\gamma = 0$ for older eggs (i.e., absence of diapausing egg bank). Parameters for the diapausing egg bank (γ and λ , the annual hatching rate) were estimated from rotifer diapausing egg banks (García-Roger, Carmona & Serra, 2006c) by adjusting them to the model described by García-Roger *et al.* (García-Roger, Carmona & Serra, 2006a).

The simulation model was implemented in C++ and based on Monte-Carlo procedures (code available at https://github.com/monpau/founder_effects). The Mersenne twister algorithm (Matsumoto & Nishimura, 1998) was used as random number generator. The logistic model was iterated numerically. 50 replicates for each parameter combination (but 100 for values of δ and recombination rate) were performed. For each replicate, a source population was randomly created by drawing from a uniform distribution the allelic frequencies of the n and s loci. After foundation of the two populations, 1000 sexual generations (4000 generations for some scenarios) were simulated.

Sampling effects were taken into account for hatching and survival of diapausing eggs if the total number of eggs in the population was lower than 1000. Selection of immigrants and gametes for mating were performed randomly regardless of the number of eggs/individuals involved.

Paralelling the procedure in an empirical study, an statistical assesment was performed Differences between F_{ST} 's values under a neutral scenario and scenarios with selective pressure and different recombination rates were analysed with an ANOVA and *a priori* contrasts. Correlations between F_{ST} and F_{STQ} at different combinations of population size, recombination rates and selective pressure were also tested using Kendall's Tau and Sperman's Rho. All statistical analyses were performed using SPSS v. 17 (SPSS Inc., Chicago, USA)

Results

The population dynamics of a newly founded population, using the parameters shown in Table 1, with $\tau = 60$ days and $r = 0.3 \text{ days}^{-1}$ – which are realistic values for both the length of the growth season (Tavernini, 2008) and the intrinsic growth rate of many aquatic invertebrates (Allan, 1976) – show that carrying capacity (K) is reached in less than two sexual generations, even in the case of the highest K (i.e. $K = 2 \cdot 10^7$ individuals). Thus, K is a good proxy of population size and we will use both terms interchangeably hereafter.

Effect of migration

The effect of the number of immigrants on genetic differentiation of neutral loci strongly depends on K (i.e. population size; Fig. 2). In both the small and the large populations, F_{ST}

decreases with increasing migration rates, as expected under a neutral scenario (Wright, 1931). For the lowest carrying capacity tested ($K=2 \cdot 10^2$ individuals; Fig. 2.A), F_{ST} decreased rapidly down to very low levels with increasing migration. By contrast, for the highest K tested ($K = 2 \cdot 10^7$ individuals; Fig. 2.B), F_{ST} was rather insensitive to the effect of migration, and populations remained highly differentiated ($F_{ST} > 0.2$) even at high levels of migration. The number of migrants needed to cause a noticeable decrease of genetic differentiation on neutral loci is in the order of 100 and 1000 individuals/sexual generation for the situation without and with diapausing eggs respectively.

Effect of population size

Carrying capacity (i.e. population size) had strong effects on F_{ST} (Fig. 3; 1000 sexual generations). In small populations (i.e. low K) populations did not differ genetically, while in large populations, F_{ST} remained as high as the values observed just after population foundation. This pattern suggests importance of migration and persistent founder effects respectively. At intermediate values of K , genetic differentiation peaked, probably due to higher drift effects. In other words, the highest F_{ST} values are found at intermediate population sizes. The $F_{ST}-K$ pattern is qualitatively similar with and without diapausing egg bank, but in absence of a egg bank a lower maximum F_{ST} at a higher K was found.

These results are robust to changes in the maximum number of sexual generations explored (results for maximum $y = 100, 500, 2000$ and 4000 generations, data not shown). However, at 100 and to a lesser extent 500 sexual generations, the peak of F_{ST} at intermediate population sizes was less pronounced than at later sexual generations. The long-term (from 1st

to the 4000th sexual generation) time course is further explored in Figure 4. In the absence of a diapausing egg bank (Fig. 4 A), F_{ST} decreases with time at low population size, and this situation is reversed when K increases, to finally become virtually constant (i.e., determined by the initial condition) at the largest population size explored ($K = 2 \cdot 10^7$). A qualitatively similar pattern is found when a diapausing egg bank is present (Fig. 4 B), although the shift to an increasing F_{ST} time course, and also to F_{ST} constancy, occurs at lower population sizes. Note that the small negative change found at $K = 2 \cdot 10^2$ (regardless whether a bank is assumed) is associated to the very low initial F_{ST} values (Fig. 4 C, D). Also note that F_{ST} values are calculated after hatching of residents and immigrants; for instance, at $y = 1$, F_{ST} value is not the value after foundation but after migration.

In summary, population size and presence or absence of a diapausing egg bank are key to predict the main force shaping the genetic structure. Decreasing F_{ST} indicates that migration is the dominant factor, while increasing values show that drift becomes dominant. The finding of stationary F_{ST} from the first generations after foundation indicates the importance of persistent founder effects on the shaping of the genetic structure of populations.

Population growth rate interacts with population size in determining the level of genetic differentiation (Fig. 5). Low growth rates result in low genetic differentiation, regardless of population size, indicating a high impact of migration. However, for population growth rates above 0.1 d^{-1} , which are common for zooplanktonic organisms, genetic differentiation becomes sensitive to variations in population size.

Effects of the number of founders

Increasing the number of population founders F results in a dramatic decrease of F_{ST} values just after foundation (Fig. 6); for instance, if compared to $F = 1$, F_{ST} is reduced by half for $F = 2$, and approaches 0 for $F = 50$. After 4000 sexual generations, the level of population differentiation still shows a negative relationship with the number of founders. Given this strong effect, we explored in further simulations how F affects the relationships between population differentiation and other factors. Our results suggest that the patterns outlined above are qualitatively maintained for $F > 1$ (data not shown).

Effect of local adaptation

Above, a selectively neutral scenario was assumed. The effect of local adaptation was explored at two levels of K ($2 \cdot 10^4$ and $2 \cdot 10^7$ individuals), which are realistic values for cladocerans and rotifers respectively. Two different selection scenarios ($\delta = 10^{-4} \text{ days}^{-1}$, light selection, and $10^{-2} \text{ days}^{-1}$, intense selection) in the presence/absence of diapausing egg bank, and six recombination rates – from complete linkage to unlinked genes – were tested (Fig. 7 summarizes the results for the scenario with diapausing egg bank; see Fig. S1, for the equivalent scenario without diapausing egg bank).

With intense selection, F_{STQ} reaches almost maximum values – i.e., populations are almost fixed for the locally adapted alleles – regardless of K (Fig. 7). In the case of low K , all F_{ST} values are statistically different from those obtained without selection (p-values < 0.05 except at 0.5 recombination rate; p-value = 0.057). However, F_{ST} values are similar irrespective of the recombination rate. In contrast, for high K , only those values of F_{ST} with complete linkage

(recombination rate = 0) are statistically different of those found without selection. This indicates that genetic hitchhiking in large populations acts only on neutral loci tightly linked to those under selection. Otherwise, linkage to the genes under selection does not reduce the persistence of founder effects.

With light selection, F_{STQ} indicates the expected result that local adaptation becomes less important than with intense selection. In large populations ($K = 2 \cdot 10^7$), F_{ST} values do not statistically differ from the neutral scenario, showing the higher importance of founder effects over local adaptation when selection is light. Moreover, F_{STQ} values also appear to be affected by persistent founder effects. In contrast to the situation with intense selection, genetic linkage does not alter differentiation at neutral loci. However, in small populations ($K = 2 \cdot 10^4$), local adaptation does play a role. Mean F_{ST} values statistically differ from the neutral scenario at all recombination rates (from 0.0 to 0.5), and the variance of the distribution of F_{ST} values is decreased (see Fig. 3 for comparison). Note that drift is the dominant factor in relatively small ($K = 2 \cdot 10^4$) populations with diapausing egg bank.

Effects of diapausing egg banks

In the presence of intense selection the effects of diapausing egg bank (see Supplementary Fig. S1) were minimal. In light selection conditions: (1) at high population density ($K = 2 \cdot 10^7$) genes under selection are less affected by persistent founder effects than when no bank is present (Fig. 7), and populations show a trend to be locally adapted; (2) at low population density ($K = 2 \cdot 10^4$), F_{ST} values at recombination rates 0.0 and 0.1 are statistically different from the neutral scenario – unlike at higher recombination rates –, which indicates that genetic hitchhiking could

be of some importance; (3) at $K = 2 \cdot 10^4$ F_{ST} and F_{STQ} had higher variance at all recombination rates than in the scenario with no diapausing egg bank (Fig. 7).

In the absence of a diapausing egg bank, populations reach maximum F_{STQ} values in about 40-50 sexual generations regardless of population size (data not shown). However, when a diapausing egg bank exists, advantageous alleles need a longer time to reach fixation (about 150 sexual generations for $K = 2 \cdot 10^4$, and about 300 generations for $K = 2 \cdot 10^7$).

We computed F_{STQ} vs. F_{ST} correlations within each tested parameter combination. Significant correlations were found only in the case of the low K ($2 \cdot 10^4$) without diapausing egg bank. Correlation coefficient is always positive, and the ranges are: Kendall's tau = 0.66-0.53 and Spearman's rho = 0.73-0.56 for intense selection; Kendall's tau = 0.68-0.32 and Spearman's rho = 0.80-0.38 for light selection.

Discussion

The understanding of the evolutionary factors responsible for the strong population structure of passively dispersed aquatic organisms in the face of potentially high gene flow has attracted considerable attention in the last decade (De Meester et al., 2002; Mills, Lunt & Gómez, 2007; Campillo et al., 2009). We have presented a specific model and, by simulation, explored the effects of genetic drift associated to population founding (founder effects), gene flow via migration and local adaptation on genetic differentiation. Our results show that the strongest effect was caused by persistent founder effects, resulting largely from the distinctive life history traits of these organisms: few population founders, high rates of population growth, large

population sizes and the presence of diapausing egg banks. These results are in agreement with those of Boileau *et al.* (Boileau, Hebert & Schwartz, 1992), who proposed that persistent founder effects are an important force shaping the genetic structure of passively dispersed aquatic organisms. Our findings show that this conclusion holds in an scenario with selection and genetic linkage, not explored by these authors. The most remarkable and novel result of our simulations is that the role of local adaptation and genetic hitchhiking on shaping genetic structure of these organisms is not significant in large populations, although it plays a significant role in small populations. This result is in agreement with recent genomic linkage data in the rotifer *B. plicatilis* indicating very low levels of genomic linkage, meaning that the opportunities for hitchhiking are limited and selection can freely act in very concrete points of the genome without affecting nearby loci (Franch Gras, 2017)

In agreement with (Boileau, Hebert & Schwartz, 1992), migration has a very limited effect on the population structure of passively dispersed aquatic organisms. For instance, a migration rate of 1000 individuals per sexual generation is needed to cause a noticeable effect on F_{ST} in a large population. Although direct estimates of the number of dispersing stages are unavailable, this extremely large value appears unlikely to occur between non-connected ponds (Cáceres & Soluk, 2002; Allen, 2007; Frisch, Green & Figuerola, 2007), and is inconsistent with estimates of the number of founders in natural populations, which are expected to be correlated with regular immigration rates (Louette *et al.*, 2007; Badosa *et al.*, 2017). However, for small

population sizes our model recovers the expected pattern for the combined effect of migration and drift under neutral genetic differentiation.

Among the factors studied in our model, population size has been shown to be largely responsible for establishing the levels of genetic differentiation observed in natural populations of aquatic organisms. In addition, this effect is strongly reinforced when a diapausing egg bank is established. Although egg banks could increase gene flow (Kaj, Krone & Lascoux, 2001; Berg, 2005), they act mainly buffering the effects of migration and reducing genetic drift, which favors the establishment of persistent founder effects. In our model, we assumed a parameter range in agreement with values reported for many aquatic organisms. Nevertheless, due to computational limitations the maximum values used for population sizes and egg bank densities had to be limited, and could underestimate those attained in many natural populations. Some estimated population sizes and diapausing egg bank densities in rotifers are one or two orders of magnitude higher than the maximum values considered here (Carmona, Gómez & Serra, 1995; Ortells, Gómez & Serra, 2003). Diapausing egg bank densities for zooplanktonic organisms are in the order of 10^3 - 10^7 eggs/m² (Hairston, 1996), although densities in the sediment layers that could provide recruits are uncertain. However, modeling larger population sizes is unlikely to change our results qualitatively; if anything, they would increase the relative impact of persistent founder effects.

Local adaptation seems to be common and has been well documented in cladocerans (Cousyn et al., 2001; De Meester et al., 2002; Decaestecker et al., 2007) and the generalist rotifer *Brachionus plicatilis* (Campillo et al., 2010; Franch-Gras et al., 2017). However, the effect

of local adaptation on genetic structure does not seem to be general, as it is weakened by neutral and demographic factors. Regarding differentiation in genetic markres, a limited role for local adaptation in continental aquatic invertebrates has been suggested (Campillo et al., 2009; Allen, Thum & Cáceres, 2010). Our results indicate that although local adaptation does occur, its effects are only apparent on population structure when population sizes and diapausing egg banks are relatively small. Given that rotifers tend to have larger population sizes than cladocerans, this would mean that the effects of local adaptation on population structure could differ between these organisms. According to our results, genetic hitchhiking appears to be of limited importance in shaping neutral genetic differentiation. We have only detected signs of its effect (1) at completely linked genes with high population size and intense selection, and (2) at intermediate population size without egg bank and light selection. The lack of observed impact does not mean that genetic hitchhiking has no importance, but that other processes are dominating the outcome. We must stress that our main question is not whether local adaptation occurs or not, but if this adaptation affects genetic differentiation in neutral markers *sensu* Nosil (Nosil, 2007). Admitadly, due to computational limitations our model simplifies the selective scenarios acting on continental aquatic invertebrates. As selection in natural populations of aquatic invertebrates is likely to be multifactorial, and fluctuating, this scenarios should be further explored.

Genetic analyses in recently established populations indicate that the number of founders is small (Haag et al., 2005; Louette et al., 2007; Ortells, Olmo & Armengol, 2011; Badosa et al., 2017), and accordingly a single founder was assumed in most simulations. By assuming a single

founder, the studied scenario corresponds, for instance, to a situation where a new region consisting of several lakes is open to colonization – e.g. after glaciation –, with few founders of any single lake, but with varying migration rates among lakes. When we relaxed the assumption of a single founder, the only remarkable observed effect was a reduction on the final value of F_{ST} .

Globally, our results show that persistent founder effects, genetic drift and local adaptation all drive population genetic structure in these organisms, but population size and the egg bank have a strong control on the dominance of each of these factors. In turn, these demographic variables can be linked to ecological features. If so, a habitat classification linking ecological factors, demographic features, and mechanisms acting on genetic structure could be possible. Therefore, in populations inhabiting permanent ponds and lakes where a low investment in diapause is generally found – as reflected in small diapausing egg banks in comparison to temporary or ephemeral ponds (Hebert, 1974a,b; García-Roger, Carmona & Serra, 2006c; Campillo et al., 2010; Montero-Pau, Serra & Gómez, 2016) –, an increased effect of local adaptation and genetic drift is expected. In contrast, in environmental conditions limiting population sizes, such as small rock pools or nutrient-poor lakes, migration can attain higher importance. If despite this, high genetic differentiation is detected, an effect of selective forces can be hypothesized. For instance, genetic hitchhiking has been suggested for a *Daphnia* metapopulation inhabiting temporal rock pools (Haag et al., 2006). Besides ecological features, our results suggest that differences can be expected between taxa differing in body size and so in their typical population sizes, and therefore differences between the smaller rotifers and the

larger cladocerans are expected. As far as our results identify a restricted number of factors driving the genetic structure, they provide insights beyond the life cycle assumed (i.e., cyclical parthenogenesis), and could be extended to organisms with similar demographic features (i.e. high growth rates, high population densities or presence of seed or egg banks). For example, populations of sexual species with high growth rates (i.e. *r* strategists) like crustaceans such *Artemia* or copepods, which produce egg banks, are also likely to benefit from a numerical inertia that will reduce the impact of migration on the genetic structure of their populations (Boileau, Hebert & Schwartz, 1992).

As we have shown, the rapid growth rate of colonists acts as a barrier against new migrants, and this is reinforced by the formation of diapausing stage banks and, in some cases, by local adaptation. This process leads to a persistent founder effect, and consequently, to a deviation from the migration-drift equilibrium. This has repercussions when interpreting phylogeographic signals (Gómez, Carvalho & Lunt, 2000; Waters, 2011). For instance patterns of ‘isolation-by-distance’ found in several aquatic organisms, regardless of their reproductive mode, have been suggested to be due to a process of sequential colonizations (Gouws & Stewart, 2007; Gómez et al., 2007; Mills, Lunt & Gómez, 2007; Muñoz et al., 2008). Our results are consistent with these proposals and suggest that caution should be applied when inferring a migration-drift mechanism of ‘isolation by distance’ from such patterns (i.e., correlation between genetic and geographical distances). Also, the establishment of persistent founder effects and competitive exclusion of closely related species can explain the phylogenetic

overdispersion in communities, given a phylogenetic limiting similarity between species (Violle et al., 2011).

During the time window from the arrival of first colonizers to the establishment of the founder effects, the genetic structure of the population is still sensitive to migration or drift. Our results point out that this period is short, as a result of the high population growth rates of most aquatic organisms. Nevertheless, we found that with relatively low population growth rates, the numerical advantage is delayed and genetic differentiation is relatively low. Inbreeding depression is expected to be larger in small populations (Lohr & Haag, 2015), and it could act favoring gene flow (Haag et al., 2002; Tortajada, Carmona & Serra, 2009, 2010). Although this factor has not been explicitly modeled here, it will act in a similar way of reducing the growth rate, which will favor effective gene flow. However, severe inbreeding could also reduce the effective population size, and increase genetic drift, which will increase genetic differentiation. A more detailed exploration of this scenario will be needed and it will depend on the relative magnitude of the purging and migration.

Other factors not implemented in our model, but likely to occur in the wild, could also counteract the high genetic differentiation. For example, processes able to reduce population size during asexual growth phase (e.g., perturbations or environmental fluctuations) could increase the impact of gene flow. In addition, it will be of interest to test the strength of persistent founder effects buffering migrants with a higher fitness than locally adapted residents. These factors – inbreeding depression, environmental fluctuations, and preadapted immigrants – were not invoked in the initial formulation of the Monopolization Hypothesis and

should be investigated in future analyses. An additional prospective is to include the effect of metapopulation structure. Walser and Haage (2012) have shown that population turnover, which is expected to have high rate in small populations, could also explain the high genetic population differentiation.

Concluding remarks

Molecular screening of natural population has uncovered an unexpectedly high genetic diversity in taxa with high dispersal potential. These findings challenged classical views of the evolutionary processes in small multicellular organisms, and when focused on aquatic invertebrates, brought to postulate a combination of processes as causal factors for that genetic differentiation, the Monopolization Hypothesis (De Meester et al., 2002). Our analysis shows that a quantitative elaboration of this multifactorial hypothesis is able to dissect the relative weights of the different factors, and their interactions. Specifically, we found that founder effects drive the genetic structure of passively dispersed aquatic organisms. We conclude that although selective factors and migration have a role in explaining genetic structure of continental aquatic invertebrates, demographic processes are dominant. By studying which factors are important in what circumstances, our analysis can help understanding relevant differences among the genetic structure of different species.

Acknowledgements

We thank Guillermo García Franco and José Gargallo Tuzón for their invaluable help with some parts of the code and programming support. We also would like to thank Luc De Meester, Raquel Ortells and M^a José Carmona for helpful comments on previous versions of this manuscript. This work was funded by a grant from the Spanish Ministerio de Ciencia e Innovación (CGL2009-07364) to MS. AG was supported by a National Environment Research Council (NERC) Advanced Fellowship (NE/B501298/1) and JM-P by a fellowship by the Spanish Ministerio de Ciencia y Tecnología (BES2004-5248).

References

- Allan JD. 1976. Life History Patterns in Zooplankton. *The American naturalist* 110:165–180.
- Allen MR. 2007. Measuring and modeling dispersal of adult zooplankton. *Oecologia* 153:135–143.
- Allen MR., Thum RA., Cáceres CE. 2010. Does local adaptation to resources explain genetic differentiation among *Daphnia* populations? *Molecular ecology* 19:3076–3087.
- Alver MO., Hagiwara A. 2007. An individual-based population model for the prediction of rotifer population dynamics and resting egg production. *Hydrobiologia* 593:19–26.
- Aparici E., Carmona MJ., Serra M. 1998. Sex Allocation in Haplodiploid Cyclical Parthenogens with Density-Dependent Proportion of Males. *The American naturalist* 152:652–657.
- Badosa A., Frisch D., Green AJ., Rico C., Gómez A. 2017. Isolation mediates persistent founder

- 509 effects on zooplankton colonisation in new temporary ponds. *Scientific reports* 7:43983.
- 510 Berg LM. 2005. Fluctuating Selection, Egg Banks and Population Genetic Structure in Cyclically
- 511 Parthenogenetic Species. *Hydrobiologia* 549:287–295.
- 512 Blackburn TM., Lockwood JL., Cassey P. 2015. The influence of numbers on invasion success.
- 513 *Molecular ecology* 24:1942–1953.
- 514 Boileau MG., Hebert PDN., Schwartz SS. 1992. Non-equilibrium gene frequency divergence:
- 515 persistent founder effects in natural populations. *Journal of evolutionary biology* 5:25–39.
- 516 Brendonck L., De Meester L. 2003. Egg banks in freshwater zooplankton: evolutionary and
- 517 ecological archives in the sediment. *Hydrobiologia* 491:65–84.
- 518 Burton OJ., Phillips BL., Travis MJ. 2010. Trade-offs and the evolution of life-histories during
- 519 range expansion. *Ecology letters* 13:1210–1220.
- 520 Cáceres CE. 1997. Temporal variation, dormancy, and coexistence: a field test of the storage
- 521 effect. *Proceedings of the National Academy of Sciences of the United States of America*
- 522 94:9171–9175.
- 523 Cáceres CE., Soluk DA. 2002. Blowing in the wind: a field test of overland dispersal and
- 524 colonization by aquatic invertebrates. *Oecologia* 131:402–408.
- 525 Campillo S., García-Roger EM., Carmona MJ., Gómez A., Serra M. 2009. Selection on life-history
- 526 traits and genetic population divergence in rotifers. *Journal of evolutionary biology*
- 527 22:2542–2553.
- 528 Campillo S., García-Roger EM., Carmona MJ., Serra M. 2010. Local adaptation in rotifer
- 529 populations. *Evolutionary ecology* 25:933–947.

- 530 Carmona MJ., Gómez A., Serra M. 1995. Mictic patterns of the rotifer *Brachionus plicatilis*
531 Müller in small ponds. In: Ejsmont-Karabin J, Pontin RM eds. *Rotifera VII*. Developments in
532 Hydrobiology. Springer Netherlands, 365–371.
- 533 Carson HL. 1968. The population flush and its genetic consequences. *Population biology and*
534 *evolution*:123–137.
- 535 Chen I-C., Hill JK., Ohlemüller R., Roy DB., Thomas CD. 2011. Rapid range shifts of species
536 associated with high levels of climate warming. *Science* 333:1024–1026.
- 537 Chesson PL. 1983. Coexistence of Competitors in a Stochastic Environment: The Storage Effect.
538 In: Freedman HI, Strobeck C eds. *Population Biology*. Lecture Notes in Biomathematics.
539 Springer Berlin Heidelberg, 188–198.
- 540 Costanzo KS., Taylor DJ. 2010. Rapid ecological isolation and intermediate genetic divergence in
541 lacustrine cyclic parthenogens. *BMC evolutionary biology* 10:166.
- 542 Cousyn C., De Meester L., Colbourne JK., Brendonck L., Verschuren D., Volckaert F. 2001. Rapid,
543 local adaptation of zooplankton behavior to changes in predation pressure in the absence
544 of neutral genetic changes. *Proceedings of the National Academy of Sciences of the United*
545 *States of America* 98:6256–6260.
- 546 Decaestecker E., Gaba S., Raeymaekers JAM., Stoks R., Van Kerckhoven L., Ebert D., De Meester
547 L. 2007. Host-parasite “Red Queen” dynamics archived in pond sediment. *Nature* 450:870–
548 873.
- 549 Declerck SAJ., Malo AR., Diehl S., Waasdorp D., Lemmen KD., Proios K., Papakostas S. 2015.
550 Rapid adaptation of herbivore consumers to nutrient limitation: eco-evolutionary

- 551 feedbacks to population demography and resource control. *Ecology letters* 18:553–562.
- 552 De Meester L. 1993. Inbreeding and outbreeding depression in *Daphnia*. *Oecologia* 96:80–84.
- 553 De Meester L., Gómez-Africa,., Okamura B., Schwenk K. 2002. The Monopolization Hypothesis
- 554 and the dispersal–gene flow paradox in aquatic organisms. *Acta Oecologica* 23:121–135.
- 555 Ebert D., Haag C., Kirkpatrick M., Riek M., Hottinger JW., Pajunen VI. 2002. A selective
- 556 advantage to immigrant genes in a *Daphnia* metapopulation. *Science* 295:485–488.
- 557 Escudero M., Vargas P., Arens P., Ouborg NJ., Luceño M. 2010. The east-west-north colonization
- 558 history of the Mediterranean and Europe by the coastal plant *Carex extensa* (Cyperaceae).
- 559 *Molecular ecology* 19:352–370.
- 560 Franch Gras L. 2017. Rotifer adaptation to environmental unpredictability.
- 561 Franch-Gras L., García-Roger EM., Serra M., José Carmona M. 2017. Adaptation in response to
- 562 environmental unpredictability. *Proceedings. Biological sciences / The Royal Society* 284.
- 563 DOI: 10.1098/rspb.2017.0427.
- 564 Freeland JR., Romualdi C., Okamura B. 2000. Gene flow and genetic diversity: a comparison of
- 565 freshwater bryozoan populations in Europe and North America. *Heredity* 85:498–508.
- 566 Frisch D., Green AJ., Figuerola J. 2007. High dispersal capacity of a broad spectrum of aquatic
- 567 invertebrates via waterbirds. *Aquatic sciences* 69:568–574.
- 568 García-Roger EM., Carmona MJ., Serra M. 2006a. A simple model relating habitat features to a
- 569 diapause egg bank. *Limnology and oceanography* 51:1542–1547.
- 570 García-Roger EM., Carmona MJ., Serra M. 2006b. Hatching and viability of rotifer diapausing
- 571 eggs collected from pond sediments. *Freshwater biology* 51:1351–1358.

- 572 García-Roger EM., Carmona MJ., Serra M. 2006c. Patterns in rotifer diapausing egg banks:
573 Density and viability. *Journal of experimental marine biology and ecology* 336:198–210.
- 574 Gómez A., Carvalho GR., Lunt DH. 2000. Phylogeography and regional endemism of a passively
575 dispersing zooplankter: mitochondrial DNA variation in rotifer resting egg banks.
576 *Proceedings of the Royal Society of London B: Biological Sciences* 267:2189–2197.
- 577 Gómez A., Montero-Pau J., Lunt DH., Serra M., Campillo S. 2007. Persistent genetic signatures
578 of colonization in *Brachionus manjavacas* rotifers in the Iberian Peninsula. *Molecular*
579 *ecology* 16:3228–3240.
- 580 Gouws G., Stewart BA. 2007. From genetic structure to wetland conservation: a freshwater
581 isopod *Paramphisopus palustris* (Phreatoicidea: Amphisopidae) from the Swan Coastal
582 Plain, Western Australia. *Hydrobiologia* 589:249–263.
- 583 Haag CR., Hottinger JW., Riek M., Ebert D. 2002. Strong inbreeding depression in a *Daphnia*
584 metapopulation. *Evolution; international journal of organic evolution* 56:518–526.
- 585 Haag CR., Riek M., Hottinger JW., Pajunen VI., Ebert D. 2005. Genetic diversity and genetic
586 differentiation in *Daphnia* metapopulations with subpopulations of known age. *Genetics*
587 170:1809–1820.
- 588 Haag CR., Riek M., Hottinger JW., Pajunen VI., Ebert D. 2006. Founder events as determinants of
589 within-island and among-island genetic structure of *Daphnia* metapopulations. *Heredity*
590 96:150–158.
- 591 Hairston NG. 1996. Zooplankton egg banks as biotic reservoirs in changing environments.
592 *Limnology and oceanography* 41:1087–1092.

- 593 Hebert PD. 1974a. Enzyme variability in natural populations of *Daphnia magna*. 3. Genotypic
594 frequencies in intermittent populations. *Genetics* 77:335–341.
- 595 Hebert PD. 1974b. Enzyme variability in natural populations of *Daphnia magna*. II. Genotypic
596 frequencies in permanent populations. *Genetics* 77:323–334.
- 597 Hedrick PW. 2011. *Genetics of Populations*. Jones & Bartlett Learning.
- 598 Kaj I., Krone SM., Lascoux M. 2001. Coalescent theory for seed bank models. *Journal of applied*
599 *probability* 38:285–300.
- 600 Kawecki TJ., Ebert D. 2004. Conceptual issues in local adaptation. *Ecology letters* 7:1225–1241.
- 601 Keller SR., Taylor DR. 2008. History, chance and adaptation during biological invasion:
602 separating stochastic phenotypic evolution from response to selection. *Ecology letters*
603 11:852–866.
- 604 Kliber A., Eckert CG. 2005. Interaction between founder effect and selection during biological
605 invasion in an aquatic plant. *Evolution; international journal of organic evolution* 59:1900–
606 1913.
- 607 Le Corre V., Kremer A. 2012. The genetic differentiation at quantitative trait loci under local
608 adaptation. *Molecular ecology* 21:1548–1566.
- 609 Lenormand T. 2002. Gene flow and the limits to natural selection. *Trends in ecology & evolution*
610 17:183–189.
- 611 Lohr JN., Haag CR. 2015. Genetic load, inbreeding depression and hybrid vigor covary with
612 population size: an empirical evaluation of theoretical predictions. *Evolution; international*
613 *journal of organic evolution*. DOI: 10.1111/evo.12802.

- 614 Louette G., Gerald L., Joost V., Raquel O., De Meester L. 2007. The founding mothers: the
615 genetic structure of newly established *Daphnia* populations. *Oikos* 116:728–741.
- 616 Makino W., Tanabe AS. 2009. Extreme population genetic differentiation and secondary contact
617 in the freshwater copepod *Acanthodiaptomus pacificus* in the Japanese Archipelago.
618 *Molecular ecology* 18:3699–3713.
- 619 Matsumoto M., Nishimura T. 1998. Mersenne twister: a 623-dimensionally equidistributed
620 uniform pseudo-random number generator. *ACM Transactions on Modeling and Computer
621 Simulation* 8:3–30.
- 622 Mills S., Lunt DH., Gómez A. 2007. Global isolation by distance despite strong regional
623 phylogeography in a small metazoan. *BMC evolutionary biology* 7:225.
- 624 Montero-Pau J., Serra M. 2011. Life-cycle switching and coexistence of species with no niche
625 differentiation. *PloS one* 6:e20314.
- 626 Montero-Pau J., Serra M., Gómez A. 2016. Diapausing egg banks, lake size, and genetic diversity
627 in the rotifer *Brachionus plicatilis* Müller (Rotifera, Monogononta). *Hydrobiologia*:1–15.
- 628 Muñoz J., Gómez A., Green AJ., Figuerola J., Amat F., Rico C. 2008. Phylogeography and local
629 endemism of the native Mediterranean brine shrimp *Artemia salina* (Branchiopoda:
630 Anostraca). *Molecular ecology* 17:3160–3177.
- 631 Nosil P. 2007. Divergent host plant adaptation and reproductive isolation between ecotypes of
632 *Timema cristinae* walking sticks. *The American naturalist* 169:151–162.
- 633 Orsini L., Mergeay J., Vanoverbeke J., De Meester L. 2013. The role of selection in driving
634 landscape genomic structure of the waterflea *Daphnia magna*. *Molecular ecology* 22:583–

- 635 601.
- 636 Ortells R., Gómez A., Serra M. 2003. Coexistence of cryptic rotifer species: ecological and
- 637 genetic characterisation of *Brachionus plicatilis*. *Freshwater biology* 48:2194–2202.
- 638 Ortells R., Olmo C., Armengol X. 2011. Colonization in action: genetic characteristics of *Daphnia*
- 639 magna Strauss (Crustacea, Anomopoda) in two recently restored ponds. *Hydrobiologia*
- 640 689:37–49.
- 641 Parmesan C., Yohe G. 2003. A globally coherent fingerprint of climate change impacts across
- 642 natural systems. *Nature* 421:37–42.
- 643 Rosenblum EB., Hickerson MJ., Moritz C. 2007. A multilocus perspective on colonization
- 644 accompanied by selection and gene flow. *Evolution; international journal of organic*
- 645 *evolution* 61:2971–2985.
- 646 Tarazona E., García-Roger EM., Carmona MJ. 2017. Experimental evolution of bet hedging in
- 647 rotifer diapause traits as a response to environmental unpredictability. *Oikos* 126:1162–
- 648 1172.
- 649 Tavernini S. 2008. Seasonal and inter-annual zooplankton dynamics in temporary pools with
- 650 different hydroperiods. *Limnologica - Ecology and Management of Inland Waters* 38:63–
- 651 75.
- 652 Templeton AR. 2008. The reality and importance of founder speciation in evolution. *BioEssays:*
- 653 *news and reviews in molecular, cellular and developmental biology* 30:470–479.
- 654 Tortajada AM., Carmona MJ., Serra M. 2009. Does haplodiploidy purge inbreeding depression in
- 655 rotifer populations? *PloS one* 4:e8195.

- 656 Tortajada AM., Carmona MJ., Serra M. 2010. Effects of population outcrossing on rotifer fitness.
657 *BMC evolutionary biology* 10:312.
- 658 Vanschoenwinkel B., Waterkeyn A., Nihwatiwa T., Pinceel T., Spooren E., Geerts A., Clegg B.,
659 Brendonck L. 2011. Passive external transport of freshwater invertebrates by elephant and
660 other mud-wallowing mammals in an African savannah habitat. *Freshwater biology*
661 56:1606–1619.
- 662 Verhoeven KJF., Macel M., Wolfe LM., Biere A. 2011. Population admixture, biological invasions
663 and the balance between local adaptation and inbreeding depression. *Proceedings.*
664 *Biological sciences / The Royal Society* 278:2–8.
- 665 Violle C., Nemergut DR., Pu Z., Jiang L. 2011. Phylogenetic limiting similarity and competitive
666 exclusion. *Ecology letters* 14:782–787.
- 667 Walser B., Haag CR. 2012. Strong intraspecific variation in genetic diversity and genetic
668 differentiation in *Daphnia magna*: the effects of population turnover and population size.
669 *Molecular ecology* 21:851–861.
- 670 Waters JM. 2011. Competitive exclusion: phylogeography’s “elephant in the room”? *Molecular*
671 *ecology* 20:4388–4394.
- 672 Waters JM., Fraser CI., Hewitt GM. 2013. Founder takes all: density-dependent processes
673 structure biodiversity. *Trends in ecology & evolution* 28:78–85.
- 674 Wright S. 1931. Evolution in Mendelian Populations. *Genetics* 16:97–159.
- 675 Xu S., Hebert PDN., Kotov AA., Cristescu ME. 2009. The noncosmopolitanism paradigm of
676 freshwater zooplankton: insights from the global phylogeography of the predatory

677 cladoceran *Polyphemus pediculus* (Linnaeus, 1761) (Crustacea, Onychopoda). *Molecular*
678 *ecology* 18:5161–5179.

679 Zierold T., Hanfling B., Gómez A. 2007. Recent evolution of alternative reproductive modes in
680 the “living fossil” *Triops cancriformis*. *BMC evolutionary biology* 7:161.

681

682

Figure legends

Figure 1. Demographic submodel.

Figure 2. Population differentiation (F_{ST}) after 1000 sexual generations plotted against migration (M) with and without a diapausing egg bank for (A) $K = 2 \cdot 10^2$, and (B) $K = 2 \cdot 10^7$ individuals. The rest of parameters were $r = 0.3 \text{ d}^{-1}$, $n = 5$, $s = 0$ and $F = 1$. Box plots are based on 50 replicate simulations. Boxes represent 25th /75th percentile and black dots the 5th/95th percentile. Thin black lines and thick gray lines in each bar represent the median and the mean, respectively. Dashed horizontal lines show the initial value of F_{ST} after foundation.

Figure 3. Population differentiation (F_{ST}) after 1000 sexual generations plotted against carrying capacity (K) with and without a diapausing egg bank. Simulation values for other parameters were $r = 0.3 \text{ d}^{-1}$, $n = 5$, $s = 0$, $F = 1$ and $M = 2$. Data is based on 50 replicate simulations. Boxes represent 25th /75th percentile and black dots the 5th/95th percentile. Thin black lines and thick gray lines in each bar represent the median and the mean, respectively. Dashed horizontal line shows the initial value of F_{ST} after foundation.

Figure 4. Effect of different carrying capacities (K) on F_{ST} along 4000 sexual generations. (A, B) Box plot of the increment of F_{ST} (ΔF_{ST}) after 4000 sexual generations (A) without and (B) with diapausing egg bank is shown. (C, D) Time course of the average F_{ST} values along 4000 generations (C) without and (D) with diapausing egg bank. Simulation conditions were $r = 0.3 \text{ d}^{-1}$, $n = 5$, $s = 0$, $F = 1$ individual and $M = 2$ individuals. Data is based on 50 replicates. Boxes represent 25th /75th percentile and black dots the 5th/95th percentile. Thin black lines and thick

gray lines in each bar represent the median and the mean, respectively. Dashed horizontal lines show the initial value of F_{ST} after foundation.

Figure 5. Contour plot showing F_{ST} values after 1000 sexual generations at different combinations to population growth rates and carrying capacity (A) without and (B) with diapausing egg bank. Simulation conditions were $n = 5$, $s = 0$, $F = 1$ and $M = 2$. Data is based on 50 replicates.

Figure 6. Time course of the average F_{ST} value along 4000 generations for different number of founders ($F = 1, 2, 5$ and 50), for $K = 2 \cdot 10^4$ (A) and $K = 2 \cdot 10^7$ (B) and $M = 2$. Solid lines: without diapausing egg bank, dotted lines: with diapausing egg bank. Average F_{ST} values obtained from 50 replicates.


Figure 7. Box plot graph of F_{ST} and F_{STQ} values after 1000 sexual generations with different recombination rates for two different values of fitness components ($\delta = 10^{-4}$ and 10^{-2} d^{-1}) and with presence of a diapausing egg bank. For each of the fitness scenario, the left panel refers to $K = 2 \cdot 10^4$ and the right panel to $K = 2 \cdot 10^7$. Other parameters were $r = 0.3 \text{ d}^{-1}$, $n = 5$, $s = 5$, $F = 1$ and $M = 2$. Data is based on 100 replicates. Boxes represent 25th /75th percentile and black dots the 5th/95th percentile. Thin black lines and thick gray lines in each bar represent the median and the mean, respectively. Dashed horizontal lines show the initial value of F_{ST} after foundation. Asterisks indicate F_{ST} statistically different from those without selection ($\delta = 0$) (**, $\alpha = 0.05$; *, $\alpha = 0.1$).

722 Table 1. Summary of model parameters and assumed values.

Parameter	Definition	Value
F	Number of founders (individuals)	1 - 50
M	Number of immigrants per sexual generation (individuals)	0 - 10^5
γ	Annual survival proportion of eggs in the egg bank	0.763 *
λ	Annual hatching proportion of diapausing eggs	0.046 *
y	Sexual generations	1000/4000
τ	Duration of the asexual growth period (days)	60
r	Clonal growth rate of each genotype (days^{-1})	0.05 - 1.00
K	Carrying capacity (individuals)	$2 \cdot 10^2$ - $2 \cdot 10^7$
m	Sexual proportion	0.7 [†]
sr	Sex ratio	0.5 [‡]
e	Diapausing egg production per sexual female	3
n	Number of neutral loci	5
δ	Additive value on r (days^{-1})	10^{-5} - 10^{-1}


723 * Calculated from (García-Roger, Carmona & Serra, 2006b) [†] (Alver & Hagiwara, 2007) [‡] (Aparici,
724 Carmona & Serra, 1998).

725




726

727 Figure 1.



728

729 Figure 2.



730

731 Figure 3.

732





Figure 4.



736

737 Figure 5.

42




Figure 6.




Figure 7.