

A peer-reviewed version of this preprint was published in PeerJ on 22 June 2018.

[View the peer-reviewed version](https://peerj.com/articles/4998) (peerj.com/articles/4998), which is the preferred citable publication unless you specifically need to cite this preprint.

Klecka J, Hadrava J, Koloušková P. 2018. Vertical stratification of plant-pollinator interactions in a temperate grassland. PeerJ 6:e4998 <https://doi.org/10.7717/peerj.4998>

Vertical stratification of plant-pollinator interactions in a temperate grassland

Jan Klecka¹, Jiří Hadrava^{1,2}, and Pavla Koloušková¹

¹Czech Academy of Sciences, Biology Centre, Institute of Entomology, České Budějovice, Czech Republic

²Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic

Corresponding author:

Jan Klecka

Email address: jan.klecka@entu.cas.cz

ABSTRACT

Visitation of plants by different pollinators depends on individual plant traits, spatial context, and other factors. A neglected aspect of small-scale variation of plant-pollinator interactions is the role of vertical position of flowers. We conducted a series of experiments to study vertical stratification of plant-pollinator interactions in a dry grassland. We observed flower visitors on cut inflorescences of *Centaurea scabiosa* and *Inula salicina* placed at different heights above ground in two types of surrounding vegetation: short and tall. Even at such a small-scale, we detected significant shift in total visitation rate of inflorescences in response to their vertical position. In short vegetation, inflorescences close to the ground were visited more frequently, while in tall vegetation, inflorescences placed higher received more visits. Moreover, we found major differences in the composition of the pollinator community on flowers at different heights. In a second experiment, we measured flower visitation rate in inflorescences of *Salvia verticillata* of variable height. Total flower visitation rate increased markedly with inflorescence height in this case. Data on seed set of individual plants provide evidence for a corresponding positive pollinator-mediated selection on increased inflorescence height. Overall, our results demonstrate strong vertical stratification of plant-pollinator interactions at the scale of mere decimetres. This may have important ecological as well as evolutionary implications.

INTRODUCTION

Interactions between plants and their pollinators play an important role in the evolution (Grant and Grant, 1965; Bronstein et al., 2006; Suchan and Alvarez, 2015) and maintenance of biodiversity (Bascompte et al., 2003, 2006; Bascompte and Jordano, 2007) in terrestrial ecosystems. However, the presence and frequency of interactions between particular plants and pollinators vary in time (Olesen et al., 2008) and space (Espíndola et al., 2011; Newman et al., 2015). Spatial variation in plant-pollinator interactions is observed from continental scales across the entire distributional range of a plant species (Espíndola et al., 2011), down to small habitat patches and individual plants (Ohashi and Yahara, 1998; Dupont et al., 2014; Akter et al., 2017). At the smallest scale, the position of an inflorescence in the context of the surrounding vegetation may affect the frequency and identity of flower visitors with consequences for plant reproduction.

Many plants show high levels of phenotypic plasticity. Inflorescence height is thus highly variable at the intraspecific level and may be important in driving visitation of individual plants. However, the importance of inflorescence height is little understood in grasslands, where the vertical distance between different flowers is rarely more than a few decimetres. In communities of multiple plants species, a few observational studies found that different bee species tend to visit flowers at different heights (Gumbert and Kunze, 1999; Hoehn et al., 2008). In addition, in a trait-based analysis of a plant-flower visitor network in a German grassland, Junker et al. (2013) found that inflorescence height was the most important species trait after phenology to explain which plant species were visited by which insects. Studies investigating the effects of inflorescence height at the intraspecific level found that inflorescence height is under significant

47 pollinator-mediated selection (Sletvold et al., 2010; Jiang and Li, 2017; Trunschke et al., 2017), although
48 these studies unfortunately did not include direct observations of flower visitors.

49 Although our knowledge of vertical stratification of plant-flower visitor interactions in grasslands is
50 limited, even less is known about how the relationship between pollination and inflorescence height is
51 modified by other environmental factors such as the structure of the surrounding vegetation. For example,
52 Sletvold et al. (2013) observed significant pollinator-mediated selection for tall *Dactylorhiza lapponica*
53 plants in tall vegetation, while there was no significant selection on plant height in short vegetation.
54 Similarly, in field experiments with *Primula farinosa*, Ehrlén et al. (2002) showed that short plants
55 were more pollen-limited than tall plants and the difference was larger in a habitat with tall vegetation.
56 However, Ågren et al. (2006) showed that removal of litter and pruning of vegetation around individual
57 *Primula farinosa* plants increased their fruit and seed production apparently because of an increase in
58 their nutritional status rather than increased pollination. The role of vegetation height for pollination thus
59 remains unclear. These studies also looked at the topic entirely from the plant's point of view and did not
60 measure the effects of vegetation height on visitation frequency or pollinator foraging behaviour. Apart
61 from vegetation height, local density of the same or other plant species (Bartkowska and Johnston, 2014),
62 distance from neighbours (Caraballo-Ortiz et al., 2011), as well as spatial variation in the proportion
63 of different morphs (Toräng et al., 2006) can also modify the effects of plant height on its reproductive
64 success.

65 Most previous research focused on the importance of inflorescence height for plant reproduction,
66 while little attention has been devoted to understanding whether and why foraging insects prefer flowers at
67 certain heights. Some insight can be gained from observations of foraging behaviour of individual insects.
68 In honeybees, "horizontal movement" characterised by a tendency of individual bees to fly between plants
69 of a similar height has been reported (Levin and Kerster, 1973; Faulkner, 1976). Preference of flowers at a
70 certain height was demonstrated also in solitary bees (Gumbert and Kunze, 1999; Hoehn et al., 2008) and
71 wasps (Peakall and Handel, 1993). Flying at a constant height may be advantageous from an energetic
72 point of view for optimally foraging flower visitors (Pyke, 1978). Also, flowers close to the ground may
73 be avoided by some insects because their visitation requires the ability to manoeuvre among plant stems,
74 which may be challenging in dense vegetation (Gumbert and Kunze, 1999).

75 We conducted a set of field experiments in a dry grassland in the Czech Republic to fill in some of
76 these knowledge gaps. Specifically, our aim was to test whether total visitation rate and the composition
77 of flower visitor assemblages depend on inflorescence height and whether the relationship is modified
78 by the height of the surrounding vegetation. Another aim was to test whether inflorescence height is
79 under pollinator-mediated selection in our system. Our field experiments with three species of plants
80 common in dry grasslands in Central Europe showed that visitation rate varied with inflorescence height,
81 moreover the relationship differed between different flower visitor taxa and was modified by the height of
82 the surrounding vegetation. We also detected significant increase in seed production with inflorescence
83 height in *Salvia verticillata*.

84 METHODS

85 Field experiments

86 We conducted two field experiments in a dry grassland near Český Krumlov, in the southern part of the
87 Czech Republic (48°49'28"N 14°18'59"E). The study site is a species rich calcareous grassland on a
88 southwest-facing slope managed by occasional pasture by cows and sheep. The area is state-owned and
89 publicly accessible. No permits were thus needed for this study.

90 In the first experiment, we observed visitation of inflorescences of two plant species, *Centaurea*
91 *scabiosa* and *Inula salicina*, at different heights above ground. To avoid confounding factors, e.g. taller
92 plants having a different size of inflorescences than shorter plants, we used inflorescences cut from plants
93 in the local population. We selected inflorescences of a similar size and general appearance and placed
94 them in 15 ml tubes with water. We attached each tube to a bamboo stick of different length and attached
95 the stick to the ground. This way, we manipulated the height of the inflorescence between 5 cm and
96 105 cm above ground. We placed the inflorescences along two short transects, each containing seven
97 inflorescences placed 50 cm apart. One transect was surrounded by short and the other by tall and dense
98 vegetation; the transects were ca. 10 m apart. The area of short vegetation was grazed by cows in the
99 spring, while the area of tall vegetation was not managed. Short vegetation was characterised by most

100 plants <10 cm tall; the average height of inflorescences of all plants growing within 50 cm from the
101 transect in all directions was 7.2 cm ($SD = 6.00$). Tall vegetation was composed of a dense layer of plants
102 reaching ca. 50 cm; plants growing within 50 cm from the transect had flowers on average 50.1 cm above
103 ground ($SD = 14.73$). We individually placed seven inflorescences of either *Centaurea scabiosa* or *Inula*
104 *salicina* in each transect at 5, 15, 25, 45, 65, 85, and 105 cm above ground in a randomised order.

105 We observed visitation of the inflorescences by insects between 10:30 and 16:00 hours, for 30
106 minutes in each transect, and identified all visitors at the species level or classified them into taxonomical
107 groups with the highest precision we could achieve without capturing the insects. Both transects were
108 observed simultaneously, one person observed each transect. After the 30 minute period, we replaced the
109 inflorescences and randomised the order of their vertical position along each transect and took another set
110 of observations. In total, we measured inflorescence visitation in 20 transects in *Inula salicina*, 10 in short
111 and 10 in tall vegetation, and 16 transects in *Centaurea scabiosa*, 8 in short and 8 in tall vegetation. In
112 total, this amounts to 18 hours of observations.

113 In the second experiment, we focused on the effect of inflorescence height for flower visitation rate
114 and its consequences for seed set in *Salvia verticillata* at the same site. In this case, we did not cut the
115 inflorescences so that we could test whether flower visitation rate varied between inflorescences within
116 the natural limits of their height above ground and to test whether variation in flower visitation rate
117 translated into differences in seed set; i.e. whether female fitness was affected by the vertical position
118 of the inflorescence. We used the following approach to minimise confounding effects, such as taller
119 plants having more resources, different display size, etc. We selected 17 plants of *Salvia verticillata* with
120 multiple ramets of approximately the same size and with the first several flowers open or with buds ready
121 to start flowering on 18 July 2017. We took advantage of the fact that the inflorescences grow on relatively
122 long and flexible stems. We bent one of them close to the ground, where it was attached to a stick so that
123 the bottom of the inflorescence was positioned just above ground. We made sure that the orientation of
124 the inflorescence remained unchanged. The second stem was attached to another stick so that it reached a
125 maximum height and the third inflorescence was positioned at an intermediate height. This way, the only
126 difference between the inflorescences was their vertical position.

127 We performed observations of flower visitation in individual *S. verticillata* plants at one of three dates
128 (20 July, 21 July, and 2 August 2017), depending on when they reached the peak of flowering. The three
129 manipulated ramets per plant were observed simultaneously during one 30 minute period and filmed using
130 three digital cameras, which gave a total of 25 hours of recordings. Afterwards, we measured the height
131 of each inflorescence as a distance of the highest open flower from the ground and counted the number of
132 open flowers. We then watched the recordings and counted and identified all flower visitors. For each
133 visitor, we also counted the number of flowers visited during each inflorescence visit.

134 We waited for the seeds of *S. verticillata* to ripen and then harvested them on 14 or 24 August 2017
135 depending on seed development in individual plants. We counted the number of developed seeds and
136 the maximum potential seed set by multiplying the number of flowers by four which is the number of
137 seeds the plant can produce per flower. We counted the flowers and seeds in individual whorls within each
138 inflorescence separately to gain data on potential differences in percentage seed set along the inflorescence
139 from the lowest to the highest whorl.

140 As already mentioned, we identified flower-visiting insects without capturing them. Naturally, we
141 could not identify all individuals to the species level, so we classified some of them into higher taxa or
142 categories, such as "small solitary bees". The most abundant flower visitors were bumblebees, *Bombus*
143 spp., some of which are difficult to identify alive. Fortunately, we have extensive collections from the
144 study site, so we know that there are three species, which we could not distinguish from *Bombus terrestris*,
145 specifically *B. lucorum*, *B. cryptarum* and *B. magnus*. However, over 85% of individuals of this species
146 group in our collections from this site belong to *B. terrestris*. Similarly, *Bombus lapidarius* could be
147 confused with *B. confusus* and *B. ruderarius*, but they have been rarely found on the site. Similar level
148 of uncertainty exists in our identification of *Bombus sylvarum*. The number of potential bumblebee
149 misidentifications during the field observations was thus low and unlikely to confound our results.

150 Data analysis

151 We tested how the total number of visits and the number and proportion of visits by individual flower
152 visitor taxa depended on inflorescence height using generalized additive models (GAM) to account for
153 the non-linear nature of these relationships. The identity of individual plants, each having three ramets

manipulated and observed, was included as a random factor in analyses of data from the experiment with *Salvia verticillata*; i.e. generalized additive mixed models (GAMM) were used in this case. Poisson distribution with overdispersion (quasipoisson distribution) was used for the number of visits, while overdispersed binomial (quasibinomial) distribution was used for data on proportions. These analyses were performed using mgcv 1.8-17 package (Wood, 2006) in R 3.2.3 (R Core Team, 2015).

To gain insights into the effects of inflorescence height on plant fitness, we tested how seed set of individual ramets depended on the number of flowers and the inflorescence height using a generalized linear model. Similarly as in analyses of selection gradients (Lande and Arnold, 1983), we standardised both predictors to have zero mean and unit variance. Partial regression coefficients then allowed us to compare whether reproductive performance (seed set) depended more strongly on the number of flowers or inflorescence height.

RESULTS

In the first experiment, we observed a strong, mostly non-linear, dependence of the total visitation rate on inflorescence height in both *Centaurea scabiosa* and *Inula salicina* (Fig. 1; raw data: Table S1). Analysis using generalised additive models (GAM, Table 1) showed that the relationship was significant in *Centaurea scabiosa* as well as in *Inula salicina* in both short and tall vegetation (Table 1). Also, there was a significant difference in the shape of the relationship between total visitation and inflorescence height in short vs. tall surrounding vegetation in both *Centaurea scabiosa* ($F = 19.27, P < 10^{-6}$) and *Inula salicina* ($F = 12.46, P = 3.10 \times 10^{-5}$). Comparison of the results presented in Fig. 1 shows that the difference between short and tall vegetation is mostly that visitation rate of inflorescences of both plant species positioned <50 cm above ground dropped in tall compared to short surrounding vegetation. Moreover, analysis of visitation rate of the most abundant flower visitors showed that different insect species had contrasting height preferences modified by the height of the surrounding vegetation (Fig. 2, Fig. 3, and Table 1). Overall, we observed 638 visits (16 taxa) on *Centaurea scabiosa* and 286 visits (13 taxa) on *Inula salicina*.

In *Centaurea scabiosa*, we observed a significant effect of inflorescence height on the number of inflorescence visits by *Bombus lapidarius*, which preferred mostly inflorescences close to the ground (Fig. 2 A., Table 1), *Halictus quadricinctus*, which preferred inflorescence high above ground (Fig. 2 C.), and small solitary bees, which visited mostly inflorescences at an intermediate height (Fig. 2 D.). The relationship was not significant in *Bombus terrestris* (Table 1). Two species, *Bombus terrestris* and *Halictus quadricinctus*, avoided the area of tall vegetation despite being frequently observed in transects surrounded by short vegetation. On the other hand, *Bombus lapidarius* was common in both habitats and showed a shift towards inflorescences higher above ground in the transects surrounded by tall vegetation (Fig. 2 A. and E.); the relationship between visitation and inflorescence height was significantly different in short and tall vegetation ($F = 21.31, P < 10^{-6}$). On the contrary, small solitary bees did not shift their visitation (Fig. 2 D. and F.).

In *Inula salicina*, only two groups of flower visitors were abundant enough for detailed analysis. Hoverflies (Diptera: Syrphidae) visited mostly inflorescences >40 cm above ground and the height of the surrounding vegetation had no effect on the relationship between the number of visits and inflorescence height (Fig. 3 A. and C.; $F = 0.04, P = 0.95$). On the other hand, small solitary bees favoured inflorescences close to the ground in short vegetation and shifted higher above ground in tall vegetation (Fig. 3 B. and D.), the relationship of visitation with inflorescence height was significantly different in short and tall vegetation ($F = 15.39, P < 10^{-6}$; Table 1).

Different flower visitors responded to inflorescence height and the height of the surrounding vegetation in a species-specific way (Fig. 4, Table 2). For example, *Bombus lapidarius* visited mostly inflorescences of *Centaurea scabiosa* positioned close to the ground when the surrounding vegetation was short, but shifted to inflorescences higher above ground when the surrounding vegetation was tall. Visitation of flowers close to the ground surrounded by tall vegetation was then dominated by small solitary bees (Fig. 4 A. and C). The composition of the flower visitor assemblage at a particular height thus differed according to the height of the surrounding vegetation.

In the second experiment, the total number of flower visitors (Fig. 5A.) and the per-flower visitation rate (Fig. 5B.) in *Salvia verticillata* significantly increased with inflorescence height (GAMM; $edf = 1.91, F = 21.04, P < 1 \times 10^{-6}$ and $edf = 1.836, F = 21.62, P < 1 \times 10^{-6}$, respectively) (raw data: Tables S2 and S3). Overall, we observed 300 visits by seven taxa.

Table 1. The effects of inflorescence height and surrounding vegetation height on the number of visits by different insects. Summary of results of generalized additive models testing the dependence of visitation of *Centaurea scabiosa* and *Inula salicina* on inflorescence height. Groups of flower visitors which had an insufficient number of observations for analysis were not analysed separately, but were included in the total visitation. *edf* = estimated degrees of freedom, which gives a measure of the complexity of the shape of the relationship (*edf* = 1 is a linear relationship). NA = cases when the number of observations was insufficient for analysis. The results are presented graphically in Figs. 1, 2, and 3.

Response	Short vegetation			Tall vegetation			Short vs. tall vegetation	
	<i>edf</i>	F	P	<i>edf</i>	F	P	F	P
Visits of <i>Centaurea scabiosa</i>								
Total visitors	3.00	10.01	$3.77 * 10^{-6}$	3.55	15.21	$< 1 * 10^{-6}$	19.27	$< 1 * 10^{-6}$
<i>Bombus lapidarius</i>	2.80	8.38	$4.17 * 10^{-5}$	3.62	6.15	0.0005	21.32	$< 1 * 10^{-6}$
<i>Bombus terrestris</i>	1.69	1.92	0.1640	NA	NA	NA	NA	NA
<i>Halictus quadricinctus</i>	2.16	6.00	0.0028	NA	NA	NA	NA	NA
small solitary bees	2.62	8.16	$6.20 * 10^{-5}$	2.62	8.16	$6.20 * 10^{-5}$	0	1
Visits of <i>Inula salicina</i>								
Total visitors	1	5.38	0.0223	2.58	6.27	0.0005	12.46	$3.10 * 10^{-5}$
small solitary bees	1.32	17.12	$4.90 * 10^{-6}$	3.65	3.75	0.0072	15.39	$< 1 * 10^{-6}$
Syrphidae	1.94	9.54	0.0001	1.94	9.54	0.0001	0.04	0.9543

208 Different groups of flower visitors also showed distinct patterns in their preference for inflorescences
 209 of different heights. *Bombus terrestris* visited mostly the highest inflorescences, followed by *Bombus*
 210 *lapidarius*, while *Bombus sylvarum* showed no significant dependence of visitation on inflorescence
 211 height and small solitary bees visited mostly inflorescences close to the ground (Fig. 6, Table 3). When
 212 expressed as the proportion of visits attributed to individual pollinator groups, our results show that plants
 213 with inflorescences closest to the ground were visited equally by *Bombus terrestris* and small solitary
 214 bees (ca. 40% each), followed by *Bombus sylvarum* (almost 20%) (Fig. 7 A., Table 3). On the other
 215 hand, visits to inflorescences high above ground were dominated solely by *Bombus terrestris* (Fig. 7 A.,
 216 Table 3). Different visitors also significantly differed in one aspect of foraging behaviour, namely in the
 217 proportion of flowers in an inflorescence probed during a visit (Table 3; GLM, $F = 5.24, P = 3.82 * 10^{-5}$).
 218 *Apis mellifera*, which was excluded from the previous analyses because it was too rare, visited on average
 219 over 60% of flowers during one inflorescence visit and the three bumblebee species over 40%. On the
 220 contrary, other visitors, which we classified as small solitary bees, Syrphidae, and other Diptera, visited
 221 less than 20% of the flowers per inflorescence visit (Fig. 7 B.).

222 Differences in visitation translated into differences in seed set, which significantly increased with both
 223 the number of flowers in an inflorescence (GLM, $F = 6.21, P = 0.0165$; Fig. 8 A.) and with inflorescence
 224 height (GLM, $F = 6.09, P = 0.0175$; Fig. 8 B.) (raw data: Table S4). Based on a comparison of
 225 partial regression coefficients, seed set depended more strongly on inflorescence height ($\beta = 0.29, SE =$
 226 0.101) than on the number of flowers ($\beta = 0.25, SE = 0.089$) (both variables were standardised to allow
 227 meaningful comparison of regression coefficients). When we included the number of inflorescence visits
 228 in the model, there was no longer any significant effect of inflorescence height ($F = 1.55, P = 0.2201$),
 229 while seed set significantly increased with the number of visits ($F = 21.46, P = 2.98 * 10^{-5}$; Fig. 8 C.).
 230 We obtained the same results when using the number of flower visits as a predictor ($F = 9.87, P = 0.0030$
 231 for the number of flower visits and $F = 1.76, P = 0.1911$ for inflorescence height; Fig. 8 C.). Hence,
 232 our results show that increased seed set of taller ramets was driven primarily by increased visitation.
 233 The importance of the vertical position of flowers is underscored by the fact that we found a significant
 234 increase in the proportional seed set of individual whorls within individual inflorescences when moving
 235 from the lowest to the highest whorl (GLM, $F = 12.80, P = 0.0004$; Fig. 9).

236 DISCUSSION

237 **The effects of inflorescence and vegetation height on visitation by potential pollinators**
 238 In the first experiment, using *Centaurea scabiosa* and *Inula salicina* as focal species, total inflorescence
 239 visitation peaked approximately at or slightly above the level of the surrounding vegetation in transects
 240 surrounded by both short and tall vegetation. These results are consistent with observations that flower

Table 2. The effects of inflorescence height and surrounding vegetation height on relative visitation by different insects. Summary of results of generalized additive models testing the dependence of the proportion of visits attributed to most abundant visitor taxa on inflorescence height in *Centaurea scabiosa* and *Inula salicina*. *edf* = estimated degrees of freedom, which gives a measure of the complexity of the shape of the relationship (*edf* = 1 is a linear relationship). NA = cases when the number of observations was insufficient for the analysis. The results are presented graphically in Fig. 4.

Response	Short vegetation			Tall vegetation			Short vs. tall vegetation	
	<i>edf</i>	F	P	<i>edf</i>	F	P	F	P
Visits of <i>Centaurea scabiosa</i>								
<i>Bombus lapidarius</i>	1	30.60	$< 1 * 10^{-6}$	2.16	4.79	0.0073	14.22	$< 1 * 10^{-6}$
<i>Bombus terrestris</i>	1	1.28	0.2640	NA	NA	NA	NA	NA
<i>Halictus quadricinctus</i>	1	21.21	$3.62 * 10^{-5}$	NA	NA	NA	NA	NA
small solitary bees	1.72	1.18	0.3045	1	16.02	0.0002	11.871	0.0005
Visits of <i>Inula salicina</i>								
small solitary bees	1	35.25	$< 1 * 10^{-6}$	1	2.16	0.1450	14.78	0.0002
Syrphidae	1.93	15.14	$1.46 * 10^{-6}$	1.93	15.14	$1.46 * 10^{-6}$	1.48	0.2333

Table 3. The effect of inflorescence height in *Salvia verticillata* on inflorescence visitation.

Summary of results of generalized additive mixed models testing the dependence of the number of visits and the proportion of visits by most frequent visitor taxa on inflorescence height in *Salvia verticillata*. *edf* = estimated degrees of freedom, which gives a measure of the complexity of the shape of the relationship (*edf* = 1 is a linear relationship). The results are plotted in Figs. 6 and 7.

Response	No. of visits / 30 min.			Proportion of visits		
	<i>edf</i>	F	P	<i>edf</i>	F	P
<i>Bombus lapidarius</i>	1.91	8.56	0.0009	1.68	0.89	0.2640
<i>Bombus terrestris</i>	1.88	19.28	$1.66 * 10^{-6}$	1	6.90	0.0118
<i>Bombus sylvarum</i>	1.21	0.03	0.8700	1	9.93	0.0029
small solitary bees	1	10.08	0.0026	1	38.48	$< 1 * 10^{-6}$

241 visiting bees tend to fly at a specific height and when they leave one inflorescence, they are more likely to
 242 fly to another one at a similar height compared to inflorescences lower or higher above ground (Levin
 243 and Kerster, 1973; Gumbert and Kunze, 1999). Flowers positioned above the level of a dense layer of
 244 vegetation are probably easier to detect and thus attract more visitors (Gumbert and Kunze, 1999). Height
 245 preferences of flower visitors may lead to vertical stratification of the plant-pollinator network if different
 246 species vary in their behaviour (Roubik, 1993; Gumbert and Kunze, 1999; Ramalho, 2004).

247 Different responses of individual species or taxonomic groups of flower visitors to inflorescence
 248 height, which we observed, mean that inflorescences at different heights varied in the composition of their
 249 flower visitor assemblages (Figs. 4 and 7). Plant-pollinator interactions thus appear stratified along the
 250 vertical dimension despite the fact that height differences between inflorescences were in the order of mere
 251 decimetres. Previous studies on vertical stratification of pollinator communities were conducted mostly
 252 in forests where the importance of the vertical dimension is more obvious. It seems there is generally a
 253 major difference in the composition of flower visitor communities between the canopy and the understory
 254 in tropical (Roubik, 1993; Nagamitsu et al., 1999; Ramalho, 2004) as well as temperate forests (Ulyshen
 255 et al., 2010), probably related to vertical distribution of flowers preferred by different species (Ramalho,
 256 2004). However, vertical stratification of plant-pollinator interactions has been less studied in grasslands
 257 where the vertical distances are limited usually to several decimetres. In one of the few available studies,
 258 (Gumbert and Kunze, 1999) observed differentiation between visitor communities on flowers below and
 259 above the dominant grass layer in a tropical wetland. Similarly, several species of bees visiting pumpkins
 260 in an agricultural landscape preferred flowers at different heights (Hoehn et al., 2008). Our results show
 261 that vertical stratification of plant-flower visitor interactions may be important also in common temperate
 262 grasslands.

263 An interesting observation is that small solitary bees were the dominant flower visitors close to the
 264 ground in all three plant species. We noticed that unlike other flower visitors, they were frequently flying
 265 among plant stems close to the ground even in dense vegetation and visiting flowers hidden there, such
 266 as flowers of *Rubus* sp., which was growing in parts of the study site. The same pattern was observed

267 by Gumbert and Kunze (1999), who suggested that small bees, which are able to manoeuvre in dense
268 vegetation, may benefit from decreased competition for floral resources because most other flower visitors
269 avoid this microhabitat. This seems to be a likely explanation for our results as well. Behaviour of small
270 solitary bees contrasted with the behaviour of larger species, such as bumblebees, which were flying
271 above the layer of dense vegetation. For example, almost all observations of *Bombus lapidarius* in tall
272 vegetation were at the height of >40 cm (Fig. 2 E.).

273 The effect of flower visitation was thus modified by the height of the surrounding vegetation because
274 most flower visitors avoided flowers within the layer of dense vegetation close to the ground. One
275 bumblebee species, *Bombus terrestris*, also visited mostly inflorescences close to the ground in short
276 vegetation (Fig. 2 B.), but almost completely avoided the area with tall vegetation, similarly to *Halictus*
277 *quadridentatus*, a solitary bee, which favoured inflorescences >60 cm above ground (Fig. 2 C.). Previous
278 studies on the effects of the structure of the surrounding vegetation for flower visitation and plant
279 reproductive success are rare and did not provide clear conclusions. For example, Ågren et al. (2006)
280 manipulated vegetation height and litter presence around individual plants of *Primula farinosa*, and found
281 that litter removal and vegetation pruning increased seed set, especially in short plants. They did not
282 report any data on visitation frequency of flowers in relation to vegetation structure. However, if we
283 assume that higher visitation leads to higher fruit or seed production, our observations of higher visitation
284 of inflorescences close to the ground in short vegetation exactly mirror these results. In tall vegetation,
285 visitation of both *Centaurea scabiosa* and *Inula salicina* peaked higher above ground, which fits the
286 results of Sletvold et al. (2013), who observed pollinator-mediated selection for taller inflorescences in
287 tall vegetation but not in short vegetation in a deceptive orchid, *Dactylorhiza lapponica*.

288 We did not measure the efficiency of different pollinators in the present study, so we cannot infer
289 consequences of the variation of visitation at different heights for the reproductive success of *Centaurea*
290 *scabiosa* and *Inula salicina*. However, data from detailed single visit experiments by other authors
291 demonstrated that different flower visitor species vary in pollen deposition by several orders of magnitude
292 (King et al., 2013). Hence, it is likely that variation in total visitation rate together with the variation
293 in visitor identity with inflorescence height affects reproductive success of plants in our system. We
294 addressed this question in a second experiment with *Salvia verticillata*.

295 **Consequences of inflorescence height for seed set and plant fitness**

296 Observations on *Salvia verticillata* were constrained by the natural range of inflorescence heights. Unlike
297 previous studies, we collected data both on flower visitation rates by pollinators and on seed set of
298 individual ramets in relation to inflorescence height. Our results are consistent with the hypothesis
299 of positive selection for inflorescence height, which other authors demonstrated in several other plant
300 species (Sletvold et al., 2010, 2013; Jiang and Li, 2017; Trunschke et al., 2017). However, we could also
301 demonstrate that inflorescences positioned higher above ground had higher total flower visitation rates.

302 Our experimental design allowed us to measure the dependence of seed set on inflorescence height,
303 because we compared similarly looking inflorescences in three ramets per plant (genet), whose height
304 was experimentally adjusted. Vertical position was thus the only apparent difference between the
305 inflorescences. This is important because taller plants usually have higher percentage seed set even
306 when they are hand-pollinated because they have more resources than shorter plants (Andersson, 1996;
307 Červenková and Münzbergová, 2014). Testing for pollinator-mediated selection on inflorescence height
308 thus requires specific experimental designs and is not possible by simply comparing plants of different
309 height. The most frequent approach is to compare selection coefficients for inflorescence height between
310 open-pollinated and hand-pollinated plants (Sletvold et al., 2010; Červenková and Münzbergová, 2014;
311 Jiang and Li, 2017; Trunschke et al., 2017). We used an alternative approach, which allowed us to skip
312 the hand-pollination treatment. We took advantage of the morphology of *Salvia verticillata*, which creates
313 multiple closely packed, relatively long, and flexible ramets, which can be easily pinned closer to the
314 ground or straightened up without causing damage. For our observations, we selected three ramets with
315 inflorescences of a similar length and general appearance per plant and randomly adjusted their vertical
316 position, so there was no known confounding factor. So, the observed positive correlation between
317 inflorescence height, visitation rate, and seed set can be interpreted as evidence for pollinator-mediated
318 selection on inflorescence height. Nevertheless, some caution is needed, because vertical position of the
319 inflorescence could also affect water transport, which is more difficult to inflorescences higher above
320 ground. Inflorescences at different heights also presumably experienced different levels of shading. In

321 additional, results of our first experiment, where we observed visitation rate on flowers of *Centaurea*
322 *scabiosa* and *Inula salicina* along a wider range of heights suggest that there may be an optimal height
323 maximising insect pollination depending on the context of the surrounding vegetation.

324 Flower visitation by different insect species can have different effects on plant fitness not only because
325 different species differ in the number of pollen grains deposited per visit (King et al., 2013), but also
326 because they differ in the relative frequency of movements between flowers on the same plant and between
327 different plants (Paton, 1993). Our observations showed that honeybees (*Apis mellifera*) visited a majority
328 of flowers within an inflorescence by moving along the inflorescence and probing one flower after another.
329 The three bumblebee species exploited slightly lower percentage of flowers, while small solitary bees and
330 Diptera usually probed only a few flowers per inflorescence (Fig. 7 B.). Different pollinators thus have
331 a different potential for geitonogamous pollination, because pollen from the previously visited plant is
332 deposited mostly on stigmas of the first few flowers and receipt of foreign pollen exponentially decreases
333 in each successive flower visit within an inflorescence (Thomson and Plowright, 1980; Gerber, 1985;
334 Morris et al., 1994). High levels of geitonogamous pollination may negatively affect plant fitness (Gerber,
335 1985; Waser and Price, 1991; de Jong et al., 1993; Ruane et al., 2013). In our population of *Salvia*
336 *verticillata*, we would expect higher level of geitonogamous pollination in inflorescences higher above
337 ground because of shifts in the flower visitor community (see Fig. 7), perhaps also in individual whorls
338 higher within an inflorescence. However, proportional seed set at the scale of entire inflorescences and
339 individual whorls increased with height, as a consequence of higher total visitation rate.

340 Inflorescences which are more attractive for pollinators may also attract higher numbers of florivores
341 and seed predators (Sletvold and Grindeland, 2008). Seed set is thus driven by a balance between
342 mutualistic and antagonistic interactions with flower visitors (Ehrlén et al., 2002; Schlinkert et al., 2016).
343 We did not notice any conspicuous evidence of florivory during observations of flower visitors or seed
344 predation when processing seeds of *Salvia verticillata*. However, in other plant species, florivory and seed
345 predation can have a large negative effect on plant fitness (Ruane et al., 2014). For example, Schlinkert
346 et al. (2016) found that abundance of both pollinators and florivores increased with plant height and
347 mutualistic and antagonistic interactions had contrasting effects on the number of seeds leading to seed
348 set being independent of plant height. The role of inflorescence height may thus be species-specific
349 and context-dependent, as shown also by our observations of inflorescence visitation in short and tall
350 vegetation.

351 **Conclusions**

352 In conclusion, we experimentally demonstrated that both total flower visitation and the composition of
353 the community of insect visitors changed with the vertical position of inflorescences in three common
354 plant species growing in a dry grassland. Moreover, we found that the dependence of visitation rate on
355 inflorescence height was mediated by the height of the surrounding vegetation. In one species, *Salvia*
356 *verticillata*, we also observed increased seed set with inflorescence height, which supports the hypothesis
357 of selection for increased inflorescence height. Overall, we detected pronounced vertical stratification of
358 plant-pollinator interactions at a scale of mere decimetres in a temperate grassland.

359 **ACKNOWLEDGMENTS**

360 This study was supported by the Czech Science Foundation (projects GP14-10035P and GJ17-24795Y
361 awarded to JK). We would like to thank Paolo Biella for help with identification of *Bombus* specimens
362 and three reviewers for their thoughtful comments.

363 **REFERENCES**

- 364 Ågren, J., Fortunel, C., and Ehrlén, J. (2006). Selection on floral display in insect-pollinated *Primula*
365 *farinosa*: effects of vegetation height and litter accumulation. *Oecologia*, 150(2):225–232.
- 366 Akter, A., Biella, P., and Klecka, J. (2017). Effects of small-scale clustering of flowers on pollinator
367 foraging behaviour and flower visitation rate. *PLoS ONE*, 12(11):e0187976.
- 368 Andersson, S. (1996). Floral variation in *Saxifraga granulata*: phenotypic selection, quantitative genetics
369 and predicted responses to selection. *Heredity*, 77(2):217–223.
- 370 Bartkowska, M. P. and Johnston, M. O. (2014). The sexual neighborhood through time: competition and
371 facilitation for pollination in *Lobelia cardinalis*. *Ecology*, 95(4):910–919.

- 372 Bascompte, J. and Jordano, P. (2007). Plant-animal mutualistic networks: the architecture of biodiversity.
373 *Annual Review of Ecology, Evolution, and Systematics*, 38:567–593.
- 374 Bascompte, J., Jordano, P., Melián, C. J., and Olesen, J. M. (2003). The nested assembly of plant–animal
375 mutualistic networks. *Proceedings of the National Academy of Sciences*, 100(16):9383–9387.
- 376 Bascompte, J., Jordano, P., and Olesen, J. M. (2006). Asymmetric coevolutionary networks facilitate
377 biodiversity maintenance. *Science*, 312(5772):431–433.
- 378 Bronstein, J. L., Alarcón, R., and Geber, M. (2006). The evolution of plant–insect mutualisms. *New*
379 *Phytologist*, 172(3):412–428.
- 380 Caraballo-Ortiz, M. A., Santiago-Valentín, E., and Carlo, T. A. (2011). Flower number and distance
381 to neighbours affect the fecundity of *Goetzea elegans* (Solanaceae). *Journal of Tropical Ecology*,
382 27(5):521–528.
- 383 Červenková, Z. and Münzbergová, Z. (2014). Pollen limitation and pollinator preferences in *Scorzonera*
384 *hispanica*. *Plant Biology*, 16(5):967–972.
- 385 de Jong, T. J., Waser, N. M., and Klinkhamer, P. G. (1993). Geitonogamy: the neglected side of selfing.
386 *Trends in Ecology & Evolution*, 8(9):321–325.
- 387 Dupont, Y. L., Trøjelsgaard, K., Hagen, M., Henriksen, M. V., Olesen, J. M., Pedersen, N. M., and
388 Kissling, W. D. (2014). Spatial structure of an individual-based plant–pollinator network. *Oikos*,
389 123(11):1301–1310.
- 390 Ehrlén, J., Käck, S., and Ågren, J. (2002). Pollen limitation, seed predation and scape length in *Primula*
391 *farinosa*. *Oikos*, 97(1):45–51.
- 392 Espíndola, A., Pellissier, L., and Alvarez, N. (2011). Variation in the proportion of flower visitors of *Arum*
393 *maculatum* along its distributional range in relation with community-based climatic niche analyses.
394 *Oikos*, 120(5):728–734.
- 395 Faulkner, G. (1976). Honeybee behaviour as affected by plant height and flower colour in brussels sprouts.
396 *Journal of Apicultural Research*, 15(1):15–18.
- 397 Gerber, M. A. (1985). The relationship of plant size to self-pollination in *Mertensia ciliata*. *Ecology*,
398 66(3):762–772.
- 399 Grant, V. and Grant, K. A. (1965). *Flower pollination in the Phlox family*. Columbia University Press,
400 New York, NY, USA.
- 401 Gumbert, A. and Kunze, J. (1999). Inflorescence height affects visitation behavior of bees—a case study of
402 an aquatic plant community in Bolivia. *Biotropica*, 31(3):466–477.
- 403 Hoehn, P., Tschartke, T., Tylianakis, J. M., and Steffan-Dewenter, I. (2008). Functional group diversity
404 of bee pollinators increases crop yield. *Proceedings of the Royal Society of London B: Biological*
405 *Sciences*, 275(1648):2283–2291.
- 406 Jiang, X.-F. and Li, Q.-J. (2017). Self-and intra-morph incompatibility and selection analysis of an
407 inconspicuous distylous herb growing on the Tibetan plateau (*Primula tibetica*). *Ecology and Evolution*,
408 7:5746–5753.
- 409 Junker, R. R., Blüthgen, N., Brehm, T., Binkenstein, J., Paulus, J., Schaefer, H. M., and Stang, M. (2013).
410 Specialization on traits as basis for the niche-breadth of flower visitors and as structuring mechanism
411 of ecological networks. *Functional Ecology*, 27(2):329–341.
- 412 King, C., Ballantyne, G., and Willmer, P. G. (2013). Why flower visitation is a poor proxy for pollination:
413 measuring single-visit pollen deposition, with implications for pollination networks and conservation.
414 *Methods in Ecology and Evolution*, 4(9):811–818.
- 415 Lande, R. and Arnold, S. J. (1983). The measurement of selection on correlated characters. *Evolution*,
416 37(6):1210–1226.
- 417 Levin, D. A. and Kerster, H. W. (1973). Assortative pollination for stature in *Lythrum salicaria*. *Evolution*,
418 27(1):144–152.
- 419 Morris, W., Price, M., Waser, N., Thomson, J., Thomson, B., and Stratton, D. (1994). Systematic increase
420 in pollen carryover and its consequences for geitonogamy in plant populations. *Oikos*, 71(3):431–440.
- 421 Nagamitsu, T., Momose, K., Inoue, T., and Roubik, D. W. (1999). Preference in flower visits and
422 partitioning in pollen diets of stingless bees in an Asian tropical rain forest. *Researches on Population*
423 *Ecology*, 41(2):195–202.
- 424 Newman, E., Manning, J., and Anderson, B. (2015). Local adaptation: Mechanical fit between floral
425 ecotypes of *Nerine humilis* (Amaryllidaceae) and pollinator communities. *Evolution*, 69(9):2262–2275.
- 426 Ohashi, K. and Yahara, T. (1998). Effects of variation in flower number on pollinator visits in *Cirsium*

- 427 *purpuratum* (Asteraceae). *American Journal of Botany*, 85(2):219–219.
- 428 Olesen, J. M., Bascompte, J., Elberling, H., and Jordano, P. (2008). Temporal dynamics in a pollination
429 network. *Ecology*, 89(6):1573–1582.
- 430 Paton, D. C. (1993). Honeybees in the Australian environment. *Bioscience*, 43(2):95–103.
- 431 Peakall, R. and Handel, S. N. (1993). Pollinators discriminate among floral heights of a sexually deceptive
432 orchid: implications for selection. *Evolution*, 47(6):1681–1687.
- 433 Pyke, G. H. (1978). Optimal foraging: movement patterns of bumblebees between inflorescences.
434 *Theoretical Population Biology*, 13(1):72–98.
- 435 R Core Team (2015). *R: A Language and Environment for Statistical Computing*. R Foundation for
436 Statistical Computing, Vienna, Austria.
- 437 Ramalho, M. (2004). Stingless bees and mass flowering trees in the canopy of atlantic forest: a tight
438 relationship. *Acta Botanica Brasilica*, 18(1):37–47.
- 439 Roubik, D. W. (1993). Tropical pollinators in the canopy and understory: field data and theory for stratum
440 “preferences”. *Journal of Insect Behavior*, 6(6):659–673.
- 441 Ruane, L. G., Hancock, L. M., Rotzin, A. T., and Luce, C. N. (2013). Pollen viability and the potential
442 for self-pollen interference in *Phlox hirsuta*, an endangered species. *International Journal of Plant
443 Sciences*, 174(9):1251–1258.
- 444 Ruane, L. G., Rotzin, A. T., and Congleton, P. H. (2014). Floral display size, conspecific density and
445 florivory affect fruit set in natural populations of *Phlox hirsuta*, an endangered species. *Annals of
446 Botany*, 113(5):887–893.
- 447 Schlinkert, H., Westphal, C., Clough, Y., Grass, I., Helmerichs, J., and Tschardt, T. (2016). Plant
448 size affects mutualistic and antagonistic interactions and reproductive success across 21 Brassicaceae
449 species. *Ecosphere*, 7(12):e01529.
- 450 Sletvold, N. and Grindeland, J. M. (2008). Floral herbivory increases with inflorescence size and local
451 plant density in *Digitalis purpurea*. *Acta Oecologica*, 34(1):21–25.
- 452 Sletvold, N., Grindeland, J. M., and Ågren, J. (2010). Pollinator-mediated selection on floral display,
453 spur length and flowering phenology in the deceptive orchid *Dactylorhiza lapponica*. *New Phytologist*,
454 188(2):385–392.
- 455 Sletvold, N., Grindeland, J. M., and Ågren, J. (2013). Vegetation context influences the strength and
456 targets of pollinator-mediated selection in a deceptive orchid. *Ecology*, 94(6):1236–1242.
- 457 Suchan, T. and Alvarez, N. (2015). Fifty years after Ehrlich and Raven, is there support for plant–insect
458 coevolution as a major driver of species diversification? *Entomologia Experimentalis et Applicata*,
459 157(1):98–112.
- 460 Thomson, J. D. and Plowright, R. (1980). Pollen carryover, nectar rewards, and pollinator behavior with
461 special reference to *Diervilla lonicera*. *Oecologia*, 46(1):68–74.
- 462 Torång, P., Ehrlén, J., and Ågren, J. (2006). Facilitation in an insect-pollinated herb with a floral display
463 dimorphism. *Ecology*, 87(8):2113–2117.
- 464 Trunschke, J., Sletvold, N., and Ågren, J. (2017). Interaction intensity and pollinator-mediated selection.
465 *New Phytologist*, 214(3):1381–1389.
- 466 Ulyshen, M. D., Soon, V., and Hanula, J. L. (2010). On the vertical distribution of bees in a temperate
467 deciduous forest. *Insect Conservation and Diversity*, 3(3):222–228.
- 468 Waser, N. M. and Price, M. V. (1991). Reproductive costs of self-pollination in *Ipomopsis aggregata*
469 (Polemoniaceae): are ovules usurped? *American Journal of Botany*, 78(8):1036–1043.
- 470 Wood, S. (2006). *Generalized Additive Models: An Introduction with R*. CRC press.




Figure 1. Inflorescence visitation in *Centaurea scabiosa* and *Inula salicina*. Visitation of inflorescences of *Centaurea scabiosa* and *Inula salicina* significantly depended on the vertical position of inflorescences above ground. This relationship was modified by the height of the surrounding vegetation as apparent from the comparison of data from transects surrounded by short (A. and B.) and tall (C. and D.) vegetation. A small amount of noise was added to the data in both x and y direction to make overlapping points visible. Summary of the statistical tests is shown in Table 1.




Figure 2. Inflorescence visitation of *Centaurea scabiosa* by the most frequent visitor taxa. The number of visits per 30 minutes in short (A. - D.) and tall (E. - H.) vegetation is plotted. A small amount of noise was added to the data in both x and y direction to make overlapping points visible. The relationship in *Bombus terrestris* (B.) is not statistically significant. Summary of the statistical tests is shown in Table 1.




Figure 3. Inflorescence visitation of *Inula salicina* by the most frequent visitor taxa. The number of visits per 30 minutes in short (A. and B.) and tall (C. and D.) vegetation is plotted. A small amount of noise was added to the data in both x and y direction to make overlapping points visible. Summary of the statistical tests is shown in Table 1.




Figure 4. Changes in relative visitation by different insects depending on inflorescence height and vegetation height. Proportions of flower visits attributed to main groups of flower visitors of *Centaurea scabiosa* and *Inula salicina* in transects surrounded by short (A. and B.) and tall (C. and D.) vegetation. The relationships were estimated using generalised additive models. A small fraction of visitors belonged to other groups omitted from the analysis because they had very low abundance. Relationships which were not statistically significant are shown in dashed lines. Summary of the statistical tests is shown in Table 2.




Figure 5. The effect of inflorescence height on visitation of *Salvia verticillata*. The number of visitors per inflorescence (A.) and the per-flower visitation rate (B.) in inflorescences of *Salvia verticillata* at different heights within the range of heights found naturally at the study site. The inflorescence height in cm refers to the top flower in each inflorescence.




Figure 6. The effect of inflorescence height on visitation of *Salvia verticillata* by different insects. The inflorescence height in cm refers to the top flower in each inflorescence. The relationships are statistically significant except in *Bombus sylvarum* (C.). Summary of the statistical tests is provided in Table 3.




Figure 7. Different insects varied in their contribution to inflorescence visitation and visited different proportions of flowers per inflorescence. (A.) Proportions of inflorescence visits attributed to main groups of visitors of *Salvia verticillata* changed significantly in relation to inflorescence height. The relationships are statistically significant except in *Bombus lapidarius*. Summary of the statistical tests is provided in Table 3. (B.) Different groups of insects differed in the proportion of flowers visited during a visit to an inflorescence of *Salvia verticillata*. The box and whiskers plot shows the median (horizontal line), interquartile range (box), and $1.5 * SD$ (whiskers) for each visitor group.




Figure 8. Seed set of *Salvia verticillata*. Seed set of individual ramets of *Salvia verticillata* increased with the number of flowers in the inflorescence (A.) and with the inflorescence height (B.). Dependence of seed set on inflorescence height could be explained by differences in visitation by pollinators in relation to inflorescence height (see Fig. 5). Seed set increased with visitation measured either as the number of inflorescence visits (C.) or as the number of flower visits (D.). Including either of these two measures of visitation rate rendered the direct effect of inflorescence height on seed set statistically non-significant.




Figure 9. Seed set of individual whorls within inflorescences of *Salvia verticillata*. The proportion of seeds produced relative to the maximum potential seeds set in individual whorls within inflorescences of *Salvia verticillata* increased with the vertical position of individual whorls. Within each inflorescence, the whorl closest to the ground was numbered as 1 and increasing numbers refer to successive whorls higher above ground.