

A peer-reviewed version of this preprint was published in PeerJ on 20 July 2018.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.5328) (peerj.com/articles/5328), which is the preferred citable publication unless you specifically need to cite this preprint.

Morsy K, Mohamed SA, Abdel-Ghaffar F, El-Fayoumi H, Abdel-Haleem H. 2018. *Rhabdias bufonis* (Rhabdiasidae) from the lung of the African common toad, *Amietophrynus regularis* (Bufonidae) in Egypt: new data on the basis of light and scanning electron microscopic study. PeerJ 6:e5328 <https://doi.org/10.7717/peerj.5328>

***Rhabdias bufonis* (Rhabdiasidae) from the lung of the African common toad, *Amietophrynus regularis* (Bufonidae) in Egypt. New data on the basis of light and scanning electron microscopic study**

Kareem Morsy^{1,2}, Sara Ali^{3*}, Fathy Abdel-Ghaffar², Hoda El-Fayoumi³, Heba Abdel-Haleem³

¹Department of Biology, College of Science, King Khalid University, Abha, Saudi Arabia

²Department of Zoology, Faculty of Science, Cairo University, Cairo, Egypt

³Department of Zoology, Faculty of Science, Beni Suef University, Beni Suef, Egypt

Corresponding author email: sara_aly_1989@yahoo.com

ABSTRACT

Background and Aims: *Rhabdias bufonis* (Rhabdiasidae) is a nematode parasite of family Rhabdiasidae infecting the lung of amphibians. The present study provides new morphological details for *R. bufonis* isolated from the lungs of the African common toad, *Amietophrynus regularis* based on observations of light and scanning electron microscopy.

Methods: Forty specimens were collected during the period from March to August 2017 from its natural habitat; the damp, moist fields and gardens at Giza governorate, Egypt. Nematode worms were isolated from the lungs, fixed and then preserved. They were examined using light and scanning electron microscopy with further line drawings.

Results: 14 specimens (35%) were found to harbor a large number of *Rhabdias*. Body of the gravid females were elongated, slender, measured 3.22–9.86 (5.64±0.03) long and 0.09–0.48 (0.23±0.02) mm wide at mid-length. Buccal capsule was with cylindrical lumen and sclerotised walls 13–31 (18 ± 2) long and 16–27 (24 ± 2) mm wide. Ovaries were almost straight. The uteri were located anterior and posterior to the vulva. Uterus filled with large number of eggs (17–42). Eggs were oval in shape, 0.120–0.132 (0.126±0.002) × 0.039–0.812 (0.517±0.02) mm and some of them were with fully developed larvae inside. The tail was comparatively short, gradually tapered, tail length 0.131–0.435 (0.320±0.02) mm long. SEM showed that worms possessed an oval anterior end with a simple, slit like oral opening. The lipless edges of mouth opening bordered with tiny cuticular elevations, radiated outwards. Two pairs of submedian cephalic papillae were observed around the oral opening as well as two amphids.

Conclusion: The current study introduced new morphological details for *R. bufonis* isolated from the Egyptian frog, which was not reported in the previous studies, of them, the morphology of the slit-like mouth opening, the two pairs of cephalic papillae, two amphids and the three pairs of cuticular elevations supporting the area around mouth opening.

Key words: *Rhabdias bufonis*, Nematoda, *Amietophrynus regularis*, Bufonidae, cuticular elevations, cephalic papillae, morphology, morphometry

INTRODUCTION

Studies on the helminth fauna of amphibians have received much attention in the recent years (Dusen & Oguz, 2010; Akani et al., 2011; Santos et al., 2013). The recent interest in amphibian parasites stems from the declining in amphibian populations (Johnson et al., 2001; Ibrahim, 2008). Nematodes of the genus *Rhabdias* Stiles et Hassall (1905) are a large group of lung-dwelling parasites and comprises about 80 nominal species of the nematodes parasitic in amphibians and reptiles (Kuzmin & Tkach, 2017). Morphologically they are females; however, hermaphroditism has been reported for some species of the genus, based on testis zones and sperm cells in the ovaries (Runey, Runey & Lauter, 1978). The life cycles of these nematodes alternate between parasitic and dioecious free-living generations. Parasitic species of *Rhabdias* are infecting the lungs of amphibians and reptiles (Anderson, 2000; Kuzmin, 2013; Tkach, Kuzmin & Snyder, 2014). Of about 39 species of *Rhabdias* parasitizing anuran hosts, eight species of them of four families have been reported from Afro tropical region (Junker et al., 2010). Twenty four species of *Rhabdias* are known from Africa (Kuzmin and Tkach 2017). Of them, 13 are parasitic in Chamaeleonidae lizards while 11 are parasites of amphibians. In Egypt, three different *Rhabdias* species were reported from amphibian hosts, *Rhabdias bufonis* Schrank (1788) (Moravec, Barus & Rysavy, 1987), *Rhabdias aegyptiaca* El-Garhy & Garo (2006) and *Rhabdias* sp. (left unnamed) from the lungs of maculated toad *Amietophrynus regularis* at Aswan governorate (Saad, Khalifa & Mostafa., 2009). Due to their similarity in the morphological characteristics such as the labial structures, the position of vulva , the tail shape and absence of males in parasitic generations, the differentiation between species belonging to the genus *Rhabdias* became complicated (Tkach, Kuzmin & Pulis, 2006; Kuzmin,Tkach & Brooks, 2007; Saad, Khalifa & Mostafa., 2009). However, the morphology of the anterior end of some *Rhabdias* species yields some characters appropriate for the species diagnostics (Kuzmin, Tkach &

Vaughan, 2005). Tkach, Kuzmin & Pulis (2006); Kuzmin , Tkach & Brooks (2007) and Martínez–Salazar, Pérez–Ponce de León & Parra–Olea (2009) suggested that, additional taxonomical tools like; molecular biology techniques, scanning electron microscopy, host specificity and geographic distribution are required to diagnose species of this genus. The African common toad, *A. regularis* Reuss (1833) is widespread in the Sub–Saharan Africa, with its range extending to the oases in Algeria and Libya as well as to northern Nilotic Egypt (Frost, 2014). According to Rödel (2000) and Ibrahim (2008) the prey of *A. regularis* often includes beetles, ants, bugs, insects, grubs, slugs, worms and other invertebrates. The present study provides new morphological details for the nematode *Rhabdias bufonis* isolated from the lungs of the African common toad, *A. regularis* by light and scanning electron microscopy.

MATERIALS & METHODS

Forty specimens of the African common toad, *Amietophrynus regularis* were collected by hand or noose from its natural habitat; the damp, moist fields and gardens at Giza governorate, Egypt (30° 2' 16.6164" N and 31° 12' 29.7360" E) from March to August 2017. Animals were subjected to euthanasia using 20% benzocaine gel (Anbesol, Pfizer, Inc., New York). Each specimen was subsequently necropsied and all organs were examined searching for helminthes using a ZEISS Compact Greenough stereomicroscope (Model Stemi 305). Where necessary, all animal procedures were carried out according to the regulatory laws regarding experimental Animal Ethics Committee, Faculty of Science, Beni–Suef University, Beni–Suef, Egypt (Ethical Approval Number: 2015/10). Nematode worms were isolated from the lungs, heat fixed in 10% neutral buffered formalin for 15 min and then preserved in 70% ethanol in 5% glycerol solution to avoid sudden drying. Finally, samples were transferred to lactophenol for clearance. The prepared samples were examined using differential interference contrast (DIC) light microscopy with digital

image analysis system (analysis auto 5.0). Drawings were made with the aid of a camera lucida. Measurements were in millimeters unless otherwise stated. For scanning electron microscopy, samples were fixed in 4% glutaraldehyde in 0.1 M sodium cacodylate buffer, washed in the same buffer, and dehydrated in a graded alcohol series (50%, 60%, 70%, 80%, 90% and 100%). Samples were then processed in a critical point drier “Bomer-900” with freon 13, sputter-coated with gold-palladium in a Technics Hummer V, and finally examined with a Jeol scanning electron microscope (Model JSM7610F).

RESULTS

Morphology of *Rhabdias bufonis* (Schränk, 1788)

Description based on 28 gravid Adults; morphometric data presented as a range followed by the mean \pm SD in parentheses.

Light microscopy

Body of the gravid females were elongated, slender (Fig.1A, B, 3A) measured 3.22–9.86 (5.64 \pm 0.03) long and 0.09–0.48 (0.23 \pm 0.02) wide at mid-length. Cuticle inflation and transverse striations (Fig. 1C) were observed, inflation widened from anterior end to level of esophagus then rapidly narrowed towards caudal region with pronounced inflation. Buccal capsule was with cylindrical lumen and sclerotised walls (Fig. 2B, C), 13–31 (18 \pm 2) long and 16–27 (24 \pm 2) wide. The anterior part of esophagus contacting posterior end of buccal capsule, esophagus was muscular, 0.27–0.63 (0.18 \pm 0.02) long and 0.25–0.5 (0.30 \pm 0.03) as a maximum width with muscular anterior third and elongated posterior bulb (Fig. 1B, D). Nerve ring surrounding esophagus and measured 0.168–0.240 from the anterior end (Fig. 3A). Anterior end of intestine connected to the esophageal bulb, it was widened at the esophageo-intestinal junction. The content of the intestine was black at their posterior part. Reproductive system was amphidephlic. Ovaries were almost straight. The uteri were located anterior and posterior to the vulva. Vulva was nearly equatorial

1.47–5.98 from the anterior end. Uterus filled with large number of eggs (17-42). Eggs were oval in shape (Fig. 1G, 3D), $0.120\text{--}0.132$ (0.126 ± 0.002) \times $0.039\text{--}0.812$ (0.517 ± 0.02) in size and some of them were with fully developed larvae inside (Fig. 3D). The tail was comparatively short, gradually tapered, tail length $0.131\text{--}0.435$ (0.320 ± 0.02) long (Figs. 1E, 2E, 3E).

Scanning electron microscopy

SEM showed that the nematode isolated in the present study possessed an oval anterior end (Figs. 2A, B, 3B) with a simple, slit like oral opening. The lipless edges of mouth opening bordered with tiny cuticular elevations, radiated outwards (Figs. 2B, C, 3B).

Two pairs of submedian cephalic papillae (two dorso-lateral and two ventro-lateral) were observed around the oral opening as well as two amphids (Figs. 2A& B, 3B). The transverse cuticle striations (Fig. 2D) and the pointed tail (Fig. 2E, 3E) were prominent by SEM.

TAXONOMIC SUMMARY

Species: Rhabdias bufonis (Schrunk, 1788)

Host: Amietophrynus regularis Reuss, 1833 (Family: Bufonidae)

Site of infection: Lungs

Locality: Giza governorate, Egypt

Prevalence: 14 out of 40 (35%) hosts were infected with an intensity of 3-5 worms per one host.

Deposition: Permanent slides as well as 70 % preserved gravid females were deposited at the Parasitology division, Zoology department museum, Faculty of Science, Cairo University, Egypt.

DISCUSSION

The differentiation among species belonging to the genus *Rhabdias* is often complicated due to their high morphological uniformity (Chu, 1936; Baker, 1978; Tkach, Kuzmin &

Pulis, 2006; Kuzmin, Tkach & Brooks, 2007). Within the species of *Rhabdias*, the shape of cephalic end represents an important character for the differentiation between species and genera of the family Rhabdiasidae (Railliet, 1915) and yields characters suitable for species diagnostics (Kuzmin, Tkach & Vaughan, 2005). Genus *Rhabdias* can be divided into three groups based on the morphology of their head end according to Baker (1978): species without lips; with six lips and with two lateral pseudolabia. Three different *Rhabdias* species were reported from amphibian hosts belonging to family Bufonidae, represented by a single genus, *Amietophrynus* (formerly included in *Bufo*), in the Afrotropical region. These were *R. bufonis* Moravec, Baruš & Ryšavý (1987), *R. picardiae* Junker et al. (2010) and *R. africanus* Kuzmin (2001). By comparing the recovered parasite with different species of the same genus previously recorded in the Afrotropical region, it was found that the present species is morphometrically similar to *Rhabdias bufonis* described previously from the same host by Moravec, Baruš & Ryšavý (1987) while many characters are different to those of the comparable species. Both species resemble each other by possessing similar head structure (absence of lips, presence of small submedian mouth papillae); the intestinal apex is broader than the esophageal base and the equatorial position of vulva. *Rhabdias bufonis* recorded in the present study has a longer body (9.86 mm vs 8.35), a shorter esophagus (270–630 vs 690–790) and a longer buccal capsule (13–41 mm vs 8–10) than *R. picardiae* and differ from *R. africanus* in the body size and the absence of two lateral pseudolabia. The present form, differs from the description of *R. bufonis* given by Travassos (1930) and Hartwich (1972) in the body length (3.22–9.86 vs 9–12) and in the size of buccal capsule which is of diagnostic importance (Lhermitte-Vallarino et al., 2008, Junker et al., 2010). The current study introduced new morphological characteristics for the nematode *Rhabdias bufonis* isolated from the lung of *Amietophrynus regularis* based on scanning electron microscopy. One of the most important features of *Rhabdias bufonis* is the cephalic end structure which has been

overlooked by previous studies. In Egypt, only three different species of the genus *Rhabdias* have been described based on the light microscopic studies and these species are: *R. bufonis* Moravec, Baruš & Ryšavý (1987), *R. aegyptiaca* El-Garhy & Garo (2006) and *Rhabdias* sp. Saad, Khalifa & Mostafa (2009). The parasite recovered in the present study differs from *Rhabdias aegyptiaca* where the later possesses six lips with unique arrangement around mouth opening, cuticle lined pores on its lateral surface and the size of the buccal capsule which is of diagnostic importance (Lhermitte–Vallarino et al., 2008; Junker et al., 2010). Also, it differs from *Rhabdias* sp. reported by Saad, Khalifa & Mostafa, 2009, where it has four developed lips and two lateral teeth on the mouth opening, the cuticular swelling that covers the whole body and the post–equatorial position of vulva. *Rhabdias* species that were recovered from a bufonid member with a sub–Saharan distribution in Egypt were compared in Table (1).

CONCLUSION

According to the current study that depended on light and scanning electron microscopy, it is concluded that there are new data were obtained and recorded for the nematode, *Rhabdias bufonis* infecting the African common toad which is the second study reported for this parasite in Egypt representing an important reference for upcoming studies.

ACKNOWLEDGEMENTS

This work was kindly supported by Zoology Departments, Faculties of Science, Cairo and Beni – Suef Universities. And we thank anonymous reviewers for reading the manuscript.

REFERENCES

- Akani GC, Luiselli L, Amuzie CC, Wokem GN. 2011.** Helminth community structure and diet of three afro topical anuran species: a test of the interactive–versus–isolationist parasite communities’ hypothesis. *Web ecology* 11:11–19.
- Anderson RC. 2000.** Nematode Parasites of Vertebrates, Their development and transmission. 2nd edn. 650 pp. CABI Publishing, Wallingford.

- Baker MR. 1978.** Morphology and taxonomy of *Rhabdias* spp. (Nematoda: Rhabdiasidae) from reptiles and amphibians of southern Ontario. *Canadian Journal of Zoology* 56: 2127–2141.
- Chu T. 1936.** A review of the status of the reptilian nematodes of the genus *Rhabdias* with a description of *Rhabdias fuscovenosa* var. *catanensis* (Rizzo, 1902) new rank. *Journal of Parasitology* 22:130–139.
- Dusen S, Oguz MC. 2010.** Metazoan endoparasites of three species of anurans collected from the middle black sea region of Turkey. *Helminthologia* 47: 226 – 232.
- El-Garhy M, Garo K. 2006.** *Rhabdias aegyptiaca* sp. n. (Nematoda: Rhabdiasidae) from the lungs of *Bufo regularis* (Amphibia: Bufonidae), in Egypt: Light and scanning electron microscopic study. *Journal of Union Arab Biology* 26: 127–137.
- Frost DR. 2014.** *Amietophrynus regularis* (Reuss, 1833). Amphibian species of the world: an Online Reference. Version 6.0. American Museum of Natural History, New York, 10/2014.
- Hartwich G. 1972.** Über *Rhabdias bufonis* (Schränk, 1788) und die Abtrennung von *Rhabdias dossei* nov. spec. (Nematoda, Rhabdiasidae). *Mitteilungen aus dem Zoologischen Museum in Berlin* 48: 401–414.
- Ibrahim MMI. 2008.** Helminth infra communities of the maculated toad *Amietophrynus regularis* (Anura: Bufonidae) from Ismailia, Egypt. *Diseases of Aquatic Organisms* 82: 19–26.
- Johnson PTJ, Lunde KB, Haight RW, Bowerman J, Blaustein AR. 2001.** *Ribeiroia ondatrae* (Trematoda: Digenea) infection induces severe limb malformations in western toads (*Amietophrynus boreas*). *Canadian Journal of Zoology* 79:370–379.
- Junker K, Lhermitte–Vallarino N, Barbuto M, Ineich I, Wanji S, Bain O. 2010.** New species of *Rhabdias* (Rhabdiasidae: Nematoda) from Afro tropical anurans, including molecular evidence and notes on biology. *Folia Parasitologica* 57: 47–61.

- Kuzmin YI. 2001.** *Rhabdias africanus* sp. nov. (Nematoda, Rhabdiasidae), a new nematode species from the South African toads (Amphibia, Bufonidae). *Acta Parasitologica* 46: 148–150.
- Kuzmin YI. 2013.** Review of Rhabdiasidae (Nematoda) from the Holarctic. *Zootaxa* 3639:1–76.
- Kuzmin Y, Tkach VV. 2017.** Rhabdias. World Wide Web electronic publication. <http://izan.kiev.ua/ppages/rhabdias/list.htm> (Last accessed on 24/08/2017)
- Kuzmin Y, Tkach VV, Brooks DR. 2007.** Two new species of *Rhabdias* (Nematoda: Rhabdiasidae) from the marine toad, *Bufo marinus* (Lissamphibia: Anura: Bufonidae), in Central America. *Journal of Parasitology* 93:159–165.
- Kuzmin Y, Tkach VV, Vaughan JA. 2005.** *Rhabdias kongmongthaensis* sp. n. (Nematoda: Rhabdiasidae) from *Polypedates leucomystax* (Amphibia: Anura: Rhacophoridae) in Thailand. *Folia Parasitologica* 52: 339–342.
- Lhermitte–Vallarino N, Barbuto M, Ineich I, Wanji S, Lebreton M, Chirio L, Bain O. 2008.** First report of *Rhabdias* (Nematoda: Rhabdiasoidea) from lungs of montane chameleons in Cameroon: description of two new species and notes on biology. *Parasite* 15: 553–564.
- Martínez–Salazar EA, Pérez–Ponce de León G, Parra Olea G. 2009. First record of the genus *Rhabdias* (Nematoda: Rhabdiasidae), endoparasite from *Scinax staufferi* (Anura: Hylidae) in Mexico. *Revista Mexicana de Biodiversidad* 80:861–865.
- Moravec F, Baruš V, Ryšavý B. 1987.** Some parasitic nematodes, excluding Heterakidae and Pharyngodonidae, from amphibians and reptiles in Egypt. *Folia Parasitologica* 34: 255–267.
- Rödel MO. 2000.** Herpetofauna of West Africa, Vol I. Amphibians of the West African savanna. Edition Chimaira, Frankfurt.

- Runey WM, Runey GL, Lauter FH. 1978.** Gametogenesis and fertilization in *Rhabdias ranae* Walton 1929: I. The parasitic hermaphrodite. *Journal of Parasitology*, 64: 1008–1014.
- Saad AI, Khalifa R, Mostafa N. 2009.** A lungworm nematode from amphibian hosts at Aswan Governorate, Egypt. *World Journal of Zoology* 4: 37–41.
- Santos VGT, Amato SB, Márcio Borges M. 2013.** Community structure of helminth parasites of the “Cururu” toad, *Rhinella icterica* (Anura: Bufonidae) from southern Brazil. *Parasitology Research* 112: 1097–1103.
- Tkach VV, Kuzmin Y, Pulis EE. 2006.** A new species of *Rhabdias* from lungs of the wood frog, *Rana sylvatica*, in North America: The last sibling of *Rhabdias ranae*?. *Journal of Parasitology* 92: 631–636.
- Tkach VV, Kuzmin Y, Snyder SD. 2014.** Molecular insight into systematics, host associations, life cycles and geographic distribution of the nematode family Rhabdiasidae. *International Journal for Parasitology* 44: 273–284.
- Travassos L. 1930. Pesquisas helminthologicas realizadas em Hamburgo. VII. Notas sobre os Rhabdiasoidea Railliet, 1916 (Nematoda). *Memórias do Instituto Oswaldo Cruz* 24: 161–181.




Figure 1 Light micrographs of *Rhabdias bufonis* showing: A. Females, left and right lateral views, scale bar 100 m. B. Anterior part, lateral view; OE, esophagus; UT, Uterus, scale bar 0.25 m. C. Transverse striations (TS) of cuticle, scale bar 0.25 m. D. Magnified anterior end, scale bar, 50 m. E. Tail region, scale bar 50 m. F. Anus (A), scale bar 0.25 m. G. Eggs (E), scale bar 0.25 m .




Figure 2 Scanning electron micrographs showing: A, B. Cephalic end, lateral (A) and Apical (B) views, four papillae (PA) and two amphids (AM) surrounding mouth opening (MO), scale bars 10 m. C. Magnified apical view for the cuticular inflation (CI), scale bars 100 m. D. Transverse striations (TS) of cuticle, scale bar 10 m. E. Tail end, scale bar 0.50 m.




Figure 3 *Rhabdias bufonis*, parasite of *Amietophrynus regularis*. A. Anterior part, lateral view, scale bar 50 m. B. Cephalic end, lateral view, scale bar 10 m. C. Loop of anterior genital tube, scale bar 50 m. D. Egg, lateral view, scale bar 0.25 m. E. Tail end, lateral view, scale bar 50 m. PA, Cephalic papillae; CI, Cuticular inflation; AM, Amphids

Table 1 *Rhabdias bufonis* (female, present study) and members of the same genus previously reported in Egypt.

Species	Length	Width	Buccal capsule	Teeth	Esophagus	Vulva	Tail length	Egg	References
<i>R. bufonis</i>	2.99–13.02	0.136–0.476	0.015	Absent	0.288–0.510	Equatorial	0.144–0.420	L: 0.117–0.144 W: 0.051–0.72	Moravec <i>et al.</i> (1987)
<i>R. aegyptiaca</i>	8–10	0.3–0.5	–	Absent	550 m	Equatorial	170–200 m	66 m in length	<i>El-Garhy and Garo (2006)</i>
<i>Rhabdias</i> sp.	5.2–12.5	0.2–0.7	0.01–0.032	Two lateral teeth	0.25–0.5 (0.3)	Post–Equatorial	0.23–0.4(0.3)	L: 0.1–0.12 W: 0.06–0.08	Saad <i>et al.</i> (2009)
<i>R. bufoni</i>	3.22–9.86 (5.64±0.03)	0.09–0.48 (0.23±0.02)	L: 13–31 (18 ± 2) W: 16–27 (24 ± 2)	Absent	L: 0.27–0.63 (0.18±0.02) W: 0.25–0.5 (0.30±0.03)	Equatorial	0.131–0.435 (0.320±0.02)	L: 0.120–0.132 (0.126±0.002) W: 0.039–0.812 (0.517±0.02)	Present study

* Measurements in mm, otherwise stated.