

A peer-reviewed version of this preprint was published in PeerJ on 26 February 2015.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.788) (peerj.com/articles/788), which is the preferred citable publication unless you specifically need to cite this preprint.

Howlin J, Cirenajwis H, Lettierio B, Staaf J, Lauss M, Saal L, Borg Å, Gruvberger-Saal S, Jönsson G. 2015. Loss of CITED1, an MITF regulator, drives a phenotype switch *in vitro* and can predict clinical outcome in primary melanoma tumours. PeerJ 3:e788
<https://doi.org/10.7717/peerj.788>

Loss of CITED1, an MITF regulator, drives a phenotype switch in vitro and can predict clinical outcome in primary melanoma tumours

CITED1 is a non-DNA binding transcriptional co-regulator whose expression can distinguish the 'proliferative' from 'invasive' signature in the *phenotype-switching* model of melanoma. We have found that CITED1 expression is repressed by TGF β in addition to other 'proliferative' signature genes while the 'invasive' signature genes are upregulated. In agreement, CITED1 positively correlates with MITF expression and can discriminate the MITF-high/pigmentation tumor molecular subtype in a large cohort (120) of melanoma cell lines. Interestingly, CITED1 overexpression significantly suppressed MITF promoter activation, mRNA and protein expression levels while MITF was transiently upregulated following siRNA mediated CITED1 silencing. Conversely, MITF siRNA silencing resulted in CITED1 downregulation indicating a reciprocal relationship. Whole genome expression analysis identified a phenotype shift induced by CITED1 silencing and driven mainly by expression of MITF and a cohort of MITF target genes that were significantly altered. Concomitantly, we found changes in the cell-cycle profile that manifest as transient G1 accumulation, increased expression of CDKN1A and a reduction in cell viability. Additionally, we could predict survival outcome by classifying primary melanoma tumors using our *in vitro* derived 'CITED1-silenced' gene expression signature. We hypothesize that CITED1 acts a regulator of MITF, functioning to maintain MITF levels in a range compatible with tumourigenesis.

1 Jillian Howlin^{*1,2}, Helena Cirenajwis¹, Barbara Lettiero¹, Johan Staaf¹, Martin
2 Lauss¹, Lao Saal¹, Åke Borg¹, Sofia Gruvberger-Saal^{1§} and Göran Jönsson^{1§}

3 1. Division of Oncology-Pathology, Lund University Cancer Center/Medicon Village,
4 Building 404:B2, Scheelevägen 2, SE-223 81 Lund, Sweden
5 2. Cell and Experimental Pathology, Department of Laboratory Medicine Malmö, Lund
6 University, Sweden.
7 [§] Equal contribution

8 ***Corresponding author:**

9 Jillian Howlin, PhD

10 Division of Oncology-Pathology, Lund University Cancer Center/Medicon Village,
11 Building 404:B2, Scheelevägen 2, SE-223 81 Lund, Sweden
12 Tel: +46 (0)709106498
13 Email: jillian.howlin@med.lu.se

14 INTRODUCTION

15 CITED1 is the founding member of the CITED (CBP/p300-Interacting
16 transactivator with glutamic acid [E]/aspartic acid [D]-rich C-terminal
17 domain) family of transcriptional co-regulators and was originally cloned from
18 a differential display screen between pigmented mouse B16 melanoma cells
19 and their dedifferentiated non-pigmented derivative, B16F10s. This led to
20 speculation that CITED1 or *msg1* (melanocyte specific gene 1) as it was
21 known at that time, was involved in the process of pigmentation(Shioda et al.
22 1996). Subsequently, Nair et al. reported that stable overexpression of
23 CITED1 increased the levels of tyrosinase, dopachrome tautomerase (Dct)
24 and melanin in B16 cells, reinforcing the idea that it had a role in
25 melanogenesis(Nair et al. 2001). By 2005, as gene expression profiling
26 became relatively commonplace, CITED1 was identified in several new
27 screens of tumors and cell lines: two studies identified CITED1 as a gene
28 whose expression distinguished nevi from primary melanoma, another found
29 CITED1 to be upregulated in advanced stage melanomas in comparison to
30 benign nevi or melanoma *in situ*, while expression profiling of an *in vitro*
31 progression model identified CITED1 among a signature of genes lost in
32 aggressive melanoma lines relative to primary melanocytes in culture (Ryu
33 et al. 2007; Haqq et al. 2005; Talantov 2005; Smith et al. 2005).

34 Based on extensive gene expression profiling of melanoma cell lines *in vitro*,
35 Hoek et al. proposed the 'phenotype-switching' model of melanoma that was
36 independent of the degree of transformation or disease progression and

sought to explain the observation that melanoma cells altered between two states: those with high proliferative potential that are less invasive and those with high metastatic potential that are less proliferative. These separate but alternating states are controlled by different transcriptional programs and can be defined by specific gene signatures (Hoek, Eichhoff, et al. 2008a). MITF expression and many of its known targets (TYR, MLANA) define the 'proliferative' group, while the 'invasive' signature group is characterized by expression of negative regulators of the Wnt signaling pathway (WNT5A, DKK1, CTGF). CITED1 expression was associated with the proliferative pathway signature and subsequently confirmed in an updated and expanded data set to be significantly correlated with the proliferative phenotype ($P < 1.00E-05$, <http://www.jurmo.ch/hopp>, accessed 19 March 2013) (Hoek et al. 2006) (Widmer et al. 2012).

Studies on CITED1 suggest that it is a non-DNA binding nuclear transcriptional co-regulator capable of influencing TGF β induced transcription mediated by ligand-induced SMAD hetero-oligomerization; estrogen-dependent transcription mediated by ER α , and Wnt/ β -Catenin-dependent transcription. These effects are dependent on CITED1-CBP/P300 binding via the conserved CITED family CR2 domain and while in the case of ER α , CITED1 is thought to act by stabilizing the CBP/P300-ER α interaction, in the case of β -Catenin it acts to repress transcription by competing for binding with CBP/P300 transcriptional co-activators (Shioda et al. 1998; Yahata 2001; Yahata 2000; Plisov 2005).

60 Microphthalmia-associated transcription factor, MITF, acts as a master-
 61 regulator of melanocyte differentiation and as a result has been intensely
 62 studied in the field of melanoma research(Widlund & Fisher 2003; Levy et al.
 63 2006). It is a basic helix-loop-helix leucine zipper transcription factor that
 64 recognizes E-box and M-box sequences in the promoter regions of its target
 65 genes. Highlighting its importance in the disease, amplification of MITF locus
 66 has been found in >15% of metastatic melanomas and germline mutations in
 67 MITF that predispose carriers to melanoma development have also been
 68 found (Garraway et al. 2005; Bertolotto et al. 2011; Yokoyama et al. 2011). In
 69 melanoma cells the target genes of MITF include most notably TYR, MCIP,
 70 DCT, MLANA involved in the process of pigmentation; cell cycle regulators
 71 such as CDK2 and CDKN1A and the more recently identified BRCA1 gene that
 72 has, with other target DNA repair genes, defined a role for MITF in the DNA
 73 damage response (DDR) (Strub et al. 2011; Beuret et al. 2011; Giuliano et al.
 74 2010).

75 The regulation of MITF is complex and tightly controlled, exhibiting both
 76 transcriptional and post-translational regulation. There are several transcript
 77 isoforms, of which MITF-M is the dominant form expressed in melanocytes.
 78 Multiple signaling pathways converge on the MITF-M specific promoter that
 79 harbors binding sites for PAX3, SOX10, CREB, FOXD3, LEF-1 and BRN2 among
 80 other transcription factors (Yokoyama & Fisher n.d.; Levy et al. 2006).
 81 Additionally, the MITF target gene CDKN1A/P21 has been shown to act as

82 reciprocal transcriptional cofactor independently of its CDK inhibitor function,
83 suggesting the existence of at least one positive feedback loop (Sestáková et
84 al. 2010).

85 MITF post-translational activity can be affected by phosphorylation,
86 sumoylation, ubiquitination and by binding with proteins that block access to
87 the DNA binding domain such as PIAS3 (Yokoyama & Fisher n.d.; Levy 2001).
88 Oncogenic BRAF (but not wildtype BRAF), which is mutated in up to 50% of
89 melanomas, also regulates MITF via simultaneously stimulating MITF
90 activation through ERK phosphorylation, which leads to its subsequent
91 degradation, and by inducing transcription of MITF via BRN2 upregulation
92 (Davies et al. 2002; Wellbrock et al. 2008).

93 The consensus regarding why the cell invests such effort in maintaining
94 control of MITF levels and why there are so many regulatory mechanisms, is
95 that melanocytes and melanoma are exquisitely sensitive to even small
96 variations in MITF expression. Ultimately its activity must be sustained within
97 the narrow window permissive for continued survival and proliferation. In this
98 study, we characterise the role of CITED1 as a novel regulator of MITF in
99 melanoma.

100 MATERIALS AND METHODS

101 **Cell lines**

Cell lines were obtained from ATCC. HT144 and SKMEL3 cells were cultured in McCoy's5A supplemented with 10% and 15% fetal bovine serum (FBS), respectively. A2058, WM852 and WM239 were cultured in RPMI 1640 supplemented with 10% FBS; A375 and HMBC cells were cultured in DMEM supplemented with 10% FBS and SKMEL5 cells were cultured in MEM media supplemented with 10% FBS. Cells were grown in the presence of penicillin and streptomycin (50 I.U./mL).

Gene expression analysis

RNA was isolated (4 replicates for each treatment) using a Qiagen RNeasy Plus mini-kit and the quality determined using a Bioanalyser (Agilent). Gene expression experiments were performed using the Illumina HT12 array covering more than 47,000 transcripts and known splice variants across the human transcriptome. The raw data was quantile normalized and Illumina control probes were removed from subsequent analysis using BASE (Vallon-Christersson et al. 2009). The data were exported to MeV, log2 transformed and gene and sample centered (Al et al. 2003). SAM (significance of microarray analysis) was performed using a two-group comparison; for the siRNA experiment the groups corresponded to siNEG vs #1 & #3 siCITED1 and for the TGF β 1 experiment the groups corresponded to cells with or without TGF β 1 treatment. In both cases there was a median false discovery risk of 10 false-positive transcripts. Hierarchical clustering was performed to visualize the data. 312 probes were found to be significantly altered in the

124 siRNA experiment (208 upregulated and 104 downregulated) while 1009
125 probes were significantly altered by TGF β 1 treatment. DAVID was used to
126 assist in functional annotation of the final gene lists (Huang et al. 2007)

127 For the publically available data cited, 120 melanoma cell lines from three
128 cohorts (PMID: 17516929, 16827748, 20406975) analyzed by Affymetrix
129 gene expression microarrays were collected, individually MAS5 normalized,
130 and merged into a single cohort. Probe sets were collapsed into single genes
131 and mean-centered across the entire cohort. Data from Harbst et al. were
132 classified using nearest centroid and pearson correlation. Survival analysis
133 and multivariate cox regression methods were performed in R.

134 **Transient transfections, promoter-reporter assay and TGF β 1-** 135 **treatment**

Transient transfections were performed using
136 Lipofectamine2000 and Opti-MEM reduced serum media (Life Technologies)
137 according to the manufactures recommendations. siRNA was purchased from
138 Applied Biosystems and the notations in the text: siNEG, #1 siCITED1 and #3
139 siCITED1 correspond to the catalogue ID numbers #4390843, #s8965 and
140 #s224062 respectively. For the MITF targeting siRNA; N, siM1 and siM3
141 correspond to the catalogue ID numbers #4390843, #s8790 and #ss8792,
142 respectively. For the luciferase reporter assay, a Dual-Luciferase Reporter
143 assay system #E1910 (Promega) was used to measure relative reporter
144 activity on a FLUOstar Omega microplate reader (BMG Labtech). A375 cells
145 were transfected with a luciferase reporter construct harboring 2.3kb of the

MITF-M specific promoter in a PGL2 vector (Wellbrock et al. 2008). A pRL-Renilla Luciferase reporter vector was used as a control for each transfection. CITED1 was overexpressed using a pRc/CMV containing a N-terminal HA-tagged human CITED1 (transcript isoform 1) referred to as pCITED1 in the text. An empty CMV-promoter expression plasmid, pcDNA3.1 (+) was used as a negative control. Recombinant human transforming growth factor- β 1 (TGF β 1), #PHG9203 was purchased from Invitrogen. For the A2058 gene expression experiment, cells were exposed to either 5 or 10ng/ml TGF β 1 in serum-free media for 24 hours. In the case of the Luciferase reporter assay, cells were serum starved the day after transfection for 3 hours and exposed to 5ng/ml TGF β 1 in serum free media for 24 hours prior to harvesting.

Antibodies and Immunoblotting

The following antibodies were used: anti-CITED1, #AB15096 from Abcam; anti-MITF (C5 clone), # MA5-14146 from ThermoScientific; anti-MITF (D5 clone) from Dako, #M3621, (used in Fig. 4c); anti-CDKN1A/P21, #2947 and anti-CDKN1C/P57, #2557 were purchased from Cell Signaling Technology and anti- β -Actin (AC-15), #A5441 from Sigma-Aldrich. Cell lysates were resolved by SDS-PAGE and transferred to 0.45 μ m PVDF membranes by electroblotting. The membranes were blocked in 5% non-fat milk in TBST prior to incubation with primary antibodies diluted 2.5% non-fat milk. The blots were probed with the appropriate secondary antibodies (Pierce

Biotechnology) in 5% non-fat milk. The membranes were developed using ECL (GE Healthcare).

Cell cycle analysis

Flow cytometry was performed on a FACSCalibur (BD Biosciences) and subsequently analysed using ModFit (Verity House Software). Briefly, following transfection, confluent cells were detached, washed in 1XPBS and fixed in 70% ethanol. Prior to analysis they were stained with a propidium iodide solution and a 20G syringe was used to obtain a homogenous single cell solution. All events were saved (up to 20,000 events per replicate) ungated, using BD Cell Quest and the data exported to ModFit where following selection of the appropriate ploidy status, a standard auto-analysis fit using autolinerarity was performed. We found that a 2-cycle aneuploid-dip/tetraploid was appropriate for HT144 and A2058 while 1-cycle diploid was suitable for A375.

Alamar Blue Assay

The Alamar blue assay reagent was purchased from Invitrogen and used according to the manufactures' instructions. Briefly, following transfection cells were seeded into 96-well plates at 5000cells/well. In each experiment, for each of the treatments i.e.: siNEG, #1 siCITED1 and #3 siCITED1, 8 wells spread over 3 rows were used. At the indicated time points (4, 72, 96 and 120 hours post-transfection), Alamar blue was added and the cells incubated at 37°C for 2 hours. Fluorescence was measured (544nm) on a FLUOstar

189 Omeaga microplate reader (BMG Labtech). The values obtained at the 4-hour
190 time point were used to normalize the fluorescence readings to account for
191 any initial cell counting error. Cells were also seeded in parallel for Western
192 blot analysis (72, 96, 120 hours) to ensure successful CITED1 down
193 regulation.

194 **Droplet digital PCR**

195 RNA was isolated from cells using a Qiagen RNeasy Plus mini-kit and
196 quantified using a Nanodrop spectrophotometer (ThermoScientific). cDNA
197 was generated from 50-100ng total RNAs using 'iScript Advanced cDNA
198 synthesis for RT-qPCR' (Bio-Rad). Bio-RAD's 'ddPCR Supermix for Probes' was
199 then used with predesigned TaqMan gene expression assays (Applied
200 Biosystems) consisting of specific primers and FAM labelled probes for MITF
201 (#Hs01117294_m1), MITF-M isoform specific transcript (Hs00165165_m1)*,
202 CITED1 (#Hs00918445_g1) and IPO8 (#Hs00183533_m1). (*There appeared
203 to be no advantage in using the MITF-M isoform specific transcript over the
204 MITF probe that could measure multiple isoforms). A manual cut-off for
205 positive/negative droplets was selected using the Biorad QuantaSoft™ data
206 analysis suite to calculate the relative copies/μl of each transcript.

207 **RESULTS**

208 **TGF β induces expression of the invasive signature genes while**
209 **suppressing a cohort of proliferative signature genes including**
210 **CITED1**

211 Hoek et al. noted that many of the genes that defined the invasive
212 phenotype were commonly TGF β -driven while at the same time only the
213 proliferative signature phenotype cells were sensitive to TGF β growth
214 inhibition *in vitro* (Hoek et al. 2006). That MITF levels increase and
215 invasiveness is enhanced in response to TGF β stimulation was also confirmed
216 subsequently (Pierrat et al. 2012; Pinner et al. 2009). In agreement, we
217 showed that the melanoma cell line A2058 upregulates WNT5A in response to
218 TGF β exposure and that exogenous Wnt-5a in turn increased their invasive
219 potential (Jenei et al. 2009). For the present study, in an effort to examine
220 what other phenotype specifying genes were directly regulated by TGF β , we
221 performed gene expression analysis and found TGF β treatment resulted in
222 both upregulation of invasive signature genes and suppression of genes
223 characterizing the proliferative phenotype (Fig. 1a). The effect is most
224 pronounced if only those signature genes that were deemed significantly
225 altered by TGF β treatment are examined. The original signature set defined
226 by Hoek et al., was redefined as more public datasets became available and
227 has a slightly different but overlapping gene profile based on the top ranked
228 differentially expressed genes (Fig. 1b). Both MITF and CITED1 are in the
229 proliferative cohort and their response to TGF β treatment was confirmed at
230 protein level in A2058 cells (Fig. 1C).

CITED1 expression positively correlates with the expression of MITF

Examination of publically available gene expression data on 120 melanoma cell lines demonstrated a consistent positive correlation between CITED1 and MITF expression ($r=0.6543$). Each cell line was assigned as either 'proliferative' or 'invasive' based on a score derived from the averaged expression values of the approximately 50 genes in each defining signature set that had matching gene symbols in our data (Fig. 2a). We also confirmed the correlation in cell lines derived from our own lab (Fig. S1). This was important as inconsistency in interlaboratory phenotype signatures has previously been reported (Widmer et al. 2012). We could additionally confirm expression at the protein level (Fig. 2b)

Gene expression analysis reveals CITED1 silencing can induce a phenotype-switch

To investigate the function of CITED1 in melanoma, we transiently downregulated its expression using CITED1 targeting siRNA. We choose the HT144 cell line as it had a relatively high level of detectable CITED1 mRNA and protein expression. A scatterplot of the 120 cell lines assigned as either 'proliferative' or 'invasive' based on the maximum matching gene signature score demonstrates the shift in phenotype that occurs following CITED1 downregulation (Fig. 3a,b). A heatmap of the expression profiles clearly illustrates that the shift is due to a general induction of the 'proliferative' and suppression of the 'invasive' cohort (Fig. 3c). It was apparent that the #3 siCITED1 siRNA was not as effective at switching the cells as the #1

254 siCITED1, this was observed consistently throughout our experiments and
255 may be due to the fact that #3 siCITED1 was not as successful at silencing
256 CITED1 (Fig. 3b, *inset*).

257 **CITED1 is a reciprocal regulator of MITF and impacts MITF target** 258 **gene expression**

259 A heatmap of only the significantly differentially induced transcripts between
260 siNEG and both #1 & #3 siCITED1 highlights the identity of the genes
261 responsible for the expression switch (Fig. 4a). Of most relevance, we found
262 MITF, a known driver of the proliferative phenotype switch and many of its
263 previously known downstream targets, these also included genes categorized
264 by Gene Ontology annotation (GO) as related to pigmentation and UV/DNA
265 damage response (Fig. 4a) (Hoek, Schlegel, et al. 2008b; McGill et al. 2006;
266 Sánchez-Martín et al. 2002; Strub et al. 2011). We could confirm that indeed
267 MITF protein levels were affected by siCITED1 in HT144 cells and that
268 conversely, overexpression of CITED1 in A2058 cells, resulted in
269 downregulation of MITF (Fig. 4b). Strub et al. identified a large number of
270 genomic targets of MITF by ChIP-seq analysis (Strub et al. 2011). A
271 comparison of the genes differentially expressed by siCITED1 compared to
272 siNEG, revealed that there was significant enrichment of these potential
273 targets (Fig. S2a). Notably, genes both up and down regulated by siCITED1
274 are represented among genes defined as having MITF-occupied promoters
275 (Fig. S2b). We also found that downregulation of MITF using siRNA in HT144
276 cells (Fig. 4c) and in WM293A, and SKMEL5 cells (Fig. S3a,b) resulted in

277 decreased protein expression of CITED1 suggesting reciprocity between these
278 factors.

279 **Induction of MITF by CITED1 silencing transiently restrains cell cycle**
280 **progression and impacts cell viability**

281 To investigate the effect of CITED1 silencing on melanoma cells behaviour we
282 analysed the cell cycle distribution following siRNA treatment, by flow
283 cytometry. In siCITED1 treated HT144 cells we saw G1 accumulation as
284 indicated by an increase in the diploid G1 fraction and a concomitant
285 reduction in the total S-phase fraction peaking at 33 hours but also observed
286 at 48 and 72 hours post-transfection in comparison to siRNA control HT144
287 cells. Again, the effect was apparent but not as pronounced using the #3
288 siCITED1 (Fig S4a). Similar effects were seen in #1 and #3 siCITED1 treated
289 A2058 and A375 cells (Fig. S4b,c).

290 Owing to the previously reported dependency of MITF-induced cell cycle
291 arrest on CDKN1A/P21 we investigated the levels of several cyclin-dependant
292 kinase inhibitors following CITED1 silencing (Carreira et al. 2005). We found
293 that CDKN1A/P21 was transiently increased in siCITED1 treated HT144 cells
294 relative to the siNEG treated HT144 cells. In contrast, in A2058 cells, which
295 do not have detectable levels of CDKN1A/P21 (Fig. S5), the levels of
296 CDKN1C/P57 were suppressed in response to CITED1 overexpression (Fig 5b).
297 We hypothesised therefore that melanoma cells can utilise either
298 CDKN1A/P21 or CDKN1C/P57 to mediate cell cycle arrest induced by MITF and

299 this is reflected in the expression levels of the alternate CDK inhibitors in
300 different melanoma cell lines (Fig. S5).

301

302 In agreement with the cell cycle data, an Alamar Blue assay revealed a
303 significant reduction in cell metabolic activity over 5 days in HT144 cells
304 treated with siCITED1 (Fig 5c). The effect was apparent but not as
305 pronounced in the #3 siCITED1 sample.

306 **The effect of CITED1 silencing on MITF is transient and mediated via** 307 **promoter activation**

308 We observed that the peak upregulation of MITF and CDKN1A/P21 protein
309 following siCITED1 treatment varied from transfection to transfection, being
310 seen between 24-48 hours post-transfection but appearing as unchanged or
311 even downregulated after this time (Fig. 6a, *upper panel*). In agreement, later
312 timepoints of the cell cycle analysis (≥ 72 hours) exhibited little or no
313 change in G1/S-phase distribution or even a reverse pattern (Fig. 5a HT144,
314 and data not shown: A2058, A375). We therefore sought to examine the
315 transcriptional dynamics more closely, map the changes in MITF following
316 CITED1 silencing and see if they corresponded to cell behaviour and changes
317 at the protein level. We used a quantitative droplet digital PCR based assay
318 (Biorad) to measure mRNA in HT144 cells transfected with siCITED1#1 and
319 siNEG as well as A2058 cells transiently overexpressing CITED1 compared to
320 an empty vector control. MITF, CITED1 and IPO8 specific primers and probes
321 were used to measure exact copies/ μ l of each mRNA from aliquots of the

322 same cDNA solution. Plots of siCITED1(copies/ μ l)/siNEG(copies/ μ l) and
 323 EV(copies/ μ l)/pCITED1(copies/ μ l) show the directional change in MITF and
 324 CITED1 relative to the housekeeper IPO8. CITED1 expression is rapidly
 325 suppressed following siCITED1 treatment of HT144 cells, concomitant with an
 326 upregulation of MITF that diminishes over time and in fact is suppressed by
 327 100 hours in accordance with observations at the protein level (Fig 6a, *lower*
 328 *panel*). In contrast, overexpression of CITED1 in A2058 cells results in
 329 transient suppression of MITF at both protein and transcript level (Fig. 6b,
 330 *upper and lower panels*).

331 The rapid MITF transcriptional response to CITED1 manipulation suggested to
 332 us that the effect could be directly mediated at the promoter level. To test
 333 this hypothesis, we over expressed an MITF-M promoter-reporter construct
 334 and CITED1 in A375 cells. We chose A375 cells, as while they had less
 335 endogenous CITED1 and MITF than HT144 or A2058 so as not to cause
 336 interference with the assay, we also knew that they could respond
 337 adequately as they had an identical G1 accumulation/S-phase decrease to
 338 both HT144 and A2058 cells following CITED1 silencing (Fig S4c). TGF β
 339 treatment was used as a positive control for repression of the MITF-M
 340 promoter. CITED1 transfection led to significant suppression of the MITF-M
 341 promoter luciferase activity relative to the empty vector control, as did TGF β
 342 treatment alone or combination with CITED1 overexpression (Fig. 6c). There
 343 did not appear to be an additive or synergistic effect using both TGF β

344 treatment and CITED1 overexpression suggesting TGF β may be dependent on
345 CITED1 for MITF suppression.

346 **The CITED1-silenced gene signature predicts outcome in primary** 347 **melanoma**

348 The 'proliferative' and 'invasive' signature phenotypes have served to define
349 the gene expression classification of melanoma cell lines. However, primary
350 tumours and metastatic lesions have also been molecularly classified into
351 several distinct groups by gene expression profiling (Harbst et al. 2012;
352 Jonsson et al. 2010). The four-class structure found in tumours consists of the
353 'pigmentation', 'proliferative', 'high-immune' and 'normal-like' subgroups
354 with a subset falling into an unclassifiable cohort (Jonsson et al. 2010). We
355 used the same tumour classification to subtype the 120 cell lines that had
356 publically available expression data and could show that the tumour
357 'pigmentation' subgroup that highly expresses MITF, corresponds to the cell
358 line 'proliferative' phenotype described by Hoek et al. Accordingly, the
359 tumour 'proliferative' and 'high-immune' subgroups comprise the cell line
360 'invasive' phenotype (Fig. 7a). It is worth noting that the names of the tumour
361 subgroups were derived from a description of the differentially expressed
362 genes that comprised each molecular classification while the 'invasive-
363 proliferative' switching phenotypes were named to reflect the *behaviour*
364 exhibited by lines classified by one or other signature. This distinction helps
365 to explain the confusing occurrence that both classifications have a group
366 referred to as 'proliferative' although they are not equivalent.

367 The overlap between the primary tumour classifying and cell line classifying
 368 systems allows us to infer that CITED1 expression is most likely restricted to
 369 a subset of MITF high 'pigmentation' subtype tumours. As the tumour
 370 subtype classification was shown to be prognostically significant in primary
 371 melanomas we were interested to know if CITED1 expression itself was
 372 independently predictive of outcome. Previously we reported on the analysis
 373 of 223 primary lesions using the Illumina WG-DASL protocol (Harbst et al.
 374 2012). As the CITED1 probe in this assay did not produce reliable data we
 375 instead derived a CITED1-silenced gene signature score based on the
 376 differentially expressed genes from the HT144 siCITED1 experiment (Fig. 3)
 377 We subsequently interrogated the gene expression data on the primary
 378 melanoma lesions using a nearest centroid approach derived from the
 379 CITED1-silenced gene signature. This revealed that primary melanomas with
 380 a gene expression signature most similar to the CITED1-silenced signature
 381 (CITED1low-class) had a significantly better outcome than those with a
 382 signature most disparate from the CITED1-silenced signature (CITED1high-
 383 class) (Fig. 7b). Importantly, the CITED1 signature classing had independent
 384 prognostic information (HR 1.85, CI 0.30-0.98, p=0.044) from the AJCC
 385 staging system (HR 5.05, CI 2.42-10.55, p=1.64x10⁻⁵). These data indirectly
 386 imply that CITED1 expression itself is a potential prognostic indicator in
 387 primary melanomas and the transcriptional program influenced by CITED1
 388 expression determines tumour behaviour *in vivo*.

DISCUSSION

One seemingly paradoxical observation from our study and previous investigations is that although CITED1 behaves as a negative regulator of MITF, both their expression levels appear positively correlated across cell lines and tumours. We maintain that this observation simply reflects the fact that where there are high levels of MITF, high levels of its negative regulator are also required. The evidence of the tight control exerted over MITF levels in melanocytes and melanoma simply speaks to the necessity of the cell to maintain a level compatible with survival and proliferation, in a type of biological 'sweet-spot' facilitating tumour progression. The cellular effects of both extremes i.e: very low or high levels of MITF, have been elegantly described by a rheostat model in order to reconcile the conflicting observations of the effects of manipulating MITF *in vitro*, and the fact that counter-intuitively, a lineage-specifying differentiation factor can behave as a potent oncogene (Hoek & Goding 2010; Carreira et al. 2006; Cheli, Giuliano, et al. 2011a). The rheostat model (Fig. S6) explains why MITF silencing can block cells in G1 and induce senescence, while it is also possible to induce a G1 arrest by MITF overexpression via CDKN2A/P16 or CDKN1A/P21 and, as we now propose, potentially also via CDKN1C/P57 (Carreira et al. 2006; Loercher et al. 2005; Carreira et al. 2005). At the extreme high end of MITF expression lies differentiated melanocytic cells, while the lowest levels can lead to senescence and irreversible cell death. Between these two extremes however it is thought that melanoma cells can oscillate from a low-MITF 'invasive' to a high-MITF 'proliferative' state via phenotype-switching.

413 We hypothesise that the role of CITED1 in melanoma is to maintain levels of
 414 MITF compatible with tumour progression and effectively tip the balance in
 415 favour of cell cycle progression rather than MITF-induced G1-arrest. This is
 416 supported by our findings that downregulation of CITED1 using siRNA results
 417 in a phenotype switch to a more pigmented state driven by increased MITF
 418 expression and concomitant upregulation of CDKN1A/P21. Conversely, we
 419 could observe that downregulation of MITF resulted in suppression of CITED1
 420 in several cell lines suggesting the existence of a classical feedback loop
 421 where low MITF levels result in inhibition of its negative regulator. MITF
 422 induced cell cycle arrest was previously shown to be dependent on
 423 CDKN1A/P21 and it was demonstrated that MITF does not cause a cell cycle
 424 arrest in CDKN1A-deficient mouse embryo fibroblasts (MEF) cells (Carreira et
 425 al. 2005). However, our data indicate that in melanoma cells deficient in
 426 CDKN1A/P21, the alternative CDK inhibitor CDKN1C/P57 is expressed and
 427 responsive to MITF.

428 Interestingly, while we observed upregulation of most MITF targets following
 429 CITED1 silencing, we found that BRCA1 and other DNA damage response
 430 (DDR) genes were suppressed, suggesting that CITED1 downregulation does
 431 not necessarily facilitate transcription of all MITF targets. It is thus tempting
 432 to speculate that rather than simply acting to induce MITF and thereby
 433 indirectly enhance transcription of its target genes, that CITED1 may also act
 434 as co-factor for MITF at various genomic locations differentially modulating

the MITF target gene response at individual promoters. One way that this might be achieved is via MITF-CITED1 competition for CBP/P300 binding as CBP/P300 is a known transcriptional coregulator for MITF, although it is not required for transcription of all MITF targets (Vachtenheim et al. 2007) (Yan et al. 2013).

As suggested by Sáez-Ayala et al., anti-cancer therapy should be ideally independent of dominant or 'driver' genetic alterations so that subclonal populations do not gain a subsequent advantage and the same holds true in the case of targeting a specific phenotype. Successful therapy will necessarily need to switch or push the subdominant phenotype into the susceptible state or eradicate the phenotype resistant to treatment. This approach was initially championed by Cheli et al., who proposed the eradication of low-MITF cells as a therapeutic strategy (Cheli, Guiliano, et al. 2011b). Indeed the idea of lineage-specific therapy has been subsequently proved in principle using methotrexate (MTX) to first activate MITF expression, in turn activating the tyrosinase enzyme, and thereby sensitising tumour cells to a tyrosinase-processed anti-folate prodrug (TMECG) (Sáez-Ayala et al. 2013). However, even without drug targeting, induction of MITF, above what is tolerated by even the highly pigmented tumor cell types, i.e: levels reaching that of melanocytes, would seem to be incompatible with melanoma progression as it inhibits cell cycle progression. Our assertion is that CITED1 acts to repress MITF in order to maintain its level in a range compatible with tumorigenesis. This assertion as a consequence naturally

suggests CITED1 as therapeutic target for genetic manipulation. Successful implementation of such a strategy would result in cell specific enhancement of MITF expression and increased susceptibility to the type of chemotherapeutic eradication demonstrated by Sáez-Ayala et al. or potentially induction of CDKN1A/p21 or CDKN1C/p57-dependent cell growth arrest even without further intervention (Fig. S6) (Sáez-Ayala et al. 2013).

Acknowledgements

We would like to thank Dr. Claudia Wellbrock (University of Manchester, UK) for the kind gift of the PG2-MITF promoter-reporter construct and Prof. Toshi Shioda (MGH, Harvard) for the human CITED1 pRc/CMV expression vector. We also would like to acknowledge Prof. Bo Baldetorp (Division of Oncology, Lund University) for his advice on cell cycle analysis and the support of Prof. Tommy Andersson (Cell and Experimental Pathology, ILMM, Lund University).

References

- Al, S. et al., 2003. TM4: a free, open-source system for microarray data management and analysis. *Biotechniques*, 32(2), pp.374-378.
- Bertolotto, C. et al., 2011. A SUMOylation-defective MITF germline mutation predisposes to melanoma and renal carcinoma. *Nature*, 480(7375), pp.94-98.
- Beuret, L. et al., 2011. BRCA1 is a new MITF target gene. *Pigment Cell & Melanoma Research*,

- 477 24(4), pp.725–727.
- 478 Carreira, S. et al., 2005. Mitf cooperates with Rb1 and activates p21Cip1 expression to
479 regulate cell cycle progression. *Nature*, 433(7027), pp.764–769.
- 480 Carreira, S. et al., 2006. Mitf regulation of Dia1 controls melanoma proliferation and
481 invasiveness. *Genes & development*, 20(24), pp.3426–3439.
- 482 Cheli, Y., Giuliano, S., et al., 2011a. Hypoxia and MITF control metastatic behaviour in mouse
483 and human melanoma cells. *Oncogene*, 31(19), pp.2461–2470.
- 484 Cheli, Y., Guiliano, S., et al., 2011b. Mitf is the key molecular switch between mouse or
485 human melanoma initiating cells and their differentiated progeny. *Oncogene*, 30(20),
486 pp.2307–2318.
- 487 Davies, H. et al., 2002. Mutations of the BRAF gene in human cancer. *Nature*, 417(6892),
488 pp.949–954.
- 489 Garraway, L.A. et al., 2005. Integrative genomic analyses identify MITF as a lineage survival
490 oncogene amplified in malignant melanoma. *Nature*, 436(7047), pp.117–122.
- 491 Giuliano, S. et al., 2010. Microphthalmia-associated transcription factor controls the DNA
492 damage response and a lineage-specific senescence program in melanomas. *Cancer
493 research*, 70(9), pp.3813–3822.
- 494 Haqq, C. et al., 2005. The gene expression signatures of melanoma progression. *Proceedings
495 of the National Academy of Sciences of the United States of America*, 102(17), pp.6092–
496 6097.
- 497 Harbst, K. et al., 2012. Molecular profiling reveals low- and high-grade forms of primary
498 melanoma. *Clinical cancer research : an official journal of the American Association for
499 Cancer Research*, 18(15), pp.4026–4036.
- 500 Hoek, K.S. & Goding, C.R., 2010. Cancer stem cells versus phenotype-switching in melanoma.
501 *Pigment Cell & Melanoma Research*, 23(6), pp.746–759.
- 502 Hoek, K.S. et al., 2006. Metastatic potential of melanomas defined by specific gene
503 expression profiles with no BRAF signature. *Pigment cell research / sponsored by the
504 European Society for Pigment Cell Research and the International Pigment Cell Society*,
505 19(4), pp.290–302.
- 506 Hoek, K.S., Eichhoff, O.M., et al., 2008a. In vivo switching of human melanoma cells between
507 proliferative and invasive states. *Cancer research*, 68(3), pp.650–656.
- 508 Hoek, K.S., Schlegel, N.C., et al., 2008b. Novel MITF targets identified using a two-step DNA
509 microarray strategy. *Pigment Cell & Melanoma Research*, 21(6), pp.665–676.
- 510 Huang, D.W. et al., 2007. DAVID Bioinformatics Resources: expanded annotation database
511 and novel algorithms to better extract biology from large gene lists. *Nucleic Acids
512 Research*, 35(Web Server), pp.W169–W175.
- 513 Jenei, V. et al., 2009. A t-butyloxycarbonyl-modified Wnt5a-derived hexapeptide functions as
514 a potent antagonist of Wnt5a-dependent melanoma cell invasion. *Proceedings of the
515 National Academy of Sciences of the United States of America*, 106(46), pp.19473–
516 19478.
- 517 Jonsson, G. et al., 2010. Gene Expression Profiling-Based Identification of Molecular Subtypes

- 518 in Stage IV Melanomas with Different Clinical Outcome. *Clinical Cancer Research*, 16(13),
519 pp.3356-3367.
- 520 Levy, C., 2001. A New Role for the STAT3 Inhibitor, PIAS3. A REPRESSOR OF
521 MICROPHTHALMIA TRANSCRIPTION FACTOR. *Journal of Biological Chemistry*, 277(3),
522 pp.1962-1966.
- 523 Levy, C., Khaled, M. & Fisher, D.E., 2006. MITF: master regulator of melanocyte development
524 and melanoma oncogene. *Trends in Molecular Medicine*, 12(9), pp.406-414.
- 525 Loercher, A. et al., 2005. MITF links differentiation with cell cycle arrest in melanocytes by
526 transcriptional activation of INK4A. *The Journal of cell biology*, 168(1), pp.35-40.
- 527 McGill, G.G. et al., 2006. c-Met Expression Is Regulated by Mitf in the Melanocyte Lineage.
528 *Journal of Biological Chemistry*, 281(15), pp.10365-10373.
- 529 Nair, S.S. et al., 2001. Over-expression of MSG1 Transcriptional Co-activator Increases
530 Melanin in B16 Melanoma Cells: A Possible Role for MSG1 in Melanogenesis. *Pigment cell
531 research / sponsored by the European Society for Pigment Cell Research and the
532 International Pigment Cell Society*, 14(3), pp.206-209.
- 533 Pierrat, M.J. et al., 2012. Expression of Microphthalmia-associated Transcription Factor (MITF),
534 Which Is Critical for Melanoma Progression, Is Inhibited by Both Transcription Factor GLI2
535 and Transforming Growth Factor. *The Journal of biological chemistry*, 287(22), pp.17996-
536 18004.
- 537 Pinner, S. et al., 2009. Intravital Imaging Reveals Transient Changes in Pigment Production
538 and Brn2 Expression during Metastatic Melanoma Dissemination. *Cancer research*,
539 69(20), pp.7969-7977.
- 540 Plisov, S., 2005. Cited1 Is a Bifunctional Transcriptional Cofactor That Regulates Early
541 Nephronic Patterning. *Journal of the American Society of Nephrology*, 16(6), pp.1632-
542 1644.
- 543 Ryu, B. et al., 2007. Comprehensive expression profiling of tumor cell lines identifies
544 molecular signatures of melanoma progression. *PLoS ONE*, 2(7), p.e594.
- 545 Sáez-Ayala, M. et al., 2013. Directed Phenotype Switching as an Effective Antimelanoma
546 Strategy. *Cancer Cell*, 24(1), pp.105-119.
- 547 Sánchez-Martín, M. et al., 2002. SLUG (SNAI2) deletions in patients with Waardenburg
548 disease. *Human molecular genetics*, 11(25), pp.3231-3236.
- 549 Sestáková, B., Ondrusová, L. & Vachtenheim, J., 2010. Cell cycle inhibitor p21/ WAF1/ CIP1 as
550 a cofactor of MITF expression in melanoma cells. *Pigment Cell & Melanoma Research*,
551 23(2), pp.238-251.
- 552 Shioda, T. et al., 1998. Transcriptional activating activity of Smad4: Roles of SMAD hetero-
553 oligomerization and enhancement by an associating transactivator. *Proceedings of the
554 National Academy of Sciences of the United States of America*, 95(17), p.9785.
- 555 Shioda, T., Fenner, M.H. & Isselbacher, K.J., 1996. msg1, a novel melanocyte-specific gene,
556 encodes a nuclear protein and is associated with pigmentation. *Proceedings of the
557 National Academy of Sciences of the United States of America*, 93(22), pp.12298-12303.
- 558 Smith, A.P., Hoek, K. & Becker, D., 2005. Whole-genome expression profiling of the
559 melanoma progression pathway reveals marked molecular differences between

- 560 nevi/melanoma in situ and advanced-stage melanomas. *Cancer Biology & Therapy*, 4(9),
561 pp.1018-1029.
- 562 Strub, T. et al., 2011. Essential role of microphthalmia transcription factor for DNA replication,
563 mitosis and genomic stability in melanoma. *Oncogene*, 30(20), pp.2319-2332.
- 564 Talantov, D., 2005. Novel Genes Associated with Malignant Melanoma but not Benign
565 Melanocytic Lesions. *Clinical Cancer Research*, 11(20), pp.7234-7242.
- 566 Vachtenheim, J., Šestáková, B. & Tuháčková, Z., 2007. Inhibition of MITF transcriptional
567 activity independent of targeting p300/CBP coactivators. *Pigment cell research /*
568 *sponsored by the European Society for Pigment Cell Research and the International*
569 *Pigment Cell Society*, 20(1), pp.41-51.
- 570 Vallon-Christersson, J. et al., 2009. BASE--2nd generation software for microarray data
571 management and analysis. *BMC bioinformatics*, 10, p.330.
- 572 Wellbrock, C. et al., 2008. Oncogenic BRAF regulates melanoma proliferation through the
573 lineage specific factor MITF. *PLoS ONE*, 3(7), p.e2734.
- 574 Widlund, H.R. & Fisher, D.E., 2003. Microphthalmia-associated transcription factor: a critical
575 regulator of pigment cell development and survival. *Oncogene*, 22(20), pp.3035-3041.
- 576 Widmer, D.S. et al., 2012. Systematic classification of melanoma cells by phenotype-specific
577 gene expression mapping. *Pigment Cell & Melanoma Research*, 25(3), pp.343-353.
- 578 Yahata, T., 2001. Selective coactivation of estrogen-dependent transcription by CITED1
579 CBP/p300-binding protein. *Genes & development*, 15(19), pp.2598-2612.
- 580 Yahata, T., 2000. The MSG1 Non-DNA-binding Transactivator Binds to the p300/CBP
581 Coactivators, Enhancing Their Functional Link to the Smad Transcription Factors. *Journal*
582 *of Biological Chemistry*, 275(12), pp.8825-8834.
- 583 Yan, G. et al., 2013. Selective inhibition of p300 HAT blocks cell cycle progression, induces
584 cellular senescence, and inhibits the DNA damage response in melanoma cells. *Journal of*
585 *Investigative Dermatology*, 133(10), pp.2444-2452.
- 586 Yokoyama, S. & Fisher, D.E., Transcriptional Regulation in Melanoma. In A. Bosserhoff, ed.
587 *Melanoma Development*. SpringerWeinNewYork.
- 588 Yokoyama, S. et al., 2011. A novel recurrent mutation in MITF predisposes to familial and
589 sporadic melanoma. *Nature*, 480(7375), pp.99-103.

Figure 1

TGF β induced gene expression in A2058 melanoma cells

(a) Distribution of the proliferative and invasive signature score genes relative to the heat map of gene expression changes induced by TGF β treatment. (b) Gene expression heatmap of the proliferative and invasive signature within those genes significantly altered by TGF β treatment (1009 transcripts following SAM, median FDR q-value=1%. '2006' refers to the signature list (motif1 and motif2) while '2012' refers to the updated signature derived from further datasets (Widmer et al. 2012). (c) Western blot of MITF and CITED showing both proteins are suppressed by TGF β treatment. β -Actin is used as a loading control.




figure 1

Figure 2

CITED1 expression correlates with MITF expression

(a) The relative MITF and CITED1 expression levels from the publically available gene expression data of 120 melanoma cell lines (Pearson correlation $r=0.6543$, $p < 0.001$). Arrows indicate the cell lines used in this study. The cell lines are further subdivided into one of either 'invasive' or 'proliferative' phenotype based on expression signature score. (b) A Western blot is shown of the relative protein expression levels of both MITF and CITED1 in our cell lines in good agreement with the transcript levels.




figure 2

Figure 3

CITED1 silencing induces a phenotype switch

(a) 120 melanoma cell lines are shown distributed on the basis of the phenotype score. The HT144 cells chosen to study the effects of CITED1 downregulation are indicated. (b) Following CITED1 downregulation a phenotype shift is observed indicated by their scatter position change according to the average expression score of genes that distinguish invasive from proliferative phenotype. For the 120 melanoma cell lines (Affymetrix platform) the expression score was derived from expression levels of 50 and 54 proliferative and invasive genes with matching genes symbols, respectively, while for the HT144 experiment (Illumina platform), 51 and 54 proliferative and invasive genes with matching genes symbols were retrieved. A Western blot of the degree of protein downregulation of CITED1 at the time of the expression analysis is also shown. β -Actin is used as a loading control (inset). (c) A heatmap comprising the 'invasive' and 'proliferative' signature genes illustrating how they are altered by CITED1 silencing; #1 and #3 denote two separate siRNAs targeting CITED1.





figure 3

Figure 4


CITED1 regulates MITF and its targets genes

(a) A heatmap showing the 312 transcripts identified as significantly changed using a SAM 2-way comparison between siNEG and siCITED1 (#1 & #3 were combined), median FDR q-value = 3%. MITF, as well as a cohort of significantly enriched MITF targets, genes associated with pigmentation, and genes involved in the UV/DNA damage response are highlighted on the right. (b) Western blot confirmation of the effect of silencing CITED1, using siRNA (#1, #3) relative to a negative control siRNA (N), on MITF protein expression in HT144 cells at 24 and 48 hours post-transfection, and the effect of overexpressing CITED1 (pCITED1) relative to an empty vector control (EV) in A2058 cells at 24 hours post-transfection. β -Actin is used as a loading control in each case. (c) Western blot showing the effect of silencing MITF using two siRNAs (siM1, siM3) on both MITF and CITED1 levels in HT144 cells at 48 hours post-transfection relative to a negative control siRNA (N). β -Actin is used as a loading control.

a



b



c




figure 4

Figure 5

CITED1 silencing restrains cell cycle progression and reduces cell viability

(a) A bar chart showing the % change in cell cycle distribution in #1 siCITED1 treated HT144 cells relative to siNEG treated HT144 cells. The reduction in the total S-phase is shown at 33 hours, 48 hours and 72 hours post-transfection in addition to the corresponding increase in the diploid G1 fraction. (b) Western blots showing upregulation of CDKN1A/P21 following CITED1 silencing in HT144 cells and suppression of CDKN1C/P57 following CITED1 overexpression in A2058 cells. (c) An Alamar Blue based metabolic assay shows a reduction in cell viability over 5 days in HT144 cells treated with siCITED1 relative to those treated with siNEG. Stars indicate significance for siNEG vs. #1 siCITED1 where *** $p \leq 0.0005$, ** $p \leq 0.005$ and * $p \leq 0.05$. In the case of siNEG vs. #1 siCITED1, the difference is significant (*) at 96 and 120 hours.




figure 5

Figure 6

CITED1 silencing transiently upregulates MITF via promoter activation

(a) A Western blot of HT144 cell lysate samples taken at the indicated time points post-transfection and showing the corresponding levels of MITF protein in #1 siCITED1 and siNEG treated cells. Shown underneath are the changes in mRNA levels of MITF-M, CITED1 and a housekeeper gene IPO8, as measured by specific ddPCR assays over a time course of 4-100 hours following transfection of HT144 cells with either siCITED1 or siNEG. (b) A Western blot of A2058 cell lysate samples taken at the indicated time points post-transfection and showing the corresponding levels of MITF protein between CITED1 overexpression (pCITED1) and empty vector (EV) control. Shown underneath are the changes in mRNA levels of MITF and a housekeeper gene IPO8, as measured by specific ddPCR assays over a time course of 33-72 hours following transfection of A2058 with either pCITED1 or an empty vector control. (c) The relative luciferase activity of the MITF-M promoter reporter measured in lysates of A375 cells transfected with the pCITED1 expression plasmid or empty vector (EV) control and treated with or without TGF β for 24 hours (*** $p \leq 0.0005$, ** $p \leq 0.005$ and *NS*= not significant).




figure 6

Figure 7

The CITED1-silenced gene signature predicts patient outcome

(a) In the leftmost panel a scatter plot of the 120 melanoma cell lines are shown distributed on the basis of their 'invasive' or 'proliferative' phenotype signature score and coloured according to the tumor molecular subtypes as defined by Jönsson et al. to illustrate the overlap between the two classification systems (Jonsson et al. 2010). In the rightmost panel the data is presented as a heat map where each gene of the proliferative or invasive signature genes is represented by a horizontal line and the 120 individual cells lines are grouped by molecular tumour subtype (coloured blocks) and shown vertically. (b) Recurrence free survival (RFS) of primary melanoma patients grouped by gene expression similarity to the CITED1 (siCITED1) silenced gene signature.




figure 7