

A peer-reviewed version of this preprint was published in PeerJ on 16 October 2014.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.631) (peerj.com/articles/631), which is the preferred citable publication unless you specifically need to cite this preprint.

Pfister CA, Gilbert JA, Gibbons SM. 2014. The role of macrobiota in structuring microbial communities along rocky shores. PeerJ 2:e631 <https://doi.org/10.7717/peerj.631>

The role of macrobiota in structuring microbial communities along rocky shores

Rocky shore microbial diversity presents an excellent system to test for microbial habitat specificity or generality, enabling us to decipher how common macrobiota shape microbial community structure. At two coastal locations in the northeast Pacific Ocean, we show that microbial composition was significantly different between inert surfaces, the biogenic surfaces that included rocky shore animals and an alga, and the water column plankton. While all sampled entities had a core of common OTUs, rare OTUs drove differences among biotic and abiotic substrates. For the mussel *Mytilus californianus*, the shell surface harbored greater alpha diversity compared to internal tissues of the gill and siphon. Strikingly, a 7-year experimental removal of this mussel from tidepools did not significantly alter the microbial community structure of microbes associated with inert surfaces when compared with unmanipulated tidepools. However, bacterial taxa associated with nitrate reduction had greater relative abundance with mussels present, suggesting an impact of increased animal-derived nitrogen on a subset of microbial metabolism. Because the presence of mussels did not affect the structure and diversity of the microbial community on adjacent inert substrates, microbes in this rocky shore environment may be predominantly affected through direct physical association with macrobiota.

1
2
3 The role of macrobiota in structuring microbial communities along rocky shores
4

5 Catherine A. Pfister^{1*}, Jack A. Gilbert^{1,2}, Sean M. Gibbons^{2,3}

6 ¹Department of Ecology and Evolution

7 and

8 ²Institute of Genomic and Systems Biology

9 Argonne National Laboratory

10 Lemont, IL 60439 USA

11 ³Biophysical Sciences Graduate Program

12 University of Chicago

13 Chicago IL 60637

14 *corresponding author: Department of Ecology & Evolution, 1101 E 57th St, University of
15 Chicago, Chicago IL 60637, tele 773-834-0071, fax 773-702-9740, cpfister@uchicago.edu
16

17 Running title: rocky shore microbial community structure
18

19 Submitted to PeerJ
20
21
22
23
24

INTRODUCTION

The dynamics and interactions of the macroscopic species on rocky shores of the northeast Pacific Ocean have been well-characterized and thus have contributed significantly to our understanding of coastal ecological processes (eg. Paine 1966; Wootton 1994; Estes and Duggins, 1995). Although some specialized symbiotic associations have been described in rocky shore species (Secord and Augustine, 2005; Bergschneider and Muller-Parker, 2008), we know little about multi-taxa microbial associations. There is increasing evidence that many marine macrobiota have surface biofilms (Grossart *et al.*, 2005; Kvennefors *et al.*, 2012) or endosymbionts (Zurel *et al.*, 2011; Wegner *et al.*, 2012) or both (e.g. Qian *et al.*, 2006; Taylor *et al.*, 2007). However, our understanding of the specificity of these associations and their functional significance remain nascent with some notable exceptions (Webster and Taylor, 2012; Fan *et al.*, 2013; Heisterkamp *et al.*, 2013). The shelf waters of the California Current Large Marine Ecosystem (CCLME) maintain diverse and unique microbial communities across upwelling areas (Bertagnolli *et al.*, 2011). The relatively high productivity of this system has been attributed to the seasonal upwelling of nitrate (Barber and Smith 1981), which can lead to significant levels of carbon fixation. The diversity of potential plant and animal 'host' species in the CCLME, their relatively large geographic range, their longevity and the provision of key resources suggest they provide unique microniches capable of increasing the diversity and function of microbial communities.

If macrobiota directly provide habitat for intertidal microbes (e.g. mussels, Pfister *et al.*, 2010; algae, Miranda *et al.*, 2013) or indirectly provide resources such as nitrogen, in the form of animal-regenerated ammonium, then our understanding of coastal biogeochemistry is incomplete without considering the contribution of macrobiota. In the northeast Pacific, Tatoosh Island, WA

shows persistent shoreward peaks of ammonium (Pfister *et al.*, 2007), while the tidepools at Second Beach, WA have animal-regenerated ammonium from the California mussel with locally enhanced algal productivity, and microbial nitrification (Pfister 2007). We employed 16S rRNA V4 amplicon sequencing to test whether microbes associated with macrobiota differ from those in the water column and on inert surfaces. We further compared the identified microbial taxa with those found when using shotgun metagenomics in a subset of samples for mussel shell biofilms. Finally, we used a manipulative field experiment of mussel presence or absence in tidepools to ask whether the increased rates of nitrogen remineralization and uptake demonstrated with mussels by tracer ammonium addition (Pather *et al.*, 2014) resulted in changed bacterial communities on inert surfaces.

MATERIALS AND METHODS

We asked whether microbial diversity and abundance were distributed differentially among intertidal microhabitats by 1) sampling distinct microhabitat types including artificial and natural surfaces at Tatoosh Island (48.32°N, 124.74°W), and 2). Sampling artificial and natural substrates in the context of an animal removal experiment at nearby Second Beach (48°, 23'N, 124, 40'W). Tatoosh Island is 0.7 km off the northwestern tip of Washington State, USA and has been well studied ecologically. Previous metagenomic analysis of biofilms associated with the shells of mussels at Tatoosh Island demonstrated a rich microbial assemblage, with the genetic capacity for nitrogen cycling (Pfister *et al.*, 2010). On 6 Aug 2009, we sampled both inert and biogenic substrates in situ, as well as artificial substrates. Biogenic substrates sampled included the surface of the red alga *Prionitis sternbergii* (n=2), the anemone *Anthopleura elegantissima* (n=2), and gill tissue (n=3) and siphon tissue (n=1) of the California mussel *Mytilus californianus*. These biogenic hosts were chosen because they are persistent members of the

community and have relatively long-lived and sessile tissues that might provide a predictable substrate. In addition to the biogenic surfaces, rocks were chipped off of bench adjacent to mussel beds. Artificial substrates (glass crucible covers, 3 cm diam, www.leco.com) were attached with epoxy on rock adjacent to mussel beds (n=4) and in tidepools (n=5) on 10 Jun 2009 and left *in situ* for 2 months. We chose glass crucible covers because they provided heterogeneous surface, yet an inert substance. Immediately after a late morning collection at low tide, samples were frozen and sent to -80°C storage at the University of Chicago prior to extraction. Environmental characteristics of the seawater, including nutrient concentrations, were recorded as part of a regular sampling program (Wootton and Pfister, 2012) and are reported in Table S1. The Upwelling Index for this latitude (48°N) was, on average, positive (www.pfeg.noaa.gov).

We tested whether the presence of mussels had entrained different microbial assemblages by sampling natural and artificial substrates in tidepools where mussels had been removed or unmanipulated since 2002 at Second Beach (Pfister, 2007), a north-facing complex of rocks 2 km east of Neah Bay, WA, USA within the Makah Tribal Reservation. For seven years prior to our microbial sample collection, mussels have been excluded from 6 tidepools that previously contained mussels by pulling them out by hand at an approximately monthly interval during the spring and summer months. Other tidepools that naturally had mussels served as controls. The biogeochemistry of these experimental and control tidepools have been characterized multiple times, allowing us to test for mussel effects on tidepool nutrient concentrations (nitrate, nitrite, ammonium, and phosphorus), and seawater pH and oxygen and how it related to microbial community structure. These parameters were measured 9 times in each tidepool during Aug 2009 and 12 times in Jul and Aug of 2010. We also estimated ammonium remineralization and

removal rates with a tracer experiment with enriched $^{15}\text{NH}_4^+$ in 2010 (Pather *et al.*, 2014). A ~ 2cm² piece of rock was chiseled from 5 pools with and 5 without mussels during morning hours on 24 Aug 2009. We also sampled from glass cover slips that had incubated for 3 months in these pools (n=7 controls, n=6 mussel removal pools) with a copper paint barrier to exclude molluscan grazers. To compare this benthic microbial assemblage to microbes in the water column, we collected 3 plankton samples on 25mm GF/F filters by pumping 300 mL of seawater on an incoming tide (1100-1130h). Due to the 0.7 um pore size, we likely excluded many free-living microbes, though high species richness still resulted (see below). All samples were frozen and sent to storage at -80°C at the University of Chicago prior to extraction.

For DNA extraction, we used the Power Soil DNA Extraction Kit (MoBio). The rock, crucible cover substrates, and mussel shells were both swabbed with sterile cotton applicators and brushed with sterile spiral dental brushes that were placed in the extraction solution. Coverslips, filters, excised mussel gill or mussel siphon, anemones, and algae were placed into the beadbeater vial and pulverized. Thus, our sampling of biogenic substrata included the potential that microbes were part of the host tissue. The PCR amplification protocol followed Caporaso *et al.* (2011) for multiplexing 16S rRNA samples. The PCR products were cleaned with MoBio™ UltraClean htp PCR Clean-up kit. We amplified the V4 variable region of the 16S rRNA gene from community DNA using bar-coded primers according to Earth Microbiome Project standard protocols (www.earthmicrobiome.org).

Sequence analysis

We used QIIME (v. 1.7.0, Quantitative Insights into Microbial Ecology; www.qiime.org) to filter reads and determine OTUs as described in Caporaso *et al.* (2010a, 2011). Briefly, we used

the open reference OTU picking workflow, where sequences were first clustered with the Greengenes (Dec 2012) reference database (McDonald *et al.*, 2012); we then allowed OTUs that did not cluster with known taxa (at 97% identity) in the database to cluster *de novo*. Singleton and chloroplast-derived sequences were removed prior to downstream analyses. Representative sequences for each OTU were aligned using PyNast, with a minimum alignment overlap of 75 bp (Caporaso *et al.*, 2010b). Alignments were used to build a phylogenetic tree (FastTree; Price, *et al.*, 2009). We computed alpha diversity metrics among substrates using the `alpha_diversity.py` script in QIIME (chao1, phylogenetic diversity and equitability), using the same sequence depth for all samples (50000 sequences per sample). We used the `beta_diversity_through_plots.py` script to compute beta diversity distances between samples (weighted UniFrac), and to construct principal component (PCoA) plots, thus accounting for both the phylogenetic composition (Lozupone *et al.*, 2011) and the relative abundance of taxa. To test for significant sample groupings based on these distance metrics, we employed PERMANOVA and PERMDISP using the `compare_categories.py` script in QIIME. We tested whether the abundance of particular OTUs differed significantly among different substrates using ANOVA analyses (Bonferonni corrected) with the `otu_category_significance.py` script. OTU networks were constructed using the `make_otu_network.py` script in QIIME. We further visualized the extent to which OTUs were shared or unique among samples using Cytoscape network layouts (www.cytoscape.org). Finally, we tested for patterns in species co-occurrence as a function of mussels with a checkerboard score (c-score) analysis using the `oecosimu` function (Vegan package) with the ‘quasiswap’ method (99 simulations) for null model construction (Barberán *et al.*, 2012).

Because nitrogen metabolism in association with animals was demonstrated in these locales (Pather *et al.* 2014, Pfister *et al.* in press), we tested whether taxa known to be involved in

nitrogen metabolic pathways were present using 3 methodologies. We first examined the taxa identified (down to the level of genus) with the Greengenes database and 16S data in each sample. From literature reports, we assigned taxa to one of 4 transformations: nitrification (either ammonia oxidation or nitrite oxidation), anammox, or nitrate reduction via DNRA or denitrification based on genera associated with each metabolism (Table 1), comparing categories in R (version 2.15, www.R-project.org).

Our second analysis of potential nitrogen cycling used PICRUSt to predict the percent of sequences associated with nitrogen metabolism in our 16s data (Langille *et al.*, 2013). Briefly, all OTUs not assigned to the Greengenes database were removed from the OTU table (closed reference), abundance was normalized by 16s rRNA read number, and PICRUSt metagenome predictions were calculated (Greengenes May 2013 release). By inferring the gene families present in our 16s data, we compared whether the different living and inert substrates on Tatoosh Island differed in the amount of OTUs associated with nitrogen metabolism or whether inert substrates in tidepools with or without mussels differed in the abundance of taxa related to nitrogen metabolism.

The discovery of nitrogen metabolizing taxa with 16S rRNA data require that those taxa are described and identical or analogous sequences are available. In contrast, shotgun metagenomics directly identifies the sequence associated with metabolic function. Two of our samples of *M. californianus* shells were also shotgun pyrosequenced (Pfister *et al.*, 2010), which allowed us to compare nitrogen metabolisms detected with shotgun metagenomics with 16S OTU reads and PICRUSt predictions.. We sequenced the 16S rRNA V4 amplicons using the Illumina platform described above from DNA archived from a previous extraction from Tatoosh Island, where microbial community biofilms associated with mussel shell surfaces were extracted and

metagenomes sequenced using 454 GS-flx pyrosequencing (Pfister *et al.*, 2010). Briefly, 6 mussels were collected on 10 April 2008 from 6 tidepools and 6 additional mussels approximately 5 m apart on an adjacent exposed rocky bench. All shells were immediately cleaned of all soft tissue. Mean shell length was 4.47 cm and 4.42 cm for tidepool and bench mussels, respectively. We thus compared the 16S rRNA V4 amplicon Illumina sequences to the shotgun metagenomic pyrosequencing to determine overlaps in key taxa. The two metagenomes from Pfister *et al.*, (2010) were reanalyzed using MG-RAST for all taxonomic matches and using the SEED Subsystems database for nitrogen metabolism with maximum e-value of $e < 10^{-5}$ (<http://metagenomics.anl.gov/>).

RESULTS

Some distinction exists among microbial assemblages associated with different substrates at Tatoosh Island

Between 54,490 and 250,432 sequences per sample were generated for 26 samples from a range of materials including inert surfaces (rock and glass crucible lids) as well as mussel shells and tissues, algal fronds, sea anemones, and the filtered plankton. All samples were rarified to 50,000 sequences per sample. OTU richness (total diversity) estimates were greatest for inert substrates and the water column, while the lowest richness was associated with mussel gill and siphon tissue (ANOVA, $F_{7,16}=4.968$, $p=0.004$). Species richness was highly correlated with other metrics of diversity including chao1 and phylogenetic diversity ($r=0.98$ to 0.99 , $p<0.001$), as was equitability (or evenness, $r=0.780$, $p<0.001$). Alphaproteobacteria, including Rhodobacteraceae and Hyphomonadaceae, dominated the algal *Prionitis* tissue and the inert substrates, while Gammaproteobacteria, especially Vibrionaceae, dominated mussel gill and siphon tissue (Fig 1).

186 The communities associated with mussel shells, anemones, and filtered plankton samples were
187 similarly dominated by Gammaproteobacteria. Mussel shells and anemones similarly had many
188 Vibrionaceae OTUs, with shells also harboring Moritellaceae. OTUs in the Psychromonadaceae
189 were prominent in the plankton. Given our use of a 0.7 μ M filter, the OTU richness may be
190 underestimated if the smallest bacteria were not retained.

191 In addition to differences in alpha diversity, the microbial community composition and
192 structure on different biotic and inert substrates showed differences in beta diversity. First, the
193 same substrates clustered in a PCoA analysis based on weighted UniFrac distances (Fig 2), e.g.
194 rock substrates clustered with the glass crucibles, while mussel gill and siphon tissue clustered
195 together. The filtered plankton samples were highly similar to each other, while the anemone and
196 *Prionitis* tissues suggest greater differences among individual hosts. Substrate differences were
197 significant with a permuted ANOVA ($F_{5,18}=6.570$, $p<0.001$) when we grouped substrates into 6
198 categories (anemone, red alga, plankton, mussel shell, mussel internal tissue, and inert
199 substrates). Analysis of the differences among those 6 categories showed that each differed
200 significantly from one or several others, except for filtered POM, which did not statistically
201 differ from any other group.

202 A second test indicating beta diversity differences among substrates was revealed in an
203 ANOVA on OTU abundance. There were 10 OTUs that differed significantly in abundance
204 (Bonferroni corrected ANOVA, $p<0.05$, Fig 3) and these distinctions came primarily from their
205 abundance in association with macrobiota. For example, *Moritella* and *Aliivibrio* were found on
206 mussel shell and gill tissue, while Cyanobacteria were primarily associated with algal fronds.

207 Further, the beta diversity differences we detected among substrates were a function of the
208 abundance of OTUs among samples. By comparing OTUs shared between substrates in a

network plot, it was evident that the abundant OTUs that were detected at least 5000 times were generally common to all substrates (Fig 4a). In contrast, when we examined shared diversity among rare OTUs (those detected only 5 to 10 times across the dataset, Fig 4b), there was strong differentiation among substrates, indicating that the rare OTUs were responsible for the majority of the compositional differences between the microbial communities associated with different substrates.

The presence of mussels has little impact on microbial assemblages in tide pools at Second Beach

In experiments performed at Second Beach, the biogeochemical parameters of tidepools were affected by the presence or absence of mussels (Supplementary Table S1). A principal components analysis that included the ammonium regeneration and removal rates (Pather *et al.*, 2014), the maximum seawater pH and dissolved oxygen, ammonium, nitrate, nitrite and phosphorus measured in the tidepools over both daytime and nighttime low tides indicated that the first principal component explained 81.7% of the variance and differed among mussel versus no mussel tidepools ($p=0.049$, Supplementary Figure S1). Rock had more than twice the microbial diversity of coverslips (Rock=3,727 OTUs, Coverslips=1,750 OTUs; $F_{1,18}=140.59$, $p<0.001$), perhaps reflecting greater time in the environment. Both types of substrata maintained more diverse and equitable community profiles in tidepools where mussels were removed, than in those with mussels present (Rock=3,282 OTUs, Coverslips=1,343 OTUs; Fig 1b, Two-way ANOVA, $F_{1,18}=12.759$, $p=0.002$). However, there was no interaction between substrate and mussel presence ($F_{1,18}=0.013$, $p=0.909$), indicating that the distinction between microbial communities associated with natural rock and glass coverslip communities did not depend upon

the presence of mussels. The equitability with which diversity was distributed was strongly correlated with total diversity ($r=0.920$, $p<0.001$), indicating that when mussels are removed, the equitability of taxa also increases. Further, the microbial communities associated with the rock substrate in tidepools at Second Beach were similar in OTU composition with rock substrate at Tatoosh Island (Fig 1a vs 1b).

A high degree of OTU sharing and community structure similarity were observed between microbial communities associated with rock surface or coverslip samples regardless of whether mussels were present (Figs 5 and 6); this was supported by the absence of OTUs with significantly different relative abundances between tidepools with or without mussels based on Bonferroni-corrected ANOVAs. However, there were fewer shared OTUs among rock samples when OTUs that were rare were considered against OTUs that were common (26.7 % versus 96.9% shared). There was no relationship in the degree of OTU sharing with mussel presence or absence. Indeed, in contrast to our comparison of Tatoosh Island substrates, both rock and coverslip samples showed no differentiation between common or rare OTUs as a function of mussels, and the clustering of nodes was highly similar. Further, the presence of mussels was not associated with any changes in the relative weight of deterministic and stochastic forces governing community assembly. OTU co-occurrence was significantly non-random on rock surfaces (c-score analysis; $p<0.01$), and this pattern did not change with mussels.

Patterns in the distribution of a subset of taxa involved in nitrogen metabolism do show responses to the presence of mussels in tide pools on Second Beach.

Nitrifying taxa were at low incidence throughout our samples, while taxa related to nitrate reduction were found in almost every sample (Table 2). Filtered plankton had the highest

incidence of nitrifying OTUs as a result of the genus *Paracoccus*. The relatively high incidence of denitrifying taxa in the anemone *Anthopleura* was primarily driven by matches in the Campylobacteraceae, the group including *Campylobacter*, a taxon that harbors *nrf* genes (Pittman and Kelly, 2005). Although most samples also had genera in the Planctomycetes, only some are known to perform anammox (Fuerst and Sagulenko, 2011), and these genera were either absent from our samples or characterized as “other Planctomycetes”. We tallied these OTUs for Table 2, but did not perform statistical analyses and interpret these with caution. OTUs associated with nitrogen metabolism, as inferred from PICRUSt, were found across all Tatoosh substrates (Table 2).

The percentage of taxa associated with nitrogen transformations and residing on mussel shells was compared between the 16s rRNA amplicon data and existing shotgun metagenomic data (Pfister *et al.*, 2010). The metagenomic data revealed a greater proportion of nitrogen metabolizing taxa, including taxa that were not observed in the amplicon data (Table 2). An analysis of SEED Subsystem functions for the two mussel shell metagenomes yielded estimates of 1.4 % and 1.3% of the 68,676 and 63,950 proteins with functions known to be related to nitrogen metabolism that were discovered in each sample. The PICRUSt analysis, which inferred functional gene presence, indicated that 0.83% of the OTUs discovered were related to nitrogen function (Table 2), a value closer to the metagenome discovery rate than our discovery analyzing only the taxa in Table 1 with 16S rRNA data, and likely larger due to the inclusion of many nitrogen-metabolizing taxa in addition to those in Table 1.

When we compared the presence of OTUs with taxa associated with certain nitrogen metabolisms (e.g. Table 1) in our experimental tidepools, we found that the presence of mussels increased the incidence of putative nitrate reducing taxa on rock substrate, but mussels had no

effect on putative nitrifiers (Table 3). The nitrogen metabolisms on inert substrates that were inferred through PICRUSt also did not differ between tidepools with or without mussels (Table 3). The maximum dissolved inorganic nitrogen in each tidepool did not correlate with the observed diversity ($r=-0.318$, $p=0.371$). Over all 46 samples from Tatoosh and Second Beach that we analyzed, the discovery rate of OTUs with known nitrogen transformations (Table 1) was unrelated to the observed diversity ($r=-0.127$, $p=0.399$, $n=46$).

DISCUSSION

All sampled substrates had microbial assemblages, which showed variation in important taxonomic and functional properties. Differences in diversity and OTU composition may reflect differences in colonization preference, temporal dynamics, the duration in the environment to accrue microbes, and host-specific interactions on the longer-lived macrobiota. The red alga, the anemone and the mussels are all long-lived and could have been present for years prior to our sampling and had ample time to accumulate a microbial assemblage. The clonal nature of the anemone *Anthopleura elegantissima* might even provide nearly immortal tissue for microbial proliferation. Relatively high diversity on rocks and crucible lids may indicate a number of micro-niches, perhaps as a result of surface irregularities on these substrates. The biotic substrates we sampled are especially likely to have a number of microbial niches, including microtopography and strong oxygen gradients (Heisterkamp et al. 2013).

Similarities between the filtered plankton and the animal tissue may have resulted from mussel and anemone tissue harboring planktonic microbiota, due to their feeding activities. Although increased sample size will be needed to quantify the extent of within-host heterogeneity, the animals we examined were dominated by Gammaproteobacteria, including the

symbiont-bearing anemone. In contrast, coral reef invertebrates with eukaryotic photosynthetic symbionts have been shown to be dominated by Alphaproteobacteria (e.g. Bourne *et al.*, 2013), suggesting differential colonization drivers in different symbiont-bearing invertebrates. Nevertheless, many specialized symbionts of marine bivalves fall within the Gammaproteobacteria (Stewart and Cavanaugh, 2006; Newton *et al.*, 2008).

The great similarity in microbial community structure on inert substrates in tidepools regardless of the presence or absence of mussels (Figure 1b, 3, 5, 6), suggests that the enhanced nitrogen regeneration and uptake that has been demonstrated with mussels (Pather *et al.*, 2014) could be due to the microbes on the mussels themselves. The only important functional difference that we found was a greater incidence of nitrate reducing taxa on rocks in the presence of mussels (Table 3), a result likely explained by the ability of mussels to temporarily reduce oxygen levels to the point where reducing processes are favored. OTUs associated with nitrification did not differ. Although mussel presence was associated with a decreased alpha diversity on inert substrates (Fig 1), this did not lead to significant differences in the network of OTUs that were shared across tidepools with and without mussels and thus did not affect beta diversity (Fig 6). Hence, the macrobiota did not drive any shifts in community structure on nearby inert substrates, even though tissue-associated microbial communities were significantly differentiated. Both 16S rRNA and shotgun metagenomic analyses (Table 2) suggest that macrobiota host OTUs that are important for nitrogen cycling. This corroborates work by Welsh and Castadelli (2004) that showed nitrogen metabolism was hosted within a related mussel species. Similarly, deep-sea mussels are known to host nitrogen-utilizing symbionts (Lee and Childress, 1995). In contrast with the shell surface microbes, those with mussel gill and siphon were less diverse (Fig 1). Thus, the elevated nitrite concentrations in tidepools with mussels,

suggesting increased nitrification (Pfister, 2007), may be the result of a direct effect of habitat provisioning for microbes by the macrobiota, rather than simply a microbial community shift on other substrates due to nutrient provisioning. Alternatively, the nitrogen increase due to mussels may not be enough to drive microbial community differences. Indeed, neither microbial community composition nor the expression of several functional genes in coastal sediments showed major changes in response to nutrient perturbations (Bowen *et al.*, 2011). Thus, further exploration of macrobiota as the repository of microbial function, not just as providers of nutrient resources, is warranted.

We note that nitrogen-transforming taxa were found with a higher incidence in the metagenomic data than the 16S rRNA V4 amplicon sequencing of mussel shells. Although there are technical differences between the two sequencing methodologies that could lead to detection differences, such as GC bias (Ross *et al.*, 2013), metagenomic data directly identify genes for nitrogen transformations that are relatively conserved and identifiable (independent of host phylogeny), even if the taxa hosting these genes are uncharacterized. Many of the taxa identified in the amplicon survey are not closely related to cultured isolates, and their N-cycling status is unknown. Although this limits our ability to infer function from phylogeny, it nevertheless is an analysis that may become increasingly insightful as our knowledge of sequence-based diversity increases. While we recognize that OTU analysis using 16S rRNA data predictions may not yet be the strongest lens to detect function, the increased characterization of taxa involved in nitrogen metabolisms (Ward *et al.* 2011, Munn 2011), and the analyses of 16S data with PICRUST (Table 2) suggests that functional inference is possible.

Marine microbial community structure has been shown to be composed of a multitude of rare taxa that have deep phylogenetic differences; the extent to which this ‘rare biosphere’ (Sogin *et*

al., 2006) drives community function is unknown, though deep sequencing efforts have revealed that there could be the equivalent of a ‘seed bank’ of rare taxa that are persistent with only relative abundance changing through time and space (Lennon and Jones, 2011; Caporaso *et al.*, 2012; Gibbons *et al.*, 2013). In some ecosystems, rare taxa have also been shown to be as metabolically active as common species (Hamasaki *et al.*, 2007), suggesting that rarity does not preclude functional importance. Although seawater and rock had the highest OTU diversity, the differences among filtered plankton and inert substrates versus biogenic substrates (mussels, seaweed, anemone) sampled *in situ* demonstrated that macrobiota enhance beta diversity by hosting unique OTUs (Fig 3, Fig 4b), presumably via the provisioning of unique habitats or resources. It is possible that particular microbial taxa end up in association only with intertidal macrobiota, though the selectivity of these associations requires further temporal and spatial sampling. Our results with these benthic macrobiota are, however, in direct contrast to analyses of seawater where the patterns of beta diversity did not differ among abundant and rare taxa (Amaral-Zettler *et al.*, 2010).

The demonstration that macrobiota host a unique microbial community compared to the water around them is supported by this study and other recent work with tadpoles (McKenzie *et al.*, 2012) and marine algae (Michelou *et al.*, 2013). Although it has been recognized for several decades that benthic invertebrates in deep sea environments host unique taxa, it may be that benthic macrobiota common to large parts of the ocean are also repositories for unique microorganisms (Grossart *et al.*, 2005; Lee *et al.*, 2011; Bengtsson *et al.*, 2012; Jackson *et al.*, 2012) and loci for important biogeochemistry (Martinez-Garcia *et al.*, 2008; Heisterkamp *et al.*, 2013). The rocky intertidal flora and fauna, though relatively well-understood in terms of the interactions among macrobiota, likely also interact and mediate productivity via a rich microbial

community that we are just beginning to describe. These intertidal macrobiota may also harbor a unique set of taxa adapted to host-associated niches, thereby promoting microbial community diversity in the coastal ocean.

ACKNOWLEDGEMENTS

S. Owens provided expertise in the lab, and G. Caporaso and J. Stombaugh provided critical code. We thank A. Olson for help in the field, O. Moulton for lab assistance, and M. Coleman for comments on the ms. We are grateful to the Makah Tribal Council for access to their lands.

REFERENCES

- Amaral-Zettler L, Artigas LP, Baross J, Loka Bharathi PA, Boetius A, Chandramohan D, Herndl G, Kogure K, Neal P, Pedros-Alio C, Ramette A, Schouten S, Stal L, Thessen A, de Leeuw J, Sogin M. 2010. A Global Census of Marine Microbes. In: McIntyre, A (ed). *Life in the World's Oceans: Diversity, Distribution, and Abundance*. Blackwell Publishing: London. pp 223-245.
- Barber RT, Smith RL. 1981. Coastal upwelling ecosystems. In: A. Longhurst (Ed.), *Analysis of Marine Ecosystems*, Academic Press: New York, pp. 31–68.
- Barberán A, Bates ST, Casamayor EO, Fierer N. 2012. Using network analysis to explore co-occurrence patterns in soil microbial communities. *International Society for Microbial Ecology Journal* 6: 343–351

- 391 Bengtsson MM, Sjøtun K, Lanzén A, Øvreås L. 2012. Bacterial diversity in relation to secondary
392 production and succession on surfaces of the kelp *Laminaria hyperborea*. *International*
393 *Society for Microbial Ecology Journal* **7**: 1452–1458.
- 394 Bergschneider, H. and Muller-Parker, G. 2008. Nutritional role of two algal symbionts in the
395 temperate sea anemone *Anthopleura elegantissima* brandt. *Biological Bulletin* **215**: 73–88.
- 396 Bertagnolli A.D., Treusch A.H., Mason O.U., Stingl U., Vergin K.L., Chan F. 2011. Bacterial
397 diversity in the bottom boundary layer of the inner continental shelf of Oregon, USA.
398 *Aquatic Microbial Ecology* **64**: 15-25.
- 399 Bourne D.G., Dennis P.G., Uthicke S, Soo R.M., Tyson G.W., Webster N. 2013. Coral reef
400 invertebrate microbiomes correlate with the presence of photosymbionts. *International*
401 *Society for Microbial Ecology Journal* **7**: 1452–1458.
- 402 Bowen, J.L., Ward B.B., Morrison H.G., Hobbie J.E., Valiela I., Deegan L.A., Sogin M.L. 2011.
403 Microbial community composition in sediments resists perturbation by nutrient enrichment.
404 *International Society for Microbial Ecology Journal* **5**: 1540-1548
- 405 Caporaso J.G., Kuczynski J., Stombaugh J., Bittinger K., Bushman F.D., Costello E.K., Fierer N,
406 Gonzalez Peña A, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE,
407 Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder M, Reeder J,
408 Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R.
409 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature*
410 *Methods* **7**:335-336.

- 411 Caporaso J.G., Bittinger K., Bushman F.D., DeSantis T.Z., Andersen G.L., Knight R. 2010.
412 PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**:
413 266-7.
- 414 Caporaso J.G., Paszkiewicz K., Field D., Knight R., Gilbert J.A. 2012. The Western English
415 Channel contains a persistent microbial seed bank. *International Society for Microbial
416 Ecology Journal* **6**:1089–1093.
- 417 DeChaine E.G., Cavanaugh C.M. 2006 Symbioses of Methanotrophs and Deep-Sea Mussels
418 (Mytilidae: Bathymodiolinae). In: Overmann, J (ed). *Molecular Basis of Symbiosis. Progress
419 in Molecular and Subcellular Biology*. Volume **41**: 227-249. Springer.
- 420 Estes, J.A., Duggins, D.O. 1995. Sea otters and kelp forests in Alaska: generality and variation in
421 a community ecological paradigm. *Ecological Monographs* **65**: 75-100.
- 422 Fan, L., Liu, M., Simister, R., Webster, N.S., and Thomas, T. 2013. Marine microbial symbiosis
423 heats up: the phylogenetic and functional response of a sponge holobiont to thermal stress.
424 *International Society for Microbial Ecology Journal* **7**: 991–1002.
- 425 Fuerst, J.A., Sagulenko, E. 2011. Beyond the bacterium: Planctomycetes challenge our concepts
426 of microbial structure and function. *Nature Reviews Microbiology* **9**: 403-413.
- 427 Gibbons, S.M., Caporaso, J.G., Pirrung, M., Field, D., Knight, R., Gilbert J.A. 2013. Evidence
428 for a persistent microbial seed bank throughout the global ocean. *Proceedings of the National
429 Academy of Sciences, USA* **110**:4651-4655.

- 430 Gilbert, J.A., Steele, J.A., Caporaso, J.G., Steinbrück, L., Reeder, J., Temperton, B., Huse S,
431 McHardy AC, Knight R, Joint I, Somerfield P, Fuhrman JA, Field D. 2012. Defining
432 seasonal marine microbial community dynamics. *International Society for Microbial Ecology*
433 *Journal* **6**: 298–308.
- 434 Grossart, H.-P., Levold, F., Allgaier, M., Simon, M., and Brinkhoff, T. 2005. Marine diatom
435 species harbour distinct bacterial communities. *Environmental Microbiology* **7**: 860–873.
- 436 Hamasaki, K, Taniguchi, A, Tada Y, Long, R. A., Azam, F. 2007. Actively growing bacteria in
437 the inland Sea of Japan, identified by combined bromodeoxyuridine immunocapture and
438 denaturing gradient gel electrophoresis. *Applied Environmental Microbiology* **73**: 2787–
439 2798.
- 440 Heisterkamp, I.M., Schramm, A., Larsen, L.H., Svenningsen, N.B., Lavik, G., de Beer, D., Stief,
441 P. 2013. Shell biofilm-associated nitrous oxide production in marine molluscs: processes,
442 precursors and relative importance. *Environmental Microbiology* **15**:1943-55.
- 443 Jackson, S.A., Kennedy, J., Morrissey, J.P., O’Gara, F., Dobson, A.D. 2012. Pyrosequencing
444 reveals diverse and distinct sponge-specific microbial communities in sponges from a single
445 geographical location in Irish waters. *Microbial Ecology* **64**: 105-16.
- 446 Kvennefors, E.C.E., Sampayo, E., Kerr, C., Vieira, G., Roff, G., Barnes, A.C. 2012. Regulation
447 of bacterial communities through antimicrobial activity by the coral holobiont. *Microbial*
448 *Ecology*. **63**: 605–618.
- 449 Langille, M.G.I., Zaneveld, J., Caporaso, J.G., McDonald, D., Knights, D., Reyes JA, Clemente
450 JC, Burkepile DE, Vega Thurber RL, Knight R, Beiko RG, Huttenhower C. 2013. Predictive

- 451 functional profiling of microbial communities using 16S rRNA marker gene sequences.
- 452 *Nature Biotechnology* **31**: 814-821.
- 453 Lee, O.O., Wang, Y., Yang, J., Lafi, F.F., Al-Suwailem, A., Qian, P.Y. 2011. Pyrosequencing
- 454 reveals highly diverse and species-specific microbial communities in sponges from the Red
- 455 Sea. *International Society for Microbial Ecology Journal* **5**: 650-64.
- 456 Lee, R.W., Childress, J.J. 1995. Assimilation of inorganic nitrogen by seep mytilid 1a, an
- 457 undescribed deep-sea mussel containing methanotrophic endosymbionts: fate of assimilated
- 458 nitrogen and the relation between methane and nitrogen assimilation. *Marine Ecology*
- 459 *Progress Series* **123**: 137-148.
- 460 Lennon, J.T., Jones, S.E. 2011. Microbial seed banks: the ecological and evolutionary
- 461 implications of dormancy. *Nature Reviews Microbiology* **9**: 119-130.
- 462 Lozupone, C., Lladser, M.E., Knights, D., Stombaugh, J., Knight, R. 2011. UniFrac: an effective
- 463 distance metric for microbial community comparison. *International Society for Microbial*
- 464 *Ecology Journal* **5**: 169–172.
- 465 Martínez-García, M., Stief, P., Díaz-Valdés, M., Wanner, G., Ramos-Esplá, A., Dubilier, N.,
- 466 Anton J. 2008. Ammonia-oxidizing Crenarchaeota and nitrification inside the tissue of a
- 467 colonial ascidian. *Environmental Microbiology* **10**: 2991-3001.
- 468 McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., DeSantis, T.Z., Probst, A., Anderson
- 469 GL, Knight RL, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit ranks
- 470 for ecological and evolutionary analyses of bacteria and archaea. *International Society for*
- 471 *Microbial Ecology Journal* **6**: 610–618.

- McKenzie, V.J., Bowers R., Fierer N., Knight R., Lauber C.L. 2012. Co-habiting amphibian species harbor unique skin bacterial communities in wild populations. *International Society for Microbial Ecology Journal* **6**: 588-596. doi: 10.1038/ismej.2011.129.
- Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E.M., Kubal, M., Paczian T, Rodriguez A, Stevens R, Wilke A, Wilkening J, Edwards RA. 2008. The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**: 386.
- Michelou, V.K., Caporaso, J.G., Knight, R., Palumbi, S.R. 2013. The ecology of microbial communities associated with *Macrocystis pyrifera*. *PLoS One* 2013 Jun 19;8(6):e67480
- Newton, I.L., Girguis, P.R., Cavanaugh, C.M. 2008. Comparative genomics of vesicomylid clam (Bivalvia: Mollusca) chemosynthetic symbionts. *BMC Genomics* **9**: 585. doi: 10.1186/1471-2164-9-585.
- Miranda, L.N., Hutchison, K., Grossman, A.R., and Brawley, S.H. 2013. Diversity and abundance of the bacterial community of the red macroalga *Porphyra umbilicalis*: Did bacterial farmers produce macroalgae? *PLoS ONE* **8**: e58269.
- Munn, C. 2011. *Marine Microbiology: Ecology and Applications*, 2nd Edition. Garland Science, New York, NY, USA.
- Paine, R.T. 1966. Food web complexity and species diversity. *American Naturalist* **100**: 65–75.
- Pather, S., Pfister, C.A., Post, D.A., Altabet, M.A. 2014. Ammonium cycling in the rocky intertidal: remineralization, removal and retention. *Limnology Oceanography* **59**: 361-372.
- Pfister, C.A. 2007. Tidepool mussels locally increase nutrients and algal growth. *Ecology* **88**: 1647–1653.

- 494 Pfister, C.A., Wootton, J.T., Neufeld, C. 2007. The relative roles of coastal and oceanic
495 processes in determining physical and chemical characteristics of an intensively sampled
496 nearshore system. *Limnology Oceanography* **52**: 1767-1775.
- 497 Pfister, C.A., Meyer, F., Antonopoulos, D.A. 2010 Metagenomic profiling of a microbial
498 assemblage associated with the California mussel, *Mytilus californianus*: a node in networks
499 of carbon and nitrogen cycling. *PLoS ONE* 5(5): e10518. doi:10.1371/journal.pone.0010518.
- 500 Pittman, M.S., Kelly D.J. 2005. Electron transport through nitrate and nitrite reductases in
501 *Campylobacter jejuni*. *Biochemical Society Transactions* **33**: 190-2.
- 502 Price, M.N., Dehal P.S., Arkin A.P. 2009. FastTree: Computing Large Minimum-Evolution
503 Trees with Profiles instead of a Distance Matrix. *Molecular Biology and Evolution* **26**: 1641-
504 1650.
- 505 Qian, P.Y., Dobretsov, S., Dahms, H.U., and Pawlik, J. 2006. Antifouling activity and microbial
506 diversity of two congeneric sponges *Callyspongia* spp. from Hong Kong and the Bahamas.
507 *Marine Ecology Progress Series* **324**: 151–165.
- 508 Ramette, A. 2007. Multivariate analyses in microbial ecology. *FEMS Microbiol Ecology* **62**:
509 142–160.
- 510 Ross, M.G., Russ C., Costello M., Hollinger A., Lennon N.J., Hegarty R., Nusbaum C, Jaffe DB.
511 2013. Characterizing and measuring bias in sequence data. *Genome Biology* **14**:R51.
512 doi:10.1186/gb-2013-14-5-r51
- 513 Secord, D. and Augustine, L. 2000 Biogeography and microhabitat variation in temperate algal-
514 invertebrate symbioses: zooxanthellae and zoochlorellae in two Pacific intertidal sea
515 anemones, *Anthopleura elegantissima* and *A. xanthogrammica*. *Invertebrate Biology* **119**:

516 139–146.

517 Sogin, M.L., Morrison, H.G., Huber, J.A., Mark Welsh, D., Huse, S.M., Neal, P.R., Arrieta JM,
518 Herndl GJ. 2006 Microbial diversity in the deep sea and the underexplored “rare biosphere”.
519 *Proceedings of the National Academy of Science, USA* **103**: 12115–12120.

520 Stahl, D.A., Lane, D.J., Olsen, G.J., Pace, N.R. 1984. Analysis of hydrothermal vent-associated
521 symbionts by ribosomal RNA sequences. *Science* **224**: 409–411.

522 Stewart, F.J., Cavanaugh C.M. 2006 Symbiosis of thioautotrophic bacteria with *Riftia*
523 *pachyptila*. *Progress in Molecular and Subcellular Biology*. **41**: 197-225.

524 Taylor, M.W., Radax, R., Steger, D., and Wagner, M. 2007. Sponge-Associated
525 microorganisms: evolution, ecology, and biotechnological Potential. *Microbiology and*
526 *Molecular Biology Reviews* **71**: 295–347.

527 Telford, R.J., Vandvik, V., Birks, H.J. 2006. Dispersal limitations matter for microbial
528 morphospecies. *Science* **312**: 1015.

529 Ward, B.B., D. J. Arp, M. J. Klotz, eds. 2011. *Nitrification*. John Wiley and Sons, Hoboken, NJ,
530 USA.

531 Webster, N.S. and Taylor, M.W. 2012. Marine sponges and their microbial symbionts: love and
532 other relationships. *Enviromental Microbiology* **14**: 335–346.

533 Wegner, K.M., Volkenborn, N., Peter, H., and Eiler, A. 2013. Disturbance induced decoupling
534 between host genetics and composition of the associated microbiome. *BMC Microbiology* **13**:
535 252.

- 536 Welsh, D.T, Castaldelli, G. 2004. Bacterial nitrification activity directly associated with isolated
537 benthic marine animals. *Marine Biology* **144**: 1029-1037.
- 538 Wootton, J.T. 1994. Predicting direct and indirect effects: An integrated approach using
539 experiments and path analysis. *Ecology* **75**: 151-165.
- 540 Zurel, D., Benayahu, Y., Or, A., Kovacs, A., and Gophna, U. 2011. Composition and dynamics
541 of the gill microbiota of an invasive Indo-Pacific oyster in the eastern Mediterranean Sea.
542 *Environmental Microbiology* **13**: 1467–1476.

545
546

FIGURE LEGENDS

547 Figure 1. The proportional representation of OTUs among the major microbial groups (colored
548 bars), with the overall mean observed OTU richness (+SE) among all substrate types at a.
549 Tatoosh Island, and b. in tidepools where natural rock substrate and coverslips were sampled in
550 the context of an experimental removal of mussels. The substrates in Fig 1a showed significant
551 differences in observed richness (ANOVA, $F_{7,16}=4.968$, $p=0.004$) with rocks ($n=3$), crucible lids
552 ($n=9$) and filtered plankton ($n=3$) showing the greatest richness while the lowest observed
553 richness was associated with mussel gill ($n=3$) and siphon ($n=1$) tissue. OTU richness of mussel
554 shell ($n=2$), anemone ($n=2$), and red algae ($n=2$) was intermediate to the others. In b. Tidepools
555 with mussels removed had greater OTU richness than those with mussels (Two-Way ANOVA,
556 $F_{1,18}=12.759$, $p=0.002$) while rock had over twice the OTU richness of coverslips ($F_{1,18}=140.59$,
557 $p<0.001$); there was no interaction between substrate and mussel presence.

558
559 Figure 2. A PCoA of the OTU beta diversity of substrates on Tatoosh Island, demonstrating the
560 clustering among the different microbial assemblages associated with each substrate. The
561 weighted UniFrac metric was used to incorporate relative abundance; the first axis explained
562 40.2% of the variance, while the second explained 14.8%. Differences among substrates were
563 significant (PERMANOVA, $F_{5,18}=6.570$, $p<0.001$), and groupings that included anemone,
564 *Prionitis*, mussel shell, mussel tissue, and inert substrates were differentiated while plankton
565 were indistinguishable from all.

566

567 Figure 3. The relative abundance of the 10 OTUs that differed among the Tatoosh Island
568 substrates (Boniferroni-corrected ANOVA, $p<0.05$).

569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593

Figure 4. Shared OTU diversity among microbes sampled from the substrate groupings as in Fig 1, 2 and 3 at Tatoosh Island and portrayed as a spring-embedded layout, where OTUs that are in common bring nodes or samples together and OTUs that are distinct repel nodes. In a., only common OTUs detected more than 5,000 times are included, while b. shows only rare OTUs that were present 5 to 10 times across the entire dataset.

Figure 5. A PCoA of the OTU diversity of tidepool rock (n=10) versus coverslip (n=12) substrates at Second Beach, demonstrating strong clustering among the microbial assemblages from the two substrates, while the presence of mussels (filled symbols) versus removal of mussels (open symbols) were not a factor for explaining beta diversity. Using weighted UniFrac, the first axis explained 46.5% of the variance, while the second explained 20.3%.

Figure 6. Shared OTU diversity among a. microbes sampled from tidepool rock versus coverslip (lighter green) substrates and b. samples distinguished by whether mussels were present (blue) or absent (red) from tidepools at Second Beach. The spring-embedded layout shows OTUs that are in common bring nodes or samples together and OTUs that are distinct repel nodes. Only common OTUs greater than >5000 are included. Analyses of relatively rare OTUs did not change the network pattern.

Table 1 (on next page)

The genera associated with different microbial nitrogen transformations.

Table 1. The genera associated with different microbial nitrogen transformations that were searched in all samples via the Greengenes database. For nitrification, any taxa associated with ammonium or nitrite oxidation are included, while nitrate reduction includes taxa for denitrification and DNRA. For anammox, we searched for taxa with “Candidatus” status.

§Due to the abundance and functional diversity of the *Vibrio* genus and the possibility that many *Vibrio* species are not involved in nitrate reduction, we did not include *Vibrio* OTUs in Tables 2 and 3.

Table 1. The genera associated with different microbial nitrogen transformations that were searched in all samples via the Greengenes database. For nitrification, any taxa associated with ammonium or nitrite oxidation are included, while nitrate reduction includes taxa for denitrification and DNRA. For anammox, we searched for taxa with “Candidatus” status. §Due to the abundance and functional diversity of the *Vibrio* genus and the possibility that many *Vibrio* species are not involved in nitrate reduction, we did not include *Vibrio* OTUs in Tables 2 and 3.

nitrification	Nitrate reduction	anammox
<i>Alcaligenes</i>	<i>Azospirillum</i>	“Candidatus”
<i>Nitrobacter</i>	<i>Campylobacter</i>	
<i>Nitrococcus</i>	<i>Desulforvibrio</i>	
<i>Nitrolancetus</i>	<i>Nitratifractor</i>	
<i>Nitrosococcus</i>	<i>Nitratiruptor</i>	
<i>Nitrosolobos</i>	<i>Paracoccus</i>	
<i>Nitrosomonas</i>	<i>Rhodobacter</i>	
<i>Nitrosopumilis</i>	<i>Sulfospirillum</i>	
<i>Nitrosovibrio</i>	<i>Wolinella</i>	
<i>Nitrospina</i>	<i>Vibrio</i> §	
<i>Nitrospira</i>		
<i>Nitrotoga</i>		
<i>Paracoccus</i>		

Table 2 (on next page)

The OTUs discovered in each substrate type at Tatoosh Island associated with 3 broad nitrogen transformations.

Table 2. The mean percent of OTUs discovered in each substrate type that is associated with each of the 3 broad nitrogen transformations (taxa listed in Table 1) or overall nitrogen metabolism (PICRUSt) at Tatoosh Island. No “Candidatus” were found in the 16S; the anammox category contains Planctomycetes as an estimate of anammox potential only. Mussel shell samples were analyzed with both the V4 region of the 16S rRNA as well as through shotgun metagenomics.

Table 2. The mean percent of OTUs discovered in each substrate type that is associated with each of the 3 broad nitrogen transformations (taxa listed in Table 1) or overall nitrogen metabolism (PICRUSt) at Tatoosh Island. No “Candidatus” were found in the 16S; the anammox category contains Planctomycetes as an estimate of anammox potential only. Mussel shell samples were analyzed with both the V4 region of the 16S rRNA as well as through shotgun metagenomics.

<u>Substrate</u>	Substrate type	Nitrification		Nitrate reduction		Anammox		N metabolism
		<u>16s</u>	<u>Meta-genome</u>	<u>16s</u>	<u>Meta-genome</u>	<u>16s</u>	<u>Meta-genome</u>	<u>PICRUSt</u>
Mussel siphon	biogenic	0.0000		0.0326		0.0008		0.8811
Mussel gill	biogenic	0.0000		0.4159		0.0009		0.9295
Mussel shell	biogenic	0.0000	0.4604	0.0049	0.9882	0.0043	0.1711	0.8344
filtered plankton	biogenic	0.0102		0.1851		0.0391		0.7636
<i>A. elegantissima</i> (anemone)	biogenic	0.0000		1.8961		0.0804		0.8658
<i>Prionitis</i> (red alga)	biogenic	0.0000		0.4278		0.0399		0.6638
rock	inert	0.0048		0.0551		0.0390		0.7119
crucible lid	artificial	0.0094		0.2120		0.0218		0.6325

Table 3(on next page)

OTUs discovered on inert substrates in experimental Second Beach tidepools associated with 3 broad nitrogen transformations.

Table 3. The mean percent of OTUs discovered on inert substrates in experimental Second Beach tidepools that were associated with each of the 3 broad nitrogen transformations (taxa listed in Table 1) or overall nitrogen metabolism (PICRUSt). P-values are listed for t-tests for a significant difference on each substrate as a function of mussel presence. The only significant contrast was the greater incidence of OTUs associated with nitrate reduction on natural rock substrate in tidepools with mussels.

Table 3. The mean percent of OTUs discovered on inert substrates in experimental Second Beach tidepools that were associated with each of the 3 broad nitrogen transformations (taxa listed in Table 1) or overall nitrogen metabolism (PICRUSt). P-values are listed for t-tests for a significant difference on each substrate as a function of mussel presence. The only significant contrast was the greater incidence of OTUs associated with nitrate reduction on natural rock substrate in tidepools with mussels.

Tidepool substrates	Nitrification		Nitrate reduction		N metabolism	
	16s	T-test	16s	T-test	PICRUSt	T-test
Natural rock substrate with mussels (n=5)	0.0001	p=0.284	0.1890	p=0.043*	0.7433	p=0.952
Natural rock substrate without mussels (n=5)	0.0011		0.0151		0.7422	
Coverslip with mussels (n=6)	0.0004	p=0.456	0.0737	p=0.471	0.7421	p=0.218
Coverslip without mussels (n=6)	0.0019		0.1785		0.7084	

Figure 1

The proportional representation of OTUs and the mean observed OTU richness among substrates sampled.

Figure 1. The proportional representation of OTUs among the major microbial groups (colored bars), with the overall mean observed OTU richness (+SE) among all substrate types at a.

Tatoosh Island, and b. in tidepools where natural rock substrate and coverslips were sampled in the context of an experimental removal of mussels. The substrates in Fig 1a showed significant differences in observed richness (ANOVA, $F_{7,16}=4.968$, $p=0.004$) with rocks ($n=3$), crucible lids ($n=9$) and filtered plankton ($n=3$) showing the greatest richness while the lowest observed richness was associated with mussel gill ($n=3$) and siphon ($n=1$) tissue. OTU richness of mussel shell ($n=2$), anemone ($n=2$), and red algae ($n=2$) was intermediate to the others. In b. Tidepools with mussels removed had greater OTU richness than those with mussels (Two-Way ANOVA, $F_{1,18}=12.759$, $p=0.002$) while rock had over twice the OTU richness of coverslips ($F_{1,18}=140.59$, $p<0.001$); there was no interaction between substrate and mussel presence.




Figure 2

A PCoA of the OTU beta diversity of substrates on Tatoosh Island.

Figure 2. A PCoA of the OTU beta diversity of substrates on Tatoosh Island, demonstrating the clustering among the different microbial assemblages associated with each substrate. The weighted UniFrac metric was used to incorporate relative abundance; the first axis explained 40.2% of the variance, while the second explained 14.8%. Differences among substrates were significant (PERMANOVA, $F_{5,18}=6.570$, $p<0.001$), and groupings that included anemone, *Prionitis*, mussel shell, mussel tissue, and inert substrates were differentiated while plankton were indistinguishable from all.




Figure 3

The relative abundance of the 10 OTUs that differed among the Tatoosh Island substrates.

Figure 3. The relative abundance of the 10 OTUs that differed among the Tatoosh Island substrates (Boniferroni-corrected ANOVA, $p < 0.05$).




Figure 4

Shared OTU diversity among microbes sampled from the substrate groupings at Tatoosh Island and portrayed as a spring-embedded layout.

Figure 4. Shared OTU diversity among microbes sampled from the substrate groupings as in Fig 1, 2 and 3 at Tatoosh Island and portrayed as a spring-embedded layout, where OTUs that are in common bring nodes or samples together and OTUs that are distinct repel nodes. In a., only common OTUs detected more than 5,000 times are included, while b. shows only rare OTUs that were present 5 to 10 times across the entire dataset.




Figure 5

A PCoA of the OTU diversity of tidepool rock versus coverslip substrates at Second Beach, WA.

Figure 5. A PCoA of the OTU diversity of tidepool rock (n=10) versus coverslip (n=12) substrates at Second Beach, demonstrating strong clustering among the microbial assemblages from the two substrates, while the presence of mussels (filled symbols) versus removal of mussels (open symbols) were not a factor for explaining beta diversity. Using weighted UniFrac, the first axis explained 46.5% of the variance, while the second explained 20.3%.




Figure 6

Shared OTU diversity among microbes sampled from tidepool rock versus coverslip substrates in tidepools at Second Beach, WA.

Figure 6. Shared OTU diversity among a. microbes sampled from tidepool rock versus coverslip (lighter green) substrates and b. samples distinguished by whether mussels were present (blue) or absent (red) from tidepools at Second Beach. The spring-embedded layout shows OTUs that are in common bring nodes or samples together and OTUs that are distinct repel nodes. Only common OTUs greater than >5000 are included. Analyses of relatively rare OTUs did not change the network pattern.

