

**A peer-reviewed version of this preprint was published in PeerJ on 12 May 2015.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.935) (peerj.com/articles/935), which is the preferred citable publication unless you specifically need to cite this preprint.

Blakeway D, Hamblin MG. 2015. Self-generated morphology in lagoon reefs. PeerJ 3:e935 <https://doi.org/10.7717/peerj.935>

## Self-generated morphology in lagoon reefs

The three-dimensional form of a coral reef develops through interactions and feedbacks between its constituent organisms and their environment. Reef morphology therefore contains a potential wealth of ecological information, accessible if the relationships between morphology and ecology can be decoded. Traditionally, reef morphology has been attributed to external controls such as substrate topography or hydrodynamic influences. Little is known about inherent reef morphology in the absence of external control. Here we use reef growth simulations, based on observations in the cellular reefs of Western Australia's Houtman Abrolhos Islands, to show that reef morphology is fundamentally determined by the mechanical behaviour of the reef-building organisms themselves—specifically their tendency to either remain in place or to collapse. Reef-building organisms that tend to remain in place, such as massive and encrusting corals or coralline algae, produce nodular reefs, whereas those that tend to collapse, such as branching *Acropora*, produce cellular reefs. The purest reef growth forms arise in sheltered lagoons dominated by a single type of reef builder, as in the branching *Acropora*-dominated lagoons of the Abrolhos. In these situations reef morphology can be considered a phenotype of the predominant reef building organism. The capacity to infer coral type from reef morphology can potentially be used to identify and map specific coral habitat in remotely sensed images. More generally, identifying ecological mechanisms underlying other examples of self-generated reef morphology can potentially improve our understanding of present-day reef ecology, because any ecological process capable of shaping a reef will almost invariably be an important process in real time on the living reef.

# Self-generated morphology in lagoon reefs

David Blakeway, Michael G Hamblin

The three-dimensional form of a coral reef develops through interactions and feedbacks between its constituent organisms and their environment. Reef morphology therefore contains a potential wealth of ecological information, accessible if the relationships between morphology and ecology can be decoded. Traditionally, reef morphology has been attributed to external controls such as substrate topography or hydrodynamic influences. Little is known about inherent reef morphology in the absence of external control. Here we use reef growth simulations, based on observations in the cellular reefs of Western Australia's Houtman Abrolhos Islands, to show that reef morphology is fundamentally determined by the mechanical behaviour of the reef-building organisms themselves—specifically their tendency to either remain in place or to collapse. Reef-building organisms that tend to remain in place, such as massive and encrusting corals or coralline algae, produce nodular reefs, whereas those that tend to collapse, such as branching *Acropora*, produce cellular reefs. The purest reef growth forms arise in sheltered lagoons dominated by a single type of reef builder, as in the branching *Acropora*-dominated lagoons of the Abrolhos. In these situations reef morphology can be considered a phenotype of the predominant reef building organism. The capacity to infer coral type from reef morphology can potentially be used to identify and map specific coral habitat in remotely sensed images. More generally, identifying ecological mechanisms underlying other examples of self-generated reef morphology can potentially improve our understanding of present-day reef ecology, because any ecological process capable of shaping a reef will almost invariably be an important process in real time on the living reef.

David Blakeway<sup>1, 2</sup> and Michael G. Hamblin<sup>3</sup>

<sup>1</sup>School of Earth and Environment, University of Western Australia, Perth, Western Australia

<sup>2</sup>Fathom 5 Marine Research, Lathlain, Western Australia

<sup>3</sup>School of Mechanical and Chemical Engineering, University of Western Australia, Perth, Western Australia

Corresponding author: David Blakeway, Fathom 5 Marine Research, 17 Staines Street, Lathlain, Western Australia, 6100, Australia, fathom5marineresearch at gmail dot com

## Introduction

Coral reefs are large organic structures created over centuries to millennia by relatively small individual organisms. The anatomy of coral reefs varies along a continuum from 'framework' reefs consisting of coral skeletons in growth position (Lowenstam 1950; Fagerstrom 1987) to 'garbage piles' of toppled corals, coral fragments and sediment (Hubbard, Miller & Scaturro 1990; Blanchon, Jones & Kalbfleisch 1997). The three-dimensional form of a reef, particularly a framework reef, is a potential repository of ecological information, because it represents a long-term integration of interactions and feedbacks between the reef-building organisms and their physical, chemical and biological environment (Roberts et al. 1975; Hopley, Smithers & Parnell 2007; Perry 2011). Reef form can perhaps also be considered a distillation of those interactions and feedbacks, in that the most influential processes in the history of the living reef could be expected to leave the greatest imprint on its morphology. Investigating the ecological mechanisms underlying aspects of reef morphology can therefore help provide a context for interpreting present-day ecological processes on reefs (Hopley, Smithers & Parnell 2007).

The primary influences on reef morphology differ across spatial scales, from the inherent forms of reef-building organisms at the small scale to the configuration of continental shelves at the large scale. Over intermediate scales (metres to kilometres) reefs exhibit a great diversity of forms, reflecting the multitude of interacting processes affecting them. But within the diversity is a subset of globally recurring forms, indicating that there are some consistent influences governing reef morphology worldwide (Wells 1957, Stoddart 1969; 1978; Goreau, Goreau & Goreau 1979; Blanchon 2011; Schlager and Purkis 2015). Traditionally, these influences have been envisaged as external controls operating at or above the scale of the morphological features, for example substrate topography (MacNeil 1954, Purdy 1974; Choi & Ginsburg 1982) or the wave field (Munk & Sargent 1954; Roberts 1974). While these factors are undoubtedly responsible for many aspects of reef morphology, they raise an interesting question: what would reefs look like in the absence of such external influences? This question brings the focus down to the reef-building organisms themselves. Because these organisms cumulatively become the reef, there is significant potential for behaviour and events at their scale to be expressed in reef morphology at the intermediate scale. Such 'emergence' of self-organised patterns from small scale processes is ubiquitous in physical and biological systems (Nicolis and Prigogine 1977; Camazine et al. 2001). While it is recognised that coral reefs are likely to exhibit this

45 trait (Drummond and Dugan 1999; Mistr and Bercovici 2003; Rietkirk and van de Koppel 2008;  
46 Blanchon 2011; Schlager and Purkis 2015), it has not been directly demonstrated.

47

48 Lagoons are the most likely setting for inherent reef growth forms to arise, as they are generally flat-  
49 floored and sheltered. Several characteristic lagoon reef forms are repeated worldwide, ranging from  
50 simple mound-like patch reefs to complex cellular<sup>1</sup> reef networks (Fig. 1; Stoddart 1969; Hopley 1982;  
51 Blanchon 2011). Patch reef development can be readily visualised in terms of expansion from a  
52 nucleus, and this mode of growth has been demonstrated repeatedly, from various nuclei including  
53 topographic highs in underlying limestones (Halley et al 1977; Mazzullo et al. 1992), sedimentary  
54 structures (Perry et al. 2012; Novak et al. 2013), or early-colonising corals (Jones 1977; Crame 1981).  
55 Cellular morphology, in contrast, is not an intuitive growth form. Cellular reefs distinctly resemble  
56 negative landforms, particularly karst terrains (terrestrial erosion landforms created in limestone  
57 through dissolution by rainwater). Based on this resemblance, and the recognition that the foundations  
58 of most reefs have been exposed to at least 100,000 years of weathering during Interglacial periods,  
59 cellular reef morphology has long been interpreted as an inheritance from underlying karst (Fairbridge  
60 1948; Purdy 1974; Guilcher 1988; Searle 1994; Macintyre, Precht & Aronson 2000; Purkis et al. 2010;  
61 Kan et al. 2015). However there has always been an opposing view attributing cellular morphology to  
62 reef growth (Dakin 1919; GIE Raro Moana 1985; Collins et al. 1993; Wyrwoll et al. 2006; Barott et al.  
63 2010; Blanchon 2011; Schlager and Purkis 2015). The growth alternative is gradually gaining  
64 acceptance, having been confirmed for the cellular reefs of Mataiva Atoll in French Polynesia (GIE  
65 Raro Moana 1985; Rossfelder 1990) and the Houtman Abrolhos Islands in Western Australia (Collins  
66 et al. 1993; Collins, Zhu & Wyrwoll 1996; 1998; Wyrwoll et al. 2006). Seismic surveys and coring at  
67 Mataiva showed the Holocene reef to be 10 to 20 m thick and demonstrated that, although the reef is  
68 underlain by a karst Tertiary limestone, the karst features are relatively small-scale and were infilled  
69 before submergence, such that the present reef morphology is independent of the substrate (Rossfelder  
70 1990). Similarly, seismic and coring in the Abrolhos lagoons recorded a Holocene reef thickness of 40  
71 m over a flat Last Interglacial grainstone, again demonstrating independence from the substrate  
72 (Collins et al. 1993; Collins, Zhu & Wyrwoll 1996; 1998).

---

<sup>1</sup> Cellular reefs (terminology after Hoskin 1963) are also widely known as reticulate reefs, from the latin *reticulum*: a network or net-like structure. However, the term reticulate has been applied to a variety of lagoon reef forms that probably develop through different mechanisms (Schlager and Purkis 2015). Therefore we consider cellular reefs to be a subdivision of reticulate reefs, distinguished by subcircular depressions as shown in Fig. 1.

73

74 While the seismic and coring work has proven beyond reasonable doubt that the cellular reefs of  
75 Mataiva and the Abrolhos have grown into their present configuration, it has not provided a generally  
76 accepted growth mechanism. Four alternative mechanisms have been proposed. The first, developed  
77 independently by GIE Raro Moana (1985) at Mataiva and Barott et al. (2010) at Millenium Atoll, is the  
78 colonisation of the lagoon floor by networks of massive corals, which are subsequently colonised by  
79 other corals and grow upward to the surface. The second, proposed by Wyrwoll et al. (2006) for the  
80 Abrolhos, is growth to sea level of isolated branching *Acropora* pinnacles and stellate (star-shaped)  
81 reefs, which subsequently extend laterally and coalesce to surround enclosed depressions. The third,  
82 proposed by Blanchon (2011), is a self-limitation mechanism based on negative feedback between reef  
83 growth and water circulation—reef growth reduces water circulation which reduces reef growth, such  
84 that the cellular depressions become self-reinforcing as the surrounding reefs grow. The fourth  
85 mechanism, proposed by Schlager and Purkis (2015) is biological self-organisation through short-range  
86 support and long-range inhibition, conceptually based on Turing's (1952) reaction-diffusion  
87 mechanism of natural pattern formation.

88

89 The alternative mechanisms outlined above are hypothetical and have not been comprehensively  
90 evaluated on real cellular reefs. In this article we use field observations and reef growth simulations to  
91 examine the process of cellular reef development in one of the type examples of cellular reefs  
92 mentioned above, those of the Houtman Abrolhos Islands. These reefs are an ideal case study site due  
93 to their flat pre-Holocene substrate, known accretion history and very pure reef-building community—  
94 cores through the Abrolhos cellular reefs consist almost entirely of branching *Acropora*, with a few  
95 tabular *Acropora* appearing as the reefs approached sea level (Collins et al. 1993; Collins, Zhu &  
96 Wyrwoll 1996; 1998). Furthermore, an apparent sequence of cellular reef development is evident in the  
97 Abrolhos lagoons (Wyrwoll et al. 2006), progressing from pinnacle reefs to stellate reefs surrounding  
98 semi-enclosed depressions to a reef platform surrounding enclosed depressions. Under the assumption  
99 that these are sequential stages of reef development, surveys of the pinnacle-stellate-platform sequence  
100 represent surveys through time. Space-for-time substitution (Darwin 1842; Maxwell 1968; Hopley  
101 1982) can therefore be applied to describe the evolution of the Abrolhos cellular reefs and, potentially,  
102 to reveal their formative mechanism.

103

## Reef survey

### Methods

We examined replicate sites of each stage in a 15 km<sup>2</sup> cellular reef complex known as the Maze in the Easter Group of the Abrolhos (Fig. 2; 28°41'S, 113°49'E). We surveyed fifteen sites in detail and many more in brief visits, including some in the Pelsaert Group to the south and the Wallabi Group to the north. At each of the fifteen Maze sites we established four transects oriented to the cardinal directions, running upslope from the deepest to shallowest habitat. Transects varied in length from 5 m at site A (maximum depth 3 m), to 75 m at site K (maximum depth 30 m). We constructed a topographic profile of each transect by recording tide-corrected depth at one metre intervals, and quantified substrate composition by filming each transect and point counting sequential still images, using five fixed points per image (English et al. 1976) and 25 benthic substrate categories (Data S1). The twenty five substrate categories were condensed into seven categories for graphical representation: tabular *Acropora*, branching *Acropora*, massive and encrusting coral, soft coral, macroalgae, dead coral, and sediment.

### Results

Underwater observations show that the different reef stages are joined in a continuous reef blanket with a distinctive undulating form, resembling the 'egg box' structure described by Kan et al. (2015) in the cellular reefs of Nagura Bay, Japan. The relationship between the shapes of the different stages can be envisaged by imagining sequentially deeper horizontal slices through a solid egg box. The initial slices contact the peaks, producing circular shapes. These reefs correspond to Wyrwoll et al.'s (2006) pinnacles but we subsequently refer to them as haystacks, based on earlier descriptions of similar *Acropora*-dominated reefs in the Caribbean (Goreau 1959; Kinzie 1973). Deeper slices reach the ridges between adjacent peaks, producing stellate shapes. Subsequent slices produce a platform surrounding enclosed depressions and eventually a solid platform. Below we describe the sequence in the three idealised stages: haystack, stellate and platform. However it should be noted that the sequence is a continuum and that sites within each stage may have features of earlier and/or later stages. Fig. 3 shows representative transect profiles from each stage and Fig. 4 is a schematic block diagram incorporating the main features of the three idealised stages.

### Haystacks



Haystack reefs occur around the margin of the Maze (e.g. sites K and M; Fig. 2). The reef surface at these sites has a sinusoidal profile, curving up over dome-shaped reef tops then descending into bowl-shaped depressions (Fig. 3). The wavelength and amplitude of the profile vary within ranges of approximately 40-100 m and 15-30 m respectively. Like the haystacks of the Caribbean (Goreau 1959; Kinzie 1973), the Abrolhos haystacks consist of loosely interlocked branching *Acropora* colonies, most in growth position but many collapsed and overturned. Adjacent haystacks are linked by saddle-shaped ridges of branching *Acropora*. The predominant *Acropora* species on the haystack reefs and ridges are *A. formosa/muricata* and *A. abrolhosensis*. Tabular *A. spicifera* is abundant at site M on the exposed northern margin of the Maze but absent from the more sheltered site K to the south. Live *Acropora* cover is 60-100% on the reef tops and ridges at both sites, decreasing to approximately 30% within the site M depressions and 0% within the more enclosed and restricted site K depressions. Dead *Acropora* branches at depth are occupied by macroalgae at both sites, predominantly *Sargassum* spp. at site M and *Lobophora variegata* at site K. Beyond the outermost haystacks, the reef surface slopes down to a flat sandy seafloor at 35 m without breaking into isolated patches (Fig. 4). Corals on these outer slopes are predominantly branching and tabular *Acropora* to approximately 25 m depth (Wilson & Marsh 1979). Below 25 m the coral community is more diverse, with a high proportion of foliose genera, particularly *Leptoseris* and *Pachyseris*.

Three islands of storm-deposited *Acropora* rubble line the eastern margin of the Maze (Collins, Zhao & Freeman 2006), and a series of submerged east-west trending linear reef banks occur on the northern margin of the Maze. Two banks can be seen in Fig. 2 and two deeper banks lie beyond them. The bank crests are 2 to 15 m deep, sloping downward to U-shaped troughs at 20 to 30m. The outermost bank reaches the seafloor at 35m. Coral cover and zonation on the banks is equivalent to that of the haystack reefs.

### Stellate reefs

In the stellate stage the haystack reef tops and ridges reach sea level, producing a network of flat-topped star-shaped reefs (sites B, E, F, J, L, N). Water circulation within the intervening depressions is further reduced and the water column is often stratified and stagnant. Live coral is consequently restricted to shallow depths, often less than 15 m in the more enclosed depressions. The shallow subtidal reef slopes and ridges remain dominated by live branching *Acropora* (Fig. 5) and occasional

foliose *Montipora*. Dead *Acropora* branches at depth are colonised by *Nephthea* soft corals and *Lobophora variegata*, and fine sediment accumulates in the bases of the depressions.

A distinct shallow coral community begins to appear in the stellate reefs, consisting of diverse massive and encrusting corals, the most abundant genera being *Montipora*, *Goniastrea*, *Favia*, *Favites*, *Merulina*, *Astreopora*, *Montastrea*, *Mycedium*, *Echinophyllia*, *Cyphastrea*, *Alveopora* and *Lobophyllia*. Several apparent developmental stages of this community are present, initiating as a discontinuous cover of small colonies on dead *Acropora* branches (Fig. 6A) and culminating in vertical or overhanging walls descending from the surface to as deep as ten metres, but typically between two and eight metres (Fig. 6B).

### Reef platform with enclosed depressions

In the platform stage (sites C, D, I, O) the trends in water quality and coral distribution that were established in the stellate reefs develop further: most depressions are rimmed by vertical walls of massive and encrusting coral, live *Acropora* cover below the walls declines rapidly with depth, the water column is usually stratified, and the depressions typically have a deep sediment fill. Late-stage enclosed depressions (A, G, H) gradually fill with sediment to the level of the surrounding reef flat. As they fill, the fringe of live *Acropora* beneath the vertical walls migrates upward, eventually overgrowing the walls and encroaching over the depression floors.

### Model

#### Rationale

Based on the survey results described above, the Abrolhos cellular reefs appear to exhibit a straightforward developmental sequence. However they provide little direct insight into the origin of cellular morphology because the cellular form is already present at the haystack stage. Given the high coral cover on the haystack reef tops and ridges, subsequent growth will inevitably enclose the depressions. Although the haystacks must have progressed through earlier stages to reach their present configuration, the existing space-for-time sequence does not extend back to the earlier stages, presumably because coral colonisation of the Last Interglacial surface ceased when it became deeply submerged and covered by sand in the mid Holocene. If this is the case, even the youngest haystacks probably initiated more than 4000 years ago. Several processes appear to be suppressing live coral

cover, and therefore accretion, within the present-day haystack reef depressions, including reduced water circulation and macroalgal colonisation. But in the absence of earlier stages in the space-for-time sequence it is impossible to determine whether these processes could have initiated the depressions or whether they are consequential. In this situation computer simulation provides a potential means of investigating the early stages of reef development.

## Methods

The model we describe below is configured as a cellular automaton: an array of identically-programmed interacting cells (Ulam 1962; von Neumann 1963; Data S2). This structure is well-suited to simulating reef growth because each cell in the array can be considered to represent a square metre of seafloor, and reefs can grow on the seafloor as three-dimensional stacks of cubic ‘corals’. Using this approach, we simulate lagoon reef development as the growth and coalescence of patch reefs from individual coral recruits on a flat seafloor. We first describe a basic model in which colonisation and growth is essentially random and unconstrained except by sea level, and subsequently introduce a parameter representing branching *Acropora*.

## Basic model

The basic model is initialised by defining the seafloor depth, the array dimensions, and the number of coral recruits. We used a default configuration of 30 m depth (static, i.e. no sea level variation), a 160 x 160 cell array (representing 160 x 160 m, or 2.56 hectares, of seafloor), and 64 randomly-spaced coral recruits, occupying 0.25% of the array. The 160 x 160 m recruitment array was centred within a larger array of 250 x 250 m, giving the reefs a 45 m margin for lateral growth. We chose 30 m as the default depth, rather than the 40 m of the Maze, because cellular reefs elsewhere appear to be thinner than the Maze; those at Mataiva for example are only 10 to 20 m thick (Rossfelder 1990). The horizontal dimensions of the array were selected to minimise computation time while still allowing adequate spatial representation of reef morphology. The colonisation density was selected such that the resulting patch reefs were close enough to eventually coalesce but not so close as to immediately coalesce. We examined variations to the default configuration, including sea level rise, and describe them later under ‘additional modifications’.

Growth from the initial coral recruits is effected by assigning two growth probabilities to each cell in the array in each iteration: a vertical probability representing the likelihood of the cell growing upward itself and a neighbour probability representing the likelihood of the cell being overgrown by a neighbouring coral (Fig. 7). The vertical probability of vacant seafloor cells is zero and the vertical probability of coral-filled cells is random. The neighbour probability of each cell is the product of a random number between zero and one and a 'neighbour value' that depends on the state of the eight surrounding cells. Cells with no shallower neighbours are assigned a neighbour value of zero; otherwise the cell's neighbour value rises incrementally for each shallower neighbour. If a cell becomes surrounded by shallower neighbours it is guaranteed to be overgrown. Otherwise growth is determined by comparing the cell's vertical and neighbour probabilities against two random numbers between zero and one. If either or both probabilities exceed their respective random numbers, the cell grows by one metre when the array is updated prior to the next iteration. Vertical accretion is halted at sea level but lateral accretion continues. The time represented by each iteration is arbitrary but we consider it to be 100 years, giving a mean vertical reef accretion rate of 7 mm/yr (the theoretical maximum rate of 10 mm/yr is not achieved because corals do not grow in every iteration).

### **Branching *Acropora* model**

Representation of branching *Acropora* was guided by the output of the basic model (Fig. 8). The basic model reefs have an irregular 'spiky' surface, with corals projecting up to four metres above the surrounding reef. Such projections cannot occur on real branching *Acropora* reefs because, due to their 'brittle tree' morphology, any branching *Acropora* colonies that grow more than a metre or two above their surroundings will inevitably collapse (Maragos 1972; Bak 1976; see Fig. 5B). This is not necessarily a disadvantage. Because broken fragments can survive and grow to form new colonies, collapse and fragmentation is recognised as an inherent and significant mode of reproduction and short range dispersal in branching *Acropora* (Gilmore and Hall 1976; Tunnicliffe 1981; Bothwell 1982; Highsmith 1982). Collapse is represented in the branching *Acropora* model by imposing a maximum height differential between neighbours (hereafter termed collapse limit) of two metres, such that corals growing to project two metres above any neighbouring cell are prevented from growing upward until the deeper cell grows. Although they cannot grow upward, projecting corals contribute to the growth probability of neighbouring cells in two ways: first, they 'support' neighbouring corals, ensuring they are unrestricted by the collapse limit, and second they may 'collapse into' deeper neighbouring cells (i.e. they increase the neighbour probability of those cells). Although this representation of collapse

involves no subtraction of height from the collapsing colony, it remains valid because it is equivalent to the projecting colony growing a metre then collapsing back a metre during the iteration. Because the imposition of the collapse limit slows reef growth, the time represented by each iteration is reduced to 40 years. This gives a mean vertical reef accretion rate of 9 mm/yr, approximating that of the Abrolhos cellular reefs (Eisenhauer et al. 1993, Collins et al. 1993).

## Additional modifications

We examined the effects of increasing the collapse limit, altering water depth, altering colonisation density, and periodically adding new coral recruits. We also simulated sea level rise and depth-dependent growth, using a simplified linear sea level rise of 10 mm/yr, stabilising at 30 m depth, and a simplified linear reduction of the coral growth rate to 10% of the surface rate at 30 m.

## Results

### Basic model

Patch reefs created with the basic model appear approximately circular in plan view and steeply conical in oblique view (Fig. 8A and B). The individual patch reefs maintain their conical form as they enlarge and coalesce with neighbouring patches (Fig. 8C and D). We use the term ‘nodular’ to describe the shapes and forms generated by the basic model. While the nodular reefs resemble many natural patch reefs (e.g. Fig. 9), they bear little resemblance to cellular reefs. In fact their shapes are the inverse of cellular reefs; nodular reefs appear convex and subcircular in plan view whereas cellular reefs are concave and stellate, surrounding subcircular depressions. However, the basic model is generic and does not intentionally represent any particular coral type.

### Branching *Acropora* model

Reefs created with the branching *Acropora* model closely resemble the Abrolhos cellular reefs (Fig. 10). The model reproduces the characteristic egg box form of the real reefs and all its corollaries including haystack reefs, stellate reefs with radiating ridges, reef platforms enclosing bowl-shaped depressions, scalloped platform margins and the presence of multiple small depressions within larger multi-lobed depressions. The 45° slopes of the model reefs are steeper than the mean of the real *Acropora* slopes (37° ± SD 6°) but within their recorded range. Fig. 11 and Movie 1 show sequential stages in the development of the branching *Acropora* reefs, demonstrating the emergence of their egg

288 box morphology. The key process is the formation of ridges between adjacent patch reefs. This process  
289 begins when the patch reefs meet, whereupon the valleys between them grow rapidly upward to  
290 become saddle-shaped ridges (Fig. 11C and D). The depressions surrounded by the reefs and ridges are  
291 initially irregular in outline but are progressively smoothed to subcircular shapes as the surrounding  
292 reef grows. Eventually the depressions become completely enclosed within the reef platform and  
293 infilled by coral (Fig. 11E).

294

## 295 **Additional modifications**

296 Increasing the collapse limit was the most influential of the additional modifications. Progressively  
297 increasing the collapse limit beyond the two metres of the branching *Acropora* model produces a  
298 transition from cellular to nodular reef forms. A three metre collapse limit creates reefs with weakly  
299 developed subcircular depressions (Fig. 12A and B) and a four metre collapse limit creates reefs with  
300 very few depressions (Fig. 12C and D). Collapse limits of more than four metres produce nodular reefs  
301 equivalent to those of the basic model.

302

303 Varying water depth also significantly influences reef morphology. Reducing depth reduces reef  
304 thickness, which constrains the morphological expression of the collapse limit such that the appearance  
305 of the branching *Acropora* reefs transforms from cellular to nodular as depth decreases (Fig. 13A and  
306 B). In the extreme case of reefs growing in only one or two metres water depth, where the collapse  
307 limit has no effect, all variants of the model produce identical nodular reefs. Increasing depth, by itself,  
308 has little influence on reef morphology (Fig. 13C and D). However, more realistic representations  
309 incorporating sea level rise and depth-dependent growth cause reef slopes to steepen significantly as  
310 depth increases (Fig. 13E and F). Variations in colonisation density and timing have relatively little  
311 effect on reef morphology, besides the expected crowding of patch reefs at high density (Fig. 14).

312

## 313 **Discussion**

### 314 **Model**

315 The resemblance in shape and form between the model reefs and real reefs suggests that the model  
316 adequately represents reality. This interpretation is supported by the model's simplicity: it has only one  
317 rule—collapse if too steep—which is intuitively reasonable and supported by field observations. Model

reef morphology is hyper-sensitive to that rule, running through a nodular to cellular spectrum as permissible steepness is reduced and collapse becomes more frequent.

The nodular reefs produced by the basic model appear straightforward and visually ‘correct’ as growth forms, because the individual patch reefs maintain their forms as they grow and merge. This straightforward morphology is indicative of pure in-situ (in place) growth. Cellular reefs are more complex because the patch reefs transform as they merge, eventually becoming linked by a network of ridges. This transformation results from the high frequency of collapse in the branching *Acropora* model. However, it is not simply the frequency of collapse that produces ridges; more important is the distribution of collapse. Because the valleys between merging patch reefs are low points in the reef structure, coral colonies in the valleys are less likely to project above their neighbours than corals elsewhere. Consequently, they are relatively unrestricted by the collapse limit and are therefore more likely to remain in place as they grow, and less likely to collapse, than colonies elsewhere (Fig. 15). The retention of in-situ colonies transforms the V-shaped valleys into U-shaped ridges that grow to sea level, enclosing depressions (Fig. 11C and D, Movie 1). The subcircular shapes of the depressions arise through the same non-uniform distribution of collapse. Colonies in the re-entrant concavities of early-stage depressions are supported by neighbours and therefore tend to remain in place while those on projecting convexities tend to collapse. Over time this creates smooth rounded shapes, the ultimate smooth shape being a circle.

### **Abrolhos cellular reefs**

The foregoing descriptions of the model branching *Acropora* reefs provide two testable predictions regarding real cellular reefs. First, their slopes should have consistent and relatively low gradients, representing the angle of repose (maximum slope stability angle) of branching *Acropora*. Second, the proportion of collapsed colonies should be lowest in the valleys and ridges and highest on reef slopes. Both predictions are supported in the Abrolhos, where *Acropora* slopes average  $37^{\circ} \pm \text{SD } 6^{\circ}$  (Fig. 3) and *Acropora* colonies in valleys and ridges are generally upright (Fig. 5A, see Fig. 10 for photo location) while those on reef slopes are often overturned (Fig. 5B). We conclude that the Abrolhos cellular reefs have developed according to the model and that Fig. 11 closely describes their morphological progression.



One significant difference between the real and model reefs is the reduced accretion rate of the real reefs once they reach sea level. Model branching *Acropora* reefs reach sea level from 30 m depth in approximately 90 iterations (3600 years) and only require 70 more iterations (2800 years) to completely fill the platform, whereas the Maze reefs, in 40 m depth, reached present sea level in approximately 4500 years (Eisenhauer et al. 1993; Collins et al 1993) but still have not filled the platform 7000 years later. The reduced accretion of the real reefs probably results from two factors not represented in the model. The first is the reduction of *Acropora* cover and vitality at depth as water circulation is restricted by the enclosure of the depressions (Wyrwoll et al. 2006). This is an example of self-limitation through negative feedback between reef growth and water circulation (Blanchon 2011). Self-limitation is therefore a significant influence on the Abrolhos cellular reefs, but operates primarily on their accretion rate not their morphology. The second factor is the colonisation of upper reef slopes by relatively slow-growing massive and encrusting corals. The steep walls created by these corals effectively exclude branching *Acropora*, because any branching *Acropora* that colonise the walls are likely to break off once they grow too large to be supported at their base. By ‘engineering’ steep walls (*sensu* Jones, Lawton & Shachak 1994), massive and encrusting corals are able to monopolise—for thousands of years—prime shallow subtidal habitat that would otherwise be occupied by fast-growing branching *Acropora*. The combination of reduced water circulation at depth and steep walls in the shallows restricts live *Acropora* to a fraction of their previous distribution, significantly slowing the overall reef accretion rate. Model cellular reefs, in contrast, rapidly fill the platform because ‘live’ *Acropora* occupy all habitats including the depression slopes and floors.

Another significant difference between the real and model reefs is the series of linear reef banks on the northern margin of the Maze. We interpret these as early to mid Holocene wave-deposited structures, resulting either from storms, cyclones (Scheffers et al. 2012) or tsunamis (Scheffers et al. 2008).

### **Application to other reefs**

The morphology of cellular reefs elsewhere appears similar enough to the Abrolhos reefs to suggest they have developed the same way, an inference supported by the abundance of branching *Acropora* in documented examples (Alacran Reef: Hoskin 1963; Solomon Islands: Morton 1974; Tétémbia Reef: de Vel and Bour 1990; Cocos-Keeling Atoll: Williams 1994; Pelican Keys: Aronson et al. 1998; Elizabeth Reef: Woodroffe et al. 2004; Pohnpei: Turak and DeVantier 2005; Tun Sakaran: Montagne et al. 2013;



380 Nagura Bay: Kan et al. 2015). At least two of these examples, the Solomon Islands (Morton 1974) and  
381 Nagura Bay (Kan et al. 2015), exhibit vertical walls of massive and encrusting corals above the  
382 *Acropora* zone, suggesting they have undergone the late-stage shallow coral community succession  
383 observed in the Abrolhos.

384  
385 We have separated Mataiva Atoll from the list above as *Porites* is abundant there and has been  
386 considered responsible for the cellular morphology (GIE Raro Moana 1985; Delesalle 1985). However,  
387 branching *Acropora* are also abundant at Mataiva (Delesalle 1985; Rossfelder 1990). We suggest that  
388 branching *Acropora* are the primary reef-builders at Mataiva and *Porites* are colonisers of the  
389 *Acropora* reef, not framework builders. Another possible exception to the rule of *Acropora* dominance  
390 is the 'Type-1' reticulate reef of the Red Sea (Purkis et al. 2010), where *Porites* is also abundant  
391 (Bruckner 2011). However, we would not classify all Type-1 Red Sea reefs as cellular because,  
392 although deep, the depressions are not always circular; more often resembling the transitional  
393 depressions produced by intermediate collapse limits (see Purkis et al.'s Fig. 2). Some Red Sea reefs  
394 are distinctly cellular and we predict those to be *Acropora*-dominated (e.g. 27°57'N, 35°13'E). Closer  
395 examination of these and other cellular reefs is required to determine whether the predominance of  
396 branching *Acropora* is universal, and whether the reef slope gradients and the distribution of collapsed  
397 colonies conform to the Abrolhos example.

398  
399 We have not classified the previously-mentioned Millenium Atoll with the cellular reefs listed above  
400 because the scale of the cellular morphology at Millenium and in many other Pacific atoll lagoons is up  
401 to an order of magnitude larger than the Abrolhos. While the large-scale cellular reefs also seem to  
402 consist predominantly of branching *Acropora* (Roy and Smith 1971; Grovhoug and Henderson 1973;  
403 Valencia 1977; Barott et al. 2010), we do not believe our model applies directly to them because it  
404 cannot produce cells of their dimensions unless it is scaled up massively, to unrealistic depths of at  
405 least 100 m. We are currently investigating the large-scale cellular morphology.

406  
407 The transitional and nodular shapes produced by increasing the collapse limit in the model also occur  
408 in real reefs (e.g. Fig. 9). The simplest interpretation of these shapes is that they indicate coral types, or  
409 mixtures of coral types, progressively less prone to collapse than branching *Acropora*. In this  
410 interpretation, transitional shapes represent reef builders that occasionally collapse, such as foliose and

tabular corals, and nodular shapes represent reef builders that rarely collapse, such as massive and encrusting corals or coralline algae. The nodular reefs of Cockatoo Island in Fig. 9 appear to conform to this interpretation, as they consist of massive and encrusting corals cemented by coralline algae (D. Blakeway, pers. obs.). However, the model indicates that transitional and nodular reef shapes are not necessarily diagnostic of coral type, because branching *Acropora* patch reefs appear nodular (i.e. circular in plan view) before they merge with adjacent patch reefs, and transitional to nodular after they merge in shallow water (Fig. 13A). This suggests that additional information on water depth, reef thickness and reef slope gradients will be required to reliably infer coral type from reef morphology in transitional and nodular reefs. Such three-dimensional data are becoming increasingly available through reef-oriented remote sensing (Zawada & Brock 2009; Zieger, Stieglitz & Kininmonth 2009; Goodman, Purkis & Phinn 2013 and references therein; Leon et al. 2013; 2015). In two-dimensional aerial images, however, the only diagnostic morphology is cellular—signifying relatively thick (>10m) reefs constructed by collapse-prone organisms.

### **Branching *Acropora***

The default collapse-prone reef builders on modern reefs are branching *Acropora*. While it seems possible for other branching coral genera or other calcareous branching invertebrates (e.g. *Millepora*) to create cellular reefs, observations worldwide (listed above) suggest it is almost exclusively *Acropora*: *A. cervicornis* in the Atlantic and multiple species in the Indo-Pacific. This is probably because branching *Acropora* have the ultimate strategy for rapid pre-emption of space in lagoon environments. *Acropora* branches not only grow quickly (up to 19 cm/yr in the Maze; Blakeway 2000), they regularly develop new branches which themselves branch and rebranch, giving them the potential for exponential expansion (Shinn 1976). Constant growth, branching and collapse produce an open three-dimensional structure that rapidly fills lagoons (Davis 1982). Our model indicates that, given adequate depth, cellular reefs are the inevitable result. Cellular reefs are essentially a phenotype of the branching *Acropora* genome(s), emerging from the innate behaviour of branching *Acropora* colonies just as colony morphology emerges from the innate behaviour of polyps.

If the relationship between cellular reef morphology and branching *Acropora* holds, the distinctive shapes of cellular reefs in remotely sensed images can potentially be used to identify and map branching *Acropora* habitat. This could be useful in reef conservation, as the sensitivity of branching

*Acropora* to environmental conditions makes them something of a canary in the coral reef coalmine (Marshall & Baird 2000; Loya et al. 2001; ABRT 2005; Roth & Dehyn 2013). However, assessing anthropogenic impacts in apparently degraded *Acropora*-dominated lagoons will rarely be straightforward, because natural self-limitation and community succession can drastically reduce *Acropora* cover and vitality in the mid to late stages of reef development. Aronson et al. (1998), Aronson (2011) and Perry & Smithers (2011) highlight the value of documenting and understanding such intrinsic trends, generated by the reef itself, before attempting to evaluate the effects of extrinsic influences imposed from outside the reef, including anthropogenic stresses.

## Conclusions

Our simulations indicate that reef morphology is fundamentally determined by the extent to which reef building organisms either remain in place or collapse. This control is best expressed in lagoons, where diminished hydrodynamic and substrate influences allow reefs to grow into their inherent forms. The purest growth forms arise in sheltered lagoons dominated by a single type of reef builder, as in the cellular reefs of the Abrolhos. In these situations reef morphology can be considered a phenotype of the predominant reef building organism.

While the propensity for collapse appears to explain the nodular to cellular spectrum of lagoon reef morphology, many more relationships between reef morphology and ecology remain to be discovered. Many of the recurrent patterns in reef morphology are likely to be ecological phenomena (Blanchon 2011; Schlager and Purkis 2015). Quantifying these patterns and identifying their underlying mechanisms can potentially improve our understanding of present-day reef ecology, because any ecological process capable of shaping a reef will almost invariably be an important process in real time on the living reef. Investigation of the relationships between reef morphology and ecology is benefiting from advances in the availability, resolution and processing of remotely sensed imagery. However, the single most important research technique remains careful and objective underwater observation. Any consistent correlations between reef morphology and underwater survey data, such as coral type, can be considered potential causal relationships warranting closer examination.

In surveying modern reefs it should be recognised that a reef's current state may not represent its formative state, particularly if the reef has reached sea level. While seismic and coring can access the

history stored within such reefs, both techniques are logistically demanding and expensive. The complementary methods we employed in the Abrolhos, space-for-time substitution and computer simulation, are relatively simple and inexpensive but can provide a comprehensive reconstruction of a reef's history and a sound basis for extrapolating its future development.

## Acknowledgements

Many thanks to field volunteers Nik Sander, Gary Watson, Tim Benfield, Nokome Bentley, Matz Berggren, Freda Blakeway, Richard Campbell, Robin Cornish, Simon Cutler, Geoff Deacon, Tim Fisher, Steven Fletcher, Erez Jacobsen, Margaret Jones, Jason Kennington, Helen Kirby, Clare Taylor and Roland Tyson. Logistic support from the mainland was provided by Freda Blakeway, Philip Blakeway and Wendy Perriam. Logistic support in the islands was provided by the Western Australian Department of Fisheries, especially Kim Nardi, Michael Byrne, Chris Chubb, Boze Hancock, David Murphy and Tony Paust. Transport to and from the islands was provided by David Kingdom, Mal McRae, David van der Oord, Ian Rowlands, Chris Shine, Fred Tucker and Geoff Whitehurst. Thanks to Justin Parker, PeerJ Academic Editor Kenneth De Baets and two anonymous reviewers for improvements to the manuscript.

## References

- Acropora* Biological Review Team (ABRT). 2005. Atlantic *Acropora* Status Review Document. Report to National Marine Fisheries Service, Southeast Regional Office. March 3, 2005.
- Aronson, RB, Precht, EF, Macintyre, IG. 1998. Extrinsic control of species replacement on a Holocene reef in Belize: the role of disease. *Coral Reefs* 17: 223-230.
- Aronson, RB. 2011. Intrinsic and extrinsic drivers on coral reefs. In Hopley, D, ed. *Encyclopedia of Modern Coral Reefs: Structure, form and process*. Dordrecht: Springer: 610-612.
- Bak, RPM. 1976. The growth of coral colonies and the importance of crustose coralline algae and burrowing sponges in relation with carbonate accumulation. *Netherlands Journal of Sea Research* 10(3): 285-337.
- Barott KL, Caselle JE, Dinsdale EA, Friedlander AM, Maragos JE, Obura, D, Rohwer, FL, Sandin, SA, Smith, JE, Zgliczynski, B. 2010. The Lagoon at Caroline/Millennium Atoll, Republic of Kiribati: Natural History of a Nearly Pristine Ecosystem. *PLoS ONE* 5(6): 1-10.
- Blakeway, DR. 2000. Geomorphology and ecology of cellular reefs in the Houtman Abrolhos Islands, Western Australia. D. Phil. Thesis, University of Western Australia, Perth.

- Blanchon, P. 2011. Geomorphic zonation. In Hopley, D, ed. *Encyclopedia of Modern Coral Reefs: Structure, form and process*. Dordrecht: Springer: 469-486.
- Blanchon, P, Jones, B, Kalbfleisch, W. 1997. Anatomy of a fringing reef around Grand Cayman: Storm rubble, not coral framework. *Journal of Sedimentary Research* 67: 1-16
- Bothwell, AM. (1982) Fragmentation, a means of asexual reproduction and dispersal in the coral genus *Acropora* (Scleractinia: Astrocoeniida: Acroporidae) – A preliminary report. *Proceedings of the 4th International Coral Reef Symposium* Volume 2: 137-144.
- Bruckner, A. (2011) Khaled bin Sultan Living Oceans Foundation Habitat Mapping and Characterization of Coral Reefs of the Saudi Arabian Red Sea: 2006-2009. *Final Report Part II, Ras Qisbah, Al Wajh, Yanbu, Farasan Banks and Farasan Islands*. Arizona: Panoramic Press.
- Camazine, S, Deneubourg, J-L, Franks, NR, Sneyd, J, Theraulaz, G, Bonabeau, E. (2001). *Self-organization in biological systems*. Princeton: Princeton University.
- Choi, DR, Ginsburg, RN. 1982. Siliciclastic foundations of Quaternary reefs in the southernmost Belize lagoon, British Honduras. *Geological Society of America Bulletin* 93: 116-126.
- Collins, LB, Zhu, ZR, Wyrwoll, K-H, Hatcher, BG, Playford, PE, Chen JH, Eisenhauer, A, Wasserburg, GJ. 1993. Late Quaternary facies characteristics and growth history of a high latitude reef complex: the Abrolhos carbonate platforms, eastern Indian Ocean. *Marine Geology* 110: 203-212.
- Collins, LB, Zhu, ZR, Wyrwoll, K-H, 1996. The structure of the Easter Platform, Houtman Abrolhos reefs: Pleistocene foundations and Holocene reef growth. *Marine Geology* 135: 1-13.
- Collins, LB, Zhu, ZR, Wyrwoll, K-H. 1998. Late Tertiary-Quaternary Geological Evolution of the Houtman Abrolhos Carbonate Platforms, Northern Perth Basin. In Purcell, R, Purcell, P. eds. *The sedimentary basins of Western Australia*, 2. Perth, Western Australia: Petroleum Exploration Society of Australia, 647-663.
- Collins, LB, Zhao, J, Freeman, H. 2006. A high-precision record of mid-late Holocene sea-level events from emergent coral pavements in the Houtman Abrolhos Islands, southwest Australia. *Quaternary International* 145-146: 78-85.
- Crame, JA. 1981. Ecological stratification in the Pleistocene coral reefs of the Kenya coast. *Palaeontology* 24(3): 609-646.
- Dakin, WJ. 1919. The Percy Sladen Trust Expedition to the Abrolhos Islands (Indian Ocean). *Journal of the Linnean Society of London (Zoology)* 34: 127-180.
- Darwin, CR. 1842. *The Structure and Distribution of Coral Reefs. Being the First Part of the Geology of the Voyage of the 'Beagle'*. London: Smith, Elder and Co.
- Davis, GE. 1982. A century of natural change in coral distribution at the Dry Tortugas: a comparison of reef maps in 1881 and 1976. *Bulletin of Marine Science* 32: 608-623.

- Delesalle, B. 1985. Environmental survey of Mataiva Atoll, Tuamotu Archipelago, French Polynesia. *Atoll Research Bulletin* 286.
- de Vel, OY, Bour, W. 1990. The structural and thematic mapping of coral reefs using high resolution SPOT data: application to the Tétémbia Reef, New Caledonia. *Geocarto International* 5: 27-34.
- Drummond, CN, Dugan, PJ. 1999. Self-organizing models of shallow-water carbonate accumulation. *Journal of Sedimentary Research* 69(4): 939-946.
- Eisenhauer, A, Wasserburg, GJ, Chen, JH, Bonani, G, Collins, LB, Zhu, ZR, Wyrwoll, KH. 1993. Holocene sea-level determination relative to the Australian continent: U/Th (TIMS) and <sup>14</sup>C (AMS) dating of coral cores from the Abrolhos Islands. *Earth and Planetary Science Letters* 114(4): 529-547.
- English, S, Wilkinson, C, Baker, V (eds.). 1997. *Survey manual for tropical marine resources*. Australian Institute of Marine Science.
- Fairbridge, RW. 1948. Notes on the geomorphology of the Pelsaert Group of the Houtman's Abrolhos Islands. *Journal of the Royal Society of Western Australia* 33: 1-43.
- GIE Raro Moana. 1985. The Phosphates from Mataiva. *Proceedings of the Fifth International Coral Reef Symposium* Volume 1: 317-319.
- Gilmore, MD, Hall, BR. 1976. Life history, growth habits, and constructional roles of *Acropora Cervicornis* in the patch reef environment. *Journal of Sedimentary Petrology* 46(3): 519-522.
- Goodman, JA, Purkis, SJ, Phinn, SR. (eds.) 2013. *Coral reef remote sensing—A guide for mapping, monitoring and management*. New York: Springer.
- Goreau, TF. 1959. The ecology of Jamaican coral reefs I. Species composition and zonation. *Ecology* 40: 67-90.
- Goreau, TF, Goreau, NI, and Goreau, TJ. 1979. Corals and coral reefs. *Scientific American* 241: 124–136.
- Grovhoug, JG, Henderson, RS. 1976. Distribution of Inshore Fishes at Canton Atoll, in SV Smith and RS Henderson (eds.) *An Environmental Survey of Canton Atoll Lagoon*, 1973, Naval Undersea Center Technical Report NUC TP 395, San Diego, California: 98-157.
- Guilcher, A. 1988. *Coral reef geomorphology*. Chichester: J. Wiley and Sons.
- Halley, R, Shinn, EA, Hudson, JH, Lidz, B. 1977 Recent and relict topography of BooBee patch reef, Belize: *Proceedings of the Third International Coral Reef Symposium* Volume 1, p. 53-96.
- Highsmith, R.C. 1982. Reproduction by fragmentation in corals. *Marine Ecology Progress Series* 7:207–226.



- 606 Hopley, D. 1982. *The geomorphology of the Great Barrier Reef: Quaternary development of coral*  
607 *reefs*. New York: Wiley-Interscience.
- 608
- 609 Hopley, D, Smithers, SG, Parnell, KE. 2007. *The Geomorphology of the Great Barrier Reef:*  
610 *development, diversity and change*. Cambridge: Cambridge University Press.
- 611
- 612 Hoskin, CM. 1963. Recent carbonate sedimentation on Alacran Reef, Yucatan, México. *National*  
613 *Academy of Science National Research Council, Foreign Field Research Program, Publication 19*.
- 614
- 615 Hubbard, DK, Miller, AI, Scaturro, D. 1990. Production and cycling of calcium carbonate in a shelf-  
616 edge reef system (St. Croix, U.S. Virgin Islands): applications to the nature of reef systems in the fossil  
617 record. *Journal of Sedimentary Petrology* 60: 335-360.
- 618
- 619 Jones, JA. 1977. Morphology and development of southeastern Florida patch reefs. *Proceedings of*  
620 *Third International Coral Reef Symposium* Volume 2: 231-235.
- 621
- 622 Jones, CG, Lawton, JH, Shachak, M. 1996. Organisms as ecosystem engineers. *Oikos* 69: 373-386.
- 623
- 624 Kan, H, Urata, K, Nagao, M, Hori, N, Fujita, K, Yokoyama, Y, Nakashima, Y, Ohashi, T, Goto, K,  
625 Suzuki, A. 2015. Submerged karst landforms observed by multibeam bathymetric survey in Nagura  
626 Bay, Ishigaki Island, southwestern Japan. *Geomorphology* 229: 112-124.
- 627
- 628 Kinzie, RA III. 1973. The zonation of West Indian gorgonians. *Bulletin of Marine Science* 23(1):93-  
629 155.
- 630
- 631 Leon, JX, Phinn, SR, Hamylton, S, Saunders, MI. 2013. Filling the ‘white ribbon’ – a multisource  
632 seamless digital elevation model for Lizard Island, northern Great Barrier Reef. *International Journal*  
633 *of Remote Sensing* 34(18): 6337-6354.
- 634
- 635 Leon, JX, Roelfsema, CM, Saunders, MI, Phinn, SR. 2015. Measuring coral reef terrain roughness  
636 using ‘Structure-from-Motion’ close-range photogrammetry. *Geomorphology*. Advance online  
637 publication, February 2015.
- 638
- 639 Loya, Y, Sakai, K, Yamazato, K, Nakano, Y, Sambali, H, van Woesik, R. (2001). Coral bleaching: the  
640 winners and the losers. *Ecology Letters* 4(2): 122-131.
- 641
- 642 Macintyre, IG, Precht, WF, Aronson, RB. 2000. Origin of the Pelican Cays Ponds, Belize. *Atoll*  
643 *Research Bulletin* 466: 1-12.
- 644
- 645 MacNeil, FS. 1954. The shape of atolls: an inheritance from subaerial erosion forms. *American Journal*  
646 *of Science* 252(7): 402-427.
- 647
- 648 Maragos, JE. 1972. A study of the ecology of Hawaiian corals. D. Phil. Thesis, University of Hawaii,  
649 Honolulu.
- 650
- 651 Marshall, PA, Baird, AH. 2000. Bleaching of corals on the Great Barrier Reef: differential  
652 susceptibilities among taxa. *Coral Reefs* 19: 155–163


- Maxwell, WGH. 1968. *Atlas of the Great Barrier Reef*. New York: Elsevier.
- Mazzullo, SJ, Anderson-Underwood, KE, Burke, CD, Bischoff, WD. 1992. Holocene Coral Patch Reef Ecology and Sedimentary Architecture, Northern Belize, Central America. *Palaios* 7(6): 591-601.
- Mistr, S, Bercovici, D. 2003. A theoretical model of pattern formation in coral reefs. *Ecosystems* 6: 61–74.
- Montagne, A, Naim, O, Tourrand, C, Pierson, B, Menier, D. 2013. Status of Coral Reef Communities on Two Carbonate Platforms (Tun Sakaran Marine Park, East Sabah, Malaysia). *Journal of Ecosystems*, Article ID 358183, 15 p.
- Morton, J. 1974. The coral reefs of the British Solomon Islands: a comparative study of their composition and ecology. *Proceedings of the Second International Coral Reef Symposium* Volume 2: 31-53.
- Munk, WH, Sargent, MC. 1954. Adjustment of Bikini Atoll to ocean waves, *U.S. Geological Survey Professional Paper* 260-C: 275–280.
- Nicolis, G, Prigogine, I. 1977. *Self Organization in Non-Equilibrium Systems*. New York: J. Wiley and Sons.
- Novak, V, Santodomingo, N, Rosler, A, Di Marino, E, Braga, JC, Taylor, PD, Johnson, KG, Renema, W. 2013. Environmental reconstruction of a late Burdigalian (Miocene) patch reef in deltaic deposits (East Kalimantan, Indonesia). *Palaeogeography, Palaeoclimatology, Palaeoecology* 374:110–122.
- Perry, CT. 2011. Carbonate budgets and reef framework accumulation. In Hopley, D, ed. *Encyclopedia of Modern Coral Reefs: Structure, form and process*. Dordrecht: Springer: 185-190.
- Perry, CT, Smithers, SG. 2011. Cycles of coral reef 'turn-on', rapid growth and 'turn-off' over the past 8500 years: a context for understanding modern ecological states and trajectories. *Global Change Biology* 17 (1):76-86.
- Perry, CT, Smithers, SG, Gulliver, P, Browne, NK. 2012. Evidence of very rapid reef accretion and reef growth under high turbidity and terrigenous sedimentation. *Geology* 40(8): 719–722.
- Purdy, EG. 1974. Reef configuration: cause and effect. In LaPorte, LF, ed. *Reefs in Time and Space*. Society of Economic Palaeontologists and Mineralogists Special Publications 18: 9-76.
- Purkis, SJ, Rowlands, GP, Riegl, BM, Renaud, PG. 2010. The paradox of tropical karst morphology in the coral reefs of the arid Middle East, *Geology*, 38: 227–230.
- Rietkerk, M, van de Koppel, J. 2008. Regular pattern formation in real ecosystems. *Trends in Ecology & Evolution* 23: 169-175.



- Roberts, HH. 1974. Variability of reefs with regard to changes of wave power around an island. *Proceedings of the Second International Coral Reef Symposium*, Volume 2: 497-512.
- Roberts, HH, Murray, SP, Suhayda, JN. 1975. Physical processes in a fringing reef system. *Journal of Marine Research*, 33: 233-260.
- Rossfelder, A.M. 1990. The submerged phosphate deposit of Mataiva Atoll, French Polynesia. In Burnett, WC and Riggs, SR, eds. *Phosphate deposits of the world. Volume 3. Neogene to Modern Phosphorites*. Cambridge: Cambridge University Press, 195-203.
- Roth, MS, Deheyn, DD. 2013. Effects of cold stress and heat stress on coral fluorescence in reef-building corals. *Scientific Reports* 3: article 1421.
- Roy KJ, Smith SV. 1971. Sedimentation and coral reef development in turbid water: Fanning Lagoon. *Pacific Science* 25(2): 234-248.
- Scheffers, SR, Scheffers, A, Kelletat, D, Bryant, EA. 2008. The Holocene paleo-tsunami history of Western Australia. *Earth and Planetary Science Letters* 270 (1-2): 137-146.
- Scheffers, AM, Scheffers, SR, Kelletat, DH, Squire, P, Collins, L, Feng, Y, Zhao, J-X, Joannes-Boyau, R, May, SM, Schellmann, G and Freeman, H. 2012. Coarse clast ridge sequences as suitable archives for past storm events? Case study on the Houtman Abrolhos, Western Australia. *Journal of Quaternary Science* 27(7): 713-724.
- Schlager, W, Purkis, S. 2015. Reticulate reef patterns: antecedent karst versus self-organization. *Sedimentology* 62(2): 501-515.
- Searle, DE. 1994. Late Quaternary morphology of the Cocos (Keeling) Islands. *Atoll Research Bulletin* 401: 1-13.
- Shinn, EA. 1976. Coral reef recovery in Florida and the Persian Gulf. *Environmental Geology* 1: 241-254.
- Stoddart, DR. 1969. Ecology and morphology of recent coral reefs. *Biological Reviews* 44(4): 433-498.
- Stoddart, DR. 1978. Descriptive reef terminology. In D.R. Stoddart and R.E. Johannes (eds.) *Coral reefs: research methods*. United Nations Educational, Scientific and Cultural Organization, 5-16.
- Tunnicliffe, V. 1981. Breakage and propagation of the stony coral *Acropora cervicornis*. *Proceedings of the National Academy of Sciences USA* 78(4): 2427-2431
- Turak, E, DeVantier, L. 2005. Reef-building corals and coral communities of Pohnpei, Federated States of Micronesia: Rapid ecological assessment of biodiversity and status. *Final report for the Conservation Society of Pohnpei, September 2005*.
- Turing, AM (1952). The chemical basis of morphogenesis. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 237(641): 37-62.

- Ulam, S. 1962. On Some Mathematical Problems Connected with Patterns of Growth and Figures. *American Mathematical Society Proceedings of Symposia in Applied Mathematics* 14: 215-224
- Valencia, MJ. 1977. Christmas Island (Pacific Ocean): Reconnaissance geologic observations. *Atoll Research Bulletin* 197: 1-17.
- von Neumann, J. 1963. *Collected Works of John von Neumann*, Taub, A.H, ed., Pergamon Press.
- Wells, JW. 1957. Coral reefs. In Hedgpeth, JW (ed.) *Treatise on Marine Ecology and Paleoecology* 1. Ecology. New York: Geological Society of America Memoir 67:609-632.
- Williams, DG. 1994. Marine habitats of the Cocos (Keeling) Islands. *Atoll Research Bulletin* 406: 1-10.
- Wilson, BR, Marsh, LM. 1979. Coral reef communities at the Houtman Abrolhos Islands, Western Australia, in a zone of biogeographic overlap. In: *Proceedings of the International Symposium on Marine Biogeography and Evolution in the Southern Hemisphere*. New Zealand Department of Scientific Information and Research, Information Series 137: 259-278.
- Woodroffe CD, Kennedy DM, Jones BG, Phipps CVG. 2004. Geomorphology and Late Quaternary development of Middleton and Elizabeth Reefs. *Coral Reefs* 23: 249-262.
- Wyrwoll, K-H, Zhu, ZR, Collins, LB, Hatcher, BG. 2006. Origin of blue hole structures in coral reefs: Houtman Abrolhos, Western Australia. *Journal of Coastal Research* 22: 202-208.
- Zawada, DG, Brock, JC. 2009. A Multiscale analysis of coral reef topographic complexity using lidar-derived bathymetry. *Journal of Coastal Research: Special Issue 53 - Coastal Applications of Airborne Lidar*: 6-15.
- Zieger S, Stieglitz TC, Kininmonth SJ. 2009. Mapping reef features from multibeam sonar data using multiscale morphometric analysis. *Marine Geology* 264: 209-217.

Cellular reefs in the Pelsaert Group lagoon, Houtman Abrolhos Islands, Western Australia (28°54'S 114°E).






Aerial photograph of the Maze in the Easter Group of the Houtman Abrolhos Islands (28°41'S 113°49'E) showing the 15 survey sites labelled A to O.




Representative transect profiles and graphs of benthic substrate composition from haystack, stellate and enclosed sites in the Maze.




Block diagram illustrating the Maze's egg-box topography and the three idealised stages of cellular reef development.

The cross-section is hypothetical but consistent with seismic and core data.






Dense in-situ (A) and collapsed (B) *Acropora* colonies on a stellate reef in the Pelsaert Group lagoon.



Inferred early (A) and late (B) stages of reef wall development in the Pelsaert Group lagoon.


The walls appear to initiate through the colonisation of dead *Acropora* branches by massive and encrusting corals (A), and subsequently grow to become vertical or overhanging (B).






Cross-section through a hypothetical model reef.

Upward-pointing arrows indicate vertical growth directions, horizontal and diagonal arrows indicate neighbour growth directions.



Reefs generated by the basic model.

A) Two-dimensional plan view of a patch reef after 80 iterations (8000 years) of growth from a single seed coral. Shading corresponds to depth—the reef top at sea level is white and the surrounding seafloor at 30m depth is black. This patch reef reached sea level in approximately 45 iterations (4500 years), and by 80 iterations has developed a 15m wide reef flat. B) Three-dimensional oblique view of the patch reef in A, showing the irregular surface morphology caused by projecting corals. The reef slopes are approximately 65°. C) Two-dimensional plan view of a coalescing patch reef system after 80 iterations. Only the uppermost 10m of the reefs is shown, simulating an aerial view with 10m water visibility. D) Three dimensional oblique view of the reefs in C.



Coalescing nodular patch reefs exposed on a low spring tide at Cockatoo Island, Western Australia, 16°4.8'S 123°35'E.

Photograph by John MacFadyen.




Plan view of model (A) and real (B) branching *Acropora* reefs.

Only the uppermost 10m of the model reef is shown, simulating an aerial view with 10m water visibility. This reef grew in 100 iterations (4000 years) from 225 corals seeded at the default colonisation density (0.25%) in a 300 x 300m array. The real reefs shown in B are those from Figure 1, in the Pelsaert Group of the Abrolhos.

A


100m



B

100m







5A → 5B



Sequential stages in the development of model branching *Acropora* reefs.

A) The set of 64 randomly spaced seed corals from which the model reefs developed. The seed coral configuration is the same as that used to create the basic model reefs in Fig. 8C and D. B) After 25 iterations (1000 years) the seed corals have developed into conical patch reefs with 45° slopes. C) After 50 iterations (2000 years) the patch reefs have enlarged and many have merged. When patch reefs meet, the valleys between them grow upward rapidly to become saddle-shaped ridges (arrowed). D) After 100 iterations (4000 years) some of the reef tops have reached sea level and the system of ridges has developed to enclose and isolate depressions, producing egg-box morphology. E) After 150 iterations (6000 years) an extensive sea level platform has developed, and most of the depressions have filled. F) Plan view showing the uppermost 10m of the reef in D, simulating an aerial view with 10m water visibility.











Influence on model reef morphology of increasing the collapse limit.

A) Plan view, 0-10m depth, of model reef with a 3m collapse limit after 90 growth iterations.

B) Oblique view of the model reef in A. The reef slopes are approximately  $55^\circ$ . C) Plan view,







0-10m depth, of model reef with a 4m collapse limit after 85 growth iterations. D) Oblique

view of the model reef in C. The reef slopes are approximately  $60^\circ$ .



Influence of water depth on the morphology of reefs created with the branching *Acropora* model (A, C, E) and the basic model (B, D, F).

These modifications were undertaken in a smaller (90 x 90 m) array, but maintained the default 0.25% colonisation density. A) Branching *Acropora* reefs grown in 10 m depth exhibit a transition toward the nodular forms of the basic model reefs shown in B). C) Branching *Acropora* reefs grown in 50 m depth retain their cellular morphology. D) basic model reefs grown in 50 m depth retain their nodular morphology. E) Branching *Acropora* reefs incorporating sea level rise and depth-dependent growth steepen to approximately 60°. The blocky appearance of these reefs is a consequence of being forced to their maximum slope, which overrides the model's randomness. F) Basic model reefs incorporating sea level rise and depth-dependent growth steepen to approximately 85°.



Influence on model reef morphology of varying the colonisation density and timing.

These plan view images show the effects of decreasing the recruitment rate from the default 0.25% to 0.125% (A —branching *Acropora* reef, 100 iterations, B —basic reef, 80 iterations), increasing the recruitment rate to 1% (C —branching *Acropora* reef, 90 iterations, D —basic reef, 60 iterations), and periodically adding new recruits during reef growth (E —branching *Acropora* reef, 100 iterations, F —basic reef, 70 iterations).






Diagram illustrating the proposed mechanism of ridge formation derived from the branching *Acropora* model.

The diagram shows a cross-section through two merging patch reefs after 50 iterations (the patch reefs arrowed in Fig. 11C). Isochrons at 20 and 40 iterations show that the patch reefs were initially conical, and that the valley between them has accreted rapidly since the patch reefs merged. Rapid accretion is attributed to the tendency for colonies in valleys to remain in place and colonies on reef slopes to collapse.

