

**A peer-reviewed version of this preprint was published in PeerJ on 17 February 2015.**

[View the peer-reviewed version](https://peerj.com/articles/775) (peerj.com/articles/775), which is the preferred citable publication unless you specifically need to cite this preprint.

Darling S, Uytman C, Allen RJ, Havelka J, Pearson DG. 2015. Body image, visual working memory and visual mental imagery. PeerJ 3:e775  
<https://doi.org/10.7717/peerj.775>

1 **Body Image, Visual Working Memory and Visual Mental Imagery.**

2 Stephen Darling<sup>1\*</sup>, Clare Uytman<sup>1</sup>, Richard J. Allen<sup>2</sup>, Jelena Havelka<sup>2</sup>, David G. Pearson<sup>3</sup>

3 <sup>1</sup> Division of Psychology and Sociology, Queen Margaret University, Edinburgh, UK

4 <sup>2</sup> School of Psychology, University of Leeds, UK

5 <sup>3</sup> School of Psychology, University of Aberdeen, UK

6 \* Corresponding author

7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21 Correspondence concerning this article should be addressed to Stephen Darling, Division of  
22 Psychology and Sociology, Queen Margaret University, Edinburgh EH21 6UU, UK.  
23

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14

## Abstract

Body dissatisfaction (BD) is a highly prevalent feature amongst females in society, with the majority of individuals regarding themselves to be overweight compared to their personal ideal, and very few self-describing as underweight. To date, explanations of this dramatic pattern have centred on extrinsic social and media factors, or intrinsic factors connected to individuals' knowledge and belief structures regarding eating and body shape, with little research examining links between BD and basic cognitive mechanisms. This paper reports a correlational study in which visual and executive cognitive processes that could potentially impact on BD were assessed. Visual memory span and self-rated visual imagery were found to be predictive of BD, alongside a measure of inhibition derived from the Stroop task. In contrast, spatial memory and global precedence were not related to BD. Results are interpreted with reference to the influential multi-component model of working memory.

## 1 Body Image, Visual Working memory and Visual Mental Imagery

2 Body image is a general term attached to a complex web of constructs by which  
3 individuals relate to their own bodies, and in particular to their appearance. In western societies  
4 the issue of weight (thinness) has been of particular significance for body image, as societal  
5 ideals have tended to emphasize unattainable degrees of thinness, especially in females  
6 (Tiggemann, 2006). These unrealistic ideals may underlie what has been termed a 'normative  
7 discontent' (Rodin, Silberstein & Striegel-Moore, 1984). There is little doubt that body  
8 dissatisfaction (BD) contributes greatly to levels of distress in the general population (Ohring,  
9 Graber, & Brooks-Gunn, 2002; Thompson, 1999), and is related to onset of pathological eating  
10 disorders (Attie & Brooks-Gunn, 1989; Killen et al., 1996). Furthermore, although females and  
11 males both show evidence of BD at roughly equivalent prevalence (e.g. Furnham & Calnan,  
12 1998; Furnham, Badmin & Sneade, 2002), the nature of dissatisfaction is qualitatively different  
13 between the sexes: the vast majority of females expressing BD express a preference for a  
14 slimmer figure, whilst for males the distribution is roughly symmetrical, with many wishing to  
15 be bulkier (Furnham & Calnan, 1998; Furnham et al, 2002; Drewnowski & Yee, 1987).

16 An increased knowledge of the predictors and correlates of BD is necessary in order to  
17 further understand this issue. A good deal is already understood about such predictors: factors  
18 such as environmental social pressure to be thin, weight-related teasing, internalization of a thin-  
19 ideal, lack of social support, individual differences in body mass and peer pressure have all been  
20 identified as specific predictors of body dissatisfaction in adolescent girls (Stice & Whitenton,  
21 2002, Presnell, Bearman and Stice, 2004).

22 Although extrinsic social forces such as those detailed above undoubtedly shape body  
23 dissatisfaction, it is important to probe intrinsic factors that may be related to negative body

1 image evaluation too. There is, for example, clear evidence for the existence of a relationship  
2 between negative affect and body dissatisfaction (Presnell, *et al* 2004; Konstanski & Gullone,  
3 2003), though longitudinal research suggests that early BD predicts later depression rather than  
4 vice versa (Stice, Hayward, Cameron, Killen & Taylor, 2000; Paxton, Neumark-Sztainer,  
5 Hannan, Eisenberg, 2006). Other intrinsic factors linked to BD include dispositional factors,  
6 personality factors and attachment (Abbate-Daga, Gramaglia, Amianto, Marzolo & Fassino,  
7 2010), alongside biological factors such as adiposity (e.g. Stice & Whitenton, 2002).

8 Information-processing factors also have the potential to impact on BD. In the last twenty  
9 years an understanding of the constraints imposed by cognitive architecture has had a profound  
10 impact on understanding of social processing (see, e.g. Fazio, Jackson, Dunton & Williams,  
11 1995; Greenwald, Nosek & Banaji, 2003). In a similar vein, a body of research has emerged  
12 focusing on information processing and BD (for a review see Lee & Shafran, 2004). This  
13 research has broadly shown that individuals with eating disorders attend differentially to  
14 information pertaining to shape, weight and eating (e.g. Vitousek & Hollon, 1990) and body  
15 components (Freeman et al., 1991); Fairburn, Shafran & Cooper, 1999). There is also a range of  
16 evidence from modifications of the Stroop task (Stroop, 1935) where participants are required to  
17 report the colour of ink in which a word is printed. In the emotional variant of the Stroop task,  
18 neutral and emotionally charged words are presented to participants who are then asked to  
19 identify the print colour of those words. It has been demonstrated that unpleasant words (e.g.,  
20 fear) produce greater slowing down of reaction times than neutral words (e.g., leaf). This pattern  
21 has also been observed in clinical populations. For example, words related to specific phobias  
22 (i.e. arachnophobia) result in slower response times when presented to individuals with those  
23 phobias (for a review see Williams, Mathews, & MacLeod, 1996), while alcohol related words

1 produce slower response times in alcoholics (e.g. Lusher, Chandler, & Ball, 2004). In relation to  
2 the present study, participants with a high 'drive for thinness' demonstrated greater impairment  
3 in ink colour naming latency when the ostensibly irrelevant text was composed of food words  
4 than control words (Perpiñá, Hemsley, Treasure & De Silva, 1993); participants who were high  
5 in dietary restraint showed a similar pattern towards food and body shape words (Green &  
6 Rogers, 1993). There is also a relationship between the retention of body-related items in  
7 memory and BD (Baker, Williamson & Sylve, 1995;). Hargreaves and Tiggeman (2002)  
8 reported that declines in girls' body satisfaction were related to the strength of cognitive  
9 schemata underlying appearance at the start of the research period, and that this was  
10 distinguishable from self-esteem. The picture emerging from this literature is that of a  
11 relationship between biases in cognition and information processing centred around concepts  
12 related to the body and food, and increased body dissatisfaction or proneness to eating disorders.

13       The cognitive factors described above, however, are intrinsically linked to the conceptual  
14 understanding of bodies and food, and are likely to reflect the content of semantic memory  
15 related to food and body related self-image. There are a further set of cognitive functions that  
16 may influence body dissatisfaction, these being more abstract cognitive processes that likely  
17 influence a range of behaviours such as, for example, planning and inhibition. There is, for  
18 example, evidence that poor inhibition, as assessed on the colour-word Stroop task, is associated  
19 with increased obesity, though not with anorexia nervosa (Fagundo et al., 2012), alongside other  
20 evidence of links between obesity and impaired executive functions such as inhibition, flexibility  
21 and planning (Boeka & Lokken, 2008, Cserjési, Luminet, Poncelet & Lénárd, 2009, Cserjési,  
22 Molnár, Luminet & Lénárd, 2007, Gunstad et al., 2007, Nederkoorn, Braet, Van Eijs, Tanghe &  
23 Jansen, 2006a, Nederkoorn, Smulders, Havermans, Roefs & Jansen, 2006b). There are also

1 relationships between eating disorders and executive functions (for reviews see: Aspen, Darcy &  
2 Lock, 2013; Dobson & Dozois, 2004; Lena, Fiocco & Leyenaar, 2004; Roberts, Tchanturia,  
3 Stahl, Southgate & Treasure, 2007).

4 Our purpose here is to focus in more detail on one particular aspect of cognition that has  
5 been largely neglected in understanding body dissatisfaction and related issues: visual cognition.  
6 The research reported here starts from a simple and relatively uncontroversial position – that  
7 there are (at least) two representative codes capable of sustaining imagery in the cognitive  
8 system, one verbal and one visuospatial in nature (Paivio, 1991). This dual coding approach  
9 underlies the highly influential Working Memory approach (Baddeley & Hitch, 1974; Baddeley,  
10 Allen & Hitch, 2011). In essence the working memory approach posits that parallel temporary  
11 verbal and visual information storage systems (known respectively as the phonological loop and  
12 visuo-spatial sketch pad) store information in a passive manner, whilst other multimodal systems  
13 can perform operations on or form links to that information. One type of multimodal system is  
14 the ‘central executive’ – a label reflecting a set of largely active processes that can manipulate  
15 the information held temporarily active in the passive stores, whilst the another, termed the  
16 ‘episodic buffer’ is more automatic in nature and involved in the formation of cohesive and  
17 unified ‘episodes’ (Baddeley, et al., 2011). When looked at from this perspective it seems likely  
18 that any representation of body image is likely to contain a visuospatial component alongside a  
19 verbal one. There is ample evidence that such visual images can be created and maintained in  
20 memory, even in the absence of explicit visual input (e.g. Quinn & McConnell, 1996; Pearson &  
21 Logie, 2004). Any task where a participant is asked to consider their own body image should  
22 invoke, at least to some extent, a visual representation. Hence, visual imagery ability should be  
23 linked to body image. One possibility is that the more realistic and veridical a participant’s

1 internal model of their own appearance is, the more resistant they should be to bias and distortion  
2 of body image driven by social and cultural factors operationalized in semantic memory.

3 Smeets, Klugkist, van Rooden, Anema, and Postma (2009) have postulated a link  
4 between visual imagery processes and body image using a variant of the Distance Comparison  
5 task (Kosslyn, Ball, & Reiser, 1978). In their study participants were asked to mentally scan  
6 across two specified body widths (e.g., hips and waist) and indicate as quickly as possible which  
7 body part was longer or shorter. The results showed females with high BD were significantly less  
8 accurate on body width comparisons involving small (less than 9 cm) differences, implying that  
9 they had a less accurate visual image of their own body in comparison to a group of low BD  
10 females.

11 However, the Distance Comparison task used by Smeets et al. is a behavioural measure  
12 that cannot directly assess a participant's subjective experience of mental imagery. Auchus,  
13 Kose and Allen (1993) report a study that, surprisingly, appears to be the only attempt to date to  
14 study the relationship between subjective self-reported visual imagery and body imagery. This  
15 study subdivided a sample on a post-hoc basis into four groups based on a test where they had to  
16 adjust the appearance of an initially distorted image of themselves until they felt it reflected  
17 reality. In a control test they adjusted the shape of an object (a tin of soup) in the same way.  
18 Participants were assessed to be either non-distorters, distorters of object but not own-body  
19 representations, distorters of own-body representations but not objects, and general distorters.  
20 Various different measures of visual imagery were taken: overall, no significant group  
21 differences were identified. The authors also adopted an extreme groups approach and claimed to  
22 observe increased ratings of imagery vividness (using the Vividness of Visual Imagery  
23 Questionnaire – VVIQ: Marks, 1973) when a subset of the participants who distorted body



1 image the most was compared against the subset that distorted the least. Such an approach is  
2 methodologically problematic when coupled with small sample size, and does not appear to have  
3 been replicated in the literature. In order to clarify and consolidate understanding of the  
4 relationship of visual imagery and body imagery, the current study adopts a correlational design  
5 in which a number of cognitive measures are adopted in order to predict BD in a female sample.  
6 We focus specifically on females because the aforementioned evidence that BD has a quite  
7 different profile in females and males would imply that separate consideration of body image  
8 issues in females and males would be appropriate.

9         The selection of measures was informed by current understandings of visual imagery and  
10 visual WM. They included a self-report measure of the vividness of visual imagery (the VVIQ),  
11 computerized versions of the Corsi Block task (Corsi, 1972; Milner, 1971) and the Visual  
12 Patterns Task (VPT: Della Sala, Gray, Baddeley & Wilson, 1997). We also included a measure  
13 of global precedence, and a standard ink-naming colour word version of the Stroop task.

14         The VVIQ allows participants to self-report their use of visual imagery. There is evidence  
15 that it does index visual working memory, but it clearly also implicates elements of long-term  
16 memory (Baddeley & Andrade, 2000). Imagery tasks also tend to implicate executive resources  
17 (Logie, 1995, 2003; Pearson, Logie & Green, 1996; Rudkin, Pearson & Logie, 2007; Quinn &  
18 Ralston, 1986). Consequently, in order to incorporate more specific measures of visuospatial  
19 working memory processing, two direct measures of visual and spatial memory span were used.  
20 There is evidence (Della Sala, Gray, Baddeley, Allamano & Wilson, 1999) that the VPT and  
21 Corsi tasks measure qualitatively different visual and spatial sub-processes of working memory,  
22 and hence together they provide the opportunity to understand relationships with BD in a more  
23 sophisticated way. Global precedence is a calculated score reflecting the degree to which a

1 person tends to view visual arrays in a global or a local manner. Global processors may  
2 remember faces better (Darling, Martin, Hellman & Memon, 2009) and process information  
3 about faces in a more configural, holistic fashion (Martin & Macrae, 2010). It is included here to  
4 see if this preference for broad configural information over small-scale local detail is related to  
5 the tendency to misrepresent ones' own body. Somewhat relatedly, Feusner, Moller, Altstein,  
6 Sugar, Bookheimer, Yoon and Hembacher (2010) observed a predisposition to local processing  
7 on a face inversion task amongst a sample of individuals with Body Dysmorphic Disorder  
8 (BDD).

9       As well as direct assessments of visual memory, we also included a measure of resistance  
10 to inhibition by irrelevant information – the classic Stroop task (Stroop, 1935). This task was  
11 included as an estimate of executive functioning, in particular indexing attention and inhibition  
12 (for a review see MacLeod, 1991), two of the so-called executive functions (e.g. Baddeley, 1996;  
13 Miyake, Freidman, Emerson, Witzki, Howerter & Wager, 2000). Although we acknowledge that  
14 executive function is a concept encompassing a diversity of elements, the Stroop task provides a  
15 simple and straightforward method of assessing at least some of these components. The classic  
16 colour-word interference version of the Stroop task used in this experiment measures the  
17 interfering effect of a colour word on the reporting of the colour of ink in which word is  
18 presented. Hence the word 'RED' might be printed in blue ink, and the participant would be  
19 required to report the ink colour as 'blue'. Successful completion of the task requires the  
20 participant to inhibit the prepotent response (reading out the colour word 'red'). This task,  
21 therefore, does not tap inhibition or orientation to particular concepts, but indexes general  
22 inhibitory functioning. Although poor inhibition control has been implicated in eating disorders  
23 and obesity, as discussed above, and although the ability to suppress undesired thoughts may be

1 useful in mitigating against body dissatisfaction itself, there is comparatively little evidence  
2 linking body dissatisfaction to Stroop performance. Hence it is an open question whether Stroop  
3 performance will relate to body dissatisfaction.

4 In summary, this report presents a study in which we sought to assess the relative  
5 importance of various visual imagery, visual memory and executive cognitive factors, with no  
6 direct link to eating behavior, in predicting BD in a sample of females. Age and obesity (assessed  
7 using BMI) were also assessed as predictors, so that an assessment of the role of cognitive  
8 factors could be made independent of age and BMI. To do this we adopted two broadly accepted  
9 measures of body dissatisfaction, a verbal questionnaire (a short form of the Body Shape  
10 Questionnaire, the BSQ-16A: Evans & Dolan, 1993) and a comparison of ideal and current  
11 visual-analogue body shape scales.

## 12 Method

### 13 *Participants*

14 The sample comprised 111 female participants who were either students or staff of Queen  
15 Margaret University, or members of the broader Edinburgh community. Participants took part  
16 either voluntarily or in return for course credit. Self-reported colour blindness was an exclusion  
17 criterion, and participants were required to have normal or corrected-to-normal vision. Mean age  
18 of participants was 26.5 years (SD = 10.2). All participants expressed informed consent in  
19 writing and the research was approved at the Psychology and Sociology Subject area level by the  
20 Queen Margaret University Research Ethics Panel.

1 *Design and Materials*

2 The following variables were assessed in each participant: Age, gender, BMI, BSQ, VPT,  
3 Corsi Blocks Span, Global Precedence, Body Dissatisfaction, VVIQ and Stroop interference  
4 score.

5 *BMI*

6 Body Mass Index was derived from mass and height using the standard formula relating  
7 weight to the square of height and was used as a proxy for degree of body fat.

8 *Body Shape Questionnaire*

9 The 16-item Body Shape Questionnaire (BSQ-16A: Evans & Dolan, 1993) is a self-  
10 report scale where participants rate the frequency of various body shape-related feelings over a  
11 period of four weeks (e.g. ‘Has worry about your shape made you diet?’). It is a shortened form  
12 of the original BSQ-34 (Cooper, Taylor, Cooper & Fairburn, 1987), which has good  
13 psychometric properties (Evans & Dolan, 1993). Ratings are expressed as a 6-point Likert scale  
14 ranging from 1 (Never) to 6 (Always), and the reported score is the total of these ratings: thus  
15 BSQ-16A scores had a potential range from 16 to 96.

16 *Visual Analogue Body Dissatisfaction (VABD)*

17 Participants were presented with a page containing two sets of 9 silhouettes of female  
18 figures ranging from severely underweight (1) to severely overweight (9): based on Stunkard,  
19 Sørensen & Schulsinger, 1983) and asked to indicate which figure most closely matched what  
20 they currently looked like on the upper set of silhouettes. They then marked, on the lower set,  
21 which item represented the shape that they would most like to look like. Dissatisfaction was  
22 assessed by subtraction of the ideal from the current rating, quantifying the discrepancy between  
23 current and ideal body representation. This method is regarded as an effective measure of BD

1 (e.g. Bulik, Wade, Heath, Martin, Stunkard & Eaves, 2001; Mciza, Goedecke, Steyn, Charlton,  
2 Puoane, Meltzer, Levitt & Lambert, 2005) and correlates with BSQ (Mciza et al, 2005). A higher  
3 value indicates a greater degree of dissatisfaction, with the current status being regarded as larger  
4 than the desired.

#### 5 *Vividness of Visual Imagery Questionnaire*

6 An unaltered version of Marks' (1973) Vividness of Visual Imagery Questionnaire  
7 (VVIQ) was used. Participants were asked to remember 4 scenarios (e.g. 'think of some relative  
8 or friend whom you frequently see ... and consider carefully the picture that comes before your  
9 mind's eye') and for each of these they were asked to attend to 4 key visual features (e.g. 'The  
10 precise carriage, length of step, etc. in walking'). For each of these 16 features, participants gave  
11 a rating of the intensity of the visual image, on a scale from 1 ('perfectly clear and as vivid as  
12 normal vision' to 5 ('no image at all – you only 'know' that you are thinking of an object'). In  
13 line with Marks' (1973) method, participants carried out the task twice, first with eyes open, and  
14 then with eyes closed. The VVIQ score was simply the total score across all 32 items: note that  
15 the scoring is counterintuitive: a lower score on the VVIQ represents more vivid imagery.

#### 16 *Visual Patterns Task*

17 The Visual Patterns Task (VPT – Della Sala, et al., 1997; Della Sala, et al., 1999) is a  
18 task designed to assess memory for static visual patterns without the need for participants to  
19 recall sequential information. It is a span task, in which the task difficulty is gradually increased  
20 until participants fail to recall a pattern. The current study did not use the original VPT, instead  
21 an electronic implementation of the VPT was presented using a standard laptop computer with a  
22 15.7" display (1024 x 768 pixels/319 x 239 mm). Participants were presented with grid arrays  
23 comprising a number of individual blank squares of side 60 pixels (19mm). Arrays ranged in size

1 from 2x1 cells to a maximum of 6 x 5. In each array, 50% of the squares (at random) were  
2 coloured blue. Arrays were visible for a total duration of 250ms per visible square (filled or  
3 unfilled), meaning larger displays were on screen for longer. Following presentation, a blank grid  
4 was shown on the screen. Participants were asked to use a standard mouse to select the squares  
5 that had previously been shown in blue. Clicked squares changed colour to indicate that they had  
6 been selected.

7 If a participant correctly reproduced an array in the first trial of a difficulty level, then the  
8 next trial increased in difficulty by increasing the number of coloured squares by 1 and the  
9 matrix size by 2. If they made any errors in reproduction, a second trial was presented at the  
10 same span size. If that pattern was recalled correctly, participants moved on to the next level of  
11 difficulty, and were again allowed 2 attempts to recall correctly at that level of difficulty if  
12 needed. However, if the second pattern was recalled incorrectly, then the procedure stopped.  
13 Participants were allocated a score that represented the maximum number of coloured squares  
14 that had been correctly remembered during the task: hence a higher score represents a higher  
15 capacity for remembering visual patterns.

#### 16 *Corsi Blocks Task*

17 The Corsi Blocks task (CBT – Corsi, 1972; Milner, 1971) requires participants to recall a  
18 sequence of locations tapped out sequentially on a board with a number of blocks randomly  
19 distributed across its surface, and is considered to be a task that indexes spatial sequential  
20 memory as distinct from visual pattern memory (Della Sala et al, 1999). In this study a  
21 computerized version of the task was used (computer was identical to that used for the VPT). As  
22 in the case of the VPT, the Corsi task was implemented as a span task, with difficulty increasing  
23 until the point that participants were no longer able to successfully reproduce the spatial pattern.

1 Participants saw a set of 9 black outlines of squares (of side 60 pixels/19mm) distributed  
2 randomly across a white square in the middle of the screen (side: 600/190mm pixels). A  
3 sequence was then traced by randomly highlighting one of these squares at a time by filling it in  
4 a green colour for 500ms. In the first trial, only one item was highlighted. If (on this or any  
5 subsequent trial) participants recalled the sequence correctly, the sequence length increased by  
6 one item. If an error was made, then a further trial of the same difficulty was presented.  
7 Successful recall of this trial enabled the participant to continue at the next level of difficulty, but  
8 the procedure stopped after two trials at any difficulty level were erroneously responded to.  
9 Participants were allocated a score representing the maximum sequence length that had been  
10 correctly remembered during the task: hence a higher score represents a higher capacity for  
11 remembering spatial sequences.

### 12 *Global Precedence*

13 This task was intended to assess participants' background inclination to process  
14 information on a global or a local basis and was based on a shortened version of the task adopted  
15 by Darling *et al* (2009). Participants saw 72 letter displays, in each of which a large letter was  
16 comprised of a number of smaller letters (e.g. a letter 'P' was formed from a number of  
17 individual letter 'U's). These displays were modeled on the stimuli used by Navon (1977).  
18 Participants were asked to respond by pressing letter keys on the keypad. On 36 trials, the letter  
19 display was preceded by an instruction to report the large letter; on the remaining 36 trials  
20 participants were told to report the small letter. The instruction used was randomized across  
21 trials. For global trials, a local-to-global conflict score was derived by subtracting mean RTs for  
22 correct responses for non-conflicting stimuli (where the global and local letters were the same)  
23 from correct RTs for conflicting stimuli, yielding a value which increases with the degree to

1 which conflicting local information impaired global processing. For local trials, a global-to-local  
2 score was derived in a similar manner reflecting the degree to which conflicting global  
3 information impaired local processing. Finally, the global precedence (GP) score was calculated  
4 by subtracting the local-to-global conflict score from the global-to-local score. This score yields  
5 an increasingly positive value the greater the degree to which global-to-local interference  
6 exceeds local-to-global interference. It provides an assessment of the degree to which irrelevant  
7 global information is automatically processed compared to irrelevant local information, and  
8 controlling for general distractibility; the higher the outcome, the more ‘global’ people are in  
9 their outlook, at least to the extent of processing irrelevant global information.

#### 10 *Stroop Interference*

11 This test was closely modeled after the task developed by Stroop (1935). Participants  
12 were shown a sheet containing a 4 x 6 grid comprising the letter string ‘XXXXXX’ represented in  
13 either blue, green, red, purple, yellow, or orange ink. Participants were asked to name all of the  
14 colours and were timed from start to finish to assess the speed with which they could correctly  
15 name all 24 colours. They were then asked to complete a similar task again, except in this second  
16 task, the ‘XXXXXX’ strings were replaced by colour names which conflicted with the ink colour  
17 that was to be reported (i.e. the word ‘ORANGE’ was depicted in blue ink. Naming time in the  
18 non-conflict condition was subtracted from naming time in the conflicting condition to give a  
19 Stroop Interference (SI) score. SI score increased with the extent to which participants’ responses  
20 were slowed down by the presentation of task-irrelevant information, and measures the degree to  
21 which participants were able to inhibit this information and attend to the ink colour. A lower  
22 score hence reflects greater ability to inhibit irrelevant information.

23



## Results

One participant was initially excluded for providing grossly outlying values (a Stroop interference score of 193, some 49 standard deviations above the mean of other participants). Two further participants did not contribute data to the global/local precedence task due to electronic data collection failures in that task. Table 1 details summary statistics of the principal body image and visual imagery measures in the study for all included participants. Correlations between the variables are reported in Table 2. Global precedence showed no correlations with any of the other variables except a small relationship with age (i.e. increased local focus with age). Focusing on relationships between the cognitive measures and measures related to body dissatisfaction, VVIQ correlated positively with BSQ and VABD, showing that higher scores (lower imagery) was related to higher body dissatisfaction. VPT correlated negatively with these measures, showing the same pattern – higher scores (higher visual memory performance) related to lower body dissatisfaction, though effect sizes were smaller than for VVIQ correlations. Correlations between Corsi performance and body dissatisfaction scores were not significant. Stroop interference correlated with both BSQ and VABD. There were also notably strong relationships between VABD, Stroop and BMI.

To address the fundamental hypotheses of this paper of links between body dissatisfaction and visuospatial working memory, the relative importance of VPT span, Corsi span, global precedence, VVIQ and Stroop interference as predictors of body image was assessed in separate regression analyses upon BSQ and VABD. Age and BMI were included in these analyses as potential covariates of body image. Table 3 reports the results of these analyses. BMI and age were directly entered into the models at Step 1: following this the remaining cognitive predictors (VVIQ, VPT, Corsi, GP and SI) were allowed to enter using a

1 stepwise procedure (probability criteria for entry/removal were  $p \leq .05 / \geq .10$ ). To avoid  
2 undue influences of individual data points on the regression model, participants with Cook's D  
3 values exceeding 0.037 (i.e.  $4/n$ ) were excluded from each analysis. The resulting models are  
4 detailed in Table 3. The models showed that BSQ scores were reliably ( $F(5,93) = 12.113, p <$   
5  $0.001: R^2 = .39$ ) predicted by a regression model including VVIQ ( $t(97) = 5.10, p < .001$ ), age  
6 ( $t(97) = -3.22, p = .002$ ), Stroop interference ( $t(97) = 2.42, p = .02$ ) and VPT ( $t(97) = -2.04, p =$   
7  $.04$ ), and that VABD scores were reliably ( $F(4,102) = 14.23, p < 0.001: R^2 = .37$ ) predicted by a  
8 regression model including BMI ( $t(101) = 4.73, p < .001$ ), VVIQ ( $t(101) = 3.95, p < .001$ ) and  
9 VPT ( $t(101) = -3.321, p = .001$ ).

## 10 Discussion

11 The principal finding of this study is of reliable relationships between some (but not all)  
12 visuospatial and cognitive measures and body dissatisfaction. The relationships identified are  
13 between body dissatisfaction and respectively visual mental imagery (vivid imagery is associated  
14 with lower dissatisfaction) and visual short-term memory (better memory is associated with  
15 lower dissatisfaction). In stark contrast to these patterns, there was no evidence of any link  
16 between BD and either spatial-sequential short-term memory or global precedence.

17 These data replicate and considerably extend the results reported by Auchus et al. (1993),  
18 who described a similar pattern based on an extreme groups analysis whereby the subset of  
19 participants who had the most distorted body image representation reported less rich imagery on  
20 the VVIQ, but that these groups did not differ in terms of their visual recall. Unfortunately, the  
21 robustness of Auchus et al.'s data is unclear, due both to the low sample size in the extreme  
22 groups analyses, and the fact that when an inclusive analysis of the data was carried out there  
23 were no differences between body image distorters on either VVIQ or visual memory. The

1 current results therefore substantively strengthen our understanding of the role of both visual  
2 imagery and memory in body imagery, firstly by convincingly replicating (with a larger sample  
3 and more appropriate analysis, therefore obtaining more satisfactory statistical evidence) the link  
4 between self-reported visual imagery (VVIQ) and BD, and secondly by establishing a link  
5 between short-term visual retention (VPT) and BD: this latter outcome is entirely novel. In doing  
6 so, the role of visual cognitive factors in body dissatisfaction is made explicit: participants with  
7 better visual memory and greater self-reported mental imagery are less likely to report being  
8 dissatisfied with their appearance. Given these results, it is necessary to ask what the  
9 mechanisms may be by which mental imagery and visual memory might influence body image:  
10 we address each concept separately.

11         Visual mental imagery reflects the conscious internal representation of visual scenes.  
12 Consequently it is a very difficult phenomenon to assess. Indeed there has been a historic debate  
13 between researchers arguing around its very existence as a separate and characteristic entity. One  
14 view holds that imagery is entirely verbal and propositional (Pylyshyn, 1981) and hence  
15 functions similarly to linguistic propositional knowledge. An alternative view contends that  
16 images depict analogue visual scenes in a direct manner (Kosslyn, 1981), based on pictorial  
17 representations. Nonetheless, there is some consensus that the VVIQ indexes a reasonably valid  
18 construct (McKelvie, 1995), and in doing so provides subjective evidence in favour of pictorial  
19 representations. Why might such a system be related to body dissatisfaction? The answer to such  
20 a question awaits future research but one possibility is that an efficient pictorial representational  
21 system may serve as a mechanism for storing a consistent and reliable visual self-percept that  
22 would be robust against biases imposed from the outside world, for example social and media  
23 pressure. In contrast, individuals who have less vivid imagery may find their internal

1 representations can be deflected more easily by external pressure and this can lead to greater  
2 inconsistency in comparing self-representation with idealized representations. This interpretation  
3 is consistent with the claim made by Smeets et al. that high BD may be associated with less  
4 accurate visual imagery for a person's body (Smeets et al., 2009).

5 Our findings can be related to a growing body of literature that demonstrates an important  
6 functional role for mental imagery in clinically-related disorders (see Pearson, Deepröse,  
7 Wallace-Hadrill, Burnett Heyes, & Holmes, 2013). For example, patients with schizophrenia and  
8 high-schizotypy controls show significantly greater vividness of imagery than low-schizotypy  
9 controls (Oertel et al., 2009). Participants with depression have also been found to be  
10 significantly slower at generating visual mental imagery than non-depressed controls (Cocude,  
11 Charlot, & Denis, 1997). Furthermore, selective dual-task interference procedures derived from  
12 the Working Memory model (Baddeley, 2000; Baddeley et al, 2011) have been applied to disrupt  
13 or reduce clinically-related mental imagery. Dynamic visual noise is a passive interference  
14 procedure known to selectively disrupt visual working memory (McConnell & Quinn, 2000;  
15 Quinn & McConnell, 2006), and Kemps and Tiggemann (2013) have recently shown that  
16 exposure to a hand-held version can significantly reduce naturally occurring food cravings linked  
17 to mental imagery. Performance of the demanding visuo-spatial game 'Tetris' has also been  
18 shown to significantly reduce the occurrence of intrusive mental images for traumatic film  
19 material (Holmes, James, Coode-Bate, & Deepröse, 2009).

20 Some caveats of the current study should be considered. One problem with assessing  
21 visual cognition using imagery questionnaires is that the task may not be particularly pure: first,  
22 reports are subjective: it is not clear to what extent the self reports are a valid reflection of  
23 internal imagery (though it is to be noted that there is evidence that VVIQ scores are affected by

1 manipulations that are known to impact on visuo-spatial working memory: Andrade, Kavanagh  
2 & Baddeley, 1997; Baddeley & Andrade, 2000). However, Pearson et al. have demonstrated  
3 participants can possess a good metacognitive understanding of their own imagery experience,  
4 with subjective ratings of image vividness being highly predictive of subsequent impact of  
5 imagery on conscious perceptual experience (Pearson, Rademaker, & Tong, 2011). Brain  
6 imaging studies have also linked differences in the introspective experience of mental imagery to  
7 differential cortical activation (Cui, Jeter, Yang, Montague, & Eagleman, 2007; Slotnick,  
8 Thompson, & Kosslyn, 2012). Second, imagery tasks are known to recruit a diverse array of  
9 cognitive components including executive functions (Logie, 1995, 2003; Pearson, et al., 1996;  
10 Rudkin et al, 2007). Consequently it is possible that the relationship of VVIQ performance and  
11 BD is due to some executive factor unrelated to the visual nature of the imagery concerned. The  
12 novel finding of an independent, consistent and statistically reliable relationship between VPT  
13 performance and BD clarifies the issue: a purer measure of visual memory is reliably related to  
14 body dissatisfaction: again this pattern is consistent with the idea that strong visual representative  
15 abilities can protect against body dissatisfaction.

16         Although superficially it may seem that VVIQ and VPT may index similar functions –  
17 one would expect strong visual representational abilities to correlate with vividness of visual  
18 imagery – in fact the correlation in our sample between VPT and VVIQ is small and non-  
19 significant ( $r(n = 110) = -.11, p = .254$ ). Bull, Pearson and Hamilton (2007) assessed  
20 correlations between VPT and VVIQ as part of a larger-scale study, and reported a slightly  
21 higher correlation between VPT and VVIQ ( $r = .22, p < .05$ ): they also noted in exploratory  
22 factor analysis that VVIQ loaded more on a measure of general executive functioning whilst  
23 VPT loaded on a measure of visual representation. On these bases, it seems reasonable to

1 speculate that differences in the profiles of body dissatisfaction relationships between VVIQ and  
2 VPT may well centre on the fact that VVIQ has a more substantive executive component,  
3 broadly consistent with our findings that VPT appears to be a stronger predictor of VABD (a  
4 visual task) than it was the questionnaire-based BSQ.

5 Stroop interference correlated with every other variable in the study apart from global  
6 precedence, highlighting its role as a measure of a general inhibitory mechanism. Some of these  
7 relationships were very high. The strong and sizeable correlation with age is of little surprise  
8 (Comalli, Wapner & Werner, 1962). A strong relationship with BMI is also consistent with  
9 previous literature (e.g. Fagundo, et al., 2012). In the context of the regression models, Stroop  
10 was a moderate predictor of BSQ but, was not a significant predictor of VABD. Looking at the  
11 high correlations between Stroop interference, BMI, VABD and BSQ it is plausible to explain  
12 this by assuming that the relationship between Stroop interference and body dissatisfaction is  
13 heavily driven by the links between obesity and inhibition: VABD is more heavily influenced by  
14 actual obesity than BSQ (given that the participant has to select a representation of their current  
15 body state) and hence the relationship to Stroop may have been substantially reduced in the  
16 regression models because they controlled explicitly for BMI, and more attenuated in VABD  
17 prediction rather than BSQ prediction. Given this it seems reasonable to hypothesize separable  
18 relationships between visual short-term memory and body dissatisfaction and between executive  
19 functions and body dissatisfaction, a testable hypothesis that should be taken up in future  
20 research. One obvious possible explanation is that increased inhibitory control may enable  
21 people to inhibit unwanted negative body-related thoughts, thus mediating dissatisfaction.

22 It is necessary to briefly consider those functions that were not apparently related to  
23 either VABD or BSQ, with due deference to the axiomatic absence of evidence not representing

1 evidence of absence. We failed to observe any relationship between spatial sequence memory  
2 and BD, nor between global and local processing and BD in our sample. Hence, whilst  
3 acknowledging that these are null findings, there is no evidence that the ‘normative discontent’  
4 (Rodin, et al., 1984) that is seen in the general population is related to intrinsic biases in  
5 processing orientation, or to individual variations in spatial sequence memory. As far as we are  
6 aware this is the first study attempting to link BD with spatial memory and global precedence.

7         This study restricted its focus to female participants. Hence we must add the caveat that  
8 these data do not necessarily generalize to males, and that research on the relationship between  
9 visual cognition and BD in males would be warranted in future. Of particular interest here may  
10 be the relatively complex patterns of gender difference in visual and spatial memory ability (e.g.  
11 Voyer, Postma, Brake & Imperato-McGinley, 2007; Cattaneo, Postma & Vecchi, 2006), and it  
12 would be worthwhile to tease out relationships between BD, gender and visual cognition in  
13 future. In the light of these gender differences, it must be made explicit here that the current data  
14 do not support the argument that females are more prone to BD or eating disorders as a  
15 consequence of gender differences in visual processing. Such a claim would be groundless, and  
16 go way beyond our current claim which is to say that we have observed evidence consistent with  
17 a role for visual processing in establishing, maintaining or protecting against BD.

18         The Working Memory model (Baddeley, 2000; Baddeley et al, 2011) was advanced  
19 earlier as a potentially useful framework for understanding the role of visual imagery in relation  
20 to BD. Given the current data, it seems likely that there is a substantive relationship between two  
21 components of working memory and BD. Efficient functioning in both the visuo-spatial sketch  
22 pad (VSSP) *and* the central executive (CE) seems to be associated with lower BD. The executive  
23 processing relationship was stronger for BSQ than for VABD, leading to the tentative conclusion

1 that the BSQ, perhaps by virtue of its use of a verbal questionnaire format, may have more  
2 executive demands. Nonetheless, an independent and reliable relationship between Stroop  
3 interference to the VABD demonstrates a relationship between CE and BD that extends beyond  
4 the nature of the stimulus materials and implicates the CE in BD. Meanwhile, a relatively small  
5 but significant independent contribution of the VPT in predicting both BSQ and VABD  
6 demonstrates the importance of visual STM to sustaining BD.

7         The present study allows a more detailed understanding of the relationship between the  
8 VSSP and BD, because only visual STM predicted BD: spatial STM had no such association.  
9 There is a reasonable consensus that the VSSP is likely to be divisible into a primarily visual and  
10 a primarily spatial system (e.g. Della Sala, et al., 1999; Logie, 1995) though there has been  
11 debate about the precise nature of these systems and how they interact (e.g. Logie, 2011;  
12 Darling, Della Sala & Logie, 2009). There is also evidence that visual and spatial imagery recruit  
13 different cognitive systems (Quinn & Ralston, 1986; McConnell & Quinn, 2000). With these  
14 patterns in mind, the present results suggest that the systems associated with BD are visual,  
15 rather than spatial, in nature: there was no evidence at all for a link between spatial sequential  
16 memory (on the Corsi task) and BD.

17         Recent models of working memory (Baddeley, 2000; Baddeley, et al., 2011) have  
18 incorporated a new component known as the episodic buffer (EB), with a role in combining  
19 information held in separate stores: the impact of this structure is less clear, and remains to be  
20 probed in future research and theorizing. The inhibition and attentional demands of the Stroop  
21 task used in the current research are more related to the Central Executive than they are clearly  
22 linked to the episodic buffer. As a hypothetical construct thought to require low levels of  
23 attention that can link information simultaneously held in different STM and LTM stores, the EB



1 might well be involved in sustaining distorted body-image related representations; for example  
2 when an individual may be thinking about their appearance it may provide access to episode-  
3 specific information which could easily be a source of biasing or negatively-valenced  
4 information. Given the role of the EB role in holding together (or ‘binding’) information held in  
5 parallel information streams in order to carry out particular task demands (e.g. Darling, Allen,  
6 Havelka, Campbell & Rattray, 2011, Allen, Havelka, Falcon, Evans & Darling, 2014), it may be  
7 the case that the efficiency of the EB may, speculatively, influence the robustness of self-image  
8 representations. However, this is speculation *in extremis*, and a fuller understanding awaits a  
9 fuller understanding of the role of the EB in imagery tasks in general, an enterprise that has not  
10 really been pursued at the present time (Pearson, 2006). We should also note that the working  
11 memory model is not the only notable model of mental imagery that separates visual and spatial  
12 processes; this explanation is also largely consistent with other approaches to imagery (Farah,  
13 Hammond, Levine & Calvanio, 1988, Kozhevnikov, Kosslyn & Shephard, 2005).

14 In summary, the results of the current study indicate that some measures of visuospatial  
15 cognitive function are predictive of body image measures. In order to understand the processes  
16 underlying body dissatisfaction, researchers would therefore be well advised to consider visual  
17 imagery and visual memory features, and we argue that the well-established working memory  
18 model proposes constructs that may help advance this understanding. It is possible that a vivid  
19 and veridical visual imagery system offers a degree of immunity to the various insults to body  
20 image that are synonymous with life in the modern world, perhaps by providing a robust parallel  
21 representation of the body that is either less or differentially affected by messages in the media  
22 and society. How this representation is established and maintained is a question for future  
23 research. At present we have identified a role for visual STM and the CE but other aspects of

1 cognition may well be involved. As a final note, we wish to make it clear that we completely  
2 acknowledge the massive importance of social and media representations to endemic, normative  
3 BD, and would not seek to underplay this. However, the finding that *visual* representation is a  
4 crucial mediator of BD is also of key importance, and we hope that this finding can be the basis  
5 for development of innovative approaches to BD that may help mitigate the pervasive and  
6 debilitating social effects of this strikingly prevalent phenomenon.

## Acknowledgements

The authors would like to acknowledge the invaluable assistance of Kirsten Millar, Kirsty Young and Catherine Leckie in collecting the data reported in this study.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## References

Abbate-Daga, G., Gramaglia, C., Amianto, F., Marzola, E., & Fassino, S. (2010). Attachment Insecurity, Personality, and Body Dissatisfaction in Eating Disorders. *The Journal Of Nervous And Mental Disease*, 198(7), 520-524. doi:10.1097/nmd.0b013e3181e4c6f7

Allen, R.J, Havelka, J., Falcon, T., Evans, S., & Darling, S. (2014). Modality Specificity and Integration in Working Memory: Insights From Visuospatial Bootstrapping. *Journal Of Experimental Psychology: Learning, Memory, And Cognition*. doi:10.1037/xlm0000058

Andrade, J., Kavanagh, D., & Baddeley, A. (1997). Eye-movements and visual imagery: A working memory approach to the treatment of post-traumatic stress disorder. *British Journal Of Clinical Psychology*, 36(2), 209-223. doi:10.1111/j.2044-8260.1997.tb01408.x

Aspen, V., Darcy, A., & Lock, J. (2013). A review of attention biases in women with eating disorders. *Cognition & Emotion*, 27(5), 820-838. doi:10.1080/02699931.2012.749777

Attie, I., & Brooks-Gunn, J. (1989). Development of eating problems in adolescent girls: A longitudinal study. *Developmental Psychology*, 25(1), 70-79. doi:10.1037//0012-1649.25.1.70

Auchus, M., Kose, G., & Allen, R. (1993). Body-image distortion and mental imagery. *Perceptual And Motor Skills*, 77(3), 719-728. doi:10.2466/pms.1993.77.3.719

Baddeley, A.D (1996). Exploring the Central Executive. *The Quarterly Journal Of Experimental Psychology A*, 49(1), 5-28. doi:10.1080/027249896392784

- 1 Baddeley, A.D. (2000). The episodic buffer: a new component of working memory?. *Trends In*  
2 *Cognitive Sciences*, 4(11), 417-423. doi:10.1016/s1364-6613(00)01538-2
- 3 Baddeley, A.D., & Andrade, J. (2000). Working memory and the vividness of imagery. *Journal*  
4 *Of Experimental Psychology: General*, 129(1), 126-145. doi: 10.1037//0096-  
5 3445.129.1.126
- 6 Baddeley, A.D., & Hitch, G.J. (1974). Working Memory . *Academic Press*, 8, 47-89.
- 7 Baddeley, A.D., Allen, R.J., & Hitch, G.J. (2011). Binding in visual working memory: The role  
8 of the episodic buffer. *Neuropsychologia*, 49(6), 1393-1400.  
9 doi:10.1016/j.neuropsychologia.2010.12.042
- 10 Baker, J., Williamson, D., & Sylve, C. (1995). Body image disturbance, memory bias, and body  
11 dysphoria: Effects of negative mood induction. *Behavior Therapy*, 26(4), 747-759.  
12 doi:10.1016/s0005-7894(05)80043-7
- 13 Boeka, A., & Lokken, K. (2008). Neuropsychological performance of a clinical sample of  
14 extremely obese individuals. *Archives Of Clinical Neuropsychology*, 23(4), 467-474.  
15 doi:10.1016/j.acn.2008.03.003
- 16 Bulik, C., Wade, T., Heath, A., Martin, N., Stunkard, A., & Eaves, L. (2001). Relating body  
17 mass index to figural stimuli: population-based normative data for Caucasians. *Int J Obes*  
18 *Relat Metab Disord*, 25(10), 1517-1524. doi:10.1038/sj.ijo.0801742

- 1 Bull, R., Pearson, D., & Hamilton, C. (2007). *Measurement and Interference of Visual-Spatial*  
2 *Memory: Full Research Report ESRC End of Award Report, RES-000-22-1507*. Swindon:  
3 ESRC. Retrieved from <http://www.esrc.ac.uk/my-esrc/grants/RES-000-22-1507/read>
- 4 Cattaneo, Z., Postma, A., & Vecchi, T. (2006). Gender differences in memory for object and  
5 word locations. *The Quarterly Journal Of Experimental Psychology*, 59(05), 904-919.  
6 doi:10.1080/02724980543000079
- 7 Comalli, P., Wapner, S., & Werner, H. (1962). Interference Effects of Stroop Color-Word Test in  
8 Childhood, Adulthood, and Aging. *The Journal Of Genetic Psychology*, 100(1), 47-53.  
9 doi:10.1080/00221325.1962.10533572
- 10 Cserjési, R., Luminet, O., Poncelet, A., & Lénárd, L. (2009). Altered executive function in  
11 obesity. Exploration of the role of affective states on cognitive abilities. *Appetite*, 52(2),  
12 535-539. doi:10.1016/j.appet.2009.01.003
- 13 Cserjési, R., Molnar, D., Luminet, O., & Lénárd, L. (2007). Is there any relationship between  
14 obesity and mental flexibility in children? *Appetite*, 49(3), 675-678.  
15 doi:10.1016/j.appet.2007.04.001
- 16 Cocude, M., Charlot, V., & Denis, M. (1997). Latency and duration of visual mental images in  
17 normal and depressed subjects. *Journal Of Mental Imagery*, 21(1-2), 127-142
- 18 Cooper, P., Taylor, M., Cooper, Z., & Fairburn, C. (1987). The development and validation of  
19 the body shape questionnaire. *International Journal Of Eating Disorders*, 6(4), 485-494.  
20 doi:10.1002/1098-108x(198707)6:4<485::aid-eat2260060405>3.0.co;2-o

- 1 Corsi, P. (1972). *Human Memory and the Medial Temporal Region of the Brain* (Ph.D.). McGill  
2 University.
- 3 Cui, X., Jeter, C., Yang, D., Montague, P., & Eagleman, D. (2007). Vividness of mental imagery:  
4 Individual variability can be measured objectively. *Vision Research*, 47(4), 474-478.  
5 doi:10.1016/j.visres.2006.11.013
- 6 Darling, S., Allen, R.J, Havelka, J., Campbell, A., & Rattray, E. (2012). Visuospatial  
7 bootstrapping: Long-term memory representations are necessary for implicit binding of  
8 verbal and visuospatial working memory. *Psychon Bull Rev*, 19(2), 258-263.  
9 doi:10.3758/s13423-011-0197-3
- 10 Darling, S., Martin, D., Hellmann, J., & Memon, A. (2009). Some witnesses are better than  
11 others. *Personality And Individual Differences*, 47(4), 369-373.  
12 doi:10.1016/j.paid.2009.04.010
- 13 Darling, S., Della Sala, S., & Logie, R. (2009). Dissociation between appearance and location  
14 within visuo-spatial working memory. *The Quarterly Journal Of Experimental Psychology*,  
15 62(3), 417-425. doi:10.1080/17470210802321984
- 16 Della Sala, S., Gray, C., Baddeley, A., & Wilson, L. (1997). *The visual patterns test: A test of*  
17 *short-term visual recall*. Bury St, Edmunds: Thames Valley Test Company.
- 18 Della Sala, S., Gray, C., Baddeley, A., Allamano, N., & Wilson, L. (1999). Pattern span: a tool  
19 for unwelding visuo-spatial memory. *Neuropsychologia*, 37(10), 1189-1199.  
20 doi:10.1016/s0028-3932(98)00159-6

- 1   Drewnowski, A., & Yee, D. (1987). Men and body image: are males satisfied with their body  
2       weight?. *Psychosomatic Medicine*, 49(6), 626-634. doi:10.1097/00006842-198711000-  
3       00008
- 4   Dobson, K., & Dozois, D. (2004). Attentional biases in eating disorders: A meta-analytic review  
5       of Stroop performance. *Clinical Psychology Review*, 23(8), 1001-1022.  
6       doi:10.1016/j.cpr.2003.09.004
- 7   Evans, C., & Dolan, B. (1993). Body shape questionnaire: Derivation of shortened “alternate  
8       forms”. *International Journal Of Eating Disorders*, 13(3), 315-321. doi:10.1002/1098-  
9       108x(199304)13:3<315::aid-eat2260130310>3.0.co;2-3
- 10   Fagundo, A., de la Torre, R., Jimenez-Murcia, S., Agüera, Z., Granero, R., & Tregue, S.  
11       et al. (2012). Executive Functions Profile in Extreme Eating/Weight Conditions: From  
12       Anorexia Nervosa to Obesity. *Plos ONE*, 7(8), e43382. doi:10.1371/journal.pone.0043382
- 13   Fairburn, C., Shafran, R., & Cooper, Z. (1999). A cognitive behavioural theory of anorexia  
14       nervosa. *Behaviour Research And Therapy*, 37(1), 1-13. doi:10.1016/s0005-  
15       7967(98)00102-8
- 16   Farah, M., Hammond, K., Levine, D., & Calvanio, R. (1988). Visual and spatial mental imagery:  
17       Dissociable systems of representation. *Cognitive Psychology*, 20(4), 439-462.  
18       doi:10.1016/0010-0285(88)90012-6



- 1 Fazio, R., Jackson, J., Dunton, B., & Williams, C. (1995). Variability in automatic activation as  
2 an unobtrusive measure of racial attitudes: A bona fide pipeline?. *Journal Of Personality  
3 And Social Psychology*, 69(6), 1013-1027. doi:10.1037//0022-3514.69.6.1013
- 4 Feusner, J., Moller, H., Altstein, L., Sugar, C., Bookheimer, S., Yoon, J., & Hembacher, E.  
5 (2010). Inverted face processing in body dysmorphic disorder. *Journal Of Psychiatric  
6 Research*, 44(15), 1088-1094. doi:10.1016/j.jpsychires.2010.03.015
- 7 Freeman, R., Touyz, S., Sara, G., Rennie, C., Gordon, E., & Beumont, P. (1991). In the eye of  
8 the beholder: Processing body shape information in anorexic and bulimic patients.  
9 *International Journal Of Eating Disorders*, 10(6), 709-714. doi:10.1002/1098-  
10 108x(199111)10:6<709::aid-eat2260100609>3.0.co;2-n
- 11 Furnham, A., & Calnan, A. (1998). Eating disturbance, self-esteem, reasons for exercising and  
12 body weight dissatisfaction in adolescent males. *European Eating Disorders Review*, 6(1),  
13 58-72. doi:10.1002/(sici)1099-0968(199803)6:1<58::aid-erv184>3.3.co;2-m
- 14 Furnham, A., Badmin, N., & Sneade, I. (2002). Body Image Dissatisfaction: Gender Differences  
15 in Eating Attitudes, Self-Esteem, and Reasons for Exercise. *The Journal Of Psychology*,  
16 136(6), 581-596. doi:10.1080/00223980209604820
- 17 Green, M., & Rogers, P. (1993). Selective attention to food and body shape words in dieters and  
18 restrained nondieters. *International Journal Of Eating Disorders*, 14(4), 515-517.  
19 doi:10.1002/1098-108x(199312)14:4<515::aid-eat2260140417>3.0.co;2-e

- 1 Greenwald, A., Nosek, B., & Banaji, M. (2003). "Understanding and using the Implicit  
2 Association Test: I. An improved scoring algorithm": Correction to Greenwald et al.  
3 (2003). *Journal Of Personality And Social Psychology*, 85(3), 481-481.  
4 doi:10.1037/h0087889
- 5 Gunstad, J., Paul, R., Cohen, R., Tate, D., Spitznagel, M., & Gordon, E. (2007). Elevated body  
6 mass index is associated with executive dysfunction in otherwise healthy adults.  
7 *Comprehensive Psychiatry*, 48(1), 57-61. doi:10.1016/j.comppsy.2006.05.001
- 8 Hargreaves, D., & Tiggemann, M. (2002). The Effect Of Television Commercials On Mood And  
9 Body Dissatisfaction: The Role Of Appearance-Schema Activation. *Journal Of Social And  
10 Clinical Psychology*, 21(3), 287-308. doi:10.1521/jscp.21.3.287.22532
- 11 Holmes, E., James, E., Coode-Bate, T., & Deepröse, C. (2009). Can Playing the Computer Game  
12 "Tetris" Reduce the Build-Up of Flashbacks for Trauma? A Proposal from Cognitive  
13 Science. *Plos ONE*, 4(1), e4153. doi:10.1371/journal.pone.0004153
- 14 Kemps, E., & Tiggemann, M. (2013). Hand-held dynamic visual noise reduces naturally  
15 occurring food cravings and craving-related consumption. *Appetite*, 68, 152-157.  
16 doi:10.1016/j.appet.2013.05.001
- 17 Killen, J., Taylor, C., Hayward, C., Haydel, K., Wilson, D., & Hammer, L. et al. (1996). Weight  
18 concerns influence the development of eating disorders: A 4-year prospective study.  
19 *Journal Of Consulting And Clinical Psychology*, 64(5), 936-940. doi:10.1037/0022-  
20 006x.64.5.936

- 1 Kozhevnikov, M., Kosslyn, S., & Shephard, J. (2005). Spatial versus object visualizers: A new  
2 characterization of visual cognitive style. *Memory & Cognition*, 33(4), 710-726.  
3 doi:10.3758/bf03195337
- 4 Kosslyn, S. (1981). The medium and the message in mental imagery: A theory. *Psychological*  
5 *Review*, 88(1), 46-66. doi:10.1037//0033-295x.88.1.46
- 6 Kosslyn, S., Ball, T., & Reiser, B. (1978). Visual images preserve metric spatial information:  
7 Evidence from studies of image scanning. *Journal Of Experimental Psychology: Human*  
8 *Perception And Performance*, 4(1), 47-60. doi:10.1037//0096-1523.4.1.47
- 9 Kostanski, M., & Gullone, E. (1999). Dieting and Body Image in the Child's World:  
10 Conceptualization and Behavior. *The Journal Of Genetic Psychology*, 160(4), 488-499.  
11 doi:10.1080/00221329909595561
- 12 Lee, M., & Shafran, R. (2004). Information processing biases in eating disorders. *Clinical*  
13 *Psychology Review*, 24(2), 215-238. doi:10.1016/j.cpr.2003.10.004
- 14 Lena, S., Fiocco, A., & Leyenaar, J. (2004). The Role of Cognitive Deficits in the Development  
15 of Eating Disorders. *Neuropsychol Rev*, 14(2), 99-113.  
16 doi:10.1023/b:nerv.0000028081.40907.de
- 17 Logie, R. (1995). *Visuo-spatial working memory*. Hove, East Sussex: L. Erlbaum Associates.
- 18 Logie, R. (2003). Spatial and visual working memory: A mental workspace. In D.E. Irwin &  
19 B.H. Ross (Eds.) *Cognitive Vision: The Psychology of Learning and Motivation Volume*  
20 42. (pp. 37-78). DOI: 10.1016/S0079-7421(03)01002-8

- 1 Logie, R. (2011). The Functional Organization and Capacity Limits of Working Memory.  
2 *Current Directions In Psychological Science*, 20(4), 240-245.  
3 doi:10.1177/0963721411415340
- 4 Lusher, J., Chandler, C., & Ball, D. (2004). Alcohol dependence and the alcohol Stroop  
5 paradigm: evidence and issues. *Drug and Alcohol Dependence*, 75(3), 225-231.
- 6 MacLeod, C. (1991). Half a century of research on the Stroop effect: An integrative review.  
7 *Psychological Bulletin*, 109(2), 163-203. doi:10.1037/0033-2909.109.2.163
- 8 Marks, D. (1973). Visual imagery differences in the recall of pictures. *British Journal of*  
9 *Psychology*, 64(1), 17-24. doi:10.1111/j.2044-8295.1973.tb01322.x
- 10 Martin, D., & Macrae, C. (2010). Processing style and person recognition: Exploring the face  
11 inversion effect. *Visual Cognition*, 18(2), 161-170. doi:10.1080/13506280902868793
- 12 McConnell, J., & Quinn, J. (2000). Interference in Visual Working Memory. *The Quarterly*  
13 *Journal Of Experimental Psychology A*, 53(1), 53-67. doi:10.1080/027249800390664
- 14 Mciza, Z., Goedecke, J., Steyn, N., Charlton, K., Puoane, T., & Meltzer, S. et al. (2005).  
15 Development and validation of instruments measuring body image and body weight  
16 dissatisfaction in South African mothers and their daughters. *Public Health Nutrition*,  
17 8(05). doi:10.1079/phn2005814
- 18 McKelvie, S. (1995). The VVIQ as a psychometric test of individual differences in visual  
19 imagery vividness: A critical quantitative review and plea for direction. *Journal Of Mental*  
20 *Imagery*, 19(3-4), 1-106.

- 1 Milner, B. (1971). Interhemispheric differences in the localization of psychological processes in  
2 man. *British Medical Bulletin*, 27(3), 272-277.
- 3 Miyake, A., Friedman, N., Emerson, M., Witzki, A., Howerter, A., & Wager, T. (2000). The  
4 Unity and Diversity of Executive Functions and Their Contributions to Complex “Frontal  
5 Lobe” Tasks: A Latent Variable Analysis . *Cognitive Psychology* , 41(1), 49-100.  
6 Retrieved from <http://www.sciencedirect.com/science/article/pii/S001002859990734X>
- 7 Navon, D. (1977). Forest before trees: The precedence of global features in visual perception.  
8 *Cognitive Psychology*, 9(3), 353-383. doi:10.1016/0010-0285(77)90012-3
- 9 Nederkorn, C., Braet, C., Van Eijs, Y., Tanghe, A., & Jansen, A. (2006). Why obese children  
10 cannot resist food: The role of impulsivity. *Eating Behaviors*, 7(4), 315-322.  
11 doi:10.1016/j.eatbeh.2005.11.005
- 12 Nederkorn, C., Smulders, F., Havermans, R., Roefs, A., & Jansen, A. (2006). Impulsivity in  
13 obese women. *Appetite*, 47(2), 253-256. doi:10.1016/j.appet.2006.05.008
- 14 Oertel, V., Rotarska-Jagiela, A., van de Ven, V., Haenschel, C., Grube, M., & Stangier, U. et al.  
15 (2009). Mental imagery vividness as a trait marker across the schizophrenia spectrum.  
16 *Psychiatry Research*, 167(1-2), 1-11. doi:10.1016/j.psychres.2007.12.008
- 17 Ohring, R., Graber, J., & Brooks-Gunn, J. (2002). Girls' recurrent and concurrent body  
18 dissatisfaction: Correlates and consequences over 8 years. *International Journal Of Eating  
19 Disorders*, 31(4), 404-415. doi:10.1002/eat.10049

- 1 Paivio, A. (1991). Dual coding theory: Retrospect and current status. *Canadian Journal Of*  
2 *Psychology/Revue Canadienne De Psychologie*, 45(3), 255-287. doi:10.1037/h0084295
- 3 Paxton, S., Neumark-Sztainer, D., Hannan, P., & Eisenberg, M. (2006). Body Dissatisfaction  
4 Prospectively Predicts Depressive Mood and Low Self-Esteem in Adolescent Girls and  
5 Boys. *Journal Of Clinical Child & Adolescent Psychology*, 35(4), 539-549.  
6 doi:10.1207/s15374424jccp3504\_5
- 7 Pearson, D.G. (2006). The episodic buffer: Implications and connections with visuo-spatial  
8 research. In: T. Vecchi (Ed.), *Imagery and Spatial Cognition: Methods, Models and*  
9 *Cognitive Assessment* (pp.139-153). Amsterdam: John Benjamins Publishing Company.
- 10 Pearson, D., & Logie, R. (2004). Effects of stimulus modality and working memory load on  
11 mental synthesis performance. *Imagination, Cognition And Personality*, 23(2), 183-191.  
12 doi:10.2190/krqb-0ced-nx6j-hq72
- 13 Pearson, D., Deepröse, C., Wallace-Hadrill, S., Heyes, S., & Holmes, E. (2013). Assessing  
14 mental imagery in clinical psychology: A review of imagery measures and a guiding  
15 framework. *Clinical Psychology Review*, 33(1), 1-23. doi:10.1016/j.cpr.2012.09.001
- 16 Pearson, D., Logie, R., & Green, C. (1996). Mental manipulation, visual working memory, and  
17 executive processes. *Psychologische Beitrage*, 38(3-4), 324-342.
- 18 Pearson, J., Rademaker, R., & Tong, F. (2011). Evaluating the Mind's Eye: The Metacognition of  
19 Visual Imagery. *Psychological Science*, 22(12), 1535-1542.  
20 doi:10.1177/0956797611417134

- 1 Perpiñá, C., Hemsley, D., Treasure, J., & De Silva, P. (1993). Is the selective information  
2 processing of food and body words specific to patients with eating disorders?.  
3 *International Journal Of Eating Disorders*, 14(3), 359-366. doi:10.1002/1098-  
4 108x(199311)14:3<359::aid-eat2260140314>3.0.co;2-g
- 5 Presnell, K., Bearman, S., & Stice, E. (2004). Risk factors for body dissatisfaction in adolescent  
6 boys and girls: A prospective study. *International Journal Of Eating Disorders*, 36(4),  
7 389-401. doi:10.1002/eat.20045
- 8 Pylyshyn, Z. (1981). The imagery debate: Analogue media versus tacit knowledge.  
9 *Psychological Review*, 88(1), 16-45. doi:10.1037//0033-295x.88.1.16
- 10 Quinn, J. (1996). Irrelevant Pictures in Visual Working Memory. *The Quarterly Journal Of*  
11 *Experimental Psychology Section A*, 49(1), 200-215. doi:10.1080/713755613
- 12 Quinn, J., & McConnell, J. (2006). The interval for interference in conscious visual imagery.  
13 *Memory*, 14(2), 241-252. doi:10.1080/09658210500210019
- 14 Quinn, J., & Ralston, G. (1986). Movement and attention in visual working memory. *The*  
15 *Quarterly Journal Of Experimental Psychology Section A*, 38(4), 689-703.  
16 doi:10.1080/14640748608401621
- 17 Roberts, M., Tchanturia, K., Stahl, D., Southgate, L., & Treasure, J. (2007). A systematic review  
18 and meta-analysis of set-shifting ability in eating disorders. *Psychological Medicine*,  
19 37(08), 1075. doi:10.1017/s0033291707009877

- 1 Rodin, J., Silberstein, L., & Striegel-Moore, R. (1984). Women and weight: A normative  
2 discontent. *Nebraska Symposium On Motivation*, 32, 267-307.
- 3 Rudkin, S., Pearson, D., & Logie, R. (2007). Executive processes in visual and spatial working  
4 memory tasks. *The Quarterly Journal Of Experimental Psychology*, 60(1), 79-100.  
5 doi:10.1080/17470210600587976
- 6 Slotnick, S., Thompson, W., & Kosslyn, S. (2012). Visual memory and visual mental imagery  
7 recruit common control and sensory regions of the brain. *Cognitive Neuroscience*, 3(1), 14-  
8 20. doi:10.1080/17588928.2011.578210
- 9 Smeets, M., Klugkist, I., Rooden, S., Anema, H., & Postma, A. (2009). Mental body distance  
10 comparison: A tool for assessing clinical disturbances in visual body image. *Acta*  
11 *Psychologica*, 132(2), 157-165. doi:10.1016/j.actpsy.2009.03.011
- 12 Stice, E., & Whitenton, K. (2002). Risk factors for body dissatisfaction in adolescent girls: A  
13 longitudinal investigation. *Developmental Psychology*, 38(5), 669-678. doi:10.1037//0012-  
14 1649.38.5.669
- 15 Stice, E., Hayward, C., Cameron, R., Killen, J., & Taylor, C. (2000). Body-image and eating  
16 disturbances predict onset of depression among female adolescents: A longitudinal study.  
17 *Journal Of Abnormal Psychology*, 109(3), 438-444. doi:10.1037//0021-843x.109.3.438
- 18 Stroop, J. (1935). Studies of interference in serial verbal reactions. *Journal Of Experimental*  
19 *Psychology*, 18(6), 643-662. doi:10.1037/h0054651



- 1 Stunkard, A., Sørensen, T., Hanis, C., Teasdale, T., Chakraborty, R., Schull, W., & Schulsinger,  
2 F. (1986). An Adoption Study of Human Obesity. *New England Journal Of Medicine*,  
3 *314*(4), 193-198. doi:10.1056/nejm198601233140401
- 4 Thompson, J. (1999). *Exacting beauty*. Washington, DC: American Psychological Association.
- 5 Tiggemann, M. (2006). The Role of Media Exposure in Adolescent Girls' Body Dissatisfaction  
6 and Drive for Thinness: Prospective Results. *Journal Of Social And Clinical Psychology*,  
7 *25*(5), 523-541. doi:10.1521/jscp.2006.25.5.523
- 8 Vitousek, K., & Hollon, S. (1990). The investigation of schematic content and processing in  
9 eating disorders. *Cognitive Therapy And Research*, *14*(2), 191-214.  
10 doi:10.1007/bf01176209
- 11 Voyer, D., Postma, A., Brake, B., & Imperato-McGinley, J. (2007). Gender differences in object  
12 location memory: A meta-analysis. *Psychonomic Bulletin & Review*, *14*(1), 23-38.  
13 doi:10.3758/bf03194024
- 14 Williams, J. M. G., Mathews, A., & MacLeod, C. (1996). The emotional Stroop task and  
15 psychopathology. *Psychological bulletin*, *120*(1), 3.
- 16  
17

1

2

Table 1. Summary statistics of principal variables

Variable	Valid <i>N</i> observations	Mean	SD	Min	Max	Skewness
BMI	110	24.25	4.53	16.85	39.76	1.42
Age	110	26.53	10.20	18	62	1.63
BSQ	110	45.18	17.90	16	89	0.32
VABD	110	1.34	1.23	-1.00	6.00	0.99
VVIQ	110	75.86	20.86	42.00	145.00	0.49
VPT	110	9.93	2.85	4	15	-0.22
Corsi	110	6.26	1.34	4	9	-0.24
Global Precedence ( <i>MS</i> )	108	-65.91	267.73	-930	758	-0.54
Stroop Interference ( <i>SI</i> )	110	6.12	3.79	-0.75	19.50	0.95

Table 2. Correlations between variables. . \* =  $p < .05$ ; \*\* =  $p < .01$ ; \*\*\* =  $p < .001$

	Age	BSQ	VABD	VVIQ	VPT	Corsi	GP	SI
<i>N</i>	108	108	108	108	108	108	106	108
BMI	.33***	.03	.53***	.12	-.07	-.07	-.06	.36***
Age		.16	.15	.03	-.11	-.22*	-.20*	.45***
BSQ			.56***	.41***	-.22*	-.03	.06	.23*
VABD				.34***	-.31**	-.14	.02	.42***
VVIQ					-.11	.03	-.11	.28**
VPT						.50***	-.01	-.32**
Corsi							.09	-.24*
GP								.06

Table 3. Summaries of the regression models predicting body dissatisfaction

Outcome	Model	Predictor	<i>b</i>	<i>SE b</i>	$\beta$	Independent Contribution to $R^{2+}$
BSQ	1 $\Delta R^2=.04$	(Constant)	39.25	9.76		
		BMI	.59	.43	.15	.01
		Age	-.38	.20	-.20	.03
	2 $\Delta R^2=.25^{***}$	(Constant)	12.67	9.56		
		BMI	.39	.37	.10	.01
		Age	-.34	.18	-.18	.03
		VVIQ	.40	.07	.51 <sup>***</sup>	.25
	3 $\Delta R^2=.07^{**}$	(Constant)	18.18	9.25		
		BMI	.22	.35	.06	<.01
		Age	-.57	.18	-.30 <sup>**</sup>	.07
		VVIQ	.34	.07	.43 <sup>***</sup>	.18
		SI	1.51	.46	.31 <sup>**</sup>	.07
	4 $\Delta R^2=.03^*$	(Constant)	30.38	10.88		
		BMI	.25	.35	.06	<.01
		Age	-.57	.18	-.30 <sup>**</sup>	.068
		VVIQ	.34	.07	.43 <sup>***</sup>	.17
SI		1.17	.48	.24 <sup>*</sup>	.04	
VPT		-1.05	.51	-.18 <sup>*</sup>	.03	
VABD	1 $\Delta R^2=.18^{***}$	(Constant)	-1.21	.52		
		BMI	.10	.02	.44 <sup>***</sup>	.17
		Age	.00	.01	-.03	<.01
	2 $\Delta R^2=.11^{***}$	(Constant)	-2.21	.55		
		BMI	.09	.02	.40 <sup>***</sup>	.14
		Age	.00	.01	-.03	<.01
		VVIQ	.02	.00	.34 <sup>***</sup>	.11
	3 $\Delta R^2=.07^{**}$	(Constant)	-1.16	.61		
		BMI	.09	.02	.40 <sup>***</sup>	.15
		Age	.00	.01	-.05	<.01
		VVIQ	.01	.00	.32 <sup>***</sup>	.10
		VPT	.00	.01	-.03 <sup>**</sup>	.11

Note., \*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$ . + derived from semipartial correlation.