

A peer-reviewed version of this preprint was published in PeerJ on 24 September 2015.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1283) (peerj.com/articles/1283), which is the preferred citable publication unless you specifically need to cite this preprint.

Sparks C. 2015. An examination of disparities in cancer incidence in Texas using Bayesian random coefficient models. PeerJ 3:e1283
<https://doi.org/10.7717/peerj.1283>

An examination of disparities in cancer incidence in Texas using Bayesian random coefficient models

Corey Sparks

Disparities in cancer risk exist between ethnic groups in the United States. These disparities often result from differential access to healthcare, differences in socioeconomic status and differential exposure to carcinogens. This study uses cancer incidence data from the population based Texas Cancer Registry to investigate the disparities in digestive and respiratory cancers from 2000 to 2008. A Bayesian hierarchical regression approach is used. All models are fit using the INLA method of Bayesian model estimation. Specifically, a spatially varying coefficient model of the disparity between Hispanic and Non-Hispanic incidence is used. Results suggest that a spatio-temporal heterogeneity model best accounts for the observed Hispanic disparity in cancer risk. Overall, there is a significant disadvantage for the Hispanic population of Texas with respect to both of these cancers, and this disparity varies significantly over space. The greatest disparities between Hispanics and Non-Hispanics in digestive and respiratory cancers occur in eastern Texas, with patterns emerging as early as 2000 and continuing until 2008.

Title: An examination of disparities in cancer incidence in Texas using Bayesian random coefficient models.

Author: Corey S. Sparks^{1,2}

¹Department of Demography

The University of Texas at San Antonio

501 West Cesar E. Chavez Blvd

San Antonio, TX 78207

Email: corey.sparks@utsa.edu

Phone: 210 458 3166

Fax: 210 458 3164

²Department of Biostatistics and Epidemiology

The University of Texas Health Science Center at San Antonio

7703 Floyd Curl Drive

San Antonio, TX 78229, USA

Keywords: health disparity, Bayesian model, cancer incidence

Abstract: Disparities in cancer risk exist between ethnic groups in the United States. These disparities often result from differential access to healthcare, differences in socioeconomic status and differential exposure to carcinogens. This study uses cancer incidence data from the population based Texas Cancer Registry to investigate the disparities in digestive and respiratory cancers from 2000 to 2008. A Bayesian hierarchical regression approach is used. All models are fit using the INLA method of Bayesian model estimation. Specifically, a spatially varying coefficient model of the disparity between Hispanic and Non-Hispanic incidence is used. Results suggest that a spatio-temporal heterogeneity model best accounts for the observed Hispanic disparity in cancer risk. Overall, there is a significant disadvantage for the Hispanic population of Texas with respect to both of these cancers, and this disparity varies significantly over space. The greatest disparities between Hispanics and Non-Hispanics in digestive and respiratory cancers occur in eastern Texas, with patterns emerging as early as 2000 and continuing until 2008.

1. Introduction

Disparities in cancer incidence and mortality exist between racial and ethnic groups in the United States and worldwide (Du et al., 2007; Elmore et al., 2005; Harper et al., 2009; Hun, Siegel, Morandi, Stock, & Corsi, 2009; McKenzie, Ellison-Loschmann, & Jeffreys, 2010; Siegel, Naishadham, & Jemal, 2012; Vainshtein, 2008). The causes of these disparities have been suggested to be rooted in different levels of socioeconomic status (SES), access to medical care, differential exposure to carcinogenic materials and differential treatment by medical staff of racial and ethnic minorities (Krieger, 2005; Sarfati, Blakely, Shaw, Cormack, & Atkinson, 2006; Schootman et al., 2010). While these causes are often non-specific in their effects of how they directly influence cancer incidence, they do allow us to conceptualize and measure key factors related inequalities in health. Furthermore, understanding disparities in cancer risk and being able to visualize the place-based differences both in the determinants of cancer inequality can be a valuable tool to both scientist and policy maker alike. The goal of this paper is to identify geographic clusters of disparities in cancer risk between the Hispanic and non-Hispanic populations of the state of Texas using current incidence data from a population based cancer registry.

Respiratory and digestive system cancers have been identified as often having direct and identifiable causal pathways associated with them, many of which are behavior or environmentally influenced. Lung cancer is perhaps the most widely recognized environmentally influenced cancer type, with strong evidence to support the effects of smoking, poor diet and direct inhalation of certain carcinogens including asbestos and other indoor air pollutants (Alberg, Ford, & Samet, 2007; Alberg & Samet, 2003; Ruano-Ravina, Figueiras, & Barros-Dios, 2003). The exposure to these carcinogens generally leads to errors in somatic cell growth, such

as chromosomal abnormalities, cellular mutations, and alterations in tumor suppressor cells. Gastrointestinal system cancers also have a variety of causes, with some consistency between the types of cancer, but other types also have distinct know etiologies. For example, hepatocellular carcinoma (primary liver cancer) has been directly linked with hepatitis infection, alcoholic cirrhosis and dietary aflotoxins (El-Serag, 2012; Stuver & Trichopoulos, 2008) while other digestive system cancers, such as colorectal cancers are heavily influenced by dietary and lifestyle factors (Chao et al., 2005). While the specific etiologies of the cancers of these two body systems sometimes have direct causal paths, they are generally thought to be influenced by both behavioral and environmental circumstances, which interact with familial and genetic pathways in complicated ways.

The state of Texas is the second most populous state in the United States, with a current population estimate of 25.7 million persons. Between 2000 and 2010, Texas was the sixth fastest growing state, and the highest in total numerical population gain (Makun & Wilson, 2011). Additionally, it is consistently in the top five fastest growing states in the nation. The Hispanic population of Texas was estimated to be 10.1 million persons, or over 38% of the population in 2013 and Texas has the second largest Hispanic population, behind only California (Makun & Wilson, 2011). In addition to being a large part of the state's population, the Hispanic population also faces socioeconomic disadvantages compared to other ethnic groups. The poverty rate for Texas Hispanics was 25.8% according to the 2010 American Community Survey, while non-Hispanic whites only had an 8.8% poverty rate (United States Department of Commerce, 2012). Likewise, Hispanics were are more likely to be employed in construction related activities (18.7% compared to 6.1% for non-Hispanic Whites), which could expose this population to more risk from air-born carcinogens.

For such a large and dynamic state, little population-based cancer disparity research has been published for Texas. In a recent study of cancer disparities in Texas counties, Phillips et. al. (2011) found that an index of socioeconomic well-being was significantly associated with county-level ratios of metastatic to non-metastatic tumors in all-cause, female genital and lung cancers. In a study of El Paso county, Collins et. al. (2011) found higher cancer risk for the Hispanic population of that area, and they go on to discuss how in El Paso, areas of the city that had the highest levels of Hispanic population who had low levels of education had six times the risk of the more educated areas, and areas with the highest proportion of Hispanic renters had seven times the risk of cancer than other, more socioeconomically advantaged areas. Using a geographically weighted regression approach, (Tian, Wilson, & Zhan, 2011) on data from the Texas Cancer Registry, found not only that that Hispanics and non-Hispanic Blacks faced disparities in breast cancer mortality, but that these disparities varied over space within the state. These studies likewise point to the place-based inequality and increased risks that minority groups, including the Hispanic population, face in certain areas within the state. This study will add to the literature on cancer disparities by employing a spatially oriented statistical analysis for the entire state over a more inclusive time period.

With respect to access-based disparities related to cancer risk, Hispanics have been shown to have lower chances of seeking preventative care (Cristancho, Garces, Peters, & Mueller, 2008; Hosain, Sanderson, Du, Chan, & Strom, 2011; Lantz et al., 2006; Shih, Zhao, & Elting, 2006; Suther & Kiros, 2009) in general, and specifically cancer screening. Reasons for not seeking care include lack of insurance, language barriers and the high cost of health care (Cristancho et al., 2008). In a study of colorectal cancer, Wan et. al. (Wan, Zhan, Lu, &

Tiefenbacher, 2012) found significant disparities for Hispanics and non-Hispanic Blacks in accessibility to care.

1.2 Visualizing disparities across space

From a methodological standpoint, testing for disparities in rates is a relatively straightforward task and a variety of statistical procedures are well suited for it. Specifically, a disparity in two rates can be measured as either a difference in total rates, or as a ratio of risks the groups being compared (Keppel et al., 2005). In terms of visualizing the disparities, this can be more of a challenge. For measuring the disparity between population subgroups, the standardized risk ratio is a useful measure, but it is often subject to noise in the underlying rates, most notably in small populations or in cases of rare disease. Maps of such relative risks, as a result of the noise caused by small populations, often lead to the reporting of unstable risk estimates. Tango (2010) describes a variety of methods for both visualizing and detecting disease clusters. Methods for mapping such risk ratios in a scan-statistic context have been described by Chen and co-authors (2008), and Bayesian disease mapping methods are also cited as being particularly good at mapping spatial disease risk (Anderson, Lee, & Dean, 2014; Choo & Walker, 2008; Earnest et al., 2010; Kim & Oleson, 2008; Lawson, 2013; Lawson et al., 2000; Lee & Mitchell, 2014; Lee & Shaddick, 2010). The Bayesian approach allows for smoothing of the relative risk by combining information across spatial units, as well as across time.

It is the purpose of this paper to investigate the spatial variation in cancer incidence disparities between Hispanic and non-Hispanic populations of the state of Texas between 2000 and 2008 using data from a population-based cancer registry. This research adds to the literature in spatial epidemiology by examining the disparities in these two populations over time and space by using a Bayesian modeling methodology, which models the variation in cancer

disparities between these two populations within the state. The Bayesian modeling framework is used to specify a series of varying coefficient models as a method of both more accurately modeling the disparity between these two populations, but also for visualizing where the disparities between the populations exist. The goal of this process is to provide a locally accurate depiction of health disparities which state and local health officials could use in combating health inequalities.

2. Data and methods

2.1 Data source

Data for this analysis come from the Texas Cancer Registry's (www.dshs.state.tx.us/tcr/) Limited-Use data file from 2000 to 2008. Access to these data was approved by the Texas Department of State Health Services IRB #12-030. These data consist of de-identified individual records of primary cancer diagnoses by oncologists in the state of Texas. For the purposes of this study, relevant variables in the data include year of diagnosis, age, sex, Hispanic ethnicity, International Catalog of Disease for Oncology (ICD-O-3) codes for cancer diagnosis site and county of residence at the time of diagnosis. Two main types of cancer were chosen: digestive system (ICD-O-3 codes C150 – C488) and respiratory system cancers (codes C300 – C399). These cancers were chosen because several of the sub-types of these cancers have been linked to environmental or behavioral influences, and several have also been shown to vary between ethnic groups in their incidence (Howe et al., 2006; Singh & Hiatt, 2006; Singh & Siahpush, 2002; Wiggins, Becker, Key, & Samet, 1993; Willsie & Foreman, 2006). These two cancers are selected for study, because they constitute 41% of all cancers in the state for this period. For the years of this study a total of n=155,652 digestive and n=124,438 respiratory system cases were in the data. The most prevalent form of digestive system cancer was colorectal cancer, with 53% of

digestive cancers, and squamous cell carcinoma of the lung was the most prevalent respiratory cancer, representing 22% of all cases.

The dependent variable in the analysis is the count of either digestive or respiratory cancers in each of the 254 counties of Texas between 2000 and 2008. The data are stratified by ethnicity into two categories Hispanic and non-Hispanic. The stratification of the cases is accomplished by using the Hispanic ethnicity variable in the registry. Thus for each year, there are two separate counts for each cancer type and for each of the 254 counties in the state. Since the dependent variables are counts, they are generally expressed as a standardized ratio of counts to expected counts. This is typically called the standardized incidence ratio (SIR), and is expressed:

$$SIR_{ijk} = y_{ijk}/e_{ijk}$$

Where y_{ijk} is the count of cases in the i^{th} county for the j^{th} year for the k^{th} ethnicity and e_{ijk} is the expected number of cases in the county for each group. Here, to estimate the expected number of cases for each county, year and ethnicity, an assumption of equal risks is used. The expected number of cases in each county, year and ethnicity, e_{ijk} , is calculated by assuming each county has the average incidence rate for the whole state for the period 2000 to 2008, or:

$$e_{ijk} = \sum n_{ijk} * r$$

, where n_{ijk} is the number of residents in each county for each ethnicity, and r is the average incidence rate for the state for the period 2000 to 2008. This is repeated for each type of cancer: digestive and respiratory. This generates a set of expected values for the Hispanic and non-Hispanic population of each county, using the statewide rate and the county population size for each group.

To control for background characteristics of the counties, and to measure proxies for factors affecting cancer risk, four independent variables are constructed. The first of these is the metropolitan status of the county, which is measured as a dummy variable indicating whether the United States Department of Agriculture's Economic Research Service considers the county metropolitan. These counties are coded as 1, and non-metro counties are coded as 0. The poverty rate in each county is calculated from the US Census Bureau's Summary File 3 for 2000, and is expressed as the proportion of all residents living below the poverty line in 1999. The proportion of the labor force in construction is used to measure a crude proxy for occupational exposure to certain carcinogens. This is again measured using the Census's Summary File 3 and expressed as a proportion. Finally, the Area Resource File (US Department of Health and Human Services, 2009) for 2008 is used to measure the number of hospitals in each county per 10,000 residents. This is used as a crude proxy for healthcare access in each county.

2.2 Statistical methods

2.2.1 Model Specification

Since the dependent variable is a count, a Poisson distribution is used to model the outcome. To model this outcome, a log-linear Poisson hierarchical regression model for each county, i , year, j , ethnicity, k , and type of cancer, C , is specified as:

$$y_{Cijk} | \theta_{Cijk} \sim \text{Poisson}(e_{Cijk} * \theta_{Cijk})$$

The relative risk function, θ_{Cijk} , can be parameterized using a number of different models, the present paper considers a Bayesian model specification.

In the Bayesian modeling paradigm all model parameters are considered to be random variables and are given a prior distribution and all inference about these parameters is made from

the posterior distribution of these parameters, given the observed data and the information given in the priors. This is generally referred to as Bayes Theorem, and typically stated as:

$$p(\theta|y) \propto p(y|\theta)p(\theta)$$

Where $p(\theta|y)$ is the posterior distribution of the model parameter of interest, $p(y|\theta)$ is the model likelihood function, here defined as a Poisson likelihood, and $p(\theta)$ is the prior distribution for the parameters in the model. Inference for all parameters is done via their posterior distribution, which can be used to derive mean values, quantiles or other descriptive statistics. One useful method for summarizing these distributions is the Bayesian Credible Interval (BCI), not unlike frequentist confidence interval, which gives the values of the posterior density for each parameter that contain $100*(1-\alpha)\%$ of the posterior density. Inference on these BCI regions usually consists of examining if the null hypothesis value of the parameter, typically zero, is contained in the interval.

Since the primary interest in this paper is the relative difference between the incidence of cancer in the Hispanic and non-Hispanic populations of each county, the simplest way to parameterize the model is as a linear difference in the incidence rates using a simple, unstructured linear predictor. This is the first model considered, and is parameterized as:

$$\begin{aligned} \ln(\theta) &= \alpha_C + \delta_C * eth_{Ci} + \sum_k \beta_{Ck} x_{ik} + u_{Ci} + v_{Ci} + t_{Cj} + \psi_{Cij} \\ \alpha_C &\sim U(-\inf, \inf) \\ \delta_C &\sim N(0, .0001) \\ \beta_{Ck} &\sim N(0, .0001) \\ v_{Ci} &\sim N(0, \tau_{Cv}) \\ u_{Ci} &\sim N\left(\frac{1}{n_j} \sum_{j=i} u_{Cj}, \tau_{Cu} / n_i\right) \\ t_{Cj} &\sim N(0, \tau_{Ct}) \\ \psi_{Cij} &\sim N(0, \tau_{\psi C}) \end{aligned} \quad (\text{Model 1})$$

, which follows the standard form for spatio-temporal disease incidence models commonly used in the literature (Blangiardo & Cameletti, 2015; Blangiardo, Cameletti, Baio, & Rue, 2013; Held, Graziano, Frank, & Rue, 2006; Knorr-Held, 2000; Lawson, 2013; Lee & Mitchell, 2014; Schrodle & Held, 2011b; Ugarte, Goicoa, Ibanez, & Militino, 2009). This model specifies the relative risk as a linear function of a grand intercept for each cancer type, α_C , a mean difference between the two ethnicities (eth) for each cancer type, δ_C , a linear predictor effect of the independent variables for each cancer type, $\sum \beta_{kC} x_{ik}$, a “convolution” spatial prior, corresponding to the Besag, York and Mollie (Besag, York, & Mollie, 1991) model, which incorporates an unstructured heterogeneity term for each county and cancer type, v_{Ci} , and a correlated heterogeneity term as a conditionally autoregressive random effect, u_{Ci} , a temporally unstructured random effect for each year and cancer type, t_{Cj} and finally a spatio-temporal interaction random effect, ψ_{Cij} , which follows the Type 1 specification in Knorr-Held (Knorr-Held, 2000). In this model there is a single parameter for measuring the disparity between Hispanics and non-Hispanics for each cancer type, and this is done on average for the entire state. This model additionally captures the underlying characteristics of the counties, the overall

¹ Other prior distributions, including a first order random walk (RW1) priors were used, but did not increase model fit in this case, so the simpler exchangeable random effect for time was used in the final model.

spatial structure of cancer risk, and the temporal variation between years in the relative risk. Priors are assigned to all parameters in a minimally informative fashion, with an improper flat prior for α_C , high variance Normal distribution priors for the δ_C and β_C and v_{Ci} , a Normal distribution prior for t_j and vague Gamma priors for the precisions of the unstructured heterogeneity, correlated heterogeneity, temporal and spatio-temporal components. For all models, the Normal distribution priors are specified in terms of their mean and precision, which is common in Bayesian modeling, with the precision being the inverse of the variance: $\tau = 1/\sigma^2$, such that low precisions equal high variances.

A second model adds more flexibility to Model 1 by including a random slope for each county's difference between Hispanic and non-Hispanic risk. This model is specified as:

$$\begin{aligned} \ln(\theta) &= \alpha_C + \delta_C * eth_{Ci} + \sum_k \beta_{Ck} x_{ik} + u_{Ci} + v_{Ci} + t_{Cj} + \psi_{Cij} \\ \alpha_C &\sim U(-\inf, \inf) \\ \delta_C &\sim \delta_{C0} + \delta_{Ci}, \delta_{Ci} \sim N(0, \tau_{C\delta}) \\ \beta_{Ck} &\sim N(0, .0001) \\ v_{Ci} &\sim N(0, \tau_{Cv}) \\ u_{Ci} &\parallel N\left(\frac{1}{n_j} \sum_{j=i} u_{Cj}, \tau_{Cu} / n_i\right) \\ t_{Cj} &\sim N(0, \tau_{Ct}) \\ \psi_{Cij} &\sim N(0, \tau_{\psi C}) \end{aligned} \quad \text{Model 2}$$

which is similar to (1), but includes a δ_{Ci} term which allows the differences between Hispanic and non-Hispanic risk to vary between counties, equivalent to an unstructured random-slopes model for the disparity. This is much like the spatially varying coefficient model discussed elsewhere (Banerjee, Carlin, & Gelfand, 2004; Gelfand, Kim, Sirmans, & Banerjee, 2003), except in this model, the random slope term is not spatially correlated.

A final model adds a correlated slope for the disparity parameter to Model 2. This model follows the example of previous authors, who model the disparity between groups as a spatially

250 autoregressive random slope (Tassone, Waller, & Casper, 2009; Wheeler, Waller, & Elliott,
251 2008). This model has the form:

$$\ln(\theta) = \alpha_c + \delta_c * eth_{Ci} + \sum_k \beta_{ck} x_{ik} + u_{Ci} + v_{Ci} + t_{Cj} + \psi_{Cij}$$

$$\alpha_c \sim U(-\inf, \inf)$$

$$\delta_{Ci} = \delta_{C0} + \delta_{Ci}, \delta_{Ci} \parallel N\left(\frac{1}{n_j} \sum_{j \sim i} \delta_{Cj}, \tau_{C\delta} / n_i\right)$$

$$\beta_{ck} \sim N(0, .0001)$$

$$v_{Ci} \sim N(0, \tau_{Cv})$$

Model 3

$$u_{Ci} \parallel N\left(\frac{1}{n_j} \sum_{j \sim i} u_{Cj}, \tau_{Cu} / n_i\right)$$

$$t_{Cj} \sim N(0, \tau_{Ct})$$

$$\psi_{Cij} \sim N(0, \tau_{\psi C})$$

253 , which smooths the disparity parameter over neighboring counties within the state.

254 For geographic modeling, neighbors are identified using a first order Queen contiguity
255 rule. Other neighbor specifications were examined, specifically a first order rook contiguity rule,
256 and the results were substantively robust to this other neighbor specification. Also, since the
257 precision terms for Bayesian hierarchical models have been shown to be sensitive to prior
258 specifications, a sensitivity analysis is performed. The models specified above all considered
259 Uniform distributions for the standard deviation of each of the precision parameters. To examine
260 the sensitivity of the models to alternative specifications, proper Gamma (.5, .0005) priors are
261 also considered for all precision terms. This prior distribution has been used by other authors,
262 and is thought of to be a sufficiently vague prior for the precision for these parameters.

263 2.3 Computing - INLA

264 The software R (R Development Core Team, 2015) and the R package R-INLA
265 (Martins, Simpson, Lindgren, & Rue, 2013; Rue, Martino, & Chopin, 2009) were used to prepare
266 data for analysis and parameter estimation. The Integrated Nested Laplace Approximation, or

INLA, approach is a recently developed, computationally simpler method for fitting Bayesian models (Rue et al., 2009), compared to traditional Markov Chain Monte Carlo (MCMC) approaches. INLA fits models that are classified as latent Gaussian models, which are applicable in many settings (Martino & Rue, 2010). In general, INLA fits a general form of additive models such as:

$$\eta = \alpha + \sum_{j=1}^{nf} f^{(j)}(u_{ij}) + \sum_{k=1}^{n\beta} \beta_k z_{ki} + \epsilon_i$$

, where η is the linear predictor for a generalized linear model formula, and is composed of a linear function of some variables u , β are the effects of covariates, z , and ϵ is an unstructured residual (Rue et al., 2009). As this model is often parameterized as a Bayesian one, we are interested in the posterior marginal distributions of all the model parameters. Rue and Martino (2007) show that the posterior marginal for the random effects (x) in such models can be approximated as:

$$\tilde{p}(x_i | y) = \sum_k \tilde{p}(x_i | \theta_k, y) \tilde{p}(\theta_k | y) \Delta_k$$

via numerical integration (Rue & Martino, 2007; Schrodle & Held, 2011a, 2011b). The posterior distribution of the hyperparameters (θ) of the model can also be approximated as:

$$\tilde{p}(\theta | y) \propto \frac{p(x, \theta, y)}{\tilde{p}G(x | \theta, y)} |_{x = x^*(\theta)}$$

, where G is a Gaussian approximation of the posterior and $x^*(\theta)$ is the mode of the conditional distribution of $p(x | \theta, y)$. Thus, instead of using MCMC to find an iterative, sampling-based estimate of the posterior, it is arrived at numerically. This method of fitting the spatio-temporal models specified above has been presented by numerous authors (Blangiardo & Cameletti, 2015;

Blangiardo et al., 2013; Lindgren & Rue, 2015; Martins et al., 2013; Schrodle & Held, 2011a, 2011b), with comparable results to MCMC.

To summarize the posterior distributions of the model parameters, posterior means and 95% credible intervals are calculated. Three models specified in 2.2.1 were examined. Model fit and improvement is assessed between the models with the Deviance Information Criterion (DIC) (Spiegelhalter, Best, Carlin, & van der Linde, 2002). The DIC measures the penalized deviance of each model, with the penalty term representing the model's estimated number of parameters. DIC for the INLA models is described in Rue et al. (2009) and uses the model deviance

$$D(\theta) = -2\log(p(y|\theta)) + pD$$

, plus a penalty component, pD , which is an approximate number of parameters in the model. DIC is used, here as a measure of relative model performance, and models with lower DIC values are preferred over those with higher DIC, analogous to the standard AIC criteria.

3. Results

3.1 Descriptive Results

Descriptive statistics for the dependent variable and the predictors are presented in Table 1.

[TABLE 1 HERE]

A gradual increase in the average number of cases per county is observed over the nine years of data. Also, many more cases of both types of cancer (on average) occur to non-Hispanics than to Hispanics. It should be noted that between 25% (2005) and 36% (2000) of counties had a zero count for Hispanic digestive cancer cases and between 38% (2003) and 46% (2002) had a zero count for Hispanic respiratory cancer cases². Also presented in Table 1 are the observed average

² The large number of zeros in the data suggests that a zero-inflated distribution be used as the model likelihood. A zero-inflated Poisson model was considered for the analysis (results

risk ratios for the state for each year. These are calculated as ratio of the observed SIR for Hispanics (SIR_H) and the observed SIR for non-Hispanics (SIR_{NH}) for each year. For digestive cancers, every year shows an elevated risk for Hispanics compared to non-Hispanics, and all years except 2000 show an elevated risk of respiratory cancer for Hispanics. Likewise, respiratory cancers show a consistent trend of higher risk in Hispanics, but not as high as for digestive cancers. With respect to the predictor variables, in 2000 nearly 18 percent of the population of Texas was in poverty, with a wide degree of variation as seen by the inter quartile range. On average there were .66 hospitals per 10,000 people in each county in the state, and there were sixty-five counties with no hospitals. Slightly over 8 percent of the work force was employed in construction, and the USDA considered thirty percent of counties in the state to be metropolitan.

3.2 Results of Bayesian models

Table 2 presents the posterior means of the regression effects for the fixed effects in the three models described above. Also, 95% Bayesian credible intervals are provided for each parameter. Model DIC values are also provided at the bottom of the table for each model. Lastly, summaries for the model hyperparameters provided.

[Table 2 HERE]

Across the three models, some of the fixed predictors show similar patterns. For digestive cancers, the poverty rate shows a negative association with overall cancer risk in Models 1 through 3. This suggests that in areas of higher poverty, the average cancer risk is lower. Respiratory cancer incidence is affected consistently by two of the predictors. The proportion of the work force in construction is positively associated with respiratory cancer risk in the three of

available from the author), but the DIC of said models suggested the Poisson model fit the data better.

the models, potentially suggesting an occupation-specific risk pattern. Likewise, a metropolitan disadvantage is seen, with higher total cancer risk in metropolitan areas. Both of these variables are in line with expectations in terms of respiratory cancer risk.

When the three models are compared using the DIC, Model 3 shows the best model fit for each cancer type, with the DIC being lowest for this model. Strong evidence is present that Model 1 is not adequate to describe the patterns of Hispanic/non-Hispanic disparities in either cancer, as every other model shows large drops in DIC. When comparing Models 2 and 3, strong evidence also exists for adding the spatially correlate random slope term temporal random effect to Model 2, again with a large drop in DIC.

Turning to the Hispanic disparity parameters, in all models, there persists a disparity between Hispanics and non-Hispanics, with the former consistently showing elevated risk for both types of disease, net of the ecological factors, and the random effects. For digestive cancers, we see an increase in risk (e^{δ}) between 5.3 and 16.4 percent, on average and between 3.8 and 20 percent when considering the 95% credible intervals, depending on the model. For respiratory cancers, we see an increase between 11.2 and 16.4 percent on average, and 9.1 and 21.1 percent when examining the credible intervals. For Models 2 and 3, the coefficients of the models are best presented graphically, as each county has an estimate for the disparity for each cancer type. These estimates are presented in Figure 1 as posterior mean estimates of the Hispanic disparity in relative risk (e^{δ_C}) for each county for Models 2 and 3.

[Figure 1 Here]

The first column of Figure 1 shows the Hispanic disparity random effect from Model 2, for respiratory and digestive cancers, respectively, when the disparity parameter was treated as unstructured. The second column of the figure shows the same parameter, when it was treated as

a spatially structured random effect (Model 3). For both respiratory and digestive system cancers, Hispanics show elevated risk in the eastern portion of the state, but they also show elevated risk in the central portion of the state for digestive system cancers, but not for respiratory cancers. The value of these figures is that the actual disparity in risk is being visualized, which shows us where within the state public health officials might try to focus activities in order to reduce the disparity in risk between these two populations.

3.3 Spatio-temporal Relative Risk Estimation

Figure 2 displays the estimated Hispanic relative risk for digestive cancers (e^θ) for each year, 2000 to 2008, estimated from Model 3.

[Figure 2 Here]

The quantity being mapped is the linear predictor of the Poisson distribution (e^θ), with all random effects included, which is interpreted as the model-based standardized incidence ratio (SIR). Each panel in the figure shows the spatial distribution for each year between 2000 and 2008. We see a general concentration of elevated Hispanic digestive cancer risk in the eastern portion of the state, as evidenced by relative risks greater than one (darker blue in color). This pattern is consistent, if not increasing over time, with more counties showing greater Hispanic relative risk over time. Lower risk ($e^\theta < 1$) for Hispanics occurs in North and Western Texas, and also along the border with Mexico, except for a few counties in extreme South Texas in the latter time periods.

[Figure 3 Here]

Figure 3 provides the complementary space-time risk map for the respiratory cancer outcome. Again, we see higher Hispanic risk in Eastern Texas, but perhaps a more concentrated pattern, compared to the digestive cancer maps. Also present is the lower risk in North and West

Texas, as seen in Figure 2 for digestive cancers. Figure 3 also highlights a consistent spatial cluster of high risk in extreme East Texas for a cluster of three to five counties located North of Harris county (city of Houston). These counties include Montgomery, Liberty, San Jacinto, Walker, Polk and Orange. These counties are quite rural and have low proportions of Hispanic residents (average of 9.3%, or about 8,900 Hispanic persons on average per county).

Finally, a sensitivity analysis of alternative priors for the model hyperparameters (all τ 's) showed very close agreement between the vague Gamma (.5, .0005) and the flat prior distributions. Since Model 3 showed evidence of being the best fitting model, the sensitivity analysis focused on its estimates. The precision point estimates for the temporal random effects (τ_t) for the digestive and respiratory cancers, respectively were 478.0 and 1538.8 from the Gamma prior and 441.5 and 1822.5 from the flat prior. The precisions for the uncorrelated heterogeneity (τ_u) were 428.7 and 923.1 for the Gamma prior and 354.0 and 1095.8 for the flat prior. The precisions for the correlated heterogeneity (τ_v) were 92.6 and 20.8 for the Gamma prior and 92.5 and 19.9 for the flat prior. The precisions for the varying disparity parameter were 15.6 and 17.9 from the Gamma and 14.9 and 17.0 from the flat prior. The precisions for the spatio-temporal random effect (τ_{ψ}) were 296.5 and 288.7 for the Gamma prior model and 298.3 and 283.8 for the flat prior model. While this is only one model, the overlap between the precisions is strong enough to validate the results. The one notable difference is the random effect for the unstructured heterogeneity (τ_u), which showed a lower precision (higher variance) in the Gamma prior model, although the parameter's 95% credible interval did show significant overlap between the two prior specifications (Figure 4).

[Figure 4 Here]

4. Discussion

This paper illustrated the application of the Bayesian varying coefficient models to the study of cancer incidence disparities between the Hispanic and non-Hispanic population of Texas over the period 2000 to 2008. This paper adds to the literature in health disparities within the state of Texas by using advanced Bayesian statistical methods to investigate the spatial non-stationarity of health disparities in two major form of cancer incidence. The primary goal of the analysis was to examine the usefulness of the spatially varying coefficient model (Banerjee et al., 2004; Gelfand et al., 2003; Tassone et al., 2009; Wheeler et al., 2008) within the Bayesian modeling framework using a variety of model specifications, including models that included interactions between space and time. Alternative model specifications modeled the disparity in incidence between the two subpopulations differently, from a fixed effect on the grand mean to a spatially varying coefficient model for each county in the state. The flexibility of the Bayesian framework also allowed for the models to be compared using standard model complexity criteria (DIC).

The model that best fit the data was the space-time model with a spatially varying slope for the disparity between Hispanics and non-Hispanics, according to the minimum DIC criteria. This suggests that the disparity between Hispanics and non-Hispanics in these two cancer types is best modeled through a spatially structured model, which allows for spatially structured variation in risk. This also suggests that there are counties within the state where the Hispanic population is at higher risk for both of these cancers, and that these counties typically occur closely to one another spatially.

Overall, a general disparity in terms of both cancers for Hispanics was found, where they face higher risk for both digestive and respiratory cancers than the non-Hispanic population of the state. Significant effects were found on cancer-specific risks consistently including the

county poverty level, metropolitan status of the county and the proportion of the workforce in construction. The labor force composition finding makes sense, as workers in construction industries often face higher levels of exposure to airborne particulates that could increase cancer risk. The finding for the county poverty rate was that in areas with higher poverty, the overall relative risk of cancer was lower, and deserves more discussion. This effect was seen for both cancer types, in all but the final model (Model 3), and is in stark contrast to findings from national data (Singh, Miller, Hankey, & Edwards, 2003) for many types of cancer, which show higher incidence and mortality in both Hispanics and non-Hispanics in areas with higher poverty. Singh et. al. did not use data from Texas, and the time period for the present study is later than those considered in their report. It is possible that the experience of the Texas population is different from the data used in their study; such local variations are common in health research.

This study had one primary limitation; the cancer incidence data had no information on residential histories of the individual cases. Any environmental exposure that could have influenced cancer risk may have come from a previous residential location. Unfortunately, the cancer registry data used in this study had no information on this subject.

Further research is needed to investigate the specifics of the counties identified in the analysis as having excess Hispanic cancer risk. This can be done by a more localized analysis of the individual-level data this analysis is derived, and by investigating housing conditions, access to healthcare and potential environmental contaminants in these areas directly. Such ecological analyses as that presented here are rarely truly informative for individual cancer diagnoses, but they can be very influential in terms of public health activities to reduce cancer disparities at the population level.

References

- Alberg, A. J., Ford, J. G., & Samet, J. M. (2007). Epidemiology of lung cancer: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest*, 132(3 Suppl), 29S-55S. doi: 10.1378/chest.07-1347
- Alberg, A. J., & Samet, J. M. (2003). Epidemiology of lung cancer. *Chest*, 123(1 Suppl), 21S-49S.
- Anderson, C., Lee, D., & Dean, N. (2014). Identifying clusters in Bayesian disease mapping. *Biostatistics*, 15(3), 457-469. doi: 10.1093/biostatistics/kxu005
- Banerjee, S., Carlin, B. P., & Gelfand, A. E. (2004). *Hierarchical modeling and analysis for spatial data*. Boca Raton: CRC/ Chapman & Hall.
- Besag, J., York, J. , & Mollie, A. (1991). Bayesian image restoration, with two applications in spatial statistics. *Annals of the Institute of Statistical Mathetmatics*, 43, 1-59.
- Blangiardo, M., & Cameletti, M. (2015). *Spatial and Spatio-temporal Bayesian Models with R-INLA*: Wiley.
- Blangiardo, M., Cameletti, M., Baio, G., & Rue, H. (2013). Spatial and spatio-temporal models with R-INLA. *Spat Spatiotemporal Epidemiol*, 7, 39-55.
- Chao, A., Thun, M. J., Connell, C. J., McCullough, M. L., Jacobs, E. J., Flanders, W. D., . . . Calle, E. E. (2005). Meat consumption and risk of colorectal cancer. *JAMA*, 293(2), 172-182. doi: 10.1001/jama.293.2.172
- Chen, J., Roth, R. E., Naito, A. T., Lengerich, E. J., & MacEachren, A. M. (2008). Geovisual analytics to enhance spatial scan statistic interpretation: an analysis of US cervical cancer mortality. *International Journal of Health Geographics*, 7.
- Choo, L., & Walker, S. G. (2008). A new approach to investigating spatial variations of disease. *Journal of the Royal Statistical Society Series a-Statistics in Society*, 171, 395-405.
- Collins, T. W., Grineski, S. E., Chakraborty, J., & McDonald, Y. J. (2011). Understanding environmental health inequalities through comparative intracategorical analysis: Racial/ethnic disparities in cancer risks from air toxics in El Paso County, Texas. *Health & Place*, 17(1), 335-344.
- Cristancho, S., Garces, D. M., Peters, K. E., & Mueller, B. C. (2008). Listening to rural Hispanic immigrants in the midwest: A community-based participatory assessment of major barriers to health care access and use. *Qualitative Health Research*, 18(5), 633-646.
- Du, X. L., Fang, S., Vernon, S. W., El-Serag, H., Shih, Y. T., Davila, J., & Rasmus, M. L. (2007). Racial disparities and socioeconomic status in association with survival in a large population-based cohort of elderly patients with colon cancer. *Cancer*, 110(3), 660-669.
- Earnest, A., Beard, J. R., Morgan, G., Lincoln, D., Summerhayes, R., Donoghue, D., . . . Mengersen, K. (2010). Small area estimation of sparse disease counts using shared component models-application to birth defect registry data in New South Wales, Australia. *Health & Place*, 16(4), 684-693.
- El-Serag, H. B. (2012). Epidemiology of Viral Hepatitis and Hepatocellular Carcinoma. *Gastroenterology*, 142(6), 1264-1273.
- Elmore, J. G., Nakano, C. Y., Linden, H. M., Reisch, L. M., Ayanian, J. Z., & Larson, E. B. (2005). Racial inequities in the timing of breast cancer detection, diagnosis, and initiation of treatment. *Medical Care*, 43(2), 141-148.

- Gelfand, A. E., Kim, H., Sirmans, C. F., & Banerjee, S. (2003). Spatial modeling with spatially varying coefficient processes. *Journal of the American Statistical Association*, 98, 387-396.
- Harper, S., Lynch, J., Meersman, S. C., Breen, N., Davis, W. W., & Reichman, M. C. (2009). Trends in Area-Socioeconomic and Race-Ethnic Disparities in Breast Cancer incidence, Stage at Diagnosis, Screening, Mortality, and Survival among Women Ages 50 Years and Over (1987-2005). *Cancer Epidemiology Biomarkers & Prevention*, 18(1), 121-131.
- Held, L., Graziano, G., Frank, C., & Rue, H. (2006). Joint spatial analysis of gastrointestinal infectious diseases. *Stat Methods Med Res*, 15(5), 465-480. doi: 10.1177/0962280206071642
- Hosain, G. M. M., Sanderson, M., Du, X. L. L., Chan, W. Y., & Strom, S. S. (2011). Racial/Ethnic Differences in Predictors of Psa Screening in a Tri-Ethnic Population. *Central European Journal of Public Health*, 19(1), 30-34.
- Howe, H. L., Wu, X. C., Ries, L. A. G., Cokkinides, V., Ahmed, F., Jemal, A., . . . Edwards, B. K. (2006). Annual report to the nation on the status of cancer, 1975-2003, featuring cancer among US Hispanic/Latino populations. *Cancer*, 107(8), 1711-1742.
- Hun, D. E., Siegel, J. A., Morandi, M. T., Stock, T. H., & Corsi, R. L. (2009). Cancer risk disparities between hispanic and non-hispanic white populations: the role of exposure to indoor air pollution. *Environ Health Perspect*, 117(12), 1925-1931. doi: 10.1289/ehp.0900925
- Keppel, K., Pamuk, E., Lynch, J., Carter-Pokras, O., Kim, Insun, Mays, V., . . . Weissman, J. S. (2005). Methodological issues in measuring health disparities. *Vital Health Stat* 2(141), 1-16.
- Kim, H., & Oleson, J. J. (2008). A Bayesian dynamic spatio-temporal interaction model: An application to prostate cancer incidence. *Geographical Analysis*, 40(1), 77-96.
- Knorr-Held, L. (2000). Bayesian modelling of inseparable space-time variation in disease risk. *Statistics in Medicine*, 19(17-18), 2555-2567. doi: 10.1002/1097-0258(20000915/30)19:17/18<2555::AID-SIM587>3.0.CO;2-# [pii]
- Krieger, N. (2005). Defining and investigating social disparities in cancer: critical issues? *Cancer Causes & Control*, 16(1), 5-14.
- Lantz, P. M., Mujahid, M., Schwartz, K., Janz, N. K., Fagerlin, A., Salem, B., . . . Katz, S. J. (2006). The influence of race, ethnicity, and individual socioeconomic factors on breast cancer stage at diagnosis. *American Journal of Public Health*, 96(12), 2173-2178.
- Lawson, A. B. (2013). *Bayesian Disease Mapping: Hierarchical Modeling in Spatial Epidemiology, Second Edition*. Boca Raton: CRC Press.
- Lawson, A. B., Biggeri, A. B., Boehning, D., Lesaffre, E., Viel, J. F., Clark, A., . . . Grp, Dis Mapping Collaborative. (2000). Disease mapping models: an empirical evaluation. *Statistics in Medicine*, 19(17-18), 2217-2241.
- Lee, D., & Mitchell, R. (2014). Controlling for localised spatio-temporal autocorrelation in long-term air pollution and health studies. *Stat Methods Med Res*, 23(6), 488-506. doi: 10.1177/0962280214527384
- Lee, D., & Shaddick, G. (2010). Spatial modeling of air pollution in studies of its short-term health effects. *Biometrics*, 66(4), 1238-1246. doi: 10.1111/j.1541-0420.2009.01376.x
- Lindgren, F., & Rue, H. (2015). Bayesian Spatial Modelling with R-INLA. *Journal of Statistical Software*, 63(19), 1-25.

- Makun, P., & Wilson, S. (2011). Population Distribution and Change: 2000 to 2010. Washington D.C.: U.S. Department of Commerce.
- Martino, S., & Rue, H. (2010). Case studies in Bayesian computation using INLA. In P. Mantovan & P. Secchi (Eds.), *Complex Data Modeling and Computationally Intensive Statistical Methods* (pp. 99-114). Milan: Springer-Verlag.
- Martins, T. G., Simpson, D., Lindgren, F., & Rue, H. (2013). Bayesian computing with INLA: New features. *Computational Statistics & Data Analysis*, 67, 68-83. doi: Doi 10.1016/J.Csda.2013.04.014
- McKenzie, F., Ellison-Loschmann, L., & Jeffreys, M. (2010). Investigating reasons for socioeconomic inequalities in breast cancer survival in New Zealand. *Cancer Epidemiology*, 34(6), 702-708.
- Philips, B. U., Gong, G., Hargrave, K. A., Belasco, E., & Lyford, C. P. (2011). Correlation of the ratio of metastatic to non-metastatic cancer cases with the degree of socioeconomic deprivation among Texas counties. *International Journal of Health Geographics*, 10.
- R Development Core Team. (2015). R: A language and environment for statistical computing (Version 3.2.0). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.r-project.org>
- Ruano-Ravina, A., Figueiras, A., & Barros-Dios, J. M. (2003). Lung cancer and related risk factors: an update of the literature. *Public Health*, 117(3), 149-156. doi: S0033-3506(02)00023-9 [pii] 10.1016/S0033-3506(02)00023-9
- Rue, H., & Martino, S. (2007). Approximate Bayesian inference for hierarchical Gaussian Markov random field models. *Journal of Statistical Planning and Inference*, 137(10), 3177-3192. doi: Doi 10.1016/J.Jspi.2006.07.016
- Rue, H., Martino, S., & Chopin, N. (2009). Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society Series B-Statistical Methodology*, 71, 319-392. doi: Doi 10.1111/J.1467-9868.2008.00700.X
- Sarfati, D., Blakely, T., Shaw, C., Cormack, D., & Atkinson, J. (2006). Patterns of disparity: ethnic and socio-economic trends in breast cancer mortality in New Zealand. *Cancer Causes & Control*, 17(5), 671-678.
- Schootman, M., Lian, M., Deshpande, A. D., Baker, E. A., Pruitt, S. L., Aft, R., & Jeffe, D. B. (2010). Temporal trends in area socioeconomic disparities in breast-cancer incidence and mortality, 1988-2005. *Breast Cancer Research and Treatment*, 122(2), 533-543.
- Schrodle, B., & Held, L. (2011a). A primer on disease mapping and ecological regression using INLA. *Computational Statistics*, 26(2), 241-258. doi: Doi 10.1007/S00180-010-0208-2
- Schrodle, B., & Held, L. (2011b). Spatio-temporal disease mapping using INLA. *Environmetrics*, 22(6), 725-734. doi: Doi 10.1002/Env.1065
- Shih, Y. C. T., Zhao, L. R., & Elting, L. S. (2006). Does Medicare coverage of colonoscopy reduce racial/ethnic disparities in cancer screening among the elderly? *Health Affairs*, 25(4), 1153-1162.
- Siegel, R., Naishadham, D., & Jemal, A. (2012). Cancer statistics for Hispanics/Latinos, 2012. *CA Cancer J Clin*, 62(5), 283-298. doi: 10.3322/caac.21153
- Singh, G. K., & Hiatt, R. A. (2006). Trends and disparities in socioeconomic and behavioural characteristics, life expectancy, and cause-specific mortality of native-born and foreign-

- born populations in the United States, 1979-2003. *International Journal of Epidemiology*, 35(4), 903-919.
- Singh, G. K., Miller, B. A., Hankey, B. F., & Edwards, B. K. (2003). Area Socioeconomic Variations in U.S. Cancer Incidence, Mortality, Stage, Treatment, and Survival, 1975–1999. *NCI Cancer Surveillance Monograph Series* (Vol. 4). Bethesda, MD: National Cancer Institute.
- Singh, G. K., & Siahpush, M. (2002). Ethnic-immigrant differentials in health behaviors, morbidity, and cause-specific mortality in the United States: An analysis of two national data bases. *Human Biology*, 74(1), 83-109.
- Spiegelhalter, D. J., Best, N., Carlin, B. P., & van der Linde, A. (2002). Bayesian measures of model complexity and fit (with discussion). *Journal of the Royal Statistical Society Series B-Methodological*, 64, 583-639.
- Stuver, S., & Trichopoulos, D. (2008). Cancer of the liver and biliary tract. In H. Adami, D. Hunter & D. Trichopoulos (Eds.), *Textbook of Cancer Epidemiology* (pp. 308-332). Oxford: Oxford University Press.
- Suther, S., & Kiros, G. E. (2009). Barriers to the use of genetic testing: A study of racial and ethnic disparities. *Genetics in Medicine*, 11(9), 655-662.
- Tango, T. (2010). *Statistical Methods for Disease Clustering*. New York: Springer.
- Tassone, E. C., Waller, L. A., & Casper, M. L. (2009). Small-area racial disparity in stroke mortality: an application of bayesian spatial hierarchical modeling. *Epidemiology*, 20(2), 234-241. doi: 10.1097/EDE.0b013e3181935aee
- Tian, N., Wilson, J. G., & Zhan, F. B. (2011). Spatial association of racial/ethnic disparities between late-stage diagnosis and mortality for female breast cancer: where to intervene? *International Journal of Health Geographics*, 10, 24. doi: 1476-072X-10-24 [pii] 10.1186/1476-072X-10-24
- Ugarte, M. D., Goicoa, T., Ibanez, B., & Militino, A. R. (2009). Evaluating the performance of spatio-temporal Bayesian models in disease mapping. *Environmetrics*, 20(6), 647-665. doi: 10.1002/env.969
- United States Department of Commerce. (2012). American Factfinder 2. Retrieved January 27, 2012 <http://factfinder2.census.gov>
- US Department of Health and Human Services. (2009). Area Resource File (ARF) 2008-2009.
- Vainshtein, J. (2008). Disparities in breast cancer incidence across racial/ethnic strata and socioeconomic status: A systematic review. *Journal of the National Medical Association*, 100(7), 833-839.
- Wan, N., Zhan, F. B., Lu, Y., & Tiefenbacher, J. P. (2012). Access to healthcare and disparities in colorectal cancer survival in Texas. *Health & Place*, 18(2), 321-329. doi: S1353-8292(11)00204-8 [pii] 10.1016/j.healthplace.2011.10.007
- Wheeler, D. C., Waller, L. A., & Elliott, J. O. (2008). Modeling epilepsy disparities among ethnic groups in Philadelphia, PA. *Stat Med*, 27(20), 4069-4085. doi: 10.1002/sim.3261
- Wiggins, C. L., Becker, T. M., Key, C. R., & Samet, J. M. (1993). Cancer Mortality among New-Mexico Hispanics, American-Indians, and Non-Hispanic Whites, 1958-1987. *Journal of the National Cancer Institute*, 85(20), 1670-1678.
- Willisie, S. K., & Foreman, M. G. (2006). Disparities in lung cancer: Focus on Asian Americans and Pacific Islanders, American Indians and Alaska Natives, and Hispanics and Latinos. *Clinics in Chest Medicine*, 27(3), 441-+.

Table 1. Descriptive statistics for dependent and independent variables used in the analysis.

Cancer Type and Year	Mean # Cases	IQR	Mean # Cases (non-Hispanic)	Mean # Cases (Hispanic)	Mean SIR _H /SIR _{NH}
Digestive Cancer Cases per County					
2000	30.9	18	49.9	12.0	0.87
2001	32.2	18	51.8	12.6	1.44
2002	32.9	19	52.6	13.2	1.18
2003	33.7	19.25	53.5	14.0	1.14
2004	34.4	22	54.0	14.8	1.31
2005	34.8	22	53.9	15.8	1.32
2006	35.2	21	54.3	16.1	1.30
2007	36.1	23	55.8	16.4	1.46
2008	36.1	20	55.1	17.0	2.06
	155,652 total cases				
Respiratory Cancer Cases per County					
2000	25.6	15	46.0	5.2	1.28
2001	26.5	17	47.2	5.8	1.42
2002	26.9	17	48.2	5.6	1.16
2003	27.8	17	49.4	6.1	1.62
2004	27.6	16.25	49.2	5.9	1.18
2005	28.1	17	49.9	6.4	1.48
2006	27.4	16	48.4	6.5	1.67
2007	27.8	16	48.7	6.8	1.61
2008	27.2	15	48.1	6.4	1.54
	123,438 total cases				
Predictors	Mean	IQR			
% in Poverty	17.76	6.58			
Hospitals/10,000 People	0.66	0.79			
% in Construction	8.11	3.15			
% Metro Counties	30.31	1.00			

n=254 counties

Table 1 (on next page)

Results from Bayesian Models

1 Table 2. Results for the alternative Bayesian model specification parameters.

	Model 1		Model 2		Model 3	
Parameter	Posterior Mean (95% Credible Interval)		Posterior Mean (95% Credible Interval)		Posterior Mean (95% Credible Interval)	
	Digestive	Respiratory	Digestive	Respiratory	Digestive	Respiratory
α	-.081 (-.119 - -.043)	-.066 (-.095 - -.037)	-.098 (-.137 - -.059)	-.074 (-.103 - -.044)	-.097 (-.136 - -.057)	-.074 (-.103 - -.044)
β						
% in Poverty	-.031 (-.052 - -.010)	.002 (-.027 - .033)	-.034 (-.057 - -.011)	.001 (-.031 - .032)	-.033 (-.057 - -.010)	.001 (-.032 - .030)
Hospitals per capita	-.016 (-.037 - .004)	-.007 (-.032 - .016)	-.015 (-.037 - .005)	-.008 (-.033 - .016)	-.016 (-.037 - .005)	-.007 (-.032 - .018)
% in Construction	-.011 (-.027 - .005)	.050 (.028 - .072)	-.009 (-.026 - .008)	.050 (.027 - .072)	-.001 (-.026 - .008)	.050 (.028 - .073)
Metro County	.023 (-.009 - .056)	.052 (.007 - .095)	.023 (-.011 - .057)	.054 (.009 - .099)	.021 (-.011 - .056)	.054 (.009 - .099)
Hispanic Disparity, δ	.052 (.038 - .066)	.107 (.087 - .126)	.138 (.106 - .171)	.146 (.109 - .184)	.152 (.122 - .183)	.152 (.112 - .192)
Model Fit						
Deviance (\bar{D})	21256.2	18625.7	20790.2	18462.5	20775.6	18436.8
DIC	21630.2	19004.4	21240.7	18888.5	21217.2	18859.9
pD	373.9	378.7	449.9	426.0	441.6	423.1
Hyperparameters						
τ_t	477.8	1552.5	478.6	1546.5	478.0	1538.8
τ_u	331.3	555.6	432.3	898.1	428.7	923.1
τ_v	133.9	24.2	93.7	20.4	92.6	20.8
τ_δ	-	-	52.3	67.5	15.6	17.9
τ_ψ	297.1	284.8	296.2	287.3	296.5	288.7


2 *Parameters in bold type represent estimates whose credible intervals do not contain 0.

3
4

1

Hispanic Relative Risk Estimated from Models 2 and 3


Hispanic Relative Risk (e^{δ}) Estimated from Models 2 and 3



2

Hispanic Fitted SIR 2000 to 2008 for Digestive Cancers


Hispanic Fitted SIR 2000 to 2008



3

Hispanic Fitted SIR 2000 to 2008 for Respiratory Cancers

Hispanic Fitted SIR 2000 to 2008



4

Marginal Densities for Model Hyperparameter

Marginal Density for Model Hyperparameter τ_u

