

The Neurophysiological Response to Manual Therapy and Its Analgesic Implications: A Narrative Review

Andrew D. Vigotsky¹

Ryan P. Bruhns^{1,2}

¹ Kinesiology Program, Arizona State University, Phoenix, Arizona, United States of America

² Department of Chemistry and Biochemistry, Arizona State University, Tempe, Arizona, United States of America

ABSTRACT

Manual therapy has long been a component of physical rehabilitation programs, especially to treat those in pain. The mechanisms of manual therapy, however, are not fully understood, and it has been suggested that its pain modulatory effects are of neurophysiological origin, and may be mediated by the descending modulatory circuit. Therefore, the purpose of this review is to examine the neurophysiological response of different types of manual therapy, in order to better understand the neurophysiological mechanisms behind each therapy's analgesic effects. It is concluded that different forms of manual therapy elicit analgesic effects via different mechanisms. Additionally, future avenues of mechanistic research pertaining to manual therapy are discussed.

Introduction

Manual therapy has been a component of physical rehabilitation programs since as early as 400 BC (Pettman 2007). Since its inception, many variations of manual therapy techniques have been developed and marketed. Each year, upwards of \$8.1 billion is spent in the US on manual therapies, including chiropractic/osteopathic manipulation and massage (Nahin et al. 2009). Despite the large annual financial expenditures on manual therapies, its mechanisms are not yet fully understood. Current research suggests that a neurophysiological response to manual therapy is responsible for clinically significant decreases in pain (Bialosky et al. 2009). Included in the neurophysiological response is the descending pain modulation circuit, which may be a principle mechanism in the analgesic effect of manual therapies.

Descending Modulation of Pain

Melzack & Wall (1965) were the first to explain the potential mechanisms of a central pain modulatory system, wherein the authors described the gate control theory of pain, which simply states that non-noxious input suppresses painful output by inhibiting dorsal root nociceptors. Numerous neurotransmitters, including serotonin (5-HT), endocannabinoids, and endogenous opioids (EO), have been shown to act on the rostral ventromedial medulla (RVM) and periaqueductal grey (PAG) in order to modulate nociceptive circuits and pain output to (Adams et al. 1986; Benedetti et al. 2013; Fields et al. 1991; Mason 1999; Nadal et al. 2013; Ossipov et al. 2010).

β -endorphins are EO peptides that have not only been shown to have a comparable analgesic effect to morphine (Gerrits et al. 2003), but are 18 to 33 times more potent (Loh et al. 1976). Diffuse noxious inhibitory control (DNIC) is the process by which afferent noxious signals are inhibited from the peripheral nervous system (PNS). Using a rat model, Le Bars et al.

(1979a); Le Bars et al. (1979b) found that neurons were inhibited by noxious stimuli (a hot bath), therein coining the term DNIC. Since then, multiple studies have suggested that EO are an underlying mechanism of DNIC (Chitour et al. 1982; Kraus et al. 1981).

Being that the analgesic effects of both human touch (Lindgren et al. 2012) and placebo (Colloca et al. 2013; Eippert et al. 2009; Morton et al. 2014; Sauro & Greenberg 2005; Wager et al. 2007; Zubieta et al. 2005) are mediated by an EO response, it is imperative that a placebo control group be utilized in research examining the neurochemical response to manual therapy, as placebo and touch alone are confounding variables. Previous reviews have noted potential descending modulatory mechanisms – an endogenous opioid response – in both physical therapy (Bender et al. 2007) and physical medicine (Crielaard et al. 1983); however, the neurochemical response to manual therapy and its implications for descending pain modulation, to the authors' knowledge, have not yet been thoroughly reviewed.

Manipulation Therapies

Through the millennia, numerous types of manipulation therapies have been developed and advocated, and have been purported to cure everything from scarlet fever and diphtheria to hearing loss (Pettman 2007). However, perhaps the most widely proclaimed outcome from manipulative therapy is pain relief, which may be modulated by neurochemicals that act on the RVM and PAG.

Osteopathic Manipulative Therapy

Degenhardt et al. (2007) recruited twenty male subjects: ten with low back pain, ten without. Four osteopathic manipulative therapy (OMT) techniques (articulatory treatment system, muscle energy, soft tissue technique, and Strain-Counterstrain) were performed on areas of subjects' "somatic dysfunction", defined as, "sites of muscle hypertonicity, tenderness, and

joint restriction” (Degenhardt et al. 2007). Blood was collected prior to (baseline), 30 minutes, and 24 hours after OMT. Increases in β -endorphin and N-palmitoylethanolamide (PEA) – an endogenous analog of arachidonylethanolamide (AEA), or anandamine, an endocannabinoid – were observed 30 minutes post treatment; at 24 hours, similar biomarker changes from baseline were found. Subjects with chronic low back pain experienced greater biomarker alterations following OMT than the control (asymptomatic) group. However, because no true control or sham group was utilized, it is not possible to distinguish whether these changes in biomarkers were due to the placebo effect, or something greater. Although, these data do show that those in pain respond differently to treatment than asymptomatic individuals.

In a blinded, randomized control trial, McPartland et al. (2005) investigated the effects of OMT on plasma endocannabinoid concentrations; that is, AEA and 2-arachidonoylglycerol (2-AG). Thirty-one subjects received either an OMT treatment (biodynamic osteopathy in the cranial field) or a sham treatment. Importantly, subjects were recruited from a patient population of an osteopath who regularly uses OMT; therefore, the patients most likely believe the treatment is efficacious. No changes were observed in 2-AG concentrations in either group. In the sham group, negligible, insignificant changes in AEA were observed (17%). The OMT group experienced a 168% increase (5.02 pmol/mL) in AEA over baseline, but this increase did not achieve statistical significance; however, this difference may certainly be clinically relevant, as indicated by changes in Drug Reaction Scale (DRS) scores. These data suggest that endocannabinoids do play a role in the analgesic effect of OMT.

Spinal Manipulation

A number of studies have investigated the pain modulation mechanisms of spinal manipulation which, as the name implies, is specific only to spinal articulation. The first to do so

were Vernon et al. (1986), who found a small but statistically significant increase in plasma β -endorphin levels in the experimental group, but not in the sham or control groups. However, two subsequent studies had findings in contrast to Vernon et al. (1986). Christian et al. (1988) and Sanders et al. (1990) both failed to find increases in plasma β -endorphin levels. Christian et al. (1988) did note a decrease in plasma cortisol levels, but this decrease also occurred in the sham groups.

Recently, Plaza-Manzano et al. (2014) compared cervical and thoracic manipulations to a control group. Both cervical and thoracic groups saw decreases in neurotensin, increases in orexin A, and decreases in oxytocin. Only the cervical group saw a decrease in cortisol.

Multiple reviews have also investigated the pain modulating mechanisms of spinal manipulation (Pickar 2002; Vernon 2000), and are in agreement that the analgesic origins are neurophysiological in nature, occurring through some type of descending pain modulation circuit. The exact circuit, however, is not fully understood, and it appears that different types of spinal manipulations, namely the velocity with which and the location at which they are performed, may elicit different neurochemical responses indicative of different descending pain modulation mechanisms.

Knee Joint Manipulation

Skyba et al. (2003) investigated the effects of knee joint manipulation in rats on monoamine, opioid, and gamma-aminobutyric acid (GABA) receptors in the spinal cord. Investigators found that the analgesic effects of knee joint manipulation were not impacted by the spinal blockade of opioid or GABA receptors, but *were* impacted by blocking the receptors of 5-HT and norepinephrine. Therefore, it was posited that descending inhibition following knee

joint manipulation may be modulated by serotonergic and noradrenergic mechanisms. These findings have yet to be replicated in humans.

Mobilization Therapies

Ankle Joint Mobilization

In mice, ankle mobilization-induced analgesia has been shown to be mediated by EO pathways (Martins et al. 2012). Importantly, researchers noted that the bottleneck in antihypersensitivity was opioid receptor availability, rather than opioid-containing leukocytes. Although opioid receptor availability may be the bottleneck in mice, this is not necessarily true for humans. These data should be replicated in human subjects, and could have large implications for those in chronic pain or those with central sensitization, as these individuals may have decreased opioid receptor availability (DosSantos et al. 2012), and therefore may not benefit as much from this technique.

Mulligan's Mobilization with Movement

Paungmali and colleagues have studied Mulligan's Mobilization with Movement (MWM) in lateral epicondylalgia (Paungmali et al. 2004; Paungmali et al. 2003). Twenty-four subjects with unilateral chronic lateral epicondylalgia were treated with MWM on six occasions at least two days apart. No significant decreases in hypoalgesic effects were seen over the treatment period (Paungmali et al. 2003). In a follow up study, Paungmali et al. (2004) failed to antagonize the hypoalgesic effects of MWM with naloxone, an opioid antagonist, and concluded that MWM works through nonopioid methods. However, as noted by Payson & Holloway (1984), naloxone by itself can produce an analgesic effect due to its inhibitory effects on inflammation and ischemia; therefore, the results of Paungmali et al. (2004) should be called into question.

Neural Mobilization

Utilizing male Wistar rats and Western blot assays of the PAG, Santos et al. (2014) examined the brains of rats following neural mobilization for mu-, delta-, and kappa-opioid receptor expression. Researchers did not find changes in delta- and mu-opioid receptor expression following neural mobilization, but kappa-opioid receptor expression underwent a significant increase, by 17%. These data indicate that neural mobilization may be modulated by EOs that work on kappa-opioid receptors, such as dynorphin.

Massage Therapies

Massage therapy is often sought for both pleasure and therapy. It has been proposed to work through the gate control theory of pain, initially described by Melzack & Wall (1965) (Field 2014). However, Field (2014) failed to note that different types of massage therapy may work via different mechanisms, nor did Field dive deeply into possible mechanisms. Therefore, a more comprehensive review of massage therapy's mechanisms is warranted.

Connective Tissue Massage

Connective tissue massage is intended to both decrease pain and increase range of motion (Threlkeld 1992). Kaada & Torsteinbo (1989) described a significant increase in plasma β -endorphin levels following connective tissue massage, similar to the time course seen in acupuncture and similar to the magnitude seen during exercise. These results are indicative of a DNIC response, which modulates pain through descending inhibition.

Acupressure

Using naloxone in rats, Trentini et al. (2005) suggested the antinociceptive effects of acupressure are mediated by EOs. Despite this, changes in plasma β -endorphin levels were not observed in follow-up research in humans (Fassoulaki et al. 2007). However, Fassoulaki et al.

(2007) only investigated the effects of one acupressure point and a sham acupressure point, both on the face. Thus, the effects of acupressure on other parts of the body remain unclear.

Conventional Massage

Regular massage, consisting of effleurage and other common techniques, has been well studied. Day et al. (1987) were the first to note there is no change in plasma β -endorphin or β -lipotropin levels following back massage; the control group also did not experience changes.

Since then, a couple of studies have found that massage increases urine concentration of dopamine and serotonin (Hernandez-Reif et al. 2001; Hernandez-Reif et al. 2004), suggesting that massage therapy's analgesic effects are mediated by dopaminergic and serotonergic pathways. A review of the mechanisms of massage therapy noted a 31% decrease in cortisol levels and a 28 and 31% increase in serotonin and dopamine levels, respectively (Field et al. 2005).

Future Research

For some therapies, such as manipulation, a minimal amount of force may be required for an analgesic effect (McLean et al. 2002), but whether a minimum force is required for descending inhibition to occur does not seem to be the case, as touch and placebo alone can trigger a descending inhibitory response. However, this may also be treatment-dependent. Being that the gate control theory of pain states that non-noxious stimuli inhibit noxious stimuli, more aggressive therapies may be too noxious to trigger a gate control response, but not noxious enough to produce a DNIC response. Thus, more research is needed to shed light on these paradoxical treatment outcomes. Future research should target therapies that have already been shown to be effective, as to prevent the wasting of resources investigating mechanisms that are not clinically meaningful, and should utilize both a control and sham group. Investigators should

185 be cautious when designing experiments that use naloxone, as it can inhibit pain via peripheral
186 mechanisms; thus, it may not be appropriate to use with those who have low back pain (Payson
187 & Holloway 1984). Lastly, it is imperative that researchers be vigilant when interpreting the
188 results of serum levels of EO, as they may not reflect levels seen in the brain or cerebral spinal
189 fluid (Wen et al. 1979).

190 **Conclusion**

191 Nearly all types of manual therapy have been shown to elicit a neurophysiological
192 response that is associated with the descending pain modulation circuit; however, it appears that
193 different types of manual therapy work through different mechanisms. For example, while
194 massage therapy appears to elicit an endogenous opioid response, spinal manipulation does not.

195 Despite the large popularity and long history of manual therapy, its mechanisms are not
196 truly understood. Understanding its mechanisms may help clinicians choose which therapy is
197 most appropriate for each patient, and may also lead to more effective therapies in the future.

199

Table 1. Findings of studies

Study	Variation	Findings
Degenhardt et al. (2007)	OMT	↑β-Endorphins ↑PEA
McPartland et al. (2005)	OMT	↑AEA
Vernon et al. (1986)	SMT	↑β-Endorphins
Christian et al. (1988)	SMT	→ β-Endorphins
Sanders et al. (1990)	SMT	→ β-Endorphins
Plaza-Manzano et al. (2014)	SMT	↑orexin A ↓ neurotensin ↓ oxytocin
Skyba et al. (2003)	Knee Manipulation	serotonin-mediated norepinephrine-mediated non-GABA-mediated
Martins et al. (2012)	Ankle Joint Mobilization	EO-mediated †
Paungmali et al. (2003)	MWM	No increase in tolerance over treatment period
Paungmali et al. (2004)	MWM	non-EO-mediated †
Santos et al. (2014)	Neural Mobilization	dynorphin-mediated
Kaada & Torsteinbo (1989)	Connective Tissue Massage	↑β-Endorphins
Trentini et al. (2005)	Acupressure	EO-mediated †
Fassoulaki et al. (2007)	Acupressure	→ β-Endorphins
Day et al. (1987)	Conventional Massage	→ β-Endorphins → β-Lipotropins
Hernandez-Reif et al. (2001)	Conventional Massage	↑dopamine ↑serotonin
Hernandez-Reif et al. (2004)	Conventional Massage	↑dopamine ↑serotonin
Field et al. (2005)*	Conventional Massage	↑dopamine ↑serotonin ↓ cortisol

* denotes review; † denotes a conclusion inferred from naloxone response

200

201

References

- Adams ML, Brase DA, Welch SP, and Dewey WL. 1986. The role of endogenous peptides in the action of opioid analgesics. *Annals of emergency medicine* 15:1030-1035.
- Bender T, Nagy G, Barna I, Tefner I, Kadas E, and Geher P. 2007. The effect of physical therapy on beta-endorphin levels. *European journal of applied physiology* 100:371-382.
- Benedetti F, Thoen W, Blanchard C, Vighetti S, and Arduino C. 2013. Pain as a reward: changing the meaning of pain from negative to positive co-activates opioid and cannabinoid systems. *Pain* 154:361-367.
- Bialosky JE, Bishop MD, Price DD, Robinson ME, and George SZ. 2009. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Manual therapy* 14:531-538.
- Chitour D, Dickenson AH, and Le Bars D. 1982. Pharmacological evidence for the involvement of serotonergic mechanisms in diffuse noxious inhibitory controls (DNIC). *Brain research* 236:329-337.
- Christian GF, Stanton GJ, Sissons D, How HY, Jamison J, Alder B, Fullerton M, and Funder JW. 1988. Immunoreactive ACTH, beta-endorphin, and cortisol levels in plasma following spinal manipulative therapy. *Spine (Phila Pa 1976)* 13:1411-1417.
- Colloca L, Klinger R, Flor H, and Bingel U. 2013. Placebo analgesia: psychological and neurobiological mechanisms. *Pain* 154:511-514.
- Crielaard JM, Bastin R, and Franchimont P. 1983. [The endorphin system and physical medicine]. *Acta Belgica Medica physica : organe officiel de la Societe royale belge de medecine physique et de rehabilitation* 6:141-145.
- Day JA, Mason RR, and Chesrown SE. 1987. Effect of massage on serum level of beta-endorphin and beta-lipotropin in healthy adults. *Physical therapy* 67:926-930.
- Degenhardt BF, Darmani NA, Johnson JC, Towns LC, Rhodes DC, Trinh C, McClanahan B, and DiMarzo V. 2007. Role of osteopathic manipulative treatment in altering pain biomarkers: a pilot study. *The Journal of the American Osteopathic Association* 107:387-400.
- DosSantos MF, Martikainen IK, Nascimento TD, Love TM, Deboer MD, Maslowski EC, Monteiro AA, Vincent MB, Zubieta JK, and DaSilva AF. 2012. Reduced basal ganglia mu-opioid receptor availability in trigeminal neuropathic pain: a pilot study. *Molecular pain* 8:74.
- Eippert F, Bingel U, Schoell ED, Yacubian J, Klinger R, Lorenz J, and Buchel C. 2009. Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron* 63:533-543.
- Fassoulaki A, Paraskeva A, Kostopanagiotou G, Tsakalozou E, and Markantonis S. 2007. Acupressure on the extra 1 acupoint: the effect on bispectral index, serum melatonin, plasma beta-endorphin, and stress. *Anesthesia and analgesia* 104:312-317.
- Field T. 2014. Massage therapy research review. *Complementary therapies in clinical practice*.
- Field T, Hernandez-Reif M, Diego M, Schanberg S, and Kuhn C. 2005. Cortisol decreases and serotonin and dopamine increase following massage therapy. *The International journal of neuroscience* 115:1397-1413.
- Fields HL, Heinricher MM, and Mason P. 1991. Neurotransmitters in nociceptive modulatory circuits. *Annual review of neuroscience* 14:219-245.

- Gerrits MA, Lesscher HB, and van Ree JM. 2003. Drug dependence and the endogenous opioid system. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 13:424-434.
- Hernandez-Reif M, Field T, Krasnegor J, and Theakston H. 2001. Lower back pain is reduced and range of motion increased after massage therapy. *The International journal of neuroscience* 106:131-145.
- Hernandez-Reif M, Ironson G, Field T, Hurley J, Katz G, Diego M, Weiss S, Fletcher MA, Schanberg S, Kuhn C, and Burman I. 2004. Breast cancer patients have improved immune and neuroendocrine functions following massage therapy. *Journal of psychosomatic research* 57:45-52.
- Kaada B, and Torsteinbo O. 1989. Increase of plasma beta-endorphins in connective tissue massage. *General pharmacology* 20:487-489.
- Kraus E, Le Bars D, and Besson JM. 1981. Behavioral confirmation of "diffuse noxious inhibitory controls" (DNIC) and evidence for a role of endogenous opiates. *Brain research* 206:495-499.
- Le Bars D, Dickenson AH, and Besson JM. 1979a. Diffuse noxious inhibitory controls (DNIC). I. Effects on dorsal horn convergent neurones in the rat. *Pain* 6:283-304.
- Le Bars D, Dickenson AH, and Besson JM. 1979b. Diffuse noxious inhibitory controls (DNIC). II. Lack of effect on non-convergent neurones, supraspinal involvement and theoretical implications. *Pain* 6:305-327.
- Lindgren L, Westling G, Brulin C, Lehtipalo S, Andersson M, and Nyberg L. 2012. Pleasant human touch is represented in pregenual anterior cingulate cortex. *Neuroimage* 59:3427-3432.
- Loh HH, Tseng LF, Wei E, and Li CH. 1976. beta-endorphin is a potent analgesic agent. *Proceedings of the National Academy of Sciences of the United States of America* 73:2895-2898.
- Martins DF, Bobinski F, Mazzardo-Martins L, Cidral-Filho FJ, Nascimento FP, Gadotti VM, and Santos AR. 2012. Ankle joint mobilization decreases hypersensitivity by activation of peripheral opioid receptors in a mouse model of postoperative pain. *Pain medicine* 13:1049-1058.
- Mason P. 1999. Central mechanisms of pain modulation. *Current opinion in neurobiology* 9:436-441.
- McLean S, Naish R, Reed L, Urry S, and Vicenzino B. 2002. A pilot study of the manual force levels required to produce manipulation induced hypoalgesia. *Clinical biomechanics* 17:304-308.
- McPartland JM, Giuffrida A, King J, Skinner E, Scotter J, and Musty RE. 2005. Cannabimimetic effects of osteopathic manipulative treatment. *The Journal of the American Osteopathic Association* 105:283-291.
- Melzack R, and Wall PD. 1965. Pain mechanisms: a new theory. *Science* 150:971-979.
- Morton DL, El-Dereby W, and Jones AK. 2014. Placebo analgesia: cognition or perception. *Handbook of experimental pharmacology* 225:71-80.
- Nadal X, La Porta C, Andreea Bura S, and Maldonado R. 2013. Involvement of the opioid and cannabinoid systems in pain control: new insights from knockout studies. *European journal of pharmacology* 716:142-157.

- Nahin R, Barnes P, Stussman B, and Bloom B. 2009. Costs of complementary and alternative medicine (CAM) and frequency of visits to CAM practitioners: United States, 2007. *National health statistics reports*:1-14.
- Ossipov MH, Dussor GO, and Porreca F. 2010. Central modulation of pain. *The Journal of clinical investigation* 120:3779-3787.
- Paungmali A, O'Leary S, Souvlis T, and Vicenzino B. 2004. Naloxone fails to antagonize initial hypoalgesic effect of a manual therapy treatment for lateral epicondylalgia. *Journal of manipulative and physiological therapeutics* 27:180-185.
- Paungmali A, Vicenzino B, and Smith M. 2003. Hypoalgesia induced by elbow manipulation in lateral epicondylalgia does not exhibit tolerance. *The journal of pain : official journal of the American Pain Society* 4:448-454.
- Payson SM, and Holloway HS. 1984. Possible complications of using naloxone as an internal opiate antagonist in the investigation of the role of endorphins in osteopathic manipulative treatment. *The Journal of the American Osteopathic Association* 84:152-156.
- Pettman E. 2007. A history of manipulative therapy. *The Journal of manual & manipulative therapy* 15:165-174.
- Pickar JG. 2002. Neurophysiological effects of spinal manipulation. *Spine J* 2:357-371.
- Plaza-Manzano G, Molina-Ortega F, Lomas-Vega R, Martinez-Amat A, Achalandabaso A, and Hita-Contreras F. 2014. Changes in biochemical markers of pain perception and stress response after spinal manipulation. *The Journal of orthopaedic and sports physical therapy* 44:231-239.
- Sanders GE, Reinert O, Tepe R, and Maloney P. 1990. Chiropractic adjustive manipulation on subjects with acute low back pain: visual analog pain scores and plasma beta-endorphin levels. *Journal of manipulative and physiological therapeutics* 13:391-395.
- Santos FM, Grecco LH, Pereira MG, Oliveira ME, Rocha PA, Silva JT, Martins DO, Miyabara EH, and Chacur M. 2014. The neural mobilization technique modulates the expression of endogenous opioids in the periaqueductal gray and improves muscle strength and mobility in rats with neuropathic pain. *Behavioral and brain functions : BBF* 10:19.
- Sauro MD, and Greenberg RP. 2005. Endogenous opiates and the placebo effect: a meta-analytic review. *Journal of psychosomatic research* 58:115-120.
- Skyba DA, Radhakrishnan R, Rohlwing JJ, Wright A, and Sluka KA. 2003. Joint manipulation reduces hyperalgesia by activation of monoamine receptors but not opioid or GABA receptors in the spinal cord. *Pain* 106:159-168.
- Threlkeld AJ. 1992. The effects of manual therapy on connective tissue. *Physical therapy* 72:893-902.
- Trentini JF, 3rd, Thompson B, and Erlichman JS. 2005. The antinociceptive effect of acupressure in rats. *Am J Chin Med* 33:143-150.
- Vernon H. 2000. Qualitative review of studies of manipulation-induced hypoalgesia. *Journal of manipulative and physiological therapeutics* 23:134-138.
- Vernon HT, Dhami MS, Howley TP, and Annett R. 1986. Spinal manipulation and beta-endorphin: a controlled study of the effect of a spinal manipulation on plasma beta-endorphin levels in normal males. *Journal of manipulative and physiological therapeutics* 9:115-123.

- 334 Wager TD, Scott DJ, and Zubieta JK. 2007. Placebo effects on human mu-opioid activity during
 335 pain. *Proceedings of the National Academy of Sciences of the United States of America*
 336 104:11056-11061.
- 337 Wen H, Ho W, Ma L, Choa G, and Ling N. 1979. The influence of electro-acupuncture on
 338 naloxone-induced morphine withdrawal: II. Elevation of immunoassayable beta-
 339 endorphin activity in the brain but not the blood. *The American journal of Chinese*
 340 *medicine* 7:237-240.
- 341 Zubieta JK, Bueller JA, Jackson LR, Scott DJ, Xu Y, Koeppe RA, Nichols TE, and Stohler CS.
 342 2005. Placebo effects mediated by endogenous opioid activity on mu-opioid receptors.
 343 *The Journal of neuroscience : the official journal of the Society for Neuroscience*
 344 25:7754-7762.
 345