

**A peer-reviewed version of this preprint was published in PeerJ on 25 August 2015.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1219) (peerj.com/articles/1219), which is the preferred citable publication unless you specifically need to cite this preprint.

Hassall C. 2015. Strong geographical variation in wing aspect ratio of a damselfly, *Calopteryx maculata* (Odonata: Zygoptera) PeerJ 3:e1219  
<https://doi.org/10.7717/peerj.1219>

# Strong longitudinal variation in wing aspect ratio of a damselfly, *Calopteryx maculata* (Odonata: Zygoptera)

Christopher Hassall

Geographical patterns in body size have been described across a wide range of species, leading to the development of a series of fundamental biological rules. However, shape variables are less well-described despite having substantial consequences for organismal performance. Wing aspect ratio (AR) has been proposed as a key shape parameter that determines function in flying animals, with high AR corresponding to longer, thinner wings that promote high manoeuvrability, low speed flight, and low AR corresponding to shorter, broader wings that promote high efficiency long distance flight. From this principle it might be predicted that populations at range edges would exhibit low AR wings. I test this hypothesis using the riverine damselfly, *Calopteryx maculata*, sampled from 34 sites across its range margin in North America. Nine hundred and seven male specimens were captured from across the 34 sites (mean=26.7  $\pm$  2.9 SE per site), dissected and measured to quantify the area and length of all four wings. Geometric morphometrics were employed to investigate geographical variation in wing shape. The majority of variation in wing shape involved changes in wing aspect ratio, confirmed independently by geometric morphometrics and wing measurements. There was a weak positive relationship between wing aspect ratio and temperature, in line with work on other insects. However, there was a strong longitudinal pattern in which western populations exhibited lower wing aspect ratio. This longitudinal pattern may be related to increasing variability in precipitation from east to west in North America. I discuss my findings in light of research of the functional ecology of wing shape across vertebrate and invertebrate taxa

2       **Strong longitudinal variation in wing aspect ratio of a damselfly, *Calopteryx***  
3                               ***maculata* (Odonata: Zygoptera)**

4  
5    Author: Christopher Hassall

6    Address: School of Biology, University of Leeds, Leeds, LS2 3JT, United Kingdom

7    Telephone: +44 0113 3435578

8    Fax: + 44 0113 3433091

9    Running title: Damselfly wing morphology

10 **ABSTRACT**

11 Geographical patterns in body size have been described across a wide range of species, leading to the  
12 development of a series of fundamental biological rules. However, shape variables are less well-described  
13 despite having substantial consequences for organism performance. Wing aspect ratio (AR) has been proposed  
14 as a key shape parameter that determines function in flying animals, with high AR corresponding to longer,  
15 thinner wings that promote high manoeuvrability, low speed flight, and low AR corresponding to shorter,  
16 broader wings that promote high efficiency long distance flight. From this principle it might be predicted that  
17 populations living in cooler areas would exhibit low AR wings to compensate for reduced muscle efficiency at  
18 lower temperatures. I test this hypothesis using the riverine damselfly, *Calopteryx maculata*, sampled from 34  
19 sites across its range margin in North America. Nine hundred and seven male specimens were captured from  
20 across the 34 sites (mean=26.7  $\pm$ 2.9 SE per site), dissected and measured to quantify the area and length of all  
21 four wings. Geometric morphometrics were employed to investigate geographical variation in wing shape. The  
22 majority of variation in wing shape involved changes in wing aspect ratio, confirmed independently by  
23 geometric morphometrics and wing measurements. There was a weak positive relationship between wing aspect  
24 ratio and temperature, in line with work on other insects. However, there was a much stronger longitudinal  
25 pattern in which western populations exhibited lower wing aspect ratio. This longitudinal pattern may be related  
26 to increasing variability in precipitation from east to west in North America. I discuss my findings in light of  
27 research of the functional ecology of wing shape across vertebrate and invertebrate taxa.

28  
29 **Keywords:** wing morphology, aspect ratio, flight, dispersal, habitat stability, precipitation, damselfly, Odonata.

## 30 INTRODUCTION

31 Powered flight has evolved independently in four different lineages: the pterosaurs, insects, birds, and bats,  
32 allowing animals to exploit novel niches and avoid predators. The adaptations that allowed each of these  
33 transitions to an aerial niche represent a suite of similar traits that can be broken down into a number of  
34 functional morphological components that influence inter- and intraspecific variation in flight performance.  
35 First, absolute body size is correlated with dispersal ability across a wide range of taxa (Jenkins et al. 2007).  
36 Second, the ratio of body mass to wing area – known as “wing loading” – has a strong influence on the amount  
37 of thrust generated per wingbeat (Dudley 2002). However, for the purposes of this study I am most interested in  
38 the third component of variation: that of wing shape. One of the principle measures of functional variation in  
39 wing shape is the length of the wing relative to the width, known as aspect ratio. In vertebrates, higher aspect  
40 ratio (longer, thinner wings) is predicted to give faster and more efficient flight (Norberg 1989) and has been  
41 shown to be associated with migratory species in birds (Mönkkönen 1995). However, there has been speculation  
42 that the benefits of high aspect ratio may be reduced or even reversed at the low Reynolds numbers experienced  
43 by insects (Ennos 1989; Wootton 1992). This fact, along with the difference in the nature of flight – number,  
44 structure and locomotory independence of wings – between birds and insects complicates the formation of  
45 hypotheses concerning the implications of variation in odonate morphology (Johansson et al. 2009). The  
46 literature on the functional relevance of insect wing morphology is heavily biased towards theory (Dudley  
47 2002), laboratory studies (Marden 1995) and observations of kinematics (Rüppell 1989; Wakeling & Ellington  
48 1997a, b, c) rather than field observations.

49  
50 Contrary to predictions for birds, a number of findings point towards lower wing aspect ratio as being beneficial  
51 for dispersal in insects. Wing aspect ratio is lower in populations of *Pararge aegeria* that have recently been  
52 founded (Hill et al. 1999). Populations of *P. aegeria* (Hughes et al. 2007; Vandewoestijne & Van Dyck 2011),  
53 *Drosophila melanogaster* (Azevedo et al. 1998), and a number of damselflies (Hassall et al. 2009; Taylor &  
54 Merriam 1995) show lower aspect ratio at higher latitudes where temperature reduces the efficiency of flight in  
55 ectotherms. Other studies have shown higher wing aspect ratio only in species of damselflies with expanding

range margins (Hassall et al. 2009), and those marginal populations exhibit wing shapes that deviate progressively away from the species average closer to the range margin (Hassall et al. 2008). Studies using common garden rearing of *Drosophila* from a range of latitudes have shown that individuals reared at lower temperatures have lower aspect ratio (Azevedo et al. 1998). While there is no clear relationship between aspect ratio and flight speed in butterflies (Berwaerts et al. 2008; but cf Berwaerts et al. 2002; Dudley 1990), species in which males "patrol" (i.e. exhibit prolonged flight) tend to have lower aspect ratios (Wickman 1992). Chironomid females have broader wings (characteristic of lower aspect ratio) to assist with flying for long periods between habitat patches (McLachlan 1986). While there are exceptions (increased fragmentation does not correlate with aspect ratio in *Plebejus argus* (Thomas et al. 1998) or *Pararge aegeria* (Merckx & Van Dyck 2006)) these findings seem to suggest that lower wing aspect ratio in insects is associated with greater dispersal.

Wing morphology in Odonata may be affected by a combination of sexual selection during intrasexual, agonistic interactions, intersexual courtship displays and dispersal (Johansson et al. 2009). In the field, intrasexual territorial contests in *C. maculata* are determined by fat reserves (Marden & Rollins 1994; Marden & Waage 1990) and contests in *Plathemis lydia* are determined by flight muscle ratio (Marden 1989). In both cases, aspect ratio was shown not to influence the outcome of the contests. Sexual selection on courtship displays focuses on patterns of pigmentation in *Calopteryx* species (Siva-Jothy 1999; Waage 1973). However, wing shape has been shown to vary with landscape structure in *C. maculata* (Taylor & Merriam 1995) and between some closely-related species of Calopterygidae in Europe (Sadeghi et al. 2009), although not all species exhibited distinct wing shapes. Based on these results, it seems that wing shape variation is under natural selection due to dispersal (within or between sites), rather than sexual selection.

Based on the reasoning presented above, I evaluate the hypothesis that, based on negative correlations between latitude and aspect ratio, a positive relationship should be found between temperature and aspect ratio to compensate for lower flight efficiency at lower temperatures. Uncertainties over the ecological role of morphology variation may stem from the partial sampling of geographical ranges (Hassall 2013). Limited

82 sampling of non-linear trends that occur over large spatial scales may produce misleading results and so I  
83 provide an analysis of wing shape variation across the entire range of the damselfly *Calopteryx maculata* in  
84 North America.

## 85 METHODS

86 A total of 907 specimens of male *C. maculata* were collected from 34 sites across the range by 25 collectors  
87 (Figure 1, Table 1). Collections took place between 13 May and 7 August 2010 and mean sample size from each  
88 site varied between 4 and 84 individuals (mean=26.7  $\pm$ 2.9 SE, details of sample sizes and mean measurements  
89 can be found in Table 1). Wings were dissected from the body as close to the thorax as possible and mounted on  
90 adhesive tape (Scotch Matte Finish Magic Tape). Wings were scanned using the slide scanner on an Epson V500  
91 PHOTO flatbed scanner with fixed exposure at 1200dpi. Wing length (the length from the costal end of the vein  
92 separating the arculus from the discoidal cell to the tip of the wing) and wing area were calculated for each of  
93 the four wings on each individual. All measurements were carried out in ImageJ (Rasband 1997-2007). During  
94 measurement, any damage to wings was noted and those measurements (length or area) which could not be  
95 accurately quantified were excluded. This resulted in the exclusion of 7 fore wing and 9 hind wing lengths, and  
96 28 fore wing and 45 hind wing areas. Aspect ratio was then calculated separately for both fore and hind wings as  
97 wingspan<sup>2</sup>/wing area (see Table 1 for summary statistics and sample sizes).

98  
99 It has been suggested that wing aspect ratio does not provide sufficient detail to be morphologically informative  
100 in butterflies (Betts & Wootton 1988) or dragonflies (Johansson et al. 2009). Therefore, in addition to  
101 calculating aspect ratio, I also use geometric morphometrics to derive descriptors of the shape of the wing. A  
102 subset of up to 10 individuals from each site were selected at random and a set of 14 landmarks were digitised  
103 on 1 fore wing and 1 hind wing (Figure 2) using tpsDig2 (v.2.12, Rohlf 2008). Mean locations for each of the 14  
104 landmarks were found for each of the 34 sites. Principal components analysis (PCA) was carried out on these  
105 landmarks after Procrustes transformation using the PAST software package (Hammer et al. 2001).  
106 Relationships between the principal components and absolute measurements were investigated using Pearson  
107 correlations. Fore and hind wings were compared using paired Hotelling's  $t^2$  tests in PAST to assess whether the  
108 two datasets could be combined.

109



110 Mean annual temperature was extracted for each site from the WORLDCLIM dataset (Haylock et al. 2008) to  
111 test the central hypothesis of the study. After initial data examination revealed a longitudinal pattern, the  
112 coefficient of variation ( $\sigma/\mu$ ) of precipitation, which exhibits strong variability from west to east, was calculated  
113 for each site using the WORLDCLIM dataset (bioclimatic variable 15, Haylock et al. 2008) in ArcGIS v9.2  
114 (ESRI 2006). Aspect ratio and the informative principal components from the shape analysis were regressed  
115 against temperature and seasonality of precipitation using linear regressions weighted by the square-root of the  
116 sample size. In each case, the models were tested with a quadratic predictor term using Akaike's information  
117 criterion (AIC) to evaluate any improvement in model fit.

118 **RESULTS**

119 Fore and hind wings vary significantly in shape ( $t^2=122500$ ,  $p<<0.001$ ) and were completely separated along the  
 120 PC1 axis which explained 80.2% of the variance in shape. As a result, fore and hind wing data are treated  
 121 separately for the rest of the analysis.

122  
 123 PCA of both fore and hind wings resulted in three components that lay before the "elbow" of the scree plot.  
 124 These explained 38.7%, 23.2% and 18.6% (total 80.5%) of the variance in fore wing shape and 44.9%, 21.4%,  
 125 and 12.6% (total 78.9%) of the variance in hind wing shape. PC1 in both cases involved a variation in the width  
 126 of the wing relative to its length, such that an increase in PC1 leads to a decrease in the width of the wing  
 127 relative to the length (Figure 3). PC1 was significantly positively correlated with aspect ratio (fore wings,  
 128  $r=0.875$ ,  $p<0.001$ ; hind wings,  $r=0.854$ ,  $p<0.001$ , Figure 4). The PC2 and PC3 involved more subtle shape  
 129 changes which were still consistent between wings. PC2 appears to involve a shortening of the pre-nodal region  
 130 and a blunting of the tip, while PC3 corresponds to a movement of wing area towards the wing tip.

131  
 132 Geographical patterns of wing aspect ratio showed strong longitudinal variation (Figure 5A), with significant  
 133 and positive relationships between longitude and aspect ratio in both fore ( $r=0.816$ ,  $p<0.001$ ) and hind wings  
 134 ( $r=0.800$ ,  $p<0.001$ ) indicating a decline in aspect ratio further west. Latitudinal patterns were significant and  
 135 positive but weaker than for longitude (fore:  $r=0.441$ ,  $p=0.010$ ; hind:  $r=0.375$ ,  $p=0.029$ ). For regression models  
 136 containing either longitude and latitude or both, models with longitude only performed best and the addition of  
 137 latitude did not produce a significant improvement in model performance ( $\Delta AIC<2$ ).

138  
 139 Regressions between aspect ratio and temperature showed no significant increase in explanatory power for the  
 140 quadratic model in the fore (linear AIC=-50.188; quadratic AIC=-50.947;  $\Delta AIC=0.76$ ) or hind wings (linear  
 141 AIC=-53.582; quadratic AIC=-53.509;  $\Delta AIC=0.073$ ) and so the quadratic term was not added. Aspect ratio was  
 142 positive and significantly related to temperature for fore wings ( $F_{1,32}=6.262$ ,  $p=0.018$ ,  $R^2=0.138$ ) but there was  
 143 no significant relationship with aspect ratio of hind wings ( $F_{1,32}=3.302$ ,  $p=0.079$ ,  $R^2=0.065$ ). Regressions

144 between aspect ratio and the seasonality of precipitation also showed no increase in explanatory power for the  
145 quadratic model in fore (linear AIC=-63.124; quadratic AIC=-64.723;  $\Delta$ AIC=1.599) or hind wings (linear AIC=-  
146 67.687; hind: AIC=-68.840;  $\Delta$ AIC=1.154) and so the quadratic term was not added. Linear regressions between  
147 aspect ratio and the seasonality of precipitation were highly significant in both fore ( $F_{1,32}=25.687$ ,  $p<0.001$ ,  
148  $R^2=0.428$ ) and hind wings ( $F_{1,32}=23.792$ ,  $p<0.001$ ,  $R^2=0.409$ , Figure 5B) and explained a high proportion of the  
149 variation in the data. Taken together, models explaining aspect ratio with seasonality of precipitation had a  
150 substantially greater explanatory power than those using temperature in both fore ( $\Delta$ AIC=13.776) and hind  
151 wings ( $\Delta$ AIC=15.331). However, the geographical distribution of aspect ratio values (Figure 1) suggests that  
152 there may be a step-change in wing shape at a certain longitude, rather than a gradual trend.

## DISCUSSION

I provide the first comprehensive assessment of intraspecific variation in wing morphology across a range in a damselfly. The use of geometric morphometrics to analyse shape confirms that changes in aspect ratio (i.e. changes in the length of the wing relative to the width) constitute the major source of variation between specimens from different sites. I demonstrate a weakly significant effect of temperature in fore wing shape, with lower wing aspect ratios at cooler temperatures as is predicted by theory. However, the dominant pattern is one of increasing aspect ratio from west to east, which has not been documented in previous studies and may be related to the greater seasonality of precipitation in western populations. This finding is consistent with the habitat-stability-dispersal hypothesis if increasing seasonality of precipitation disrupts flow regimes in *Calopteryx* breeding sites. It is worth noting that the nature of this study is such that I cannot disentangle the effects of selection from those of phenotypic plasticity. Indeed, previous studies have demonstrated that while some flight morphological parameters are under genetic control, wing aspect ratio shows a plastic response to the environment in *Drosophila* (Azevedo et al. 1998).

The literature on the functional relevance of aspect ratio has produced conflicting findings. While the pattern observed in the present study is consistent with a range of studies that suggest lower aspect ratio favours more efficient or longer flight, higher aspect ratios have been associated with fragmented habitat (Taylor & Merriam 1995) and range expansion (Hassall et al. 2009) in calopterygid damselflies in other studies. The sites involved in Taylor and Merriam's study were all north of Ottawa, Canada, where my results indicate aspect ratios are high. It may be that the increase in distance between foraging and reproductive habitat that was associated with morphological change in their study operates on a different scale to the patterns describe in the present study. I suggest that the findings of previous studies are limited by their consideration of a relatively small area and number of populations.

It is generally considered that higher aspect ratios provide a benefit for longer-distance (Mönkkönen 1995) or efficient, gliding flight (Ennos 1989), although this is equivocal in Lepidoptera (Betts & Wootton 1988).

179 However, as I argue in the introduction, the opposite appears to be true for the majority of studies of insects  
180 where smaller Reynolds numbers operate. Indeed, this study provides support for a positive association between  
181 aspect ratio and temperature, which may be selected for because lower aspect ratio is more efficient at lower  
182 temperatures (Vandewoestijne & Van Dyck 2011). There remains a gap in the literature that needs to be filled  
183 with flight laboratory experiments of the functional implications of aspect ratio variation in odonates and other  
184 insects as have been carried out in some butterflies (Berwaerts et al. 2008; Berwaerts et al. 2002; Davis et al.  
185 2012). In particular, a test of the hypothesis that lower aspect ratios enhance flight efficiency at lower  
186 temperatures is warranted given the increasing evidence for the correlation between aspect ratio and  
187 temperature.

188  
189 The findings here are consistent with selection for dispersal in habitats with a greater variability in precipitation  
190 which causes greater variability in streamflow. The range of potential oviposition sites is limited by fluctuations  
191 in rainfall which may raise or lower flow-rates, moving them outside of the narrow range of values that *C.*  
192 *maculata* prefer (Gibbons & Pain 1992), such that a habitat that is suitable in one year may not remain so in the  
193 next. It is also conceivable that seasonality of precipitation increases the fragmentation of these suitable sites:  
194 sites which are only marginally suitable become unsuitable during extreme precipitation events, while a subset  
195 of sites which are resilient to flow rate variation remain suitable. Such resilience may be provided by  
196 geomorphological characteristics of the river channel (Goldstein et al. 2007) or the presence of microhabitats  
197 which are less susceptible to variations in flow rate (Brooks et al. 2005). This elimination of marginal larval  
198 habitats effectively fragments the landscape resulting in spatial variability.

199  
200 Habitat selection in Odonata is a hierarchical process, with selection steps occurring at the level of the  
201 landscape, the habitat, and the oviposition site (Buchwald 1995; Wildermuth 1994). At least the first two of  
202 these steps involves visual cues which include the presence of conspecifics (Corbet 1999), linear polarised light  
203 reflected from water bodies (Wildermuth 1998) and structural features, including vegetation (Wildermuth 1992).  
204 Calopterygid damselflies also exhibit very specific habitat requirements relating to the depth and flow rate of the

205 rivers which form the larval habitat (Abbott 2005). Male *Calopteryx splendens* and *Calopteryx xanthostoma* land  
206 on the water to demonstrate the flow-rate of the stream in their territory, and successful courtship is associated  
207 with increased flow-rate up to  $0.15\text{m.s}^{-1}$  (Gibbons & Pain 1992). The benefits of this flow-rate to the resulting  
208 offspring are higher embryonic development rates and lower accumulation of encrusting algae (Siva-Jothy et al.  
209 1995). Changes in flow rates may result in either (i) variation in larval mortality rates, or (ii) variation in the  
210 tendency for females to oviposit. It is well-established that variability in precipitation produces  
211 disproportionately large variations in streamflow (Dettinger & Diaz 2000). Given the apparent sensitivity of  
212 Calopterygidae to variations in flow rates (Gibbons & Pain 1992; Siva-Jothy et al. 1995), an assessment of  
213 regional variations in population sizes with respect to extremes of precipitation (and, hence, streamflow) might  
214 be warranted. Extremes of temperature and precipitation resulting from climate change are likely to manifest  
215 long before the gradual increase in long-term mean climate (Karl & Trenberth 2003; Palmer & Raisanen 2002)  
216 and may pose a particular threat to this family.

217  
218 Previous studies have questioned the use of aspect ratio as a single numerical metric describing wing shape in  
219 insects, due to its inability to represent the complexity of wing morphology (Betts & Wootton 1988; Johansson  
220 et al. 2009). However, I find that a complex method of shape analysis using geometric morphometrics yields  
221 patterns that strongly resemble variation in the simpler concept of aspect ratio. However, it is clear from the  
222 explanatory power of those principal components that correlate with aspect ratio (38.7% and 44.9%) that there is  
223 a great deal of variability in addition to this dimension. It is worth noting that insects exhibit a great deal of  
224 variation in aspect ratio. Odonates have high aspect ratios compared to some other insects, for example  
225 *Drosophila virilis* with an aspect ratio of 2 (Vogel 1957), and *Bombus terrestris* with an aspect ratio of 6.4.  
226 However, butterflies show higher aspect ratios of 9.8-10.5 in *Pararge aegeria* (Berwaerts et al. 2008; Berwaerts  
227 et al. 2002). The data presented here show aspect ratios of hind wings between 5.61 and 7.79 and of forewings  
228 between 5.70 and 7.56. *Aeshna cyanea*, a large odonate, exhibits aspect ratio of 8.4 and 11.6 for hind and fore  
229 wings, respectively (Ellington 1984). What makes the odonate wing very different is the extent of the venation

230 in odonate wings compared to other taxa. This venation may be associated with the pleating of the wing, which  
231 enhances aerodynamic performance relative to a smooth with of the same shape (Vargas et al. 2008).

232

233 The results presented here demonstrate clear geographical variation in flight morphology in a damselfly across  
234 its entire range. While the other studies investigating geographical variation in odonate morphology have  
235 focused on north-south transects (Johansson 2003), there are clearly important patterns occurring along the east-  
236 west axis of the range highlighting the need to consider range-wide surveys to understand macroecological and  
237 macroevolutionary patterns (Hassall 2013, 2014). Furthermore, although habitat fragmentation is associated with  
238 higher aspect ratios in *C. maculata* at local scales, lower aspect ratios may be associated with lower habitat  
239 stability at continental scales. From the survey of studies that have included aspect ratio, it is clear that  
240 laboratory studies are needed to clarify the relationship between form and function in odonate wing shape.

241 **ACKNOWLEDGEMENTS**

242 I am extremely grateful to Arne Iserbyt, Mary Burnham, Chris Lewis, Shari Sokay, Darrin OBrien, Fred Sibley,  
243 Giff Beaton, George Harp, George Sims, Harris Luckham, John Abbott, Joseph Carson, Jeni Eggers and Elliott  
244 Porter, Jeffrey Willers, Michael Blust, Marion Dobbs, Mark Musselman, Pat Heithaus, Rick Abad, Ryan  
245 Spafford, Steve Hummel, Sarah Richer, Timothy Sesterhenn, William Lamp and Wade Worthen for giving so  
246 graciously of their time to assist with collections. Carley Centen provided valuable assistance in the field and  
247 Tom Langen provided assistance with logistics. The study was funded by a British Ecological Society Small  
248 Ecological Project Grant and I was supported by a Government of Canada Postdoctoral Fellowship and an  
249 Ontario MRI Fellowship.



250 **REFERENCES**

- 251 Abbott JC. 2005. *Dragonflies and Damselflies of Texas and the South Central United States*. Princeton:  
252 Princeton University Press.
- 253 Azevedo RBR, James AC, McCabe J, and Partridge L. 1998. Latitudinal variation of wing:thorax size ratio and  
254 wing-aspect ratio in *Drosophila melanogaster*. *Evolution* 52:1353-1362.
- 255 Berwaerts K, Matthysen E, and Van Dyck H. 2008. Take-off flight performance in the butterfly *Pararge aegeria*  
256 relative to sex and morphology: a quantitative genetic assessment. *Evolution* 62:2525-2533.
- 257 Berwaerts K, Van Dyck H, and Aerts P. 2002. Does flight morphology relate to flight performance? An  
258 experimental test with the butterfly *Pararge aegeria*. *Functional Ecology* 16:484-491.
- 259 Betts CR, and Wootton RJ. 1988. Wing shape and flight behaviour in butterflies (Lepidoptera: Papilionoidea and  
260 Hesperioidea): a preliminary analysis. *Journal of Experimental Biology* 138:271-288.
- 261 Brooks AJ, Haeusler TIM, Reinfelds I, and Williams S. 2005. Hydraulic microhabitats and the distribution of  
262 macroinvertebrate assemblages in riffles. *Freshwater Biology* 50:331-344.
- 263 Buchwald R. 1995. Structure and floristic composition of vegetation: what is the significance for the occurrence  
264 of dragonfly species? 13th International Symposium on Odonatology. Essen.
- 265 Corbet PS. 1999. *Dragonflies: Behaviour and Ecology of Odonata*. Colchester: Harley Books.
- 266 Davis AK, Chi J, Bradley C, and Altizer S. 2012. The redder the better: wing color predicts flight performance  
267 in monarch butterflies. *PLoS ONE* 7:e41323.
- 268 Dettinger MD, and Diaz HF. 2000. Global characteristics of stream flow seasonality and variability. *Journal of*  
269 *Hydrometeorology* 1:289-310.
- 270 Dudley R. 1990. Biomechanics of insect flight in neotropical butterflies: morphometrics and kinematics. *Journal*  
271 *of Experimental Biology* 150:37-53.
- 272 Dudley R. 2002. *The Biomechanics of Insect Flight*. Princeton: Princeton University Press.
- 273 Ellington CP. 1984. The aerodynamics of hovering insect flight: II. Morphological parameter. *Philosophical*  
274 *Transactions of the Royal Society: Series B (Biological Sciences)* 305:17-40.

275 Ennos AR. 1989. The effect of size on the optimal shapes of gliding insects and seeds. *Journal of Zoology*  
 276 219:61-69.

277 ESRI. 2006. *ArcGIS v.9.2*. Redlands: Environmental Systems Research Institute, Inc.

278 Gibbons DW, and Pain D. 1992. The influence of river flow rate on the breeding behaviour of *Calopteryx*  
 279 damselflies. *Journal of Animal Ecology* 61:283-289.

280 Goldstein RM, Meador MR, and Ruhl KE. 2007. Relative influence of streamflows in assessing temporal  
 281 variability in stream habitat. *Journal of the American Water Resources Association* 43:642-650.

282 Hammer Ø, Harper DAT, and Ryan PD. 2001. PAST: Paleontological Statistics Software Package for Education  
 283 and Data Analysis. *Paleontologica Electronica* 4:9.

284 Hassall C. 2013. Time stress and temperature explain continental variation in damselfly body size. *Ecography*  
 285 36:894-903.

286 Hassall C. 2014. Continental variation in wing pigmentation in *Calopteryx* damselflies is related to the presence  
 287 of heterospecifics. *PeerJ* 2:e438.

288 Hassall C, Thompson DJ, and Harvey IF. 2008. Wings of *Coenagrion puella* vary in shape at the northern range  
 289 margin (Odonata: Coenagrionidae). *International Journal of Odonatology* 11:35-41.

290 Hassall C, Thompson DJ, and Harvey IF. 2009. Variation in morphology between core and marginal populations  
 291 of three British damselflies. *Aquatic Insects* 31:187-197.

292 Haylock MR, Hofstra N, Klein Tank AMG, Klok EJ, Jones PD, and New M. 2008. A European daily high-  
 293 resolution gridded dataset of surface temperature and precipitation. *Journal of Geophysical Research Series D,*  
 294 *Atmospheres* 113.

295 Hill JK, Thomas CD, and Blakeley DS. 1999. Evolution of flight morphology in a butterfly that has recently  
 296 expanded its geographic range. *Oecologia* 121:165-170.

297 Hughes CL, Dytham C, and Hill JK. 2007. Modelling and analysing evolution of dispersal in populations at  
 298 expanding range boundaries. *Ecological Entomology* 32:437-445.

299 Jenkins DG, Brescacin CR, Duxbury CV, Elliott JA, Evans JA, Grablow KR, Hillegass M, Lyon BN, Metzger  
 300 GA, Olandese ML, Pepe D, Silvers GA, Suresch HN, Thompson TN, Trexler CM, Williams GE, Williams NC,  
 301 and Williams SE. 2007. Does size matter for dispersal distance? *Global Ecology and Biogeography* 16:415.  
 302 Johansson F. 2003. Latitudinal shifts in body size of *Enallagma cyathigerum* (Odonata). *Journal of*  
 303 *Biogeography* 30:29-34.  
 304 Johansson F, Söderquist M, and Bokma F. 2009. Insect wing shape evolution: independent effects of migratory  
 305 and mate guarding flight on dragonfly wings. *Biological Journal of the Linnean Society* 97:362-372.  
 306 Karl TR, and Trenberth KE. 2003. Modern global climate change. *Science* 302:1719-1723.  
 307 Marden JH. 1989. Bodybuilding dragonflies: costs and benefits of maximizing flight muscle. *Physiological*  
 308 *Zoology* 62:505-521.  
 309 Marden JH. 1995. Large-scale changes in thermal sensitivity of flight performance during adult maturation in a  
 310 dragonfly. *Journal of Experimental Biology* 198:2095-2102.  
 311 Marden JH, and Rollins RA. 1994. Assessment of energy reserves by damselflies engaged in aerial contests for  
 312 mating territories. *Animal Behaviour* 48:1023-1030.  
 313 Marden JH, and Waage JK. 1990. Escalated damselfly contests are energetic wars of attrition. *Animal Behaviour*  
 314 39:954-959.  
 315 McLachlan AJ. 1986. Sexual dimorphism in midges: strategies for flight in the rain-pool dweller *Chironomus*  
 316 *imicola* (Diptera: Chironomidae). *Journal of Animal Ecology* 55:261-267.  
 317 Merckx T, and Van Dyck H. 2006. Landscape structure and phenotypic plasticity in flight morphology in the  
 318 butterfly *Pararge aegeria*. *Oikos* 113:226-232.  
 319 Mönkkönen M. 1995. Do migrant birds have more pointed wings? A comparative study. *Evolutionary Ecology*  
 320 9:520-528.  
 321 Norberg UM. 1989. *Vertebrate Flight, Mechanics, Physiology, Morphology, Ecology and Evolution*. Berlin:  
 322 Springer-Verlag.  
 323 Palmer TN, and Raisanen J. 2002. Quantifying the risk of extreme seasonal precipitation events in a changing  
 324 climate. *Nature* 415:512-514.

325 Rasband WS. (1997-2007) *ImageJ*. <<http://rsb.info.nih.gov/ij/>>.

326 Rohlf FJ. 2008. *tpsDig, digitize landmarks and outlines, version 2.12*. State University of New York at Stony

327 Brook: Department of Ecology and Evolution.

328 Rüppell G. 1989. Kinematic analysis of symmetrical flight manoeuvres of Odonata. *Journal of Experimental*

329 *Biology* 144:13-42.

330 Sadeghi S, Adriaens D, and Dumont HJ. 2009. Geometric morphometric analysis of wing shape variation in ten

331 European populations of *Calopteryx splendens* (Harris, 1782) (Zygoptera: Odonata). *Odonatologica* 38:343-360.

332 Siva-Jothy MT. 1999. Male wing pigmentation may affect reproductive success via female choice in a

333 calopterygid damselfly (Zygoptera). *Behaviour* 136:1365-1377.

334 Siva-Jothy MT, Gibbons DW, and Pain D. 1995. Female oviposition site preference and egg hatching success in

335 the damselfly *Calopteryx splendens xanthostoma*. *Behavioral Ecology and Sociobiology* 37:39-44.

336 Taylor PD, and Merriam G. 1995. Wing morphology of a forest damselfly is related to landscape structure.

337 *Oikos* 73:43-48.

338 Thomas CD, Hill JK, and Lewis OT. 1998. Evolutionary consequences of habitat fragmentation in a localized

339 butterfly. *Journal of Animal Ecology* 67:485-497.

340 Vandewoestijne S, and Van Dyck H. 2011. Flight morphology along a latitudinal gradient in a butterfly: do

341 geographic clines differ between agricultural and woodland landscapes? *Ecography* 34:876-886.

342 Vargas A, Mittal R, and Dong H. 2008. A computational study of the aerodynamic performance of a dragonfly

343 wing section in gliding flight. *Bioinspiration & Biomimetics* 3:1-13.

344 Vogel S. 1957. Flight in *Drosophila*: III. Aerodynamic characteristics of fly wing and wing models. *Journal of*

345 *Experimental Biology* 46:431-443.

346 Waage JK. 1973. Reproductive behavior and its relation to territoriality in *Calopteryx maculata* (Beauvois)

347 (Odonata: Calopterygidae). *Behaviour* 47:240-256.

348 Wakeling JM, and Ellington CP. 1997a. Dragonfly flight. II. Velocities, accelerations and kinematics of flapping

349 flight. *Journal of Experimental Biology* 200:557-582.

- 350 Wakeling JM, and Ellington CP. 1997b. Dragonfly flight. III. Lift and power requirements. *Journal of*  
351 *Experimental Biology* 200:583-600.
- 352 Wakeling JM, and Ellington CP. 1997c. Dragonfly flight: I Gliding flight and steady-state aerodynamic forces.  
353 *Journal of Experimental Biology* 200:543-556.
- 354 Wickman PO. 1992. Sexual selection and butterfly design - a comparative study. *Evolution* 46:1525-1536.
- 355 Wildermuth H. 1992. Visual and tactile stimuli in choice of oviposition substrates by the dragonfly *Perithemis*  
356 *mooma* Kirby (Anisoptera: Libellulidae). *Odonatologica* 21:309-321.
- 357 Wildermuth H. 1994. Habitatselektion bei Libellen. *Advances in Odonatology* 6:223-257.
- 358 Wildermuth H. 1998. Dragonflies recognise the water of rendezvous and oviposition sites by horizontally  
359 polarised light: a behavioural field test. *Naturwissenschaften* 85:297-302.
- 360 Wootton RJ. 1992. Functional morphology of insect wings. *Annual Review of Entomology* 37:113-140.

## 363 TABLES

364 Table 1 - Sampling site locations, sample sizes and aspect ratios of wings of male *Calopteryx maculata*.

365 “Measurements” gives the sample size for the total number of measured specimens, “Geo Morph” gives the

366 sample sizes used in the geometric morphometric analysis ( $N_{\text{fore}}$  = sample size for fore wings,  $N_{\text{hind}}$  = sample size

367 for hind wings).

							Measurements			Geo Morph	
Region	Site	Latitude	Longitude	Date	Fore wing	Hind wing	N <sub>total</sub>	N <sub>fore</sub>	N <sub>hind</sub>	N <sub>fore</sub>	N <sub>hind</sub>
					aspect ratio (±SE)	aspect ratio (±SE)					
Ontario	Blakeney Falls	45.268	-76.250	31/05/10	6.845 (±0.044)	6.392 (±0.037)	23	23	23	10	10
Ontario	Dorset	45.271	-78.960	31/07/10	7.053 (±0.075)	6.564 (±0.069)	7	6	7	6	7
Ontario	Heber Down	43.941	-78.988	08/06/10	6.845 (±0.034)	6.380 (±0.034)	20	20	20	10	10
Ontario	Lucknow	43.954	-81.497	28/07/10	7.018 (±0.041)	6.578 (±0.040)	20	20	19	10	10
Ontario	North Bay	44.947	-79.471	20/06/10-21/06/10	6.811 (±0.019)	6.372 (±0.019)	84	84	84	10	10
Ontario	Peterborough	44.315	-78.343	15/06/10	6.792 (±0.048)	6.352 (±0.052)	20	20	20	10	10
Ontario	Ridgetown	42.439	-81.831	11/07/10	6.707 (±0.048)	6.280 (±0.039)	18	18	18	10	10
Ontario	Sault Ste Marie	46.582	-84.300	24/06/10-26/06/10	6.651 (±0.025)	6.231 (±0.023)	60	60	59	10	10
Ontario	Serena Gundy Park	43.716	-79.353	15/07/10	6.772 (±0.042)	6.378 (±0.040)	25	25	25	10	10
Quebec	Dunany	45.758	-74.304	25/06/10	6.925 (±0.036)	6.457 (±0.040)	15	14	15	10	10
Quebec	Shawinigan	46.514	-72.679	27/06/10	6.857 (±0.032)	6.491 (±0.059)	33	26	25	10	10
Arkansas	Smithville	36.235	-91.470	22/05/10-07/08/10	6.382 (±0.027)	6.014 (±0.028)	35	35	33	10	10
Florida	8 Mile Creek	30.483	-87.326	26/06	6.653 (±0.045)	6.278 (±0.039)	20	19	19	10	10
Georgia	Conyers Monastery	33.584	-84.073	04/08	6.755 (±0.049)	6.331 (±0.045)	11	11	11	10	10
Georgia	Rome	34.443	-85.150	18/06/10-27/06/10	6.651 (±0.041)	6.221 (±0.036)	20	19	15	10	10
Illinois	Rockford	42.211	-88.976	17/07/10	6.332 (±0.040)	5.956 (±0.040)	20	20	20	10	10
Iowa	Gateway Hills Park	42.008	-93.647	24/06/10	6.298 (±0.037)	5.879 (±0.035)	20	20	20	10	10
Iowa	Odebolt	42.274	-95.129	15/07/10	6.391 (±0.025)	6.040 (±0.024)	73	73	73	10	10
Kentucky	Fossil Creek	37.773	-84.561	07/06/10	6.757 (±0.046)	6.265 (±0.036)	25	25	25	10	10
Maryland	Folly Quarter Creek	39.255	-76.927	13/07/10	6.603 (±0.029)	6.247 (±0.031)	33	32	32	10	10
Michigan	Johnson Creek	42.399	-83.528	19/06/10-26/06/10	6.826 (±0.041)	6.405 (±0.038)	24	23	21	10	10
Mississippi	Starkville	33.567	-89.041	05/07/10	6.580 (±0.035)	6.190 (±0.031)	26	26	24	10	10
Missouri	Eleven Point River	36.793	-91.331	05/06/10	6.279 (±0.047)	5.885 (±0.042)	12	12	12	10	10
Missouri	White River	36.654	-92.230	05/06/10	6.273 (±0.028)	5.903 (±0.028)	25	24	21	10	10
Nebraska	Chappell	41.083	-102.467	30/06/10	6.408 (±0.065)	6.070 (±0.061)	6	6	6	6	6

Nebraska	Kimball	41.232	-103.843	01/07/10	6.401 ( $\pm 0.030$ )	6.038 ( $\pm 0.030$ )	32	32	32	10	10
Nebraska	Leigh	41.701	-97.247	21/06/10	6.359 ( $\pm 0.034$ )	5.963 ( $\pm 0.034$ )	25	23	22	10	10
Ohio	Mt Vernon	40.405	-82.487	16/06/10	6.748 ( $\pm 0.023$ )	6.300 ( $\pm 0.025$ )	40	39	39	10	10
South Carolina	Four Holes Swamp	33.212	-80.348	14/07/10	6.782 ( $\pm 0.059$ )	6.445 ( $\pm 0.046$ )	21	21	21	10	10
South Carolina	Little Creek	34.842	-82.402	15/07/10	6.777 ( $\pm 0.040$ )	6.529 ( $\pm 0.050$ )	29	28	28	10	10
Texas	Powderly	33.753	-95.605	13/05/10	6.287 ( $\pm 0.033$ )	5.929 ( $\pm 0.030$ )	22	19	18	10	10
Vermont	Lamoille River	44.681	-73.068	18/06/10	6.873 ( $\pm 0.123$ )	6.473 ( $\pm 0.112$ )	4	4	4	4	4
Vermont	West Haven	43.624	-73.362	24/07/10	6.688 ( $\pm 0.037$ )	6.277 ( $\pm 0.035$ )	17	11	10	10	10
Vermont	Winooski River	46.352	-72.571	04/07/10-18/07/10	6.895 ( $\pm 0.034$ )	6.477 ( $\pm 0.028$ )	42	42	41	10	10




Figure 1 –

Geographic distribution of *Calopteryx maculata* (light area) and 34 sampling locations (symbol size is proportional to aspect ratio, see legend in lower left).






Figure 2 – Locations of 14 landmarks on the wing of *Calopteryx maculata*.




Figure 3 – Deformation plots showing the effect of increasing the value of each principal component on the relative locations of wing landmarks. Arrows indicate the direction and extent of change. Percentages are the percentage of variation explained by each principal component for fore and hind wings, respectively.




Figure 4 – Relationship between aspect ratio and the first principal component describing variation in wing shape for fore (closed symbols, solid line) and hind wings (open symbols, dotted line). Points are mean values from each of 34 sampling sites for both variables.





Figure 5 – Relationships between (A) aspect ratio and longitude for fore (closed symbols, solid line) and hind wings (open symbols, dotted line), and (B) aspect ratio and seasonality of precipitation (only fore wing data are shown). Points are mean values from each of 34 sampling sites for both variables.


## Map of sampling sites

Figure 1 - Geographic distribution of *Calopteryx maculata* (light area) and 34 sampling locations (symbol size is proportional to aspect ratio, see legend in lower left).




## Geometric morphometric landmarks

Figure 2 - Locations of 14 geometric morphometric landmarks on the wing of *Calopteryx maculata*.




### *Calopteryx maculata* wing deformation plots

Figure 3 - Deformation plots showing the effect of increasing the value of each principal component on the relative locations of wing landmarks. Arrows indicate the direction and extent of change. Percentages are the percentage of variation explained by each principal component for fore and hind wings, respectively.



## Relationship between aspect ratio and wing shape principal component

Figure 4 - Relationship between aspect ratio and the first principal component describing variation in wing shape for fore (closed symbols, solid line) and hind wings (open symbols, dotted line). Points are mean values from each of 34 sampling sites for both variables.





Relationship between longitude, seasonality of precipitation and aspect ratio in *Calopteryx maculata*.

Figure 5 - Relationships between (A) aspect ratio and longitude for fore (closed symbols, solid line) and hind wings (open symbols, dotted line), and (B) aspect ratio and seasonality of precipitation (only fore wing data are shown). Points are mean values from each of 34 sampling sites for both variables.

